

This is a repository copy of *The N2-Src neuronal splice variant of C-Src has altered SH3 domain ligand specificity and a higher constitutive activity than N1-Src*.

White Rose Research Online URL for this paper:  
<https://eprints.whiterose.ac.uk/95181/>

Version: Submitted Version

---

**Article:**

Keenan, Sarah, Lewis, Philip A, Wetherill, Sarah J et al. (2 more authors) (2015) The N2-Src neuronal splice variant of C-Src has altered SH3 domain ligand specificity and a higher constitutive activity than N1-Src. *FEBS Letters*. pp. 1995-2000. ISSN 0014-5793

<https://doi.org/10.1016/j.febslet.2015.05.033>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

Figure 1




Figure 2






Figure 3



# Figure 4





Figure 5

