

**Original citation:**

Arduini, Arduino and Zammit, Victor A. (2016) A tennis lesson : sharp practice in the science behind the Sharapova case. Postgraduate Medical Journal .

**Permanent WRAP URL:**

<http://wrap.warwick.ac.uk/80281>

**Copyright and reuse:**

The Warwick Research Archive Portal (WRAP) makes this work of researchers of the University of Warwick available open access under the following conditions.

This article is made available under the Creative Commons Attribution 4.0 International license (CC BY 4.0) and may be reused according to the conditions of the license. For more details see: <http://creativecommons.org/licenses/by/4.0/>

**A note on versions:**

The version presented in WRAP is the published version, or, version of record, and may be cited as it appears here.

For more information, please contact the WRAP Team at: [wrap@warwick.ac.uk](mailto:wrap@warwick.ac.uk)

# A tennis lesson: sharp practice in the science behind the Sharapova case

Arduino Arduini,<sup>1</sup> Victor A Zammit<sup>2</sup>

Maria Sharapova (and hundreds of other elite athletes) took meldonium, a drug developed at the time of the USSR for the treatment of heart attack and stroke, though it has never been approved for use anywhere outside of the former Soviet Union. Meldonium is an inhibitor of  $\gamma$ -butyrobetaine hydroxylase, an enzyme involved in the carnitine biosynthetic pathway.<sup>1</sup> Intake results in a reduction of tissue carnitine content, including the heart and skeletal muscles. Carnitine plays a critical role in transferring long-chain fatty acids across mitochondrial inner membrane into the mitochondrial matrix, to enable entry of the fatty acid moiety into the oxidation pathway, to synthesise ATP aerobically.<sup>2</sup> In cases when intracellular carnitine availability is compromised, such as in neonatal primary carnitine deficiency,<sup>3</sup> those affected experience hypoglycaemia, fatigue, seizures and cardiomyopathy.<sup>3</sup> Carnitine-deficient individuals are sensitive to prodrugs containing the pivaloyl moiety, as pivalic acid is a prodrug able to induce carnitine deficiency (like meldonium); this results in lethal cardiac arrhythmias.<sup>4</sup> Meldonium administered orally at 'recommended' doses (500 mg, twice daily)<sup>5</sup> would not have such a drastic effect. However, even a moderate reduction of muscle carnitine content in healthy volunteers treated with pivaloyl-conjugated antibiotics for 54 days reduces work at maximal oxygen uptake ( $VO_2$ max), maximal heart rate,<sup>6</sup> reduced interventricular septum thickness and left ventricular mass.<sup>7</sup> The oral administration of meldonium to healthy volunteers (500 mg, twice daily) for 4 weeks leads to a significant 18% reduction of plasma carnitine.<sup>5</sup> Even though we are not aware of the meldonium dosage prescribed to Ms Sharapova, it is safe to assume that, in the absence of oral carnitine supplementation, her muscle carnitine content would have been, at most, only modestly decreased, and would have resulted in a minimal

impact on her muscle oxidative capacity, as evidenced by her continued performance at the highest levels in her sport. The medical team looking after Ms Sharapova appears to have confounded the tangible beneficial effects of depressing acylcarnitine formation in *ischaemic* heart (which may improve glucose use, eg, during ischaemia-reperfusion) with the requirement of normal cardiac muscle for high rates of carnitine-dependent fatty acid oxidation. A recent study in rodents has shown that even a moderate, silent carnitine-deficient phenotype may reveal its fragility when subjected to strong adrenergic stress.<sup>8</sup> Therefore, unless meldonium has unidentified off-target pharmacological effects, one has to be thankful that it was not administered in high enough doses so as to damage the athletes who continued to take it.

Another aspect of meldonium pharmacology is the inhibition of carnitine acetyltransferase (CAT) activity,<sup>9</sup> an enzyme that reversibly transfers the acetyl moiety of acetyl coenzyme A to carnitine in mitochondria and peroxisomes. Due to its capacity to buffer mitochondrial acetyl coenzyme A levels, CAT plays a prominent role in intermediary metabolism and muscle bioenergetics.<sup>2</sup> As expected, in rodent models, a lowered muscle CAT activity results in muscle fatigue and exercise intolerance.<sup>10</sup> This would again strongly argue against the use of meldonium as a performance-enhancing drug, particularly if such CAT-inhibitory activity is combined with carnitine depletion.

The World Anti-Doping Agency (WADA) may have decided to add meldonium to the Prohibited List, not necessarily because of any illusory performance-enhancing action in healthy athletes, but because of the real danger that it carries for the overenthusiastic athlete who may overdose on it and lower intracellular carnitine to pathological levels. A comprehensive literature search of several databases failed to unearth any cogent evidence of the potential meldonium effect in enhancing physical performance in healthy subjects, with only one clinical trial conducted on a small cohort of old (average >60 year) patients with *angina pectoris*, to evaluate exercise tolerance. On such minimal findings is the dangerous extrapolation to an assumed performance-enhancing

effect of meldonium in healthy elite athletes based. It is not an overstatement to emphasise that the understanding of metabolism is a postmodernist discipline in which neither the inadequately trained scientists (Metabolism is a Cinderella subject in most universities) nor the editors of journals or the clinicians who peruse their publications know how to interpret the intricacies of control in metabolism. This is an era in which metabolic science is overrun by over-emphasis on gross changes in gene expression, and too little on the reductionist discipline of the study of the function of proteins, their interactions, and their effects on metabolic fluxes. Athletes chasing the shadows of marginal gains are the unwitting victims. Hats off to WADA who may have recognised that, even in the absence of any performance-enhancing properties of meldonium, they should ban it for health reasons. If anything, Ms Sharapova and other ill-advised athletes may not have attained their true athletic potential as a result of having taken meldonium.

**Contributors** AA contributed to the idea of publishing an Editorial and to the writing of the text. VAZ contributed to the idea of publishing an Editorial and to the writing of the text.

**Competing interests** None declared.

**Provenance and peer review** Not commissioned; externally peer reviewed.



## OPEN ACCESS

**Open Access** This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

**To cite** Arduini A, Zammit VA. *Postgrad Med J* Published Online First: [please include Day Month Year] doi:10.1136/postgradmedj-2016-134124

Received 29 March 2016

Accepted 18 April 2016

*Postgrad Med J* 2016;0:1–2.

doi:10.1136/postgradmedj-2016-134124

## REFERENCES

- 1 Simkhovich BZ, Shutenko ZV, Meirëna DV, et al. 3-(2,2,2 Trimethylhydrazinium)propionate (THP)—a novel  $\gamma$ -butyrobetaine hydroxylase inhibitor with cardioprotective properties. *Biochem Pharmacol* 1988;37:195–202.
- 2 Zammit VA, Ramsay RR, Bonomini M, et al. Carnitine, mitochondrial function and therapy. *Adv Drug Deliv Rev* 2009;61:1353–62.
- 3 Longo N, Amat di San Filippo C, Pasquali M. Disorders of carnitine transport and the carnitine

<sup>1</sup>CoreQuest, Manno, Switzerland; <sup>2</sup>Translational and Experimental Medicine, Warwick Medical School, Coventry, UK

**Correspondence to** Professor Victor A Zammit, Translational Medicine, Warwick Medical School, Coventry CV4 7AL, UK; [v.a.zammit@warwick.ac.uk](mailto:v.a.zammit@warwick.ac.uk)

## Editorial

- cycle. *Am J Med Genet C Semin Med Genet* 2006;142C:77–85.
- 4 Rasmussen J, Nielsen OW, Lund AM, *et al.* Primary carnitine deficiency and pivalic acid exposure causing encephalopathy and fatal cardiac events. *J Inherit Metab Dis* 2013;36:35–41.
- 5 Liepinsh E, Konrade I, Skapare E, *et al.* Mildronate treatment alters  $\gamma$ -butyrobetaine and l-carnitine concentrations in healthy volunteers. *J Pharm Pharmacol* 2011;63:1195–201.
- 6 Abrahamsson K, Eriksson BO, Holme E, *et al.* Pivalic acid-induced carnitine deficiency and physical exercise in humans. *Metab Clin Exp* 1996;45:1501–7.
- 7 Abrahamsson K, Mellander M, Eriksson BO, *et al.* Transient reduction of human left ventricular mass in carnitine depletion induced by antibiotics containing pivalic acid. *Br Heart J* 1995;74: 656–9.
- 8 Giudice PL, Bonomini M, Arduini A. A moderate carnitine deficiency exacerbates isoproterenol-induced myocardial injury in rats. *Cardiovasc Drugs Ther* 2016;30:119–27.
- 9 Jaudzems K, Kuka J, Gutsaits A, *et al.* Inhibition of carnitine acetyltransferase by mildronate, a regulator of energy metabolism. *J Enzyme Inhib Med Chem* 2009;24:1269–75.
- 10 Seiler SE, Koves TR, Gooding JR, *et al.* Carnitine Acetyltransferase Mitigates Metabolic Inertia and Muscle Fatigue during Exercise. *Cell Metab* 2015;22:65–76.



# A tennis lesson: sharp practice in the science behind the Sharapova case

Arduino Arduini and Victor A Zammit

*Postgrad Med J* published online June 1, 2016

---

Updated information and services can be found at:

<http://pmj.bmj.com/content/early/2016/06/01/postgradmedj-2016-134124>

---

*These include:*

## References

This article cites 10 articles, 1 of which you can access for free at: <http://pmj.bmj.com/content/early/2016/06/01/postgradmedj-2016-134124#BIBL>

## Open Access

This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

## Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

---

## Topic Collections

Articles on similar topics can be found in the following collections

- [Open access](#) (37)
- [Drugs: cardiovascular system](#) (353)
- [Cardiomyopathy](#) (26)
- [Ischaemic heart disease](#) (139)
- [Diabetes](#) (136)
- [Drugs: infectious diseases](#) (216)
- [Epilepsy and seizures](#) (29)
- [Interventional cardiology](#) (110)
- [Metabolic disorders](#) (213)
- [Occupational and environmental medicine](#) (36)
- [Stroke](#) (134)

---

## Notes

---

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>