

This is a repository copy of *Tree encroachment may lead to functionally-significant changes in peatland testate amoeba communities*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/99318/>

Version: Accepted Version

Article:

Payne, Richard J., Creevy, Angela, Malysheva, Elena et al. (9 more authors) (2016) Tree encroachment may lead to functionally-significant changes in peatland testate amoeba communities. *Soil Biology and Biochemistry*. pp. 18-21. ISSN 0038-0717

<https://doi.org/10.1016/j.soilbio.2016.04.002>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Tree encroachment may lead to functionally-significant changes in peatland testate amoeba communities

Richard J Payne^{a,b}, Angela Creevy^{c,d}, Elena Malysheva^b, Joshua Ratcliffe^{c,e}, Roxane Andersen^c, Andrey N Tsyganov^b, James G Rowson^d, Katarzyna Marcisz^f, Małgorzata Zielińska^f, Mariusz Lamentowicz^f, Elena D Lapshina^g, Yuri Mazei^{b,h}

^a Environment, University of York, Heslington, York YO10 5DD, United Kingdom. E-mail richard.payne@york.ac.uk

^b Department of Zoology and Ecology, Penza State University, Krasnaya str. 40, 440026 Penza, Russia.

^c Environmental Research Institute, North Highland College, University of the Highlands and Islands, Castle Street, Thurso, Caithness, KW14 7JD, Scotland, United Kingdom.

^d Geography, Edge Hill University, St Helens Road, Ormskirk, Lancashire L39 4QP, United Kingdom.

^e Science & Engineering, University of Waikato, Private Bag 3105, Hamilton 3240, New Zealand.

^f Laboratory of Wetland Ecology and Monitoring & Department of Biogeography and Palaeoecology, Faculty of Geographical and Geological Science, Adam Mickiewicz University in Poznań, ul. Dziegielowa 27, 61-680 Poznań, Poland.

^g Center for Environmental Dynamics and Climate Change, Yugra State University, Chekhova str. 16, Khanty-Mansiysk 628012, Russia.

^h Department of Hydrobiology, Lomonosov Moscow State University, Leninskiye gory, 1, Moscow 119991, Russia.

ABSTRACT

Climate change is likely to cause increased tree recruitment on open peatlands but we currently have little idea what consequences this vegetation change may have below-ground. Here we use transects across forested to open bog ecotones at three Russian peatland complexes to assess potential changes in the most abundant group of peatland protists – the testate amoebae. We show that the testate amoeba communities of forested and open bog are markedly different with a very abrupt boundary at, or near, the vegetation ecotone. Changes along our transects suggest that tree encroachment may reduce the trophic level of testate amoeba communities and reduce the contribution of mixotrophic testate amoebae to primary production. Our study strongly suggests that increased tree recruitment on open peatlands will have important consequences for both microbial biodiversity and microbially-mediated ecosystem processes.

KEYWORDS: Peatland; Forest; Tree; Protist; Climate; Carbon

Climate change is causing treelines to move pole-wards and to higher altitudes around the world (Harsch et al., 2009). Peatland ecosystems are a globally-significant carbon store (c.4-600 GtC (Yu,

2012; Loisel et al., 2014)) with two alternative stable states as unforested 'open' bogs and forested bogs (often known by their Russian name 'ryam')(Agnew et al., 1993). Modelling and experimental evidence show that climatic warming and water table drawdown can lead to a switch between these states with the establishment of trees on formerly-open peatland (Heijmans et al., 2013; Limpens et al., 2014; Holmgren et al., 2015). It is reasonable to expect that rising temperatures and increased drought frequency, combined with ongoing anthropogenic drainage, will lead to increasing areas of boreal peatland switching to tree domination. There is both palaeoecological evidence for greater forest cover in peatland-dominated boreal regions during warmer phases of the Holocene (MacDonald et al., 2008) and monitoring data suggesting recent increases in tree and shrub recruitment in many peatland regions (Esper and Schweingruber, 2004; Shiyatov et al., 2005; Berg et al., 2009). The incursion of trees into previously open bogs has the potential to create positive feedback loops through increased transpiration, interception and shade leading to drying of the bog surface (Waddington et al., 2015). The consequences for microbial communities of such a switch in vegetation are largely unknown but may have important implications for carbon and nutrient cycling. Here we use a space-for-time substitution approach (Blois et al., 2013) to assess possible changes in communities of the key eukaryotic microbes of peatlands – the testate amoebae – with tree establishment. Testate amoebae are the most abundant protists in peatlands (Gilbert et al., 1998) and play important roles in carbon and nutrient cycling (Wilkinson and Mitchell, 2010).

We positioned ten transects of 30-45m length across the forested to open bog ecotone at three peatland complexes in western Siberia (Mukhrino 60.9°N, 68.7°E), the Karelia region of northwest Russia ('Black River' 66.5°N, 32.9°E) and the Penza region of European Russia (Morsovo 53.8°N, 42.3°E). Transects (3-4 per site) were labelled based on a qualitative judgement of the centre point of the vegetation ecotone (0m), with positive numbers representing the open bog and negative numbers the forested bog ends of the transect respectively. Samples were extracted every 5m by removing the upper 5cm of bryophytes and any litter (c. 25cm³), giving a total of 74 samples across all sites. Testate amoebae were extracted, identified and enumerated under the microscope using standard methods (Booth et al., 2010)(Supplementary Table 1). The mean count total was 211 tests per sample and the minimum was 100, sufficient to adequately characterise the community (Payne and Mitchell, 2009). To assess community change along the transects we summarised the datasets using first axis scores from an NMDS ordination on Bray-Curtis dissimilarity (Bray and Curtis, 1957). Results (Fig. 1B) show large changes in most transects with a very abrupt shift between low and high axis scores. To identify points along the transects which maximised community difference we applied non-parametric change-point analysis (nCPA; (Qian et al., 2003)) (Fig. 1A). Identified change-points were all immediately adjacent to the centre-point of the vegetation ecotone as judged in the field, with two exceptions from the Morsovo site where the vegetation change is more gradual. Redundancy analysis (accounting for the transect structure) showed a highly significant difference between the amoeba communities either side of these points (Hellinger-transformed data; 17.8% variance, P=0.001). This difference remained highly significant when accounting for water table depth (measured in Mukhrino only; 7.5% variance, P=0.002) and more variance was explained by the difference across the change points than by tree stem density for the only site where this was recorded (Mukhrino only; 29.9% variance, P=0.001 versus 19.2% variance, P=0.001).

We used Indicator Value analysis (IndVal (Dufrêne and Legendre, 1997)) to identify taxa typifying samples on either side of the change points. These groups have clear differences (Fig. 1A). Taxa

78 typifying the open bog end of the transects are generally larger and mixotrophic (e.g. *Hyalosphenia*
79 *papilio*; *Archerella flavum*) while taxa typifying the forested end are often smaller and bacterivorous
80 (e.g. *Assulina muscorum*; *Corythion dubium*). We used Bray-Curtis dissimilarity to assess pairwise
81 mean similarity to communities of mineral soil forests (111 analyses from across Siberia (Malysheva,
82 2011)) and open peatlands (68 analyses (Lamentowicz et al., 2015b)). These results showed some
83 variability but communities at the open bog end of the transects showed greater similarity to other
84 analyses from open peatland and differed strongly ($p < 0.01$; Supplementary Fig. 1) from communities
85 at the forested end, which were more similar to mineral soil forest.

86 To assess the possible functional significance of these changes we considered two key functional
87 traits of testate amoeba communities: the proportion of mixotrophic taxa and the community-
88 weighted mean aperture diameter, which is increasingly used as a metric of trophic position
89 (Fournier et al., 2012; Lamentowicz et al., 2013; Fournier et al., 2015; Lamentowicz et al., 2015a).
90 Aperture diameter data were taken from the literature and direct observations and assigned to five
91 size classes (1=0-10 μ m; 2=10-20 μ m; 3=20-30 μ m; 4=30-40 μ m; 5=>40 μ m) using a mean of the
92 maximum and minimum dimensions where reported. Results show an abrupt increase in the
93 proportion of mixotrophs from forested to open bog (although mixotrophs were rare in Morsovo).
94 Testate amoebae from the open bog end of the transects generally had larger apertures, suggesting
95 larger prey and higher trophic position. Both these differences were highly significant ($P < 0.01$; Fig.
96 1C & Fig. 1D).

97 Differences in testate amoeba communities between forested and open bog are not particularly
98 surprising but we are not aware of any previous study which has directly demonstrated this. More
99 surprising is the scale of the difference and the abruptness of the change demonstrated by our data,
100 which has the non-linear characteristics of an ecological threshold (Groffman et al., 2006). An open
101 question is whether this response reflects alternative stable states within the testate amoeba
102 community *itself* or whether this is driven by similarly abrupt changes in plant communities and
103 environmental conditions. Changes along the transects may be partly driven by moisture availability,
104 paralleling evidence from many previous studies, but variance partitioning suggests that other
105 factors must also be involved. The most distinct change along the transects is a loss of mixotrophic
106 testate amoebae with tree cover, suggesting that reduced light penetration is an important factor
107 driving the change in community (*cf.* Marcisz et al., 2014). Recent research has suggested that
108 mixotrophic testate amoebae may make a non-trivial contribution to peatland primary production
109 (Jassey et al., 2015). Although microbially-fixed carbon is likely to be a relatively small input
110 compared to plant production, greater lability may make this pool disproportionately important in
111 driving change in the microbial food-web. Our data also show that testate amoebae in open
112 peatland tend to have larger apertures than those of forested peatland. This may reflect differences
113 in available food sources with larger food items such as microalgae and ciliates likely to be more
114 abundant in the open peatland. It is possible that tree encroachment might lead to greater grazing
115 pressure from testate amoebae on bacteria due to the loss of alternative food types, with
116 consequences for the biogeochemical processes those bacteria control.

117 Our study does not allow us to assess how quickly testate amoeba communities respond to tree
118 establishment or whether these changes are reversible, but opens the way to palaeoecological

119 studies, which could address this question. Our dataset also provides a baseline for reassessments of
120 these sites in the future.

121 On the basis of our results it seems probable that tree encroachment may lead to large changes in
122 testate amoeba community with important implications for ecosystem processes such as microbial
123 primary production and predation.

124 ACKNOWLEDGEMENTS

125 Work in European Russia and data analysis was supported by the Russian Scientific Fund, grant 14-
126 14-00891 to YuM. Fieldwork in Siberia was funded by the European Union Seventh Framework
127 Programme under grant agreement no. 262693 [INTERACT] as part of the PEATEDGE (primary data
128 collection), CliMireSiber (comparison dataset) and DAVOCCAP projects (participation of RA). UK-
129 Russia research cooperation was supported by a Royal Society International Exchange grant
130 (IE150173). During the research JR was supported by the Carnegie Trust for the Universities of
131 Scotland through grant LG13STIR007 and AC was supported by a Natural Talent Apprenticeship from
132 The Conservation Volunteers, funded by the Heritage Lottery Fund. RJP and RA are funded by the
133 Leverhulme Trust through project RPG-2015-162. The comparison dataset was partially funded by
134 grant PSPB-013/2010 (CLIMPEAT) through the Swiss Contribution to the enlarged European Union.

135 Author contributions: RJP, JR and YuM conceived the study. RJP, AC, JR, RA, EM, ANT, YuM, ML and
136 EDL conducted fieldwork. AC, EM, KM and MZ analysed testate amoebae. RJP, YuM, ML, JGR and RA
137 obtained funding and/or supervised research students. RJP conducted the data analysis and wrote
138 the first draft of the paper. All authors commented on the draft and provided interpretation.

139

140

141


- 143 Agnew, A.D.Q., Wilson, J.B., Sykes, M.T., 1993. A Vegetation Switch as the Cause of a Forest/Mire
144 Ecotone in New Zealand. *Journal of Vegetation Science* 4, 273-278.
- 145 Berg, E.E., Hillman, K.M., Dial, R., DeRuwe, A., 2009. Recent woody invasion of wetlands on the Kenai
146 Peninsula Lowlands, south-central Alaska: a major regime shift after 18 000 years of wet Sphagnum-
147 sedge peat recruitment. *Canadian Journal of Forest Research* 39, 2033-2046.
- 148 Blois, J.L., Williams, J.W., Fitzpatrick, M.C., Jackson, S.T., Ferrier, S., 2013. Space can substitute for
149 time in predicting climate-change effects on biodiversity. *Proceedings of the National Academy of
150 Sciences* 110, 9374-9379.
- 151 Booth, R.K., Lamentowicz, M., Charman, D.J., 2010. Preparation and analysis of testate amoebae in
152 peatland palaeoenvironmental studies. *Mires & Peat* 7.
- 153 Bray, J.R., Curtis, J.T., 1957. An ordination of the upland forest communities of southern Wisconsin.
154 *Ecological monographs* 27, 325-349.
- 155 Dufrêne, M., Legendre, P., 1997. Species assemblages and indicator species: the need for a flexible
156 asymmetrical approach. *Ecological monographs* 67, 345-366.
- 157 Esper, J., Schweingruber, F.H., 2004. Large-scale treeline changes recorded in Siberia. *Geophysical
158 Research Letters* 31.
- 159 Fournier, B., Lara, E., Jassey, V.E., Mitchell, E.A., 2015. Functional traits as a new approach for
160 interpreting testate amoeba palaeo-records in peatlands and assessing the causes and consequences
161 of past changes in species composition. *The Holocene*, 0959683615585842.
- 162 Fournier, B., Malysheva, E., Mazei, Y., Moretti, M., Mitchell, E.A.D., 2012. Toward the use of testate
163 amoeba functional traits as indicator of floodplain restoration success. *European Journal of Soil
164 Biology* 49, 85-91.
- 165 Gilbert, D., Amblard, C., Bourdier, G., Francez, A.J., 1998. The microbial loop at the surface of a
166 peatland: structure, function, and impact of nutrient input. *Microbial ecology* 35, 83-93.
- 167 Groffman, P., Baron, J., Blett, T., Gold, A., Goodman, I., Gunderson, L., Levinson, B., Palmer, M.,
168 Paerl, H., Peterson, G., Poff, N.L., Rejeski, D., Reynolds, J., Turner, M., Weathers, K., Wiens, J., 2006.
169 *Ecological Thresholds: The Key to Successful Environmental Management or an Important Concept
170 with No Practical Application?* *Ecosystems* 9, 1-13.
- 171 Harsch, M.A., Hulme, P.E., McGlone, M.S., Duncan, R.P., 2009. Are treelines advancing? A global
172 meta-analysis of treeline response to climate warming. *Ecology letters* 12, 1040-1049.
- 173 Heijmans, M.M., Knaap, Y.A., Holmgren, M., Limpens, J., 2013. Persistent versus transient tree
174 encroachment of temperate peat bogs: effects of climate warming and drought events. *Global
175 change biology* 19, 2240-2250.
- 176 Holmgren, M., Lin, C.-Y., Murillo, J.E., Nieuwenhuis, A., Penninkhof, J., Sanders, N., van Bart, T., van
177 Veen, H., Vasander, H., Vollebregt, M.E., Limpens, J., 2015. Positive shrub–tree interactions facilitate
178 woody encroachment in boreal peatlands. *Journal of Ecology* 103, 58-66.
- 179 Jassey, V.E., Signarbieux, C., Hättenschwiler, S., Bragazza, L., Buttler, A., Delarue, F., Fournier, B.,
180 Gilbert, D., Laggoun-Défarge, F., Lara, E., 2015. An unexpected role for mixotrophs in the response of
181 peatland carbon cycling to climate warming. *Scientific reports* 5.
- 182 Lamentowicz, M., Bragazza, L., Buttler, A., Jassey, V., Mitchell, E., 2013. Seasonal patterns of testate
183 amoeba diversity, community structure and species–environment relationships in four Sphagnum-
184 dominated peatlands along a 1300 m altitudinal gradient in Switzerland. *Soil Biology and
185 Biochemistry* 67, 1-11.
- 186 Lamentowicz, M., Gałka, M., Obremaska, M., Köhl, N., Lücke, A., Jassey, V., 2015a. Reconstructing
187 climate change and ombrotrophic bog development during the last 4000 years in northern Poland
188 using biotic proxies, stable isotopes and trait-based approach. *Palaeogeography, Palaeoclimatology,
189 Palaeoecology* 418, 261-277.
- 190 Lamentowicz, M., Słowiński, M., Marcisz, K., Zielińska, M., Kaliszan, K., Lapshina, E., Gilbert, D.,
191 Buttler, A., Fiałkiewicz-Kozieł, B., Jassey, V.E., 2015b. Hydrological dynamics and fire history of the

192 last 1300years in western Siberia reconstructed from a high-resolution, ombrotrophic peat archive.
193 Quaternary Research 84, 312-325.
194 Limpens, J., Egmond, E., Li, B., Holmgren, M., 2014. Do plant traits explain tree seedling survival in
195 bogs? Functional ecology 28, 283-290.
196 Loisel, J., Yu, Z., Beilman, D.W., Camill, P., Alm, J., Amesbury, M.J., Anderson, D., Andersson, S.,
197 Bochicchio, C., Barber, K., 2014. A database and synthesis of northern peatland soil properties and
198 Holocene carbon and nitrogen accumulation. The Holocene, 0959683614538073.
199 MacDonald, G., Kremenetski, K., Beilman, D., 2008. Climate change and the northern Russian
200 treeline zone. Philosophical Transactions of the Royal Society B: Biological Sciences 363, 2283-2299.
201 Malysheva, E., 2011. Testate amoeba community structure in the contact zones of different
202 biotopes. Penza State University, Penza.
203 Marcisz, K., Lamentowicz, Ł., Słowińska, S., Słowiński, M., Muszak, W., Lamentowicz, M., 2014.
204 Seasonal changes in Sphagnum peatland testate amoeba communities along a hydrological gradient.
205 European journal of protistology 50, 445-455.
206 Payne, R.J., Mitchell, E.A.D., 2009. How many is enough? Determining optimal count totals for
207 ecological and palaeoecological studies of testate amoebae. Journal of Paleolimnology 42, 483-495.
208 Qian, S.S., King, R.S., Richardson, C.J., 2003. Two statistical methods for the detection of
209 environmental thresholds. Ecological Modelling 166, 87-97.
210 Shiyatov, S., Terent'Ev, M., Fomin, V., 2005. Spatiotemporal dynamics of forest-tundra communities
211 in the Polar Urals. Russian Journal of Ecology 36, 69-75.
212 Waddington, J., Morris, P., Kettridge, N., Granath, G., Thompson, D., Moore, P., 2015. Hydrological
213 feedbacks in northern peatlands. Ecohydrology 8, 113-127.
214 Wilkinson, D.M., Mitchell, E.A., 2010. Testate amoebae and nutrient cycling with particular reference
215 to soils. Geomicrobiology Journal 27.
216 Yu, Z., 2012. Northern peatland carbon stocks and dynamics: a review. Biogeosciences 9, 4071-4085.

217


218

220 Figure 1. Testate amoeba community change along forested to open bog transects. A) nCPA change-
 221 points (vertical lines) and significant indicator species as identified by IndVal showing taxa IV>50 and
 222 P<0.05. B) Community change as summarised by first axis scores from an NMDS ordination using
 223 Bray-Curtis dissimilarity. C) Proportion of mixotrophic taxa. D) Community weighted mean (CWM)
 224 aperture diameter in five size classes (1=0-10µm; 2=10-20µm; 3=20-30µm; 4=30-40µm; 5=>40µm).
 225 In all plots lines represent individual transects from Mukhrino (shades of red); Morsovo (shades of
 226 green) and Black River (shades of blue). In the bottom right of each plot are results of one way
 227 repeated measures analysis of variance (RM-ANOVA) contrasting data either side of the nCPA
 228 change-point.



230 Supplementary Table 1. Species abundance across the three study sites.

231 Supplementary Figure 1. A) Pairwise mean Bray-Curtis dissimilarity relative to testate amoebae
232 communities of open bog (Lamentowicz et al. 2015b). B) Pairwise mean Bray-Curtis dissimilarity
233 relative to testate amoeba communities of conifer forests on mineral soil (Malysheva, 2011). Details
234 as for Figure 1.



235