Huntjens, B. & O'Donnell, C. (2006). Diabetes mellitus and the eye: axial length. Paper presented at the 42nd EASD Meeting, 14-09-2006 - 17-09-2006, Copenhagen (Denmark).



City Research Online

Original citation: Huntjens, B. & O'Donnell, C. (2006). Diabetes mellitus and the eye: axial length. Paper presented at the 42nd EASD Meeting, 14-09-2006 - 17-09-2006, Copenhagen (Denmark).

Permanent City Research Online URL: http://openaccess.city.ac.uk/5601/

Copyright & reuse

City University London has developed City Research Online so that its users may access the research outputs of City University London's staff. Copyright © and Moral Rights for this paper are retained by the individual author(s) and/ or other copyright holders. All material in City Research Online is checked for eligibility for copyright before being made available in the live archive. URLs from City Research Online may be freely distributed and linked to from other web pages.

Versions of research

The version in City Research Online may differ from the final published version. Users are advised to check the Permanent City Research Online URL above for the status of the paper.

Enquiries

If you have any enquiries about any aspect of City Research Online, or if you wish to make contact with the author(s) of this paper, please email the team at publications@city.ac.uk.

ABSTRACT EASD (poster)

AUTHORS

Byki Huntjens Clare O'Donnell

TITLE

Diabetes Mellitus and the Eye: the Axial Length.

Background and aims: The refractive error of the eye is dependent on its axial length. Refractive error is known to fluctuate significantly in poorly-controlled diabetic patients. Recently it has been reported that human eyes fluctuate in axial length during the day. However, this change is not detectable in all subjects, suggesting physiological influences such as diet. The purpose of this study was to investigate fluctuations in axial length and blood glucose levels (BGLs) in diabetic patients and control subjects, using partial coherence interferometry.

Material and methods: Periodic axial length measurements were taken with an IOLMaster (Zeiss, Oberkochen, Germany) in 21 type 2 diabetic subjects (age 56 ± 11 years), 19 type 1 diabetic subjects (age 38 ± 15 years) and 18 non-diabetic controls (age 46 ± 22 years). Additionally, BGLs were measured using a HemoCue Beta glucose analyser (HemoCue, Ängelholm, Sweden). Measurements were taken between 8AM and 8PM at approximately two hourly intervals. The ocular length variability was mapped against time of day and related to BGLs, age, and the degree of diabetic retinopathy.

Results: Axial length variations were found to be significant with time (ANOVA, P=.003) and there was a significant effect for the interaction between group and time of day (ANOVA, P<.0005). DM type 1 and control subjects showed a significantly increased axial length in the afternoon compared to in the morning (t-test, P<.0005). No significant change was observed among DM type 2 subjects. Axial length fluctuated sinusoidally with time of day. A three-parameter sinusoidal curve was used to fit each set of axial length measurements, which resulted in a correlation coefficient $R^2 \ge 0.7$ in 62% of type 2 diabetic subjects, 78% of type 1 diabetic subjects and 72% of control subjects. Multiple regression analysis established that neither BGLs, age, nor the degree of diabetic retinopathy had a significant effect on the fluctuations of the axial length (P>.05)

Conclusion: Axial length varies throughout the day in diabetic and control subjects; however, this change does not appear to be significant in every individual. The data suggest a maximum axial length value in the afternoon. Changes in axial length from baseline did not correlate with BGLs, age, or the degree of diabetic retinopathy. Therefore, blood glucose levels may not be the main physiological influence in axial length fluctuations.

Acknowledgements: Supported by a grant awarded by Lein Applied Diagnostics.