

Dhanani, S., Asaria, T. S., Barbur, J. L. & Huntjens, B. (2012). Relationship between Macular Pigment and Foveal Anatomic Architecture in an Asian Population. Paper presented at the Association for Research in Vision and Ophthalmology, 06-05-2012 - 10-05-2012, Fort Lauderdale, USA.



**CITY UNIVERSITY
LONDON**

[City Research Online](#)

Original citation: Dhanani, S., Asaria, T. S., Barbur, J. L. & Huntjens, B. (2012). Relationship between Macular Pigment and Foveal Anatomic Architecture in an Asian Population. Paper presented at the Association for Research in Vision and Ophthalmology, 06-05-2012 - 10-05-2012, Fort Lauderdale, USA.

Permanent City Research Online URL: <http://openaccess.city.ac.uk/5606/>

Copyright & reuse

City University London has developed City Research Online so that its users may access the research outputs of City University London's staff. Copyright © and Moral Rights for this paper are retained by the individual author(s) and/ or other copyright holders. All material in City Research Online is checked for eligibility for copyright before being made available in the live archive. URLs from City Research Online may be freely distributed and linked to from other web pages.

Versions of research

The version in City Research Online may differ from the final published version. Users are advised to check the Permanent City Research Online URL above for the status of the paper.

Enquiries

If you have any enquiries about any aspect of City Research Online, or if you wish to make contact with the author(s) of this paper, please email the team at publications@city.ac.uk.

Purpose: : The extent to which reduced macular pigment optical density (MPOD) contributes to the prevalence of age related macular degeneration (ARMD) in Caucasians compared to other ethnicities has often been questioned. Foveal architecture may be related to MPOD levels and hence be a contributing factor. Previous studies have reported race-linked differences in peak MPOD and its central spatial distribution. This study investigates the relationship between MPOD and foveal architecture in an adult Asian population.

Methods: : The spatial profile of MPOD was assessed in 55 healthy Asian subjects (mean age 21 ± 4 years) using heterochromatic flicker photometry (Ophthalmic Physiol Opt. 30:470-483,2010). High-resolution macular thickness maps were obtained using a spectral-domain optical coherence tomography (Spectralis OCT, Heidelberg, Germany). The following relationships were investigated: 1) Peak MPOD (at 0° eccentricity) and minimal foveal thickness (MFT) measured manually at the point of sharpest foveal reflex; 2) Peak MPOD and central foveal thickness (CFT: average retinal thickness within central 1mm circle of the ETDRS grid); 3) Peak MPOD and foveal width (FW: measured from crest to crest); 4) MFT and FW; 5) Average MPOD (OD_{av} : over an area subtending $\pm 2.8^\circ$ centred at the fovea) and FW; and 6) OD_{av} and CFT.

Results: : The peak MPOD values (mean 0.56 ± 0.17 log units) and the corresponding MFT showed good correlation ($R^2 = 0.34$ $p < 0.0005$). A weaker correlation was found between peak MPOD and CFT ($R^2 = 0.12$; $p = 0.01$). A moderately significant negative correlation was found between FW and CFT ($R^2 = 0.22$; $p < 0.0005$) and between FW and MFT ($R^2 = 0.1$; $p = 0.03$). In contrast, no correlation was found between FW and peak MPOD at 0° ($p = 0.84$), OD_{av} and FW ($p = 0.41$), or OD_{av} and CFT ($p = 0.59$).

Conclusions: : The current findings suggest that minimal foveal thickness (measured manually at the point of sharpest foveal reflex) is a better predictor value for MPOD compared to the central foveal thickness given by the OCT. The expectation that a wider foveal width may contain more macular pigment because of longer cone axon fibres is not supported by our findings in the Asian subject group. Our results suggest that differences in foveal architecture can explain some of the measured variations in MPOD.