

Bender, C., Moshe, M. & Sarkar, S. (2012). PT-symmetric interpretation of double-scaling. Journal of Physics A: Mathematical and Theoretical, 46(10), e102002. doi: 10.1088/1751-8113/46/10/102002



**CITY UNIVERSITY
LONDON**

[City Research Online](#)

Original citation: Bender, C., Moshe, M. & Sarkar, S. (2012). PT-symmetric interpretation of double-scaling. Journal of Physics A: Mathematical and Theoretical, 46(10), e102002. doi: 10.1088/1751-8113/46/10/102002

Permanent City Research Online URL: <http://openaccess.city.ac.uk/13275/>

Copyright & reuse

City University London has developed City Research Online so that its users may access the research outputs of City University London's staff. Copyright © and Moral Rights for this paper are retained by the individual author(s) and/ or other copyright holders. All material in City Research Online is checked for eligibility for copyright before being made available in the live archive. URLs from City Research Online may be freely distributed and linked to from other web pages.

Versions of research

The version in City Research Online may differ from the final published version. Users are advised to check the Permanent City Research Online URL above for the status of the paper.

Enquiries

If you have any enquiries about any aspect of City Research Online, or if you wish to make contact with the author(s) of this paper, please email the team at publications@city.ac.uk.

\mathcal{PT} -Symmetric Interpretation of Double-Scaling

Carl M Bender^{a,‡}, Moshe Moshe^{b,§}, Sarben Sarkar^{c,||}

^aDepartment of Physics, Washington University, St. Louis MO 63130, USA

^bDepartment of Physics, Technion – Israel Institute of Technology, 32000 Haifa, Israel

^cDepartment of Physics, King's college London, Strand, London WC2R 2LS, UK

Abstract. The conventional double-scaling limit of an $O(N)$ -symmetric quartic quantum field theory is inconsistent because the critical coupling constant is negative. Thus, at the critical coupling the Lagrangian defines a quantum theory with an upside-down potential whose energy appears to be unbounded below. Worse yet, the integral representation of the partition function of the theory does not exist. It is shown that one can avoid these difficulties if one replaces the original theory by its \mathcal{PT} -symmetric analog. For a zero-dimensional $O(N)$ -symmetric quartic vector model the partition function of the \mathcal{PT} -symmetric analog is calculated explicitly in the double-scaling limit.

PACS numbers: 11.15.Pg, 11.30.Er, 03.65.Db

Submitted to: *J. Phys. A: Math. Gen.*

1. Introduction

The techniques of \mathcal{PT} quantum mechanics have been used to solve several long-standing problems, such as the violation of unitarity in the Lee model [1] and the appearance of ghosts in the Pais-Uhlenbeck model [2] and in other field-theory models [3]. In this paper we clarify a serious problem with the double-scaling limit in quantum field theory that in our opinion has not yet been satisfactorily addressed; namely, that the critical coupling constant g_{crit} is *negative*. Since $g_{\text{crit}} < 0$, the double-scaling limit appears to be unphysical because near g_{crit} the potential is upside-down and the energy appears to be unbounded below. However, we show that if we approach the critical theory in a \mathcal{PT} -symmetric fashion, the resulting correlated limit gives a physically acceptable quantum theory. Here, we study the double-scaling limit of a zero-dimensional $O(N)$ -symmetric quartic vector field-theoretic model. (In a more detailed paper we will show that the same approach works for theories in higher dimension [4].)

‡ email: cmb@wustl.edu

§ email: moshe@technion.ac.il

|| email: sarben.sarkar@kcl.ac.uk

To explain the mathematical problem addressed in this paper we consider the partition function $Z(g)$ for a toy zero-dimensional quartic quantum field theory model:

$$Z(g) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dx e^{-x^2/2 - gx^4/4}, \quad (1)$$

where the coupling constant g is assumed at first to be positive. The weak-coupling series for $Z(g)$ is obtained by expanding the integrand of (1) in powers of g and then integrating term by term. (This is a trivial application of Watson's Lemma [5].) The result is the (divergent) asymptotic-series representation for $Z(g)$:

$$Z(g) \sim \sum_{n=0}^{\infty} (-g)^n \frac{(4n-1)!!}{4^n n!} \quad (g \rightarrow 0^+). \quad (2)$$

This series alternates in sign when $g > 0$ and it can be Borel summed to recover the integral representation (1).

The integral representation (1) for $Z(g)$ ceases to exist when g is negative. Thus, to study the behavior of $Z(g)$ for negative g we evaluate the integral for positive g in terms of a parabolic cylinder function [6]:

$$Z(g) = (2g)^{-1/4} e^{1/(8g)} D_{-1/2} \left(1/\sqrt{2g} \right). \quad (3)$$

We conclude from this formula that $Z(g)$ is a multiple-valued function of g ; it is defined on a four-sheeted Riemann surface and is complex when g is negative. Thus, to evaluate $Z(g)$ for $g < 0$ we must define precisely the path from positive to negative g . We obtain four possible values for $Z(g)$ depending on how we rotate in the complex- g plane from $g > 0$ to $g < 0$.

Even though $Z(g)$ is a four-valued function of g , each term in the asymptotic series (2) is single valued. The resolution of this apparent discrepancy lies in identifying the wedge-shaped region in which a series is asymptotic to the function that it represents; this region is called a *Stokes wedge*. (For a detailed discussion of Stokes wedges see Refs. [5, 7].) The angular opening of the Stokes wedge for $D_\nu(x)$ for $x \gg 1$ is $|\arg x| < 3\pi/4$ [5]. Thus, in the complex- g plane the Stokes wedge is centered about the real- g axis and the full opening angle is 3π . This means that the series (2) is asymptotic to $Z(g)$ in (3) on the negative- g axes $\arg g = \pm\pi$ as $|g| \rightarrow 0$. Thus, we have the paradoxical result that on the negative axes (2) is real while $Z(g)$ is complex. This happens because the imaginary part of $Z(g)$ is *subdominant* (exponentially small); it is of order $e^{1/g}$, as can be determined by a saddle-point calculation.

In this paper we are interested in the partition function for negative coupling constant [8]. However, we will not be interested in the complex partition function that one obtains by rotating $x^2 + gx^4$ from positive g to negative g in the complex- g plane. Rather, we will keep the coupling constant g fixed and perform the \mathcal{PT} -symmetric limit of $x^2 + gx^2(ix)^\varepsilon$ as ε goes from 0 to 2 [9]. The resulting partition function is

$$Z(g) = \frac{1}{\sqrt{2\pi}} \int_C dx e^{-x^2/2 + gx^4/4}, \quad (4)$$

where C is a complex contour of integration. The original integration path lies on the real axis when $\varepsilon = 0$. The path rotates downward into the complex plane as ε increases. When $\varepsilon = 2$, the final contour C comes inward from $x = \infty$ in the 45° wedge $-7\pi/8 < \arg x < -5\pi/8$ and goes back out to $x = \infty$ in the 45° wedge $-3\pi/8 < \arg x < -\pi/8$. As a consequence, this partition function is *real*. This is the N -dimensional version of the \mathcal{PT} -symmetric theory that is studied in this paper.

This paper is organized as follows. Correlated limits are explained in Sec. 2 using several illustrative examples. Then the uncorrelated and correlated expansions of an $O(N)$ -symmetric vector theory in zero dimensions are investigated in Secs. 3 and 4 and problems with implementing the double-scaling limit are described. To solve these problems the vector theory is reformulated as a \mathcal{PT} -symmetric model in Sec. 5, and the uncorrelated and correlated expansions of this \mathcal{PT} -symmetric model are derived in Secs. 6 and 7. Brief concluding remarks are given in Sec. 8.

2. Correlated limits

Correlated limits arise frequently in physical problems when there are two parameters, say ε and α , and ε is treated as small ($\varepsilon \ll 1$) so that it plays the role of a perturbation parameter. If we treat α as fixed, the solution $\mathcal{S}(\varepsilon, \alpha)$ to the problem is a formal perturbation series in powers of ε : $\mathcal{S}(\varepsilon, \alpha) \sim \sum_{n=0}^{\infty} a_n(\alpha)\varepsilon^n$. Usually, the perturbation series is a *divergent* asymptotic series in which each term in the series is negligible relative to the previous term as ε tends to 0; that is, $a_n(\alpha)\varepsilon^n \ll a_{n-1}(\alpha)\varepsilon^{n-1}$ as $\varepsilon \rightarrow 0$ for all n . A correlated limit occurs when we do *not* treat α as fixed, but rather allow it to tend to a limit as $\varepsilon \rightarrow 0$; that is, we take α to depend on ε : $\alpha = \alpha(\varepsilon)$.

A nontrivial correlated limit arises if we choose the functional dependence so that *all* terms in the perturbation series become comparable as $\varepsilon \rightarrow 0$. When this happens, the series undergoes a transmutation in which it depends on just *one parameter*, which we call γ . In this correlated limit the perturbation series still diverges, but we sum the series for $\mathcal{S}(\gamma)$ by using Borel summation. Correlated limits are remarkable in that $\mathcal{S}(\gamma)$ is a *universal* function that reveals the essential features of the problem while being insensitive to specific details. Often, $\mathcal{S}(\gamma)$ is *entire* (analytic for all γ).

This paper examines the correlated limit of $O(N)$ -symmetric quantum field theories. Such theories have been used to model a variety of physical phenomena because the large- N expansion in powers of $1/N$ often reveals the phase structure of the theory; quantities such as masses and Green's functions can be expressed as asymptotic series having the form $\sum_{k=0}^{\infty} a_k N^{-k}$ [10]. In the *double-scaling* limit $N \rightarrow \infty$ and $g \rightarrow g_{\text{crit}}$ in a correlated fashion in which all terms in the $1/N$ expansion are of comparable size. In the correlated limit the sum of the series is no longer dominated by early terms; the N^{-k} power in the k th term is compensated by a sizable coefficient a_k [11]. Furthermore, this limit is characterized by a *universal* function of the parameters that describes the correlated limit [11]. In the correlated limit $O(N)$ -symmetric vector models represent discretized branched polymers. Additionally, dynamically triangulated random surfaces

summed on different topologies can be represented by matrix models in the double-scaling limit [7, 12].

We have devised three elementary examples to illustrate correlated limits:

Example 1: *Behavior of a nonuniformly convergent Fourier sine series near the boundary of its interval of convergence.* Here, the number of terms N in the partial sum of the Fourier series is correlated with the distance x to the boundary. This limit is described by the Gibbs function $G(\gamma) = \text{Si}(2\gamma)$ (the sine-integral function), where $N \rightarrow \infty$, $x \rightarrow 0$, and $\gamma \equiv Nx$. The Gibbs function is entire and it is *universal* because it describes the behavior of the Fourier sine series for *any* differentiable function $f(x)$ such that $f(0) \neq 0$ and/or $f(\pi) \neq 0$. To explain the famous *universal* 18% overshoot exhibited by all nonuniformly convergent Fourier series at the boundary, we simply verify that $G'(\pi/2) = 0$ and that $G(\pi/2) = 1.18\dots$

Example 2: *Laplace's method for the asymptotic expansion of integrals.* To find the large- N behavior of the Laplace integral

$$Z(N) = \int_0^\infty dr e^{-NS(r)}, \quad (5)$$

we assume that $S'(r) > 0$ for all $r \geq 0$ and use repeated integration by parts to obtain the complete asymptotic expansion of $Z(N)$ as $N \rightarrow \infty$ [5]:

$$Z(N) \sim e^{-NS(0)} \sum_{k=1}^\infty N^{-k} \left[\frac{1}{S'(r)} \frac{d}{dr} \right]^{k-1} \frac{1}{S'(r)} \Big|_{r=0}. \quad (6)$$

This is an uncorrelated large- N expansion.

Laplace's method emerges as a correlated limit of integration by parts: Suppose now that $S'(0)$ is small [but that the higher derivatives of $S(r)$ are not small at $r = 0$]. As $S'(0) \rightarrow 0$, the k th term in (6) is approximated by

$$N^{-k} [-2S''(0)]^{k-1} [S'(0)]^{1-2k} \Gamma(k - 1/2) / \Gamma(1/2) \quad (7)$$

because this has the greatest number of powers of $S'(0)$ in the denominator. Consider the correlated limit $N \rightarrow \infty$, $S'(0) \rightarrow 0$, where $\gamma^2 \equiv N[S'(0)]^2/S''(0)$ is a fixed parameter. [We assume that $S''(0) > 0$ so that $\gamma^2 > 0$.] In this limit (6) becomes

$$Z(\gamma) \sim \frac{e^{-NS(0)}}{\sqrt{NS''(0)}} \sum_{k=0}^\infty (-2)^k \gamma^{-2k-1} \frac{\Gamma(k + 1/2)}{\Gamma(1/2)}. \quad (8)$$

This series diverges, but we can obtain its Borel sum in terms of the parabolic cylinder function $D_\nu(z)$ [6]:

$$Z(\gamma) \sim e^{-NS(0)} \exp(\gamma^2/4) D_{-1}(\gamma) / \sqrt{NS''(0)}. \quad (9)$$

The function $Z(\gamma)$ is entire. Also, it is universal because it depends only on the two numbers $S(0)$ and $S''(0)$, and thus it applies *universally* to all functions $S(r)$ with these particular values. [In contrast, the uncorrelated series (6) depends on *all* derivatives of $S(r)$ at $r = 0$.]

For the special value $\gamma = 0$, $D_{-1}(0) = \sqrt{\pi/2}$ gives the famous result

$$Z(N) \sim e^{-NS(0)} \sqrt{\pi/[2NS''(0)]} \quad (N \rightarrow \infty) \quad (10)$$

of Laplace's method applied to (5). The asymptotic formula (10) is a limiting case of the correlated limit (9) for which $S'(0) = 0$ and $S''(0) > 0$. Thus, (9) describes the approach of $Z(\gamma)$ to Laplace's asymptotic formula (10). To be precise, the asymptotic expansion of the Laplace integral is controlled by the function $S(r)$ at its Laplace point (that is, its maximum point). Laplace's formula (10) gives the asymptotic behavior of the integral for the special case in which $S(r)$ is *level* (has a vanishing derivative) at its Laplace point. Thus, the correlated limit describes in a smooth and *universal* fashion what happens as the derivative of $S(r)$ approaches 0 at the Laplace point, just as the Gibbs function describes in a smooth and universal fashion how a nonuniformly convergent Fourier series for $f(x)$ behaves as x approaches the boundary of the interval.

Example 3: *Transition in a quantum-mechanical wave function between a classically allowed region and a classically forbidden region.* This transition is described by the universal Airy function $\text{Ai}(\gamma)$, which is obtained by summing the divergent WKB series in the correlated limit $\hbar \rightarrow 0$, $x \rightarrow 0$ (where x is the distance to the turning point), with the ratio $\gamma = x^{3/2}/\hbar$ held fixed. Consider the one-turning-point problem for the Schrödinger equation $\hbar^2 \phi''(x) = Q(x)\phi(x)$, where $Q(x)$ is linear in x near the turning point at $x = 0$: $Q(x) \sim ax$ ($x \rightarrow 0$). When $x \neq 0$ the WKB approximation to $\phi(x)$ is

$$\phi_{\text{WKB}}(x) = \exp \left[\frac{1}{\hbar} \int_0^x ds \sum_{n=0}^{\infty} \hbar^n S_n(s) \right] \quad (\hbar \rightarrow 0), \quad (11)$$

where $S_0(x) = \pm \sqrt{Q(x)}$, $S_1(x) = -Q'(x)/[4Q(x)]$, and $S_n(x)$ obeys the recursion relation $2S_0(x)S_n(x) = -S'_{n-1}(x) - \sum_{j=1}^{n-1} S_j(x)S_{n-j}(x)$ for $n \geq 2$.

The WKB approximation $\phi_{\text{WKB}}(x)$ is invalid at the turning point because it blows up at $x = 0$ while the exact solution $\phi(x)$ remains finite. To see how the WKB solution behaves as x approaches 0, we choose the negative solution for S_0 : $S_0(x) \sim -\sqrt{ax}$. Then, $S_1(x) \sim -1/(4x)$ and the functions $S_n(x)$ have the asymptotic form

$$S_n(x) \sim -4^{-n} a^{1/2-n/2} x^{1/2-3n/2} s_n \quad (x \rightarrow 0), \quad (12)$$

where $s_n = (4 - 3n)s_{n-1} - \frac{1}{2} \sum_{j=1}^{n-1} s_j s_{n-j}$ for $n \geq 2$ and $s_0 = s_1 = 1$. The numerical coefficients s_n become complicated as n increases, but an easy way to understand the WKB series is to exponentiate it:

$$\exp \left[\sum_{n=2}^{\infty} \varepsilon^{n-1} \int dx S_n(x) \right] = \sum_{k=0}^{\infty} (-1)^k g_k \gamma^{-k}, \quad (13)$$

where $\gamma = a^{1/2} x^{3/2}/\hbar$. The numerical coefficients g_n in the exponentiated series (13) now have the simple form $g_k = \pi^{-1/2} \Gamma(3k + 1/2) 9^{-k} / \Gamma(2k + 1)$. The series on the right side of (13) is just the asymptotic expansion of the Airy function $\text{Ai}(z)$ as $z \rightarrow +\infty$:

$$\text{Ai}(z) \sim \frac{z^{-1/4}}{2\sqrt{\pi}} e^{-\frac{2}{3}z^{3/2}} \sum_{k=0}^{\infty} (-1)^k g_k z^{-3k/2}. \quad (14)$$

This series diverges but it is Borel summable; in the correlated limit $\hbar \rightarrow 0$, $x \rightarrow 0$, γ fixed, the solution to the Schrödinger equation is $\phi(\gamma) = c\text{Ai}(\gamma)$, where c is an arbitrary constant. This shows that the solution to the famous one-turning-point problem is a

correlated limit. The Airy function is an *entire* function of γ and it is universal because it is valid for all potentials $Q(x)$ that vanish linearly at the turning point.

3. Uncorrelated large- N series for quantum field theory in zero dimensions

The partition function

$$Z = \int d^{N+1}x \exp \left[-\frac{1}{2} \sum_{n=1}^{N+1} x_n^2 - \frac{\lambda}{4} \left(\sum_{n=1}^{N+1} x_n^2 \right)^2 \right] \quad (15)$$

represents a zero-dimensional quartic quantum field theory having $O(N+1)$ symmetry. The coupling constant λ is assumed to be *positive* so that the integral converges. To derive an uncorrelated large- N expansion of Z , we exploit the rotational symmetry by introducing the radial variable r , $\sum_{n=1}^{N+1} x_n^2 = Nr^2$, and we let $\lambda = g/N$. The partition function now takes the one-dimensional form

$$Z = \mathcal{A}_{N+1} \int_0^\infty dr e^{-NL(r)}, \quad (16)$$

where $\mathcal{A}_N = 2\pi^{N/2}/\Gamma(N/2)$ is the surface area of an N -dimensional sphere of radius 1 and $L(r) = r^2/2 + gr^4/4 - \log r$. We emphasize that we must assume that g is positive so that the integral representation (16) for the partition function Z converges.

The integral (16) is a Laplace integral and the correlated expansion of the Laplace integral (5) was considered earlier. However, in contrast with $Z(N)$ in (5), $L(r)$ is not positive. We will see that as a result, the correlated limit of (16) lies in a different universality class and is characterized by a different universal function; specifically, the universal function for (16) is an Airy function rather than a parabolic cylinder function.

The standard procedure (Laplace's method) for finding the large- N asymptotic behavior of the integral (16) begins by locating the Laplace points, which are the zeros of $L'(s) = r + gr^3 - 1/r$. Just one Laplace point $r_0 = \sqrt{(G-1)/(2g)}$, where $G \equiv \sqrt{1+4g}$, lies in the range of integration $0 \leq r < \infty$. Laplace's method relies on the crucial fact that if r_0 is a *global minimum* of $L(r)$, then as $N \rightarrow \infty$ the entire asymptotic expansion of the integral is determined by the behavior of $L(r)$ on the infinitesimal region $r_0 - \varepsilon < r < r_0 + \varepsilon$ containing r_0 . [The Laplace point r_0 is a global minimum because $L''(r) = 1 + 3gr^2 + 1/r^2$ and $L''(r_0) = 2G > 0$.] We thus Taylor expand $L(r)$ about r_0 and conclude that for large N ,

$$Z \sim \mathcal{A}_{N+1} e^{-NL(r_0)} \int_{r_0-\varepsilon}^{r_0+\varepsilon} dr e^{-NG(r-r_0)^2} \exp \left[-\sum_{k=3}^{\infty} \frac{N}{k!} L^{(k)}(r_0) (r-r_0)^k \right].$$

To evaluate this integral we translate the integration region by substituting $t = r - r_0$ and then we perform the scaling $t = u/\sqrt{NG}$. Because the sum in square brackets begins at $k = 3$, all terms in the sum are small as $N \rightarrow \infty$. Thus, we can expand the exponential in powers of u . Making only transcendentally small errors, we

extend the range of integration to $-\infty < u < \infty$, and perform the Gaussian integrals term-by-term. This yields an infinite series in powers of $1/N$:

$$Z \sim \frac{\mathcal{A}_{N+1} e^{-NL(r_0)}}{\sqrt{NG/\pi}} \sum_{k=0}^{\infty} a_k N^{-k} \quad (N \rightarrow \infty). \quad (17)$$

This is the full asymptotic expansion of Z and it is the *uncorrelated* large- N expansion of the partition function (15). Only integer powers of N^{-1} appear in this expansion; there are no half-odd-integer powers because Gaussian integrals over odd powers of u vanish. The first three coefficients in this series are explicitly

$$a_0 = 1, \quad a_1 = \frac{5 - 6G^2 - G^3}{24G^3}, \quad a_2 = \frac{385 - 924G^2 - 10G^3 + 684G^4 + 12G^5 - 143G^6}{1152G^6}.$$

4. Correlated limit of the large- N expansion (17)

Proceeding formally, we now attempt to construct a correlated large- N limit of the expansion in (17). For all terms in the expansion to have the same order of magnitude, the correlated limit must be $N \rightarrow \infty$ and $g \rightarrow g_{\text{crit}} = -1/4$ (that is, $G \rightarrow 0$) with $\gamma \equiv NG^3/2$. In this limit the asymptotic approximation (17) becomes

$$Z \sim \frac{\mathcal{A}_{N+1} e^{-NL(r_0)}}{\sqrt{NG/\pi}} \left(1 + \frac{5}{48\gamma} + \frac{385}{4608\gamma^2} + \dots \right). \quad (18)$$

We recognize that the series in (18) is the asymptotic expansion for large γ of $\text{Bi}(\gamma^{2/3})\sqrt{\pi}e^{-2\gamma/3}\gamma^{1/6}$. [This is like the series in (13) for the Airy function Ai , but it does not alternate in sign.] Thus, we are tempted to conclude that the correlated limit of the series (17) is

$$Z \sim \mathcal{A}_{N+1} e^{NL(\sqrt{2})} 2^{-1/6} \pi N^{-1/3} \text{Bi}(\gamma^{2/3}) e^{-2\gamma/3}. \quad (19)$$

However, this correlated limit is invalid because it requires that $g < 0$. Furthermore, to obtain the Airy function Bi in (19), we have had to sum a *nonalternating* divergent series; such a series is not Borel summable.

5. \mathcal{PT} -symmetric reformulation of the theory

In place of (15) we consider the $O(N+1)$ -symmetric partition function $Z = \text{Re} \int d^{N+1}x e^{-L}$, where we assume that N is an *even* integer. The Lagrangian L has the form

$$L = \frac{1}{2} \sum_{j=1}^{N+1} x_j^2 + \frac{\lambda i^\varepsilon}{2 + \varepsilon} \left(\sum_{j=1}^{N+1} x_j^2 \right)^{1+\varepsilon/2}. \quad (20)$$

The multiple integral above is taken on the real axis and it converges if $\varepsilon < 1$.

We let $\lambda = gN^{-\varepsilon/2}$ and again introduce the radial variable r by $\sum_{n=1}^{N+1} x_n^2 = Nr^2$. The crucial assumption that N is *even* allows us to extend the radial integral to the entire real- r axis:

$$Z = \frac{1}{2} \mathcal{A}_{N+1} \int_{-\infty}^{\infty} dr e^{-N\mathcal{L}(r)}, \quad (21)$$




Figure 1. The four saddle points [zeros of $\mathcal{L}'(r)$] for the case $g < 1/4$ with the up and down directions indicated by the letters U and D. The steepest-descent path is \mathcal{PT} -symmetric (left-right symmetric) and runs along the real axis between the distant pair of saddle points. Then it turns downward, curves, and asymptotes at $-\pi/4$ and at $-3\pi/4$. For large N the contribution to the integral is localized at the pair of saddle points nearest the origin.

where $\mathcal{L} = r^2/2 + gr^2(ir)^\varepsilon/(2 + \varepsilon) - \log r$. This Laplace integral is real because the integrand is \mathcal{PT} symmetric; that is, the integrand is invariant under $r \rightarrow -r$ and $i \rightarrow -i$ [9]. Without its logarithm term, \mathcal{L} has a standard \mathcal{PT} -symmetric structure that has been studied in great detail [9]. The logarithm in the exponent does not make the integral complex. [We can make \mathcal{L} explicitly \mathcal{PT} symmetric by including an additive constant $\log r \rightarrow \log(ir)$ and taking the branch cut to lie on the negative-imaginary axis. The additive constant $\log i$ has no effect on the forthcoming steepest-descent analysis.] To achieve the \mathcal{PT} -symmetric partition function (21) we have had to work in a space of *odd* dimension $N + 1$. This requirement can be traced to the absence of a distinct parity operator in even-dimensional space, where the sign of \vec{x} can be changed by a rotation.

To obtain a quartic theory as $\varepsilon \rightarrow 2$, we must redefine the boundary conditions on the integral (21) accordingly: For any $\varepsilon \geq 0$ the integral converges if the integration contour lies inside a pair of \mathcal{PT} -symmetric Stokes wedges centered about $-\pi\varepsilon/(4 + 2\varepsilon)$ and $-(4\pi + \pi\varepsilon)/(4 + 2\varepsilon)$. The wedges have angular opening $\pi/(2 + \varepsilon)$ and contain the real- r axis if $\varepsilon < 1$. As ε increases above 1, the wedges rotate downward into the complex plane and become narrower. At $\varepsilon = 2$ the wedges are centered about $-\pi/4$ and $-3\pi/4$ and have angular opening $\pi/4$.

6. Uncorrelated limit of the \mathcal{PT} -symmetric theory (21)

To find the large- N asymptotic behavior of Z in (21) when $\varepsilon = 2$, we use the *method of steepest descents* [5]. The saddle points are the zeros of $\mathcal{L}'(r) = r - gr^3 - 1/r$. There are four saddle points, which are the roots of $r_0^2 = (1 \pm \sqrt{1 - 4g})/(2g)$. If $g < 1/4$, the saddle points are all real and are shown in Fig. 1. (If $g > 1/4$, the saddle points are all imaginary, but this does not affect the asymptotic analysis of the double-scaling limit.)

We determine the *directions* of the saddle points by calculating $\mathcal{L}''(r) = 1 - 3gr^2 + 1/r^2$. At the saddle points $\mathcal{L}'' = \mp 2\sqrt{1 - 4g}$. For the minus (plus) sign the steepest-descent path moves away from the saddle point in the imaginary (real) direction. Thus, the complete steepest-descent path, as shown in Fig. 1, follows the real- r axis until it reaches the distant pair of saddle points. Then it turns downward and curves off at the angles $-\pi/4$ and $-3\pi/4$. The entire contribution to the uncorrelated limit comes from an infinitesimal region surrounding the two saddle points closest to $r = 0$.

7. Correlated limit of the \mathcal{PT} -symmetric theory

In the correlated limit the critical point is determined by requiring that $\mathcal{L}''(r) = 0$ in addition to $\mathcal{L}'(r) = 0$. The critical value of the coupling constant is $g_{\text{crit}} = 1/4$. At this value of g there is a coalescence of the two uncorrelated saddle points at $r_{\text{crit}} = \pm\sqrt{2}$, as shown in Fig. 2. The steepest curve is shown in Fig. 2 and the contribution to the integral in the double-scaling limit comes from *four* infinitesimal regions at the saddle points, two horizontal and two at 60° angles. We must now evaluate four *convergent* integrals of the form $\int_0^L dr e^{ar^2+br^3}$, where $L = \infty$ or $L = \infty e^{-i\pi/3}$. Each of these integrals separately gives a combination of Airy functions and a ${}_2F_2$ hypergeometric function. However, we will now demonstrate that when the four integrals are combined there is a dramatic simplification and the result is identical to that in (18).




Figure 2. Correlated limit of Fig. 1. At $g = g_{\text{crit}}$ the quadratic saddle points coalesce to form cubic saddle points with up and down directions indicated by U and D. The steepest path now runs along the real axis between the saddle points. At the saddle points it veers off at 60° angles, curves, and asymptotes at $-\pi/4$ and at $-3\pi/4$.

We begin with the integral $\int dr e^{-r^2+r^3/(3\sqrt{\gamma})}$, where $\gamma = G^3 N/2$ and $G = \sqrt{1-4g}$. Note that the sign of λ is reversed in (21) compared with (15). There are four saddle points of cubic type. This gives rise to two integrals for each saddle point:

$$\int_{-\infty}^0 dr + \int_0^{\infty e^{-i\pi/3}} dr + \text{complex conjugate}.$$

In the first integral we replace $r \rightarrow -r$ and in the second we replace $r \rightarrow r e^{-i\pi/3}$. The first integral then becomes $3^{1/3}\gamma^{1/6} \int_0^\infty e^{fu^2-u^3}$, where $f = -3^{2/3}\gamma^{1/3}$, and the second becomes $3^{1/3}\gamma^{1/6} e^{-i\pi/3} \int_0^\infty e^{fu^2-u^3}$, where $f = e^{i\pi/3} 3^{2/3}\gamma^{1/3}$. Next, we use the identity

$$\int_0^\infty e^{fu^2-u^3} = 2\pi e^{2f^3/27} 3^{-4/3} \text{Bi}(3^{-4/3} f^2) + \frac{f}{3} {}_2F_2\left(\frac{1}{2}, 1; \frac{2}{3}, \frac{4}{3}; \frac{4f^3}{27}\right), \quad (22)$$

where the generalized hypergeometric function is defined by the Taylor series

$${}_2F_2(a, b; c, d; z) \equiv \frac{\Gamma(c)\Gamma(d)}{\Gamma(a)\Gamma(b)} \sum_{n=0}^{\infty} \frac{\Gamma(a+n)\Gamma(b+n)}{n!\Gamma(c+n)\Gamma(d+n)} z^n.$$

For both the first integral and the second integral, $z = -4\gamma/3$. Thus, the two hypergeometric functions are real and identical, their sum exactly cancels, and only the Bi functions remain. There are four such Airy functions:

$$2\pi e^{-2\gamma/3} 3^{-4/3} \left[2\text{Bi}(\gamma^{2/3}) - e^{2i\pi/3} \text{Bi}(e^{2i\pi/3} \gamma^{2/3}) - e^{-2i\pi/3} \text{Bi}(e^{-2i\pi/3} \gamma^{2/3}) \right].$$

We simplify this expression by using the identity $\text{Bi}(z) + \omega \text{Bi}(\omega z) + \omega^2 \text{Bi}(\omega^2 z) = 0$ [6]. The final result for the integration is therefore $2\pi e^{-2\gamma/3} 3^{-1/3} \text{Bi}(\gamma^{2/3})$. This concludes the demonstration.

8. Final remarks

We have shown that the key to constructing and interpreting the double-scaling limit is continuing in ε rather than in g . In the past the way to deal with a negative-quartic potential has been to imagine an analytic continuation from positive g to negative g . However, this procedure would give complex-energy eigenvalues because the path from positive to negative g crosses branch cuts in the coupling-constant plane, which is a multsheeted Riemann surface [13]. We have used a different analytic continuation, $\varepsilon : 0 \rightarrow 2$, which does not involve the coupling constant. To do so, we have written the potential in manifestly *PT*-symmetric form. The complex potentials for all $\varepsilon \geq 0$ give real partition functions and at $\varepsilon = 2$ we have made contact with the quartic theory obtained in the conventional double-scaling limit.

CMB was supported by the U.K. Leverhulme Foundation and the U.S. Department of Energy.

- [1] C. M. Bender, S. F. Brandt, J.-H. Chen, and Q. Wang, Phys. Rev. D **71**, 025014 (2005); G. Källén and W. Pauli, Mat.-Fys. Medd. **30**, No. 7, 1-23 (1955).
- [2] C. M. Bender and P. D. Mannheim, Phys. Rev. Lett. **100**, 110402 (2008).
- [3] T. Curtright, E. Ivanov, L. Mezincescu, and P. K. Townsend, JHEP 0704:020 (2007); T. Curtright and A. Veitia, J. Math. Phys. **48**, 102112 (2007); E. A. Ivanov and A. V. Smilga, JHEP 0707:036 (2007).
- [4] We extend our analysis to one-dimensional space time (quantum mechanics) by using an $O(N)$ -symmetric Schrödinger equation (C. M. Bender, M. Moshe, and S. Sarker, in preparation). The problem here is that in the double-scaling limit the resulting potential violates the boundary conditions imposed on the original potential. For earlier papers on $O(N)$ -symmetric quantum-mechanical systems see C. M. Bender, L. D. Mlodinow, and N. Papanicolaou, Phys. Rev. D **25**, 1305 (1982); T. C. Germann, D. R. Herschbach, M. Dunn, and D. K. Watson, Phys. Rev. Lett. **74**, 658 (1995); D. K. Watson and D. Z. Goodson, Phys. Rev. A **51**, R5 (1995).
- [5] See, for example, C. M. Bender and S. A. Orszag, *Advanced Mathematical methods for Scientists and Engineers* (McGraw-Hill, New York, 1978), chap. 6.
- [6] I. S. Gradshteyn and I. M. Ryzhik, *Tables of Integrals, Series, and Products* (Academic Press, San Diego, 2000), 6th ed.
- [7] See, for example, M. Mariño, “Lectures on non-perturbative effects in large N theory, matrix models and topological strings,” Notes of lectures held at the conference of the Research Programm on Topological String Theory, Modularity & non-perturbative Physics, ESI, Vienna; Notes available under <http://www.th.physik.uni-bonn.de/People/rauch/viennamarino.pdf> and M. Mariño, JHEP **0812** (2008) 114 [arXiv:0805.3033 [hep-th]]. See also C. M. Bender and S. P. Klevansky, Phys. Rev. Lett. **105**, 031601 (2010).
- [8] Z. Ahmed, C. M. Bender, and M. V. Berry, J. Phys. A: Math. Gen. **38**, L627-L630 (2005); H. F. Jones and J. Mateo, Phys. Rev. D **73**, 085002 (2006); C. M. Bender, D. C. Brody, J.-H. Chen, H. F. Jones, K. A. Milton, and M. C. Ogilvie, Phys. Rev. D **74**, 025016 (2006).
- [9] C. M. Bender, Rep. Prog. Phys. **70**, 947 (2007).
- [10] K. G. Wilson, Phys. Rev. D **7**, 2911 (1973). The large- N limit was first used for spin models by H. E. Stanley, Phys. Rev. **176**, 718 (1968), who showed its relation to the Berlin-Kac spherical

- model (Phys. Rev. **86**, 821 (1952). See also M. Moshe and J. Zinn-Justin, Phys. Rept. **385**, 69 (2003).
- [11] S. Nishigaki and T. Yoneya, Nucl. Phys. B **348**, 787 (1991); P. Di Vecchia, M. Kato, and N. Ohta, Nucl. Phys. B **357**, 495 (1991); J. Zinn-Justin, Phys. Lett. B **257**, 335 (1991); P. Di Vecchia, M. Kato, and N. Ohta, Int. J. Mod. Phys. A **7**, 1391 (1992); H. J. Schnitzer, Mod. Phys. Lett. A **7**, 2449 (1992); P. Di Vecchia and M. Moshe, Phys. Lett. B **300**, 49 (1993); G. Eyal, M. Moshe, S. Nishigaki, and J. Zinn-Justin, Nucl. Phys. B **470**, 369 (1996).
- [12] M. Douglas and S. Shenker, Nucl. Phys. B **335**, 635 (1990); E. Brézin and V. Kazakov, Phys. Lett. B **236**, 144 (1990); D. Gross and A. A. Migdal, Phys. Rev. Lett. **64**, 127 (1990) and Nucl. Phys. B **340**, 333 (1990). For reviews see P. Di Francesco, P. Ginsparg, and J. Zinn-Justin, Phys. Rep. **254**, 1 (1995); M. C. Ogilvie and P. N. Meisinger, J. Phys. A: Math. Theor. **41**, 244021 (2008).
- [13] C. M. Bender and T. T. Wu, Phys. Rev. **184**, 1231 (1969).