

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Tahita, MC; Tinto, H; Yarga, S; Kazienga, A; Traore/Coulibaly, M; Valea, I; Van Overmeir, C; Rosanas-Urgell, A; Ouedraogo, JB; Guiguemde, RT; van Geertruyden, JP; Erhart, A; D'Alessandro, U (2015) Ex vivo anti-malarial drug susceptibility of *Plasmodium falciparum* isolates from pregnant women in an area of highly seasonal transmission in Burkina Faso. *Malar J*, 14 (1). p. 251. ISSN 1475-2875 DOI: <https://doi.org/10.1186/s12936-015-0769-1>

Downloaded from: <http://researchonline.lshtm.ac.uk/2212687/>


DOI: [10.1186/s12936-015-0769-1](https://doi.org/10.1186/s12936-015-0769-1)

Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by/2.5/>

Additional file: Figure S1 Scatter of the IC50 values with geometric mean and the resistance cutoff (where available) of each drug



— Geometric means IC 50

— Resistance threshold