

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Peltier, J; Shaw, HA; Couchman, EC; Dawson, LF; Yu, L; Choudhary, JS; Kaever, V; Wren, BW; Fairweather, NF (2015) Cyclic-di-GMP regulates production of sortase substrates of *Clostridium difficile* and their surface exposure through ZmpI protease-mediated cleavage. *The Journal of biological chemistry*. ISSN 0021-9258 DOI: 10.1074/jbc.M115.665091

Downloaded from: <http://researchonline.lshtm.ac.uk/2281267/>

DOI: [10.1074/jbc.M115.665091](https://doi.org/10.1074/jbc.M115.665091)

Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by/2.5/>

Cyclic-di-GMP regulates production of sortase substrates of *Clostridium difficile* and their surface exposure through ZmpI protease-mediated cleavage*

Johann Peltier¹, Helen A. Shaw¹, Edward C. Couchman¹, Lisa F. Dawson², Lu Yu³, Jyoti S. Choudhary³, Volkhard Kaever⁴, Brendan W. Wren² and Neil F. Fairweather¹

Supplemental material

Table S1. Strains, plasmids and oligonucleotides used for the present study.

Strain, plasmid or primer	Characteristic/description	Source/construction
Strains		
<i>E. coli</i> NEB5 α	General cloning	New England Biolabs
<i>E. coli</i> Rosetta	Protein expression	Novagen
<i>E. coli</i> CA434	Conjugation donor	(1)
<u><i>C. difficile</i> strains</u>		
630	Wild type Erm ^R . Virulent and multidrug-resistant PCR ribotype 01	(2)
630 Δ <i>erm</i>	Erythromycin-sensitive derivative of wild-type <i>C. difficile</i> 630	(3)
Δ <i>srtB</i>	630 isogenic <i>srtB</i> deletion mutant	This study
Δ <i>zmpI</i>	630 isogenic <i>zmpI</i> deletion mutant	This study
Δ <i>srtB</i> - Δ <i>zmpI</i>	630 isogenic double mutant with deletions in the aforementioned genes	This study
<i>P</i> _{ter} - <i>srtB</i>	Insertion of the P _{ter} promoter upstream <i>srtB</i> of the 630 strain by allele exchange	This study
Δ <i>srtB</i> _{comp}	Δ <i>srtB</i> complemented on the chromosome by allele exchange	This study
Plasmids		
pET28a	<i>E. coli</i> expression vector. Kan ^R	Novagen
pUC19	<i>E. coli</i> cloning vector. Amp ^R	New England Biolabs
pMTL-SC7215	<i>E. coli</i> - <i>C. difficile</i> shuttle vector for <i>codA</i> -mediated allele exchange mutagenesis. Tm ^R	(4)
pRPF144	<i>E. coli</i> - <i>C. difficile</i> shuttle vector for protein expression. <i>P</i> _{cwp2} - <i>gusA</i> , Tm ^R	(5)
pRPF185	<i>E. coli</i> - <i>C. difficile</i> shuttle vector for protein expression. <i>P</i> _{ter} - <i>gusA</i> , Tm ^R	(5)
pHAS027	CD2831 (32-290)-His ₆ in pET28a	This study
pHAS042	His ₆ -CD27L (1-179) in pET28a	This study
pJKP001	SrtB (27-225)-His ₆ in pET28a	This study
pJKP019	pUC19 with construct for <i>srtB</i> deletion	This study
pJKP022	pMTL-SC7215 with construct for <i>srtB</i> deletion	This study

pJKP029	pMTL-SC7215 with construct for <i>P_{ter}-srtB</i> strain creation	This study
pJKP041	pRFP185 derivative carrying <i>P_{ter}-CD2831</i> for inducible CD2831 expression	This study
pJKP070	pRFP185 derivative carrying <i>P_{ter}-CD3392</i> with addition of a HA-tag encoding sequence into <i>CD3392</i> and a functional RBS upstream <i>CD3392</i> for inducible CD3392 _{HA} expression	This study
pJKP088	pMTL-SC7215 with construct for <i>zmpI</i> deletion	This study
pJKP095	pJKP041 with addition of R-HA-tag encoding sequence into <i>CD2831</i>	This study
pJKP096	pMTL-SC7215 with construct for <i>srtB</i> complementation	This study
pECC12	pRFP144 derivative carrying <i>P_{cwp2-}dccA</i> with addition of His-tag encoding sequence into 3'-extremity of <i>dccA</i> for constitutive expression of DccA.	This study
pECC17	pRFP185 derivative carrying <i>P_{ter}-dccA</i> with addition of His-tag encoding sequence into 3'-extremity of <i>dccA</i> for inducible expression of DccA.	This study

Primer	Sequence (5' to 3')*	Characteristics or use
NF1614	GGGCCATGGATTTCAGAATTAG GAGAGAATAGTCAGATTC	Amplify 5' extremity of CD2831 for pET28a cloning
NF1615	GGGCTCGAGTTCTATTTTTCC AGTAGTTTCAATATGTATTTT TG	Amplify 5' extremity of CD2831 for pET28a cloning
NF1968	ATATCCATGGAACCTTACCAAA TACAATCATGATACTA	Amplify <i>srtB</i> deleted in the N-terminal hydrophobic domain for pET28a cloning
NF1969	TGAGCTCGAGAATCAATCTAC CATGAATCACCAT	Amplify <i>srtB</i> deleted in the N-terminal hydrophobic domain for pET28a cloning
NF2126	GATCGAGCTCGTATTTTATTT TGGAGAAATTAATATGTTTAA AG	Amplify <i>dccA</i> with the addition of a C-terminal His-tag encoding sequence for pRFP144 or pRFP185 cloning.
NF2155	GATAGAATTCCCATTTTTAAT ATCTGTACTAACTTATATTAC	Construction of pJKP019 for <i>srtB</i> deletion
NF2156	GATAGGATCCGAGAAGACCT GTCCTTAGTAC	Construction of pJKP019 for <i>srtB</i> deletion
NF2157	GCCAGAATGGTGATTCATGG	Construction of pJKP019 for <i>srtB</i>

NF2158	CTGTAAAGTATTACTAATACT AGTATAATGT	deletion Construction of pJKP019 for <i>srtB</i> deletion
NF2169	GGATTTACATTTGCCGTTTT GTAAAC	Screen for insert into pMTL- SC7215
NF2170	GATCTTTTCTACGGGGTCTGA C	Screen for insert into pMTL- SC7215
NF2171	GCTACATCTTATATCATATCA TATTATTTACATC	Screen for single and double crossover for <i>srtB</i> deletion
NF2172	CGGTCCCTATCCGTCGC	Screen for single and double crossover for <i>srtB</i> deletion
NF2199	GATCGGATCCTTAATGGTGAT GGTGATGGTGCTCGAGATAAT CATTTTTATCAAATTTTTTCTT GTTTTTC	Amplify <i>dccA</i> with the addition of a C-terminal His-tag encoding sequence for pRFP144 or pRFP185 cloning.
NF2214	AAACTCCTTTTTGATAATCTC ATGACC	Linearize pMTL-SC7215 by inverse PCR at the PmeI site
NF2215	AAACTTAGGGTAACAAAAAA CACCG	Linearize pMTL-SC7215 by inverse PCR at the PmeI site
NF2254	gTTTTgttaccctaagtttGTAATATTTA CATTGTGCAAAGTTG	Construction of pJKP029 for creation of P_{tet} - <i>srtB</i> strain
NF2257	atgtagttaaGGAGCTCAGATCTGT TAAC	Construction of pJKP029 for creation of P_{tet} - <i>srtB</i> strain
NF2258	tgagctccTAAACTACATTTAACT CGAGGTG	Construction of pJKP029 for creation of P_{tet} - <i>srtB</i> strain
NF2259	gattatcaaaaaggagtttCAATCTACCA TGAATCACCATTC	Construction of pJKP029 for creation of P_{tet} - <i>srtB</i> strain
NF2260	CACTCAAAACCTTCACTCCTT AAATC	Screen for single and double crossover for creation of P_{tet} - <i>srtB</i> strain
NF2261	CAGTTTAAAATTATTGTTTTA AACTGTGTCTG	Screen for single and double crossover for creation of P_{tet} - <i>srtB</i> strain
NF2266	cagaattcgTATATCTATATTACAC TATTTTTGAATAAATTTAAAT TTTTAG	Construction of pJKP029 for creation of P_{tet} - <i>srtB</i> strain
NF2267	atatagatataCGAATTCTGCATCAA GCTAG	Construction of pJKP029 for creation of P_{tet} - <i>srtB</i> strain
NF2378	GATAGAGCTCAAAATGAAAG GAGCATTAGATTTATG	Amplify <i>CD3392</i> for pRFP185 cloning
NF2379	GATAGGATCCTTATGATTTCT TCATTTTACGGC	Amplify <i>CD3392</i> for pRFP185 cloning
NF2451	GGATCCTATAAGTTTTAATAA AACTTTAAATAG	Remove <i>CD3392</i> coding sequence by inverse PCR
NF2501	GATAGAGCTCGGAAGATTAG ATTATAAATTATAGGTAAC	Amplify <i>CD2831</i> for pRFP185 cloning
NF2502	GATAGGATCCTAATTTGTATT TTTATTTCTTCTTAATACGATA AG	Amplify <i>CD2831</i> for pRFP185 cloning
NF2726	TTCCAGATTATGCTGGTCAGA	Add a sequence encoding for HA

	CACCAGATGTAGA	tag into the CD3246 coding sequence by inverse PCR
NF2727	CATCATATGGATAATTTGATT GTTCTTCTCCATTAAGTATTG	Add a sequence encoding for HA tag into the CD3246 coding sequence by inverse PCR
NF2858	AAATCTAATGCTCCTTTCATT TTGAGC	Remove CD3392 coding sequence by inverse PCR
NF2860	gaaaggagcattagatttTTGAAAACAA AAATTA AAAAATCAAGTATA ATC	Amplify <i>CD3246</i> for pRFP185 cloning
NF2861	attaaaacttataggatccTTAATTTCTT TTTATTTTACATTGATACGC	Amplify <i>CD3246</i> for pRFP185 cloning
NF2922	gtttttgttaccctaagtttCATACCATAT TATCCAATATGCT	Construction of pJKP088 for <i>zmpI</i> deletion
NF2923	atcaatattcCAAACCTTATAGCTTTT TGCAA AATTTAG	Construction of pJKP088 for <i>zmpI</i> deletion
NF2924	tataagtttgGAATATTGATATTAT AGCTATTAATAATTTTTTACTT G	Construction of pJKP088 for <i>zmpI</i> deletion
NF2925	gattatcaaaaaggagtttGGGGAGACT ATACGAAATG	Construction of pJKP088 for <i>zmpI</i> deletion
NF2926	CCTGTGCAAAGTTGGGTTAC	Screen for single and double crossover for <i>zmpI</i> deletion
NF2956	CTTTACCAGCATTGAAAAAT TAGCC	Screen for single and double crossover for <i>zmpI</i> deletion
NF2958	CAGATTATGCTGTAAATCCAC CAGTACCACC	Add a sequence encoding for the R-HA tag into the CD2831 encoding gene of pJKP041 by inverse PCR
NF2959	GAACATCATATGGATACCGCA AAGTATCATCTTTAACTTTAT TATTTTTTTC	Add a sequence encoding for the R-HA tag into the CD2831 encoding gene of pJKP041 by inverse PCR
NF2966	gtttttgttaccctaagtttCGTCTTCTTC TCTCTGATAG	Construction of pJKP096 for complementation of <i>srtB</i> mutant
NF2967	tcaatctacCATGAATCACCATTCT GGC	Construction of pJKP096 for complementation of <i>srtB</i> mutant
NF2968	tgattcatgGTAGATTGATTTAGAA ACTCACAGACAC	Construction of pJKP096 for complementation of <i>srtB</i> mutant
NF2969	gattatcaaaaaggagtttCGGAAAGAC CCCATGGAG	Construction of pJKP096 for complementation of <i>srtB</i> mutant
NF2970	GAAGGTCTTTATATAGGTTTA AAACC	Screen for single and double crossover for complementation of <i>srtB</i> mutant
NF2971	CGGATACCTGCGACAGG	Screen for single and double crossover for complementation of <i>srtB</i> mutant

*Underlined bases indicate engineered restriction sites; lowercase bases indicate overlapping sequences

REFERENCES

1. Purdy, D., O'Keeffe, T. A., Elmore, M., Herbert, M., McLeod, A., Bokori-Brown, M., Ostrowski, A., and Minton, N. P. (2002) Conjugative transfer of clostridial shuttle vectors from *Escherichia coli* to *Clostridium difficile* through circumvention of the restriction barrier. *Mol Microbiol* **46**, 439-452
2. Sebahia, M., Wren, B. W., Mullany, P., Fairweather, N. F., Minton, N., Stabler, R., Thomson, N. R., Roberts, A. P., Cerdeno-Tarraga, A. M., Wang, H., Holden, M. T., Wright, A., Churcher, C., Quail, M. A., Baker, S., Bason, N., Brooks, K., Chillingworth, T., Cronin, A., Davis, P., Dowd, L., Fraser, A., Feltwell, T., Hance, Z., Holroyd, S., Jagels, K., Moule, S., Mungall, K., Price, C., Rabbinowitsch, E., Sharp, S., Simmonds, M., Stevens, K., Unwin, L., Whithead, S., Dupuy, B., Dougan, G., Barrell, B., and Parkhill, J. (2006) The multidrug-resistant human pathogen *Clostridium difficile* has a highly mobile, mosaic genome. *Nat Genet* **38**, 779-786
3. Hussain, H. A., Roberts, A. P., and Mullany, P. (2005) Generation of an erythromycin-sensitive derivative of *Clostridium difficile* strain 630 (630 Δ erm) and demonstration that the conjugative transposon Tn916E enters the genome of this strain at multiple sites. *J Med Microbiol* **54**, 137-141
4. Cartman, S. T., Kelly, M. L., Heeg, D., Heap, J. T., and Minton, N. P. (2012) Precise Manipulation of the *Clostridium difficile* Chromosome Reveals a Lack of Association Between tcdC Genotype and Toxin Production. *Appl Environ Microbiol* **78**, 4683-4690
5. Fagan, R. P., and Fairweather, N. F. (2011) *Clostridium difficile* has two parallel and essential Sec secretion systems. *J Biol Chem* **286**, 27483-27493