

RESEARCH ARTICLE

Associations between Common Variants in Iron-Related Genes with Haematological Traits in Populations of African Ancestry

Wanjiku N. Gichohi-Wainaina^{1,2*}, Toshiko Tanaka³, G. Wayne Towers⁴, Hans Verhoef^{5,6,7}, Jacobien Veenemans^{8,9}, Elise F. Talsma^{1,10}, Jan Harryvan¹, Mark V. Boekschoten¹¹, Edith J. Feskens¹, Alida Melse-Boonstra¹

1 Division of Human Nutrition, Wageningen University, Wageningen, the Netherlands, **2** International institute of Tropical Agriculture (IITA), Dar es Salaam, Tanzania, **3** Translational Gerontology Branch, National Institute on Aging, Baltimore, MD, United States of America, **4** Centre of Excellence for Nutrition, North-West University (Potchefstroom campus), Potchefstroom, South Africa, **5** Cell Biology and Immunology Group, Wageningen University, Wageningen, The Netherlands, **6** Medical Research Council (MRC) International Nutrition Group, London School of Hygiene and Tropical Medicine, London, United Kingdom, **7** Medical Research Council Banjul, The Gambia, **8** Laboratory for Microbiology and Infection Control, Amphia Hospital, Breda, The Netherlands, **9** Department of Medical Microbiology and Immunology, Admiraal De Ruyter Hospital, Goes The Netherlands, **10** HarvestPlus, International Center for Tropical Agriculture (CIAT), Cali, Colombia, **11** Nutrition, Metabolism and Genomics group, Division of Human Nutrition, Wageningen University, Wageningen, The Netherlands

* wngichohi@gmail.com; W.gichohi@cgiar.org



CrossMark
click for updates

 OPEN ACCESS

Citation: Gichohi-Wainaina WN, Tanaka T, Towers GW, Verhoef H, Veenemans J, Talsma EF, et al. (2016) Associations between Common Variants in Iron-Related Genes with Haematological Traits in Populations of African Ancestry. PLoS ONE 11(6): e0157996. doi:10.1371/journal.pone.0157996

Editor: David C. Samuels, Vanderbilt University Medical Center, UNITED STATES

Received: January 14, 2016

Accepted: June 8, 2016

Published: June 22, 2016

Copyright: This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the [Creative Commons CC0](https://creativecommons.org/licenses/by/4.0/) public domain dedication.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: The Kenyan cohort research was conducted within the framework of INSTAPA Project, funded by the European Union's Seventh Framework Programme [FP7/2007-2013] under grant agreement nr 211484. The Tanzanian cohort research was supported through funding from the Netherlands Organisation for Scientific Research (NWO)/WOTRO [W93-413, WAO93-441, as well as WIZ93-465, UNICEF, Cornelis Visser Foundation and Wageningen University (Interdisciplinary Research

Abstract

Background

Large genome-wide association (GWA) studies of European ancestry individuals have identified multiple genetic variants influencing iron status. Studies on the generalizability of these associations to African ancestry populations have been limited. These studies are important given interethnic differences in iron status and the disproportionate burden of iron deficiency among African ancestry populations.

Methods

We tested the associations of 20 previously identified iron status-associated single nucleotide polymorphisms (SNPs) in 628 Kenyans, 609 Tanzanians, 608 South Africans and 228 African Americans. In each study, we examined the associations present between 20 SNPs with ferritin and haemoglobin, adjusting for age, sex and CRP levels.

Results

In the meta analysis including all 4 African ancestry cohorts, we replicated previously reported associations with lowered haemoglobin concentrations for rs2413450 ($\beta = -0.19$, $P = 0.02$) and rs4820268 ($\beta = -0.16$, $P = 0.04$) in *TMPRSS6*. An association with increased ferritin concentrations was also confirmed for rs1867504 in *TF* ($\beta = 1.04$, $P = <0.0001$) in the meta analysis including the African cohorts only.

and Education Fund). The PURE study was supported through grants provided by the South African Medical Research Council (Self-Initiated Research Grants), Cape Town, South Africa, with further support from a fellowship provided by Unilever R&D, Vlaardingen, The Netherlands, and from the European Union (EU) funded staff exchange European-South African Partnership in Nutrition Research program (grant MC-IRSES-2008-230786). The BLSA was supported through funding by the Intramural Research Program of the NIH, National Institute on Aging. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Conclusions

In all meta analyses, we only replicated 4 of the 20 single nucleotide polymorphisms reported to be associated with iron status in large GWA studies of European ancestry individuals. While there is now evidence for the associations of a number of genetic variants with iron status in both European and African ancestry populations, the considerable lack of concordance highlights the importance of continued ancestry-specific studies to elucidate the genetic underpinnings of iron status in ethnically diverse populations.

Introduction

Body iron status is influenced by a combination of genes as well as environmental factors such as diet, blood loss, pregnancy, alcohol intake and infections [1, 2]. In order to maintain iron levels within normal ranges and thus prevent iron excess or iron deficiency, iron homeostasis has evolved as a complex trait and a tightly coordinated process of interaction between genes and environmental factors. Whether genetic influences on iron metabolism may influence iron status at the public health level, is a subject of recent interest. This is of interest since segments of populations that are genetically prone to iron deficiency can be specifically targeted with interventions.

Understanding the genetic underpinnings of low iron status is particularly relevant for African ancestry populations, who experience a disproportionately high burden of low iron status. In particular, it has been observed that in the US, African-Americans have lower haemoglobin concentrations and serum transferrin saturation (TS), and higher serum ferritin (SF) concentrations compared to individuals of European Ancestry [3]. Furthermore, according to WHO regional prevalence estimates, Africa has the highest proportion of individuals affected by anaemia worldwide [4].

Previous studies on iron status report heritability estimates ranging from of 20% to 30%, supporting a genetic component to iron status [5, 6]. Genome-wide association (GWA) studies have identified common variants that are associated with iron status in largely European ancestry populations [7–10]. It is questionable however whether these findings can be generalized to African ancestry populations. Given the greater levels of linkage disequilibrium among European-ancestry individuals [11], it is expected that index single nucleotide polymorphisms (SNPs) tag larger regions of the genome than they would among African ancestry individuals. Therefore index SNPs in European populations may not necessarily tag causal variants within the African population. Moreover, African populations are exposed to very different environmental factors that further influence iron status. Recently, genetic studies on iron status have been conducted in the African American population [12–16]. This population has varying degrees of admixture that may complicate transferability of these results to African populations [17]. Conclusive inferences about the potential generalization of previous findings to African ancestry populations have been precluded by the lack of data from African cohorts. In part this has been due to the fact that it is challenging to find African cohorts which can be utilized in genetic epidemiologic studies. In the present study, we utilized data from four African ancestry cohorts that were initiated with different primary hypotheses but had collected the necessary phenotypic and genetic data to address the question regarding the genetic architecture of iron metabolism. We investigated 20 SNPs previously reported to be associated with iron status in GWA studies [7, 8, 10, 14, 18] with haematological indicators in several African ancestry populations.

Methods

Study area and population

Four cohorts of African ancestry were selected; from Kenya (n = 628), Tanzania (n = 609); South Africa (n = 686) and the United States of America (n = 228). The four cohorts differed largely in age, gender and burden of inflammation as described in more detail below.

Kenyan cohort. Samples were obtained from both a cross sectional study and from the baseline data of a randomized trial (*Cassavita study*) designed to assess the efficacy of bio-fortified cassava to improve vitamin A deficiency [19, 20]. The cross sectional study was conducted in June 2010. The randomized trial (*Cassavita study*) was conducted from May until November 2012 in 3 primary schools in Kibwezi District, Kenya. Haematological markers were measured prior to the start of the trial. As determined by self-report in both studies children were of Akamba (Bantu) ancestry. Blood samples for the cross-sectional study were collected from children aged 6–12 years (n = 375) who were located in the Kibwezi and Makindu Districts in Eastern Kenya, which is a low malarial transmission area [21]. Blood samples for the randomized trial were collected from children aged 5–13 years (n = 342) in three primary schools located in the Kibwezi District. Children with lowest retinol-binding protein concentrations were included in the trial to select for those with the lowest vitamin A status. All subjects (for both the trial and cross sectional study) were apparently healthy and without fever (ear drum temperature < 37.5°C) upon examination by the research physician and children with inflammation (C-reactive protein concentration > 8mg/L), low haemoglobin concentrations (<70 g/L), or *Plasmodium* infection were excluded.

The cross sectional study as well as the trials were approved by ethical committees in Kenya and the Netherlands (ClinicalTrials.gov:NCT01614483). Written informed consent was obtained from parents or guardians, and assent from children.

Tanzanian cohort. Children in this cohort were part of a randomized placebo controlled trial (*MACH study*) [22]. Data were collected between February 2008 and March 2009. Haematological markers as were measured in samples collected prior to the start of the trial and ancestry was self-reported. People in this area originally belonged to the Wazigua and Wabondei Bantu tribes, but settlement of migrant plantation workers in the past has resulted in the admixture of many tribes with different origins. Samples were obtained from children aged 6–60 months (n = 612) with height-for-age z-score ≤ -1.5 SD in a rural area of Handeni District in North-Eastern Tanzania, which has intense year-round *P. falciparum* transmission. Children with severe wasting (weight-for-age z-score < -3 SD), haemoglobin concentration < 70 g/L, *Plasmodium* infection or signs of chronic illness were excluded.

The study was approved by the Ethical Review Committee of Wageningen University, The Netherlands (ClinicalTrials.gov: NCT00623857) and the National Health Research Ethics Review sub-Committee, Dar es Salaam, Tanzania. Written individual informed consent was obtained from parents or primary caretakers.

South African cohort. This study population was a subset of Tswana speaking pre- and post-menopausal women (n = 686) enrolled as part of the South African arm of the Prospective Urban and Rural Epidemiological (PURE) study. The PURE study began in 2005. Haematological measures utilized in this study were collected in a follow up study conducted in 2010. The participants were from the North-West Province in South Africa, aged 35 years or older and were mainly of Tswana (Bantu) ancestry as determined by self-report. Women were only eligible for this sub-study if they were apparently healthy, not pregnant, non-lactating and were at the time of blood collection not diagnosed with any chronic disease or taking drugs for any chronic disease. Additionally, we included only participants with all outcome measures determined and who were HIV negative. HIV testing was undertaken with the appropriate pre- and

post-test counselling as required by South African law. This study was approved by the Ethics Committee of the North-West University and signed informed consent forms were obtained from all participants.

African American cohort. This cohort consisted of participants from the Baltimore Longitudinal Study on Aging (BLSA). The BLSA is a population-based study aimed at evaluating contributors of healthy aging in the older population residing predominantly in the Baltimore-Washington DC area [23]. Started in 1958, participants aged 17–94 years were examined every one to four years depending on their age. Currently, there are approximately 1,100 active participants enrolled in the study. For the present study, 228 subjects of African ancestry with complete data on both genetic and haematological traits were available and were used for the analysis. The BLSA has continuing approval from the Institutional Review Board (IRB) of Medstar Research Institute, Johns Hopkins, Bayview Medical Centre, and the University of Maryland. Additionally, BLSA study participants provide written informed consent to participate in both direct and ancillary studies related to their collected data.

Iron related measurements

Kenyan and Tanzanian cohorts. For the cross sectional study (Kenyan cohort), serum ferritin concentrations were determined by enzymatic immunoassay (Ramco Laboratories, Stafford Texas) and haemoglobin concentration (Hb) was assessed using a hematology analyzer (Celltac- α , MEK-6410K, Nihon Kohden Corporation, Tokyo, Japan). Serum concentrations of CRP were determined by immunoturbidometric assay on a Cobas Integra 800 system (Roche Diagnostics, Mannheim, Germany). For the Kenyan randomized trial as well as the Tanzanian cohort, concentrations of ferritin and concentrations of CRP were measured on a Beckman Coulter UniCel DxC 880i analyzer as per manufacturer's instructions.

South African cohort. Serum ferritin was determined using an enzyme immunoassay procedure (Ramco Laboratories, Inc, Stafford Texas). Hb concentrations were determined spectrophotometrically from whole blood using the Hemocue[®] HB201 haemoglobin meter (Hemocue AB, Angelholm Sweden). Serum concentrations of CRP were measured from serum via a particle-enhanced immunoturbidometric assay using Sequential Multiple Analyser Computer (SMAC) using the Konelab20i[™] auto analyser (Thermo Fisher Scientific Oy, Vantaa, Finland). Whole blood was used for the determination of HIV status making use of the First Response (PMC Medical, India) rapid HIV Card Test. If tested positive, the test was repeated with the Pareeshak (BHAT Bio-tech India) card test.

African American cohort. Serum ferritin was measured using an immunoassay type two-stage sandwich method using two antiferritin antibodies (Advia Centaur, Bayer). Haemoglobin was assessed using SYSMEX XE-series (Sysmex Corporation). Serum concentrations of CRP were measured using the BNII nephelometer from Dade Behring utilizing a particle enhanced immunonephelometric assay.

SNP selection, genotyping and quality control procedures

SNP selection. To set up the assay for genotyping it was required to develop a list of reference SNPs (rs) numbers from Genbank to be included in the analysis. This list was set up by performing a literature search of Pubmed, Sciencedirect, Ebscohost, HUGE navigator and Google Scholar using the following keywords:

1. Iron, ferritin, transferrin
2. Anaemia
3. Haemochromatosis

4. Human
5. SNP, Polymorphism, Mutation
6. Genetic

All abstracts were scanned for appropriateness of use, that is, association or linkage of a SNP to the haematological parameters or key biomarkers of iron status (Hb, SF, serum transferrin receptor (sTfR) and body iron (BI)). If appropriate, the reference SNP (rs) number of the polymorphism was recorded and included in the list to be sent to Illumina for verification of use in the array. Based on this, we selected 32 SNPs for genotyping. Once SNPs were selected, genotyping was performed for the Kenyan, Tanzanian and South African cohorts. These data were complemented with the data we obtained from the African American cohort in the BLSA.

Genotyping and quality control procedures. For the Kenyan, Tanzanian and South African cohorts, all 32 rs numbers were submitted to Illumina Technical Support for evaluation using the Assay Design Tool. Each SNP was scored (varying from 0 to 1) by the same, based on its compatibility for use in a GoldenGate genotyping assay. SNPs with a score above 0.4 were chosen for genotyping. No SNP fell out based on these criteria. Genotyping for Kenyan (efficacy trial only), South African and Tanzanian cohorts was conducted using the BeadXpress™ platform. Once genotyping was performed, several quality control procedures were applied. Lack of a detectable signal after analysis was designated as a failed SNP. SNP clustering was also assessed visually to determine success of genotyping. SNPs with a GenCall Score of >0.5 and samples with a call rate of ≥ 0.95 were included in the final analysis. Genotyping was performed for only three SNPs in *TMPRSS6* (rs2413450, rs4820268, rs228918), for the Kenyan pilot study samples using Applied Biosystems TaqMan Assays. Data obtained with this method were merged with the data from the Kenyan efficacy trial. Eventually, we considered only 20 SNPs that had passed all quality control procedures in all four cohorts for association analyses (S1 Table).

For the African American cohort, genome-wide genotyping was conducted using the Illumina Infinium HumanHap 550K. Genotyping was completed for 284 subjects, and imputation of ~2.5 million HapMap SNPs was conducted using 501,764 SNPs that passed quality control (minor allele frequency $> 1\%$, genotyping completeness $> 99\%$, and Hardy Weinberg-equilibrium > 0.0001). Imputation was conducted with MACH using the HapMap YRI sample (HapMap data release rel#23a NCBI B36) as a reference. The candidate iron-homeostasis SNPs that were either directly genotyped or imputed (with an imputation quality of greater than $r^2 > 0.3$) were used for the present analysis.

Statistical analyses

Outcome variables (haemoglobin and serum ferritin concentrations) were determined to be continuous quantitative traits. Anaemia was defined using different cut offs for each cohort as cut offs are age dependent [24]. Cut offs were as follows: Hb concentration < 11.5 g/dL (Kenyan cohort), < 11 g/dL (Tanzanian cohort), < 12 g/dL for the South African and African American cohorts, while iron deficiency was defined as serum ferritin concentration < 15 μ g/L for Kenyan, South African and African American samples, and plasma ferritin concentration < 12 μ g/L for Tanzanian samples. \log_{10} transformations were applied to the ferritin outcomes to improve their fit to the normal distribution. Results from associations involving ferritin were then presented as back-transformed data. Potential disturbances in the distributions of the genotypes were tested for deviation from the assumptions of Hardy-Weinberg equilibrium (HWE) by using the Chi-square test. To compare minor allele frequencies (MAFs) across cohorts, a spider web graph was generated using Microsoft Excel (Version 2010), including European ancestry

populations from the 1000 Genomes Project [20]. Linear regression was used to assess associations between SNPs and outcome measures. For association testing, genotypes were coded as 0, 1, and 2 reflecting the number of copies of the minor allele (additive genetic model), while imputed genotypes were coded using fractional counts between 0 and 2 according to the estimated number of copies of each minor allele. All associations were adjusted for age and gender. Additionally, associations involving ferritin concentrations were adjusted for inflammation by including CRP concentrations as a covariate in the model. Statistical analyses were performed using version 20 of the Statistical Package for Social Sciences (SPSS Inc Chicago III) or SAS version 9.2 (Cary, NC). All reported P-values were two-tailed, and adjustment for multiple testing was performed using the false discovery rate (FDR) method. The R program for statistical computing version 2.15.2 [25] was used to perform all meta-analyses and chi square tests conducted to test for heterogeneity between studies. We performed the meta analyses in order to increase the sample size and obtain a summary statistic. Comparisons of effect sizes and directions were made within and across populations, and meta-analyses were performed to generate summary effect estimates with and without including the African American cohort. Statistical significance for all analyses was defined as $\alpha < 0.05$, while the q values cut-off was set at 0.10.

Results

There were large differences in age between the cohorts, as the studies from Kenya and Tanzania were pre- and school going children, while the studies from South Africa and the United States were of adults (Table 1). With the exception of the South African cohort which was all women, the sex distribution was approximately equal. Prevalence of inflammation was below 5% in the Kenyan and African American cohort and 33% and 43% in the Tanzanian and South African cohort, respectively. Iron deficiency was highest in the Kenyan cohort (37%) and lowest in the African American cohort (3%). Anaemia was highest in the Tanzanian cohort (32%) and lowest in the Kenyan cohort at (7%).

In comparing minor allele frequencies across cohorts, the SNPs genotyped for the present study were generally observed to be more common in European ancestry cohorts as reported in the 1000 Genomes Project database (Fig 1). For example, for the commonly investigated rs4820268 SNP in the *TMPRSS6* gene, the frequency of the allele associated with lower iron

Table 1. General characteristics of the study participants in the four cohorts.

	<i>n</i>	Kenya	<i>n</i>	Tanzania	<i>N</i>	South Africa	<i>N</i>	USA (African American)
Age, y	628	9.0 (2.1)	609	2.7 (1.3)	686	50.0 (10.0)	228	67.4 (11.0)
Gender (% female)	628	52	609	51	686	100	228	59
Haemoglobin (g/dL)	626	13.9 (1.6)	607	10.3 (12.8)	481	13.8 (1.6)	228	13.0 (1.4)
Ferritin (µg/L) [§]	623	17.8 (17.0, 18.6)	603	31.6 (31.6, 39.8)	686	85.1 (77.6, 93.3)	221	103.4 (96.6)
Inflammation (%)	620	3	609	33	290	43	228	5
Anaemia (%)	626	7	609	32	481	8	227	19
Iron deficiency (%)	623	37	606	18	686	11	227	3

[§] Values are geometric means (confidence intervals) or otherwise means (SD); Cut-offs for inflammation: C-reactive protein >5mg/L; Iron deficiency: Serum ferritin <15µg/l (Kenyan, South African and African American cohorts), and Plasma ferritin <12µg/L (Tanzanian cohort); Anaemia: Haemoglobin <11.5 g/dL (Kenyan cohort), <11 g/dL (Tanzanian cohort), or <12 g/dL (South African and African American cohorts).




Fig 1. Minor allele frequency of common genetic variants related to iron metabolism in four African ancestry populations in comparison to the African and European population in the 1000 Genomes Project database. MAF 1000 A, Minor allele frequency for Africans-1000 Genomes Project; MAF 1000 E, Minor allele frequency for Europeans- 1000 Genomes Project.

doi:10.1371/journal.pone.0157996.g001

status was 27%, 22%, 18%, and 30% in the Kenyan, Tanzanian, South African and African American cohorts, respectively. According to the 1000 Genomes Project database, this is comparable to the 31% reported for Africans, but substantially lower than the 41% reported for Europeans. Similarly, for rs2413450 (*TMPRSS6*), MAFs were ~12% in the Kenyan, Tanzanian and South African cohorts and 24% in the African American cohort, which is comparable to the MAF for Africans (17%), but again lower than that for Europeans (42%) reported in the 1000 Genomes Project database. On the other hand, rs1421312 (*TMPRSS6*) was more common in the African cohorts (Kenya, 58%; Tanzania, 66%; South Africa, 68%; 1000 Genomes, 58%) compared to the Europeans (41%). For the same SNP (rs1421312), the African American cohort had a MAF of 43%.

In linear regression analyses with haemoglobin concentrations as the dependent outcome the minor allele (A) of rs10904850 in *CUBN* was associated with lower Hb concentrations of 0.38 g/dL ($P = 0.02$). Lower Hb concentrations were also observed for the minor allele (G) of rs4820268 in *TMPRSS6* (0.28 g/dL; $P = 0.01$) in the Kenyan children (Table 2). Conversely, the minor allele (A) of rs10421768 in *HAMP* was positively associated with increased Hb concentrations of 0.37 g/dL ($P = 0.01$) in the Kenyan population.

After meta-analyses of data from all three African cohorts, the minor allele (T) of rs3811658 in *TF* was associated with a higher Hb concentration of 0.36 g/dL ($P = 0.02$) while minor alleles of both rs2413450 (A) and rs4820268 (G) in *TMPRSS6* were associated with lower Hb

Table 2. Associations of single nucleotide polymorphism with haemoglobin in four cohorts of African ancestry with meta-analyses.

SNP	Minor Allele	Ke		Tz		SA†		Met_African		AA		Met_All	
		β (se)	P	β (se)	P	β (se)	P	β (se)	P (Het)	β (se)	P	β (se)	P
<i>TF</i>													
rs3811658	T	0.01(0.13)	0.95	1.57(0.99)	0.11	0.10(0.04)	0.49	0.36(0.04)	0.02	0.08(0.16)	0.61	0.09(0.03)	0.01
<i>CUBN</i>													
rs10904850	A	-0.38(0.16)	0.02	0.91(1.19)	0.44	0.07(0.11)	0.56	-0.11(0.10)	0.14	0.03(0.16)	0.86	-0.07(0.14)	0.63
<i>HAMP</i>													
rs10421768	A	0.37(0.19)	0.01	0.46(0.94)	0.62	-0.01(0.15)	0.93	0.19(0.18)	0.29	0.14(0.12)	0.12	0.17(0.10)	0.09
<i>TMPRSS6</i>													
rs2413450	A	-0.20(0.12)	0.10	-0.65(1.13)	0.57	-0.23(0.18)	0.20	-0.23(0.10)	0.02	-0.09(0.14)	0.55	-0.19(0.08)	0.02
rs4820268	G	-0.28(0.11)	0.01	-0.26(0.87)	0.76	-0.05(0.14)	0.74	-0.18(0.07)	0.01	-0.09(0.13)	0.48	-0.16(0.07)	0.04

Differences in haemoglobin concentrations are in g/dL. SNP = Single nucleotide polymorphism; Ke = Kenyan cohort; Tz = Tanzanian cohort; SA = South African cohort; AA = African American cohort; Met-African = meta analyses of beta and SE values from African cohorts; Met_All = meta analyses of beta and SE values from all cohorts. Results are presented as mean (SE) change per copy of the minor allele from regression analysis using an additive genetic model and with adjustment for age and sex. † Values not adjusted for gender as all participants were women. P (Het) = Cochran's Q to measure heterogeneity between cohorts with a value of 0.10 as a cut-off for significance. Values in bold are significant before FDR correction for multiple testing. Explained variance ranged from <1% to 3%, with the largest explained variance being 3% for rs10421768 in the Kenyan population.

doi:10.1371/journal.pone.0157996.t002

concentrations of -0.23 g/dL ($P = 0.02$) and -0.18 g/dL ($P = 0.01$) respectively. The associations of rs3811658, rs2413450 and rs4820268 with Hb remained significant and in the same direction after including the African American cohort in the meta-analysis. There was no heterogeneity between studies for these SNPs.

The minor allele (A) of rs1799852 in *TF* was associated with lower SF concentrations in the South African cohort (-0.62 $\mu\text{g/L}$, $P = 0.01$). On the other hand the minor allele (A) of rs1867504 also in *TF* was associated with increased serum ferritin concentrations (1.04 $\mu\text{g/L}$, $P = 0.06$), in the meta analyses of data from the three African populations.

The minor alleles (C) of rs7385804 in *TFR2* was associated with decreased serum ferritin concentrations in the Kenyan population of -0.86 $\mu\text{g/L}$ ($P = 0.02$). Lowered serum ferritin concentrations were also observed for rs1800629 (A) in *TNF* (-0.76 $\mu\text{g/L}$, $P = 0.03$, respectively) within the African American cohort (Table 3). No significant associations were observed in the meta analyses including the African American cohort though associations of the minor alleles of rs1525892 ($P = 0.05$), rs1867504 ($P = 0.06$) and rs 3811647 ($P = 0.05$) in *TF* and rs10421768 in *HAMP* ($P = 0.06$) with ferritin concentrations were nearly significant (S1 Table). Complete information on associations of all 20 SNPs with concentrations of haemoglobin and ferritin can be found in the supplementary table (S1 Table).

None of the associations for either Hb or SF as outcome measures were significant after FDR correction for multiple testing.

Discussion

We report here one of the few candidate gene association studies focussing on the genetics of iron status in various African ancestry populations. In the meta analyses involving all four cohorts, we have observed small differences in Hb concentrations between genotype groups: slightly lower Hb concentrations were associated with the minor alleles A of rs2413450 and G of rs4820268, both in the *TMPRSS6* gene and higher Hb concentrations involving the minor allele T in rs3811658 in the *TF* gene. In similar investigations involving ferritin concentrations, the minor allele A of rs1867504 in *TF* was associated with higher ferritin concentrations. Additionally, there were marginally significant associations with higher ferritin concentrations for the minor alleles of rs1525892, and rs3811647 in *TF*, rs7385804 in *TFR2* and rs10421768 in *HAMP*. We further observed that many of the candidate SNPs genotyped in our cohorts, were more common in the European ancestry cohort in the 1000 Genomes Project as compared to our African ancestry cohorts.

The most consistent association in terms of direction as well as statistical significance was observed for rs2413450 and rs4820268 in the *TMPRSS6* gene, both associated with lower Hb concentration. Similar observations have previously been made within populations of European and Asian ancestry [7, 10, 26, 27]. The exact mechanism through which *TMPRSS6* action is exerted is still under investigation. It has generally been hypothesised that *TMPRSS6* polymorphisms affect hepcidin antimicrobial peptide (HAMP) transcription, thereby influencing hepcidin concentrations in response to systemic iron concentrations [28–30]. However, two recent studies [31, 32] did not confirm an intermediate role for hepcidin in the SNP-iron status parameter associations. Currently, it is proposed that matriptase-2 could regulate hepcidin expression by cleaving HJV to decrease BMP-SMAD signalling [33].

Several *TF* variants (rs3811658, rs1867504 and rs1799852) were also significantly associated with various iron status measures. In contrast to what is known from other studies, we observed associations between rs3811658 in the *TF* gene with higher Hb concentrations. This SNP is in linkage disequilibrium with rs3811647 that has previously been negatively associated with serum iron, body iron, transferrin saturation and serum ferritin concentrations and

Table 3. Associations of single nucleotide polymorphism with ferritin in four cohorts of African ancestry with meta-analyses.

SNP	Minor Allele	Ke		Tz		SA†		Met_African		AA		Met_All		
		β (se)	P	β (se)	P	β (se)	P	β (se)	P	β (se)	P	β (se)	P	
TF														
rs1799852	A	-0.98(1.09)	0.80	1.20(1.15)	0.18	-0.62(1.2)	0.01	-0.15(0.68)	0.41	1.23(1.17)	0.34	0.21	0.18(0.60)	0.76
rs1867504	A	1.04(1.06)	0.62	1.05(1.07)	0.52	1.03(1.08)	0.69	1.04(0.06)	0.00*	1.05(1.10)	1.00	0.62	1.04(0.54)	0.06
TFR2														
rs7385804	C	0.86(1.07)	0.02	0.95(1.08)	0.48	1.15(1.07)	0.06	0.99(0.62)	0.11	1.07(1.10)	0.98	0.52	1.01(0.54)	0.06
TNF														
rs1800629	A	1.02(1.10)	0.85	1.05(1.12)	0.62	1.03(1.09)	0.74	1.03(0.64)	0.10	-0.76(1.15)	1.00	0.03	0.61(0.56)	0.27

Ferritin concentrations are in µg/L. Ferritin values are Log₁₀ back transformed data. SNP = Single nucleotide polymorphism; Ke = Kenyan cohort; Tz = Tanzanian cohort;

SA = South African cohort; AA = African American cohort; Met-African = meta analyses of beta and SE values from African cohorts; Met_All = meta analyses of beta and SE values from all cohorts. Results are presented as mean (SE) change per copy of the minor allele from regression analysis using an additive genetic model and with adjustment for age and sex.

† Values not adjusted for gender as all participants were women.

* = P < 0.0001; P (Het) = Cochran's Q to measure heterogeneity between cohorts with a value of 0.10 as a cut-off for significance. Values in bold are significant before FDR correction for multiple testing. Explained variance ranged from < 1% to 3%, with the largest explained variance being 3% for rs10421768 in the Kenyan population.

doi:10.1371/journal.pone.0157996.t003

positively associated with serum transferrin concentrations [9, 34]. Also, the SNP has been associated with increased risk of iron deficiency in a population of men and women of European descent [35]. As we would have expected to see negative associations between rs3811658 with haemoglobin concentrations, our finding requires further investigation.

The variant rs1867504 in *TF* has previously been associated with increased serum transferrin concentrations in a cohort of European ancestry [9] but not with serum ferritin as seen in our study. It is possible that this variant has pleiotropic effects on ferritin as has previously been observed for other *TF* SNPs [9].

Previously, three variants in *TF* (rs3811647, rs1799852 and rs2280673) plus the HFE C282Y mutation have been observed to explain approximately 40% of genetic variation in serum transferrin [9]. We only observed a significant association of rs1799852 with serum ferritin in the South African cohort. Furthermore, we did not have information on rs2280673 and HFE C282Y. We therefore could not replicate this previously observed association.

TF is a crucial biological carrier of iron in blood plasma [36]. Severe mutations in the transferrin gene lead to transferrinaemia [37] and have been determined to cause microcytic anaemia accompanied by hepatic accumulation of iron in humans [38] and mice [39]. Further investigations into the influence of various *TF* variants on haematological traits are therefore required.

A large portion of systemic iron deficiency undoubtedly results from non-genetic factors. Environmental factors such as diets low in bioavailable iron [40] as well as a high inflammation burden [41], are probably the foremost factors predisposing to iron deficiency. High inflammation burden causes iron to be retained in the reticuloendothelial system, resulting in iron being unavailable for erythropoiesis [41]. Diets in African populations are typically low in bioavailable iron [42]. We corrected for dietary iron intake in the African American as well as South African cohorts (not presented), but the difference between the crude and adjusted regression models was negligible. To account for other environmental and individual influences on iron status, we adjusted our regression models for major biological confounders that are known to affect Hb and ferritin concentrations including age, gender [43] and the inflammatory acute phase protein CRP [41] to increase precision of our estimates. Age and gender are known to influence iron status [44, 45]. Although we corrected the associations with serum ferritin as the outcome for CRP, we may not have captured the inflammation at later convalescence.

It is noteworthy that we did observe population specific significant associations that were not replicated in all cohorts. This was particularly observed in the Kenyan cohort that consisted of school going children (mean age 9.0 years). In this cohort we specifically highlight the association of the A allele of rs10421768 in *HAMP* with Hb and the C allele of rs7385804 in *TFR2*.

The *HAMP* gene encodes hepcidin, which is a 25-amino acid peptide produced and secreted mainly by hepatocytes and is a major regulator of systemic iron homeostasis [33]. As far as we know, no functional studies have investigated the effect of rs10421768 on dysregulation of hepcidin expression in humans. Based on our data, we speculate that this SNP down regulates *HAMP*, thereby increasing the amount of iron in circulation available for erythropoiesis. Indeed, we observed rs10421768 in *HAMP* to be consistently associated with higher serum ferritin concentrations across all study populations (S1 Table).

The SNP rs7385804 has previously been associated with reduced serum iron, serum transferrin and transferrin saturation concentrations in a cohort of elderly Chinese women [26]. *TFR2* polymorphisms have also been reported to be associated with red blood cell count and mean cell volume [46–48]. In terms of function, *TFR2* is believed to be involved in the hepcidin regulation pathway [49]. The fact that SNPs directly linked to hepcidin were observed to be associated with various iron status measures underscores the key role that hepcidin could play in determining iron status in the Kenyan cohort. The children in this cohort did not reside in a

malaria endemic area and based on CRP concentrations did not have a high prevalence of inflammation, both of which are likely to impact significantly on hepcidin concentrations. Hepcidin however responds to signals from diverse physiological processes: it can be induced by infection, inflammation and iron loading and suppressed by iron deficiency, hypoxia, and expanded erythropoiesis [50]. We hypothesise this observation is because the typical diet of this cohort of children is low in iron which would impact on iron loading thereby having an impact on hepcidin response.

Within the African American population, rs1800629 in *TNF* was significantly associated with lower plasma ferritin concentrations. This SNP has previously been associated with iron deficiency as part of a haplotype together with other *TNF* SNPs [51], as well as with an increased risk of severe anemia in low-birth-weight children in Kenya [52] and increased risk of severe malarial anemia in Gambian children [53]. It has been suggested previously that this haplotype would only influence iron status due to pressure of infectious diseases [51]. However, our African American cohort consisted of apparently healthy elderly subjects who are not at risk of developing malaria. Nevertheless, there may still have been undetected low grade inflammation from undiagnosed infections or diseases in this elderly population. Disease induced inflammatory processes may therefore underlie this observation.

A limitation inherent to our study is that age and sex distributions differed radically between some of the cohorts analyzed and therefore simple adjustments may not have eliminated all confounding by these variables. In future studies, including cohorts that are more similar in aspects such as age and sex would be desirable. Additionally, our cohorts were smaller than those used in GWA studies and therefore may have lacked power to confirm previously observed associations. Indeed none of the associations with either Hb or SF as outcome measures were significant after FDR correction for multiple testing. The sample size for detecting associations between disease and SNP markers is affected by prevalence of the condition under study, allele frequency, linkage disequilibrium (LD), inheritance models (e.g., additive, dominant, and multiplicative models), and effect size of the genetic variants (e.g., odds ratio, relative risk, etc.) [54–56]. We conducted a retrospective sample size calculation in which we assumed an additive model. From the calculation, 780 individuals are required to conduct association analyses for a SNP explaining 1% of the variation in the outcome measure, with $\alpha = 0.05$ and power of 80%. If the genetic variant is one of 20 independent genetic variants tested, adjustment for multiple testing using a Bonferroni correction would require a significance level of $0.05/20$ or 0.0025 as the threshold. To detect this same association at this adjusted significance level, with 80% power would require a large increase in sample size.

We observed large differences in MAF of investigated SNPs between our study cohorts and those of the Europeans included in the 1000 Genomes project. This may be due to selective pressure caused by environmental conditions thereby altering the frequency of genetic variants between different populations [57]. Theoretically, these differences in MAF between populations could result in dissimilarities in the prevalence of a disease outcome, for example iron deficiency, and in the case of strong gene-disease associations call for population specific strategies to address this.

All of our cohorts were not specifically designed to study the genetics of iron metabolism and therefore not all of the factors taken into account in sample size calculation were considered. However our study does offer an initial indication of what genetic loci would be important for haematological traits in African ancestry cohorts. Another important consideration not taken into account in our analyses is the linkage disequilibrium structure in African ancestry populations, which is different from that of Europeans resulting from the greater variation within the genome of African ancestry populations [58]. It is thus possible that actual disease markers tagged by the investigated SNPs in European populations are not in LD in our African

cohorts, and could explain why only a few significant associations were observed. A recent genome wide admixture and association study indeed discovered novel loci to be associated with iron status in an African-American population [59].

In conclusion, we have attempted to replicate reported associations between genetic loci and iron status indicators from European ancestry populations in various African ancestry populations with limited success. This may partly be due to the greater genetic variation within the African ancestry populations [60] which limits the transferability of findings from European ancestry populations. Additionally, populations vary by ethnicity in terms of allele frequencies and biological adaptations due to various environmental factors. For these reasons, population specific genomic studies are warranted. GWA studies in African ancestry populations can potentially reveal novel genetic variants that are specific to these populations. In addition, functional studies are required to further explore the mechanisms involved in the regulation of iron metabolism.

Supporting Information

S1 Table. Associations of single nucleotide polymorphism with iron status parameters.

SNP = Single nucleotide polymorphism; Ke = Kenyan cohort; Tz = Tanzanian cohort; SA = South African cohort; AA = African American cohort; P(Het) = P value of heterogeneity testing; Met- = meta analyses of beta and SE values from African cohorts; Met_All = meta analyses of beta and SE values from all cohorts (African+American); Direction = direction of association based on beta value Hb = Haemoglobin, Values in bold are significant. Results are presented as mean (SE) change per copy of the minor allele from regression analysis using additive genetic model and with adjustment for age and sex for associations involving Hb; and for age, sex and CRP concentrations for associations involving ferritin. † Values not adjusted for gender as all participants were women. Values in bold are significant before FDR correction for multiple testing. Explained variance ranged from < 1% to 3%, with the largest explained variety being 3% for rs10421768 in the Kenyan population.

(DOC)

Acknowledgments

Kenyan cohort (Cassavita study) acknowledgements

We thank the participating children, caretakers, teachers, and the KEMRI/CPHR team without whom the execution of the study would not have been possible. Dr. Alice Mwangi, Gloria Mbera, Christopher Masyuki, Julius Musyoki and Jasper van der Hout are duly acknowledged for their roles in study execution as well as participant data organization.

Tanzanian cohort (MACH study) acknowledgements

We thank the MACH study participants, district medical officers of Handeni, field staff, village elders, community volunteers, other local collaborators and students for magnificent contributions. QuikRead tests and trimagnesium dicitrate were donated by Orion Diagnostica, Espoo, Finland and Jungbunzlauer, Ladenburg, Germany, respectively; supplements were manufactured and donated by Medisan, Heerenveen, The Netherlands.

South African cohort (PURE study) acknowledgements

We thank the PURE study participants, local collaborators of the PURE study North-West Department of Health, tribal chiefs, community leaders, employers, and mayors. In addition Tinashe Chikowore and Herman Myburgh, Centre of Excellence for Nutrition, North West University, Potchefstroom Campus, South Africa, are duly acknowledged for assistance in sample preparation.

African American cohort (BLSA study) acknowledgements

We thank the BLSA study participants without whom this work would not have been possible.

We thank Richard Crooijmans and the Animal Breeding Group of Wageningen University, The Netherlands, for conducting the genotyping work for the Kenya, Tanzania and South African cohorts.

Author Contributions

Conceived and designed the experiments: WNGW TT GWT MB EJF AMB. Performed the experiments: WNGW MB JH. Analyzed the data: WNGW. Wrote the paper: WNGW TT GWT MB EJF HV JV ET AMB JH.

References

1. Andrews NC. Iron homeostasis: insights from genetics and animal models. *Nature reviews Genetics*. 2000; 1(3):208–17. Epub 2001/03/17. doi: [10.1038/35042073](https://doi.org/10.1038/35042073) PMID: [11252750](https://pubmed.ncbi.nlm.nih.gov/11252750/).
2. Whitfield JB, Zhu G, Heath AC, Powell LW, Martin NG. Effects of alcohol consumption on indices of iron stores and of iron stores on alcohol intake markers. *Alcoholism, clinical and experimental research*. 2001; 25(7):1037–45. Epub 2001/08/16. PMID: [11505030](https://pubmed.ncbi.nlm.nih.gov/11505030/).
3. Beutler E, West C. Hematologic differences between African-Americans and whites: the roles of iron deficiency and alpha-thalassemia on hemoglobin levels and mean corpuscular volume. *Blood*. 2005; 106(2):740–5. Epub 2005/03/26. doi: [10.1182/blood-2005-02-0713](https://doi.org/10.1182/blood-2005-02-0713) PMID: [15790781](https://pubmed.ncbi.nlm.nih.gov/15790781/); PubMed Central PMCID: [PMC1895180](https://pubmed.ncbi.nlm.nih.gov/PMC/PMC1895180/).
4. World Health Organization. *Worldwide prevalence of anaemia 1993–2005*. 2008.
5. Njajou OT, Alizadeh BZ, Aulchenko Y, Zillikens MC, Pols HA, Oostra BA, et al. Heritability of serum iron, ferritin and transferrin saturation in a genetically isolated population, the Erasmus Rucphen Family (ERF) Study. *Human heredity*. 2006; 61(4):222–8. Epub 2006/08/01. doi: [10.1159/000094777](https://doi.org/10.1159/000094777) PMID: [16877869](https://pubmed.ncbi.nlm.nih.gov/16877869/).
6. Whitfield JB, Cullen LM, Jazwinska EC, Powell LW, Heath AC, Zhu G, et al. Effects of HFE C282Y and H63D polymorphisms and polygenic background on iron stores in a large community sample of twins. *American journal of human genetics*. 2000; 66(4):1246–58. Epub 2000/03/31. doi: [10.1086/302862](https://doi.org/10.1086/302862) PMID: [10739755](https://pubmed.ncbi.nlm.nih.gov/10739755/); PubMed Central PMCID: [PMC1288192](https://pubmed.ncbi.nlm.nih.gov/PMC/PMC1288192/).
7. Tanaka T, Roy CN, Yao W, Matteini A, Semba RD, Arking D, et al. A genome-wide association analysis of serum iron concentrations. *Blood*. 2010; 115(1):94–6. Epub 2009/11/03. doi: [10.1182/blood-2009-07-232496](https://doi.org/10.1182/blood-2009-07-232496) PMID: [19880490](https://pubmed.ncbi.nlm.nih.gov/19880490/); PubMed Central PMCID: [PMC2803694](https://pubmed.ncbi.nlm.nih.gov/PMC/PMC2803694/).
8. Benyamin B, Ferreira MA, Willemsen G, Gordon S, Middelberg RP, McEvoy BP, et al. Common variants in TMPRSS6 are associated with iron status and erythrocyte volume. *Nature genetics*. 2009; 41(11):1173–5. Epub 2009/10/13. doi: [10.1038/ng.456](https://doi.org/10.1038/ng.456) PMID: [19820699](https://pubmed.ncbi.nlm.nih.gov/19820699/); PubMed Central PMCID: [PMC3135421](https://pubmed.ncbi.nlm.nih.gov/PMC/PMC3135421/).
9. Benyamin B, McRae AF, Zhu G, Gordon S, Henders AK, Palotie A, et al. Variants in TF and HFE explain approximately 40% of genetic variation in serum-transferrin levels. *American journal of human genetics*. 2009; 84(1):60–5. Epub 2008/12/17. doi: [10.1016/j.ajhg.2008.11.011](https://doi.org/10.1016/j.ajhg.2008.11.011) PMID: [19084217](https://pubmed.ncbi.nlm.nih.gov/19084217/); PubMed Central PMCID: [PMC2668053](https://pubmed.ncbi.nlm.nih.gov/PMC/PMC2668053/).
10. Chambers JC, Zhang W, Li Y, Sehmi J, Wass MN, Zabaneh D, et al. Genome-wide association study identifies variants in TMPRSS6 associated with hemoglobin levels. *Nature genetics*. 2009; 41(11):1170–2. Epub 2009/10/13. doi: [10.1038/ng.462](https://doi.org/10.1038/ng.462) PMID: [19820698](https://pubmed.ncbi.nlm.nih.gov/19820698/); PubMed Central PMCID: [PMC3178047](https://pubmed.ncbi.nlm.nih.gov/PMC/PMC3178047/).
11. Need AC, Goldstein DB. Next generation disparities in human genomics: concerns and remedies. *Trends in genetics: TIG*. 2009; 25(11):489–94. Epub 2009/10/20. doi: [10.1016/j.tig.2009.09.012](https://doi.org/10.1016/j.tig.2009.09.012) PMID: [19836853](https://pubmed.ncbi.nlm.nih.gov/19836853/).
12. McLaren CE, McLachlan S, Garner CP, Vulpe CD, Gordeuk VR, Eckfeldt JH, et al. Associations between single nucleotide polymorphisms in iron-related genes and iron status in multiethnic populations. *PloS one*. 2012; 7(6):e38339. Epub 2012/07/05. doi: [10.1371/journal.pone.0038339](https://doi.org/10.1371/journal.pone.0038339) PMID: [22761678](https://pubmed.ncbi.nlm.nih.gov/22761678/); PubMed Central PMCID: [PMC3382217](https://pubmed.ncbi.nlm.nih.gov/PMC/PMC3382217/).
13. Lo KS, Wilson JG, Lange LA, Folsom AR, Galarnau G, Ganesh SK, et al. Genetic association analysis highlights new loci that modulate hematological trait variation in Caucasians and African Americans. *Human genetics*. 2011; 129(3):307–17. Epub 2010/12/15. doi: [10.1007/s00439-010-0925-1](https://doi.org/10.1007/s00439-010-0925-1) PMID: [21153663](https://pubmed.ncbi.nlm.nih.gov/21153663/); PubMed Central PMCID: [PMC3442357](https://pubmed.ncbi.nlm.nih.gov/PMC/PMC3442357/).

14. Chen Z, Tang H, Qayyum R, Schick UM, Nalls MA, Handsaker R, et al. Genome-wide association analysis of red blood cell traits in African Americans: the COGENT Network. *Human molecular genetics*. 2013; 22(12):2529–38. Epub 2013/03/01. doi: [10.1093/hmg/ddt087](https://doi.org/10.1093/hmg/ddt087) PMID: [23446634](https://pubmed.ncbi.nlm.nih.gov/23446634/); PubMed Central PMCID: PMCPmc3658166.
15. Ding K, de Andrade M, Manolio TA, Crawford DC, Rasmussen-Torvik LJ, Ritchie MD, et al. Genetic variants that confer resistance to malaria are associated with red blood cell traits in African-Americans: an electronic medical record-based genome-wide association study. *G3 (Bethesda, Md)*. 2013; 3(7):1061–8. Epub 2013/05/23. doi: [10.1534/g3.113.006452](https://doi.org/10.1534/g3.113.006452) PMID: [23696099](https://pubmed.ncbi.nlm.nih.gov/23696099/); PubMed Central PMCID: PMCPmc3704235.
16. Li J, Glessner JT, Zhang H, Hou C, Wei Z, Bradfield JP, et al. GWAS of blood cell traits identifies novel associated loci and epistatic interactions in Caucasian and African-American children. *Human molecular genetics*. 2013; 22(7):1457–64. Epub 2012/12/25. doi: [10.1093/hmg/dds534](https://doi.org/10.1093/hmg/dds534) PMID: [23263863](https://pubmed.ncbi.nlm.nih.gov/23263863/); PubMed Central PMCID: PMCPmc3657475.
17. Parra EJ, Marcini A, Akey J, Martinson J, Batzer MA, Cooper R, et al. Estimating African American admixture proportions by use of population-specific alleles. *American journal of human genetics*. 1998; 63(6):1839–51. Epub 1998/12/05. doi: [10.1086/302148](https://doi.org/10.1086/302148) PMID: [9837836](https://pubmed.ncbi.nlm.nih.gov/9837836/); PubMed Central PMCID: PMC1377655.
18. Kullo IJ, Ding K, Jouni H, Smith CY, Chute CG. A genome-wide association study of red blood cell traits using the electronic medical record. *PLoS one*. 2010; 5(9). Epub 2010/10/12. doi: [10.1371/journal.pone.0013011](https://doi.org/10.1371/journal.pone.0013011) PMID: [20927387](https://pubmed.ncbi.nlm.nih.gov/20927387/); PubMed Central PMCID: PMC2946914.
19. Talsma EF V H, Brouwer ID, Mburu-de Wagt AS, Hulshof PJM, Melse-Boonstra A. Proxy markers of serum retinol concentration, used alone and in combination, to assess population vitamin A status in Kenyan children: a cross-sectional study. *BMC Med* 2015.
20. Patel A, Pusdekar Y, Badhoniya N, Borkar J, Agho KE, Dibley MJ. Determinants of inappropriate complementary feeding practices in young children in India: secondary analysis of National Family Health Survey 2005–2006. *Maternal & child nutrition*. 2012; 8 Suppl 1:28–44. Epub 2011/12/30. doi: [10.1111/j.1740-8709.2011.00385.x](https://doi.org/10.1111/j.1740-8709.2011.00385.x) PMID: [22168517](https://pubmed.ncbi.nlm.nih.gov/22168517/).
21. WHO. [cited 2015 4-2-2015]. Available from: <http://www.who.int/malaria/en/>.
22. Veenemans J, Milligan P, Prentice AM, Schouten LR, Inja N, van der Heijden AC, et al. Effect of supplementation with zinc and other micronutrients on malaria in Tanzanian children: a randomised trial. *PLoS medicine*. 2011; 8(11):e1001125. Epub 2011/12/02. doi: [10.1371/journal.pmed.1001125](https://doi.org/10.1371/journal.pmed.1001125) PMID: [22131908](https://pubmed.ncbi.nlm.nih.gov/22131908/); PubMed Central PMCID: PMC3222646.
23. Shock NW, Greulich RC, Andres R, Arenberg D, Paul T, Costa J, Lakata EG, et al. *Normal Human Aging: The Baltimore Longitudinal Study of Aging*. 1984.
24. Vitamin and Mineral Nutrition Information System. Geneva WHO, 2011 (WHO/NMH/NHD/MNM/11.1). Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity
25. Team RC. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. 2013.
26. An P, Wu Q, Wang H, Guan Y, Mu M, Liao Y, et al. TMPRSS6, but not TF, TFR2 or BMP2 variants are associated with increased risk of iron-deficiency anemia. *Human molecular genetics*. 2012; 21(9):2124–31. Epub 2012/02/11. doi: [10.1093/hmg/dds028](https://doi.org/10.1093/hmg/dds028) PMID: [22323359](https://pubmed.ncbi.nlm.nih.gov/22323359/).
27. Gan W, Guan Y, Wu Q, An P, Zhu J, Lu L, et al. Association of TMPRSS6 polymorphisms with ferritin, hemoglobin, and type 2 diabetes risk in a Chinese Han population. *The American journal of clinical nutrition*. 2012; 95(3):626–32. Epub 2012/02/04. doi: [10.3945/ajcn.111.025684](https://doi.org/10.3945/ajcn.111.025684) PMID: [22301935](https://pubmed.ncbi.nlm.nih.gov/22301935/).
28. Finberg KE, Heeney MM, Campagna DR, Aydinok Y, Pearson HA, Hartman KR, et al. Mutations in TMPRSS6 cause iron-refractory iron deficiency anemia (IRIDA). *Nature genetics*. 2008; 40(5):569–71. Epub 2008/04/15. doi: [10.1038/ng.130](https://doi.org/10.1038/ng.130) PMID: [18408718](https://pubmed.ncbi.nlm.nih.gov/18408718/); PubMed Central PMCID: PMCPmc3104019.
29. Du X, She E, Gelbart T, Truksa J, Lee P, Xia Y, et al. The serine protease TMPRSS6 is required to sense iron deficiency. *Science (New York, NY)*. 2008; 320(5879):1088–92. Epub 2008/05/03. doi: [10.1126/science.1157121](https://doi.org/10.1126/science.1157121) PMID: [18451267](https://pubmed.ncbi.nlm.nih.gov/18451267/); PubMed Central PMCID: PMCPmc2430097.
30. De Falco L, Sanchez M, Silvestri L, Kannengiesser C, Muckenthaler MU, Iolascon A, et al. Iron refractory iron deficiency anemia. *Haematologica*. 2013; 98(6):845–53. Epub 2013/06/05. doi: [10.3324/haematol.2012.075515](https://doi.org/10.3324/haematol.2012.075515) PMID: [23729726](https://pubmed.ncbi.nlm.nih.gov/23729726/); PubMed Central PMCID: PMCPmc3669438.
31. Galesloot TE, Geurts-Moespot AJ, den Heijer M, Sweep FC, Fleming RE, Kiemeny LA, et al. Associations of common variants in HFE and TMPRSS6 with iron parameters are independent of serum hepcidin in a general population: a replication study. *Journal of medical genetics*. 2013. Epub 2013/06/25. doi: [10.1136/jmedgenet-2013-101673](https://doi.org/10.1136/jmedgenet-2013-101673) PMID: [23794717](https://pubmed.ncbi.nlm.nih.gov/23794717/).
32. Traglia M, Girelli D, Biino G, Campostrini N, Corbella M, Sala C, et al. Association of HFE and TMPRSS6 genetic variants with iron and erythrocyte parameters is only in part dependent on serum

- hepcidin concentrations. *Journal of medical genetics*. 2011; 48(9):629–34. Epub 2011/07/26. doi: [10.1136/jmedgenet-2011-100061](https://doi.org/10.1136/jmedgenet-2011-100061) PMID: [21785125](https://pubmed.ncbi.nlm.nih.gov/21785125/).
33. Meynard D, Babitt JL, Lin HY. The liver: conductor of systemic iron balance. *Blood*. 2014; 123(2):168–76. Epub 2013/11/10. doi: [10.1182/blood-2013-06-427757](https://doi.org/10.1182/blood-2013-06-427757) PMID: [24200681](https://pubmed.ncbi.nlm.nih.gov/24200681/); PubMed Central PMCID: PMCPmc3888285.
 34. McLaren CE, Garner CP, Constantine CC, McLachlan S, Vulpe CD, Snively BM, et al. Genome-wide association study identifies genetic loci associated with iron deficiency. *PLoS one*. 2011; 6(3):e17390. Epub 2011/04/13. doi: [10.1371/journal.pone.0017390](https://doi.org/10.1371/journal.pone.0017390) PMID: [21483845](https://pubmed.ncbi.nlm.nih.gov/21483845/); PubMed Central PMCID: PMCPmc3069025.
 35. Blanco-Rojo R, Toxqui L, Lopez-Parra AM, Baeza-Richer C, Perez-Granados AM, Arroyo-Pardo E, et al. Influence of diet, menstruation and genetic factors on iron status: a cross-sectional study in Spanish women of childbearing age. *International journal of molecular sciences*. 2014; 15(3):4077–87. Epub 2014/03/26. doi: [10.3390/ijms15034077](https://doi.org/10.3390/ijms15034077) PMID: [24663082](https://pubmed.ncbi.nlm.nih.gov/24663082/); PubMed Central PMCID: PMCPmc3975385.
 36. Lee PL, Halloran C, Trevino R, Felitti V, Beutler E. Human transferrin G277S mutation: a risk factor for iron deficiency anaemia. *British journal of haematology*. 2001; 115(2):329–33. Epub 2001/11/13. PMID: [11703331](https://pubmed.ncbi.nlm.nih.gov/11703331/).
 37. Aslan D, Crain K, Beutler E. A new case of human atransferrinemia with a previously undescribed mutation in the transferrin gene. *Acta haematologica*. 2007; 118(4):244–7. Epub 2007/12/22. doi: [10.1159/000112726](https://doi.org/10.1159/000112726) PMID: [18097132](https://pubmed.ncbi.nlm.nih.gov/18097132/).
 38. Beutler E, Gelbart T, Lee P, Trevino R, Fernandez MA, Fairbanks VF. Molecular characterization of a case of atransferrinemia. *Blood*. 2000; 96(13):4071–4. Epub 2000/12/09. PMID: [11110675](https://pubmed.ncbi.nlm.nih.gov/11110675/).
 39. Trenor CC 3rd, Campagna DR, Sellers VM, Andrews NC, Fleming MD. The molecular defect in hypotransferrinemic mice. *Blood*. 2000; 96(3):1113–8. Epub 2000/07/27. PMID: [10910930](https://pubmed.ncbi.nlm.nih.gov/10910930/).
 40. Jonker FA, Boele van Hensbroek M. Anaemia, iron deficiency and susceptibility to infections. *The Journal of infection*. 2014. Epub 2014/09/30. doi: [10.1016/j.jinf.2014.08.007](https://doi.org/10.1016/j.jinf.2014.08.007) PMID: [25264159](https://pubmed.ncbi.nlm.nih.gov/25264159/).
 41. Trey JE, Kushner I. The acute phase response and the hematopoietic system: the role of cytokines. *Critical reviews in oncology/hematology*. 1995; 21(1–3):1–18. Epub 1995/11/01. PMID: [8822494](https://pubmed.ncbi.nlm.nih.gov/8822494/).
 42. Zimmermann MB, Chaouki N, Hurrell RF. Iron deficiency due to consumption of a habitual diet low in bioavailable iron: a longitudinal cohort study in Moroccan children. *The American journal of clinical nutrition*. 2005; 81(1):115–21. Epub 2005/01/11. PMID: [15640469](https://pubmed.ncbi.nlm.nih.gov/15640469/).
 43. Coad J, Conlon C. Iron deficiency in women: assessment, causes and consequences. *Current opinion in clinical nutrition and metabolic care*. 2011; 14(6):625–34. Epub 2011/09/22. doi: [10.1097/MCO.0b013e32834be6fd](https://doi.org/10.1097/MCO.0b013e32834be6fd) PMID: [21934611](https://pubmed.ncbi.nlm.nih.gov/21934611/).
 44. Garn SM, Ryan AS, Owen GM, Abraham S. Income matched black-white hemoglobin differences after correction for low transferrin saturations. *The American journal of clinical nutrition*. 1981; 34(9):1645–7. Epub 1981/09/01. PMID: [7282587](https://pubmed.ncbi.nlm.nih.gov/7282587/).
 45. Yip R, Johnson C, Dallman PR. Age-related changes in laboratory values used in the diagnosis of anemia and iron deficiency. *The American journal of clinical nutrition*. 1984; 39(3):427–36. Epub 1984/03/01. PMID: [6695842](https://pubmed.ncbi.nlm.nih.gov/6695842/).
 46. Ganesh SK, Zakai NA, van Rooij FJ, Soranzo N, Smith AV, Nalls MA, et al. Multiple loci influence erythrocyte phenotypes in the CHARGE Consortium. *Nature genetics*. 2009; 41(11):1191–8. Epub 2009/10/29. doi: [10.1038/ng.466](https://doi.org/10.1038/ng.466) PMID: [19862010](https://pubmed.ncbi.nlm.nih.gov/19862010/); PubMed Central PMCID: PMCPmc2778265.
 47. Pichler I, Minelli C, Sanna S, Tanaka T, Schwienbacher C, Naitza S, et al. Identification of a common variant in the TFR2 gene implicated in the physiological regulation of serum iron levels. *Human molecular genetics*. 2011; 20(6):1232–40. Epub 2011/01/07. doi: [10.1093/hmg/ddq552](https://doi.org/10.1093/hmg/ddq552) PMID: [21208937](https://pubmed.ncbi.nlm.nih.gov/21208937/); PubMed Central PMCID: PMCPmc3043660.
 48. Soranzo N, Spector TD, Mangino M, Kuhnel B, Rendon A, Teumer A, et al. A genome-wide meta-analysis identifies 22 loci associated with eight hematological parameters in the HaemGen consortium. *Nature genetics*. 2009; 41(11):1182–90. Epub 2009/10/13. doi: [10.1038/ng.467](https://doi.org/10.1038/ng.467) PMID: [19820697](https://pubmed.ncbi.nlm.nih.gov/19820697/); PubMed Central PMCID: PMCPmc3108459.
 49. Weiss G. Genetic mechanisms and modifying factors in hereditary hemochromatosis. *Nature reviews Gastroenterology & hepatology*. 2010; 7(1):50–8. Epub 2009/11/18. doi: [10.1038/nrgastro.2009.201](https://doi.org/10.1038/nrgastro.2009.201) PMID: [19918260](https://pubmed.ncbi.nlm.nih.gov/19918260/).
 50. Ganz T. Heparin and iron regulation, 10 years later. *Blood*. 2011; 117(17):4425–33. Epub 2011/02/25. doi: [10.1182/blood-2011-01-258467](https://doi.org/10.1182/blood-2011-01-258467) PMID: [21346250](https://pubmed.ncbi.nlm.nih.gov/21346250/); PubMed Central PMCID: PMCPmc3099567.
 51. Atkinson SH, Rockett KA, Morgan G, Bejon PA, Sirugo G, O'Connell MA, et al. Tumor necrosis factor SNP haplotypes are associated with iron deficiency anemia in West African children. *Blood*. 2008; 112

- (10):4276–83. Epub 2008/08/22. doi: [10.1182/blood-2008-06-162008](https://doi.org/10.1182/blood-2008-06-162008) PMID: [18716131](https://pubmed.ncbi.nlm.nih.gov/18716131/); PubMed Central PMCID: [PMCPmc2581977](https://pubmed.ncbi.nlm.nih.gov/PMCPmc2581977/).
52. Aidoo M, McElroy PD, Kolczak MS, Terlouw DJ, ter Kuile FO, Nahlen B, et al. Tumor necrosis factor-alpha promoter variant 2 (TNF2) is associated with pre-term delivery, infant mortality, and malaria morbidity in western Kenya: Asembo Bay Cohort Project IX. *Genetic epidemiology*. 2001; 21(3):201–11. Epub 2001/10/23. doi: [10.1002/gepi.1029](https://doi.org/10.1002/gepi.1029) PMID: [11668577](https://pubmed.ncbi.nlm.nih.gov/11668577/).
 53. McGuire W, Hill AV, Allsopp CE, Greenwood BM, Kwiatkowski D. Variation in the TNF-alpha promoter region associated with susceptibility to cerebral malaria. *Nature*. 1994; 371(6497):508–10. Epub 1994/10/06. doi: [10.1038/371508a0](https://doi.org/10.1038/371508a0) PMID: [7935762](https://pubmed.ncbi.nlm.nih.gov/7935762/).
 54. Scherag A, Muller HH, Dempfle A, Hebebrand J, Schafer H. Data adaptive interim modification of sample sizes for candidate-gene association studies. *Human heredity*. 2003; 56(1–3):56–62. Epub 2003/11/14. PMID: [14614239](https://pubmed.ncbi.nlm.nih.gov/14614239/).
 55. Gordon D, Levenstien MA, Finch SJ, Ott J. Errors and linkage disequilibrium interact multiplicatively when computing sample sizes for genetic case-control association studies. *Pacific Symposium on Biocomputing Pacific Symposium on Biocomputing*. 2003:490–501. Epub 2003/02/27. PMID: [12603052](https://pubmed.ncbi.nlm.nih.gov/12603052/).
 56. Pfeiffer RM, Gail MH. Sample size calculations for population- and family-based case-control association studies on marker genotypes. *Genetic epidemiology*. 2003; 25(2):136–48. Epub 2003/08/14. doi: [10.1002/gepi.10245](https://doi.org/10.1002/gepi.10245) PMID: [12916022](https://pubmed.ncbi.nlm.nih.gov/12916022/).
 57. Fumagalli M, Sironi M, Pozzoli U, Ferrer-Admetlla A, Pattini L, Nielsen R. Signatures of environmental genetic adaptation pinpoint pathogens as the main selective pressure through human evolution. *PLoS genetics*. 2011; 7(11):e1002355. Epub 2011/11/11. doi: [10.1371/journal.pgen.1002355](https://doi.org/10.1371/journal.pgen.1002355) PMID: [22072984](https://pubmed.ncbi.nlm.nih.gov/22072984/); PubMed Central PMCID: [PMCPmc3207877](https://pubmed.ncbi.nlm.nih.gov/PMCPmc3207877/).
 58. Tishkoff SA, Kidd KK. Implications of biogeography of human populations for 'race' and medicine. *Nature genetics*. 2004; 36(11 Suppl):S21–7. Epub 2004/10/28. doi: [10.1038/ng1438](https://doi.org/10.1038/ng1438) PMID: [15507999](https://pubmed.ncbi.nlm.nih.gov/15507999/).
 59. Li J, Lange LA, Duan Q, Lu Y, Singleton AB, Zonderman AB, et al. Genome-wide admixture and association study of serum iron, ferritin, transferrin saturation and total iron binding capacity in African Americans. *Human molecular genetics*. 2015; 24(2):572–81. Epub 2014/09/17. doi: [10.1093/hmg/ddu454](https://doi.org/10.1093/hmg/ddu454) PMID: [25224454](https://pubmed.ncbi.nlm.nih.gov/25224454/); PubMed Central PMCID: [PMCPmc4334839](https://pubmed.ncbi.nlm.nih.gov/PMCPmc4334839/).
 60. Gomez F, Hirbo J, Tishkoff SA. Genetic variation and adaptation in Africa: implications for human evolution and disease. *Cold Spring Harbor perspectives in biology*. 2014; 6(7):a008524. Epub 2014/07/06. doi: [10.1101/cshperspect.a008524](https://doi.org/10.1101/cshperspect.a008524) PMID: [24984772](https://pubmed.ncbi.nlm.nih.gov/24984772/).