

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Cauchi, D; Glonti, K; Petticrew, M; Knai, C (2016) Environmental components of childhood obesity prevention interventions: an overview of systematic reviews. *Obesity reviews*, 17 (11). pp. 1116-1130. ISSN 1467-7881 DOI: <https://doi.org/10.1111/obr.12441>

Downloaded from: <http://researchonline.lshtm.ac.uk/2664376/>

DOI: [10.1111/obr.12441](https://doi.org/10.1111/obr.12441)

Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

Environmental components of childhood obesity prevention interventions: an overview of systematic reviews

Daniel Cauchi¹, Ketevan Glonti¹, Mark Petticrew², Cécile Knai¹

¹ Department of Health Services Research and Policy, Faculty of Public Health and Policy, London School of Hygiene and Tropical Medicine, , 15-17 Tavistock Place, WC1H 9SH, London, United Kingdom

² Department of Social and Environmental Health Research, Faculty of Public Health and Policy, London School of Hygiene and Tropical Medicine, , 15-17 Tavistock Place, WC1H 9SH, London, United Kingdom

Keywords: Overview; environmental interventions; childhood obesity; obesity prevention

Running title: Environmental interventions for childhood obesity

Acknowledgements: This study was partly supported by a grant from the Malta Government Scholarship Scheme. The funding agreement ensured the authors' independence in designing the study, interpreting the data, writing, and publishing the report.

Corresponding author: Daniel Cauchi

Email: Dcauchi@gmail.com; Daniel.cauchi@lshtm.ac.uk

Postal address: LG13, LSHTM

15-17 Tavistock Place

London WC1H 9SH

United Kingdom

Conflicts of interest: Partial financial support for this study was provided to DC by a PhD scholarship grant from the Malta Government Scholarship Scheme. The funding agreement ensured the authors' independence in designing, writing, and publishing this study. KG, CK and MP have no conflict of interest to declare.

Abbreviations: % BF, percentage body fat; AMSTAR, Assessment of Multiple Systematic Reviews; BMI, body mass index; EPPI-Centre, Evidence for Policy and Practice Information and Co-ordinating Centre; ES, Effect Sizes; HSSS, High in Fat, Sugar and Salt, OR, odds ratio; MVPA, Moderate-Vigorous Physical Activity; PA, Physical Activity; PE, Physical Education; SE, standard error; SFT, skin fold thickness; SR, Systematic Review; SSB, Sugar Sweetened Beverage; WC, waist circumference

Abstract

Childhood obesity has a complex multi-factorial aetiology grounded in environmental and individual level factors that affect behaviour and outcomes. An ecological, systems-based approach to addressing childhood obesity is increasingly being advocated. The primary aim of this review is to summarise the evidence reported in systematic reviews on the effectiveness of population-level childhood obesity prevention interventions which have an environmental component. We conducted a systematic review of reviews published since 1995, employing a standardized search strategy in nine databases. Inclusion criteria required that reviews be systematic and evaluated at least one population-level, environmental intervention in any setting aimed at preventing or reducing obesity in children (5-18 years). Sixty-three reviews were included, ten of which were of high quality. Results show modest impact of a broad range of environmental strategies on anthropometric outcomes. Systematic reviews vary in methodological quality, and not all relevant primary studies may be included in each review. To ensure relevance of our findings to practice, we also report on relevant underlying primary studies, providing policy-relevant recommendations based on the evidence reviewed. Greater standardization of review methods and reporting structures will benefit policymakers and public health professionals seeking informed decision-making.

Introduction

Childhood obesity is a global public health challenge due to concerns about increasing prevalence^{1,2}, the likelihood of obesity tracking into adolescence and adulthood³⁻⁵, and its association with a range of adverse health outcomes^{6,7}. Increasingly, an ecological, systems-based approach to addressing

obesity that acknowledges its complex multi-factorial aetiology and recognizes the policy, environmental, and individual level factors that influence behaviour and outcomes is being advocated^{8,9}. This entails recognition of a broad range of physical, socio-cultural, economic and political dimensions within which individuals are embedded, as well as attributes and behaviours of individuals themselves¹⁰. Education-based interventions to address childhood obesity have had little success^{11,12} as changing human behaviour within an ‘obesogenic’ environment that does not support healthy choices is difficult to achieve and sustain¹³. Multi-component interventions where several environmental aspects are addressed simultaneously may lead to more sustainable results^{8,14}. Childhood and adolescence are particularly vulnerable life stages requiring protective social and public health policies¹⁵. This is an additional challenge for decision-makers often seeking easy solutions for rapid implementation during their relatively short term in office, and represents an all the more pressing rationale for providing them with up-to-date evidence on the range and effectiveness of environmental, population-level interventions to prevent or help reverse childhood overweight and obesity. We focus on synthesising systematic reviews as they can be useful decision-making tools that objectively summarize large amounts of information in a format that is of relevance to health practitioners, policy makers and researchers in order to guide health policy, clinical guidelines and research efforts^{16,17}. The applicability of review findings to practice has also been called into question, with reviews often being criticized for their narrow scope and for lacking context-specific details that are essential for knowledge-translation into policies^{18,19}. To ensure relevance of our findings to practice, we also investigate and report on relevant underlying primary studies, providing policy-relevant recommendations based on the evidence reviewed.

Existing systematic reviews

Several reviews summarising the effects of built or policy environmental change on obesity outcomes have been published²⁰⁻²⁷. The Evidence for Policy and Practice Information and Co-ordinating Centre (EPPI-Centre) has mapped out systematic reviews (SRs) on social and environmental interventions to address childhood obesity²⁸, and a number of overviews of systematic reviews^{29,30} on school-based

childhood obesity interventions ^{18,31–33} are available. Overviews on the impact of the built environment on physical activity have also been conducted ^{34,35}, however, to the best of our knowledge, no overview of reviews focusing specifically on environmental interventions to prevent or reduce excess weight in children exists.

Aim of this review

The primary aim of this review is to summarise the evidence reported in SRs on the range and effectiveness of population-level interventions aimed at preventing overweight or obesity in children which have an environmental component. We adopted a broad interpretation of what is meant by an ‘environmental intervention’, defined as a strategy that involves changing the physical surroundings and social, economic or organizational systems to facilitate healthy choices and enable people to adopt healthy behaviours without requiring significant motivation on the part of the individual. ‘Effectiveness’ was defined as achieving a beneficial or protective, statistically significant, anthropometric outcome. A secondary aim is to assess and critique the methodological quality of included systematic reviews.

Methods

An overview of systematic reviews was conducted. Reviews were judged to be systematic if they synthesised peer-reviewed articles; explicitly reported pre-defined objectives, search strategy details and inclusion and exclusion criteria; and clearly identified all included studies. The full text of potentially eligible environmental interventions reported upon in included SRs was also reviewed, and relevant study data extracted.

Search strategy

We conducted a search of free text terms and subject headings from January 1995 to May 2015, using the Population, Intervention, Comparison and Outcome model ³⁶, to describe the target population (healthy children and adolescents), intervention (population-level prevention of obesity), comparison (SRs), and outcome (anthropometric outcomes). The initial search string was developed in Medline

(Appendix S1) and further refined for use in the different databases: Database of Abstracts of Reviews of Effects (DARE), Medline, Embase, PsychINFO, CINAHL PLUS, SCOPUS, Social Policy and Practice Database, Database of promoting health effectiveness reviews (DoPHER), and CENTRAL. Reference lists of identified SRs were manually searched to identify any additional reviews. A grey literature search in TRIP and Google Scholar was performed. Project websites and those of collaborative groups that conduct SRs of public health interventions (e.g. EPPI-Centre ²⁸; the Community Guide ³⁷; Health Systems Evidence ³⁸; Health Evidence Network ³⁹; Agency for Healthcare Research and Quality ⁴⁰; Centre for Reviews and Dissemination ⁴¹; and the National Institute for Health and Care Excellence ⁴²) were also searched for relevant publications, and their bibliography reviewed. No language restrictions were applied.

Inclusion and exclusion criteria

Systematic reviews were required to be: published between January 1995 and March 2015; reviewing interventions to prevent obesity and overweight in children and adolescents aged 5 - <18 years; assessing at least one population-level intervention with an environmental component; and reporting anthropometric outcome data on the effectiveness of interventions (i.e. using a standardized or accepted measure of obesity such as body mass index (BMI), BMI z-score, waist circumference (WC), skin fold thickness (SFT), percentage body fat (% BF), overweight or obesity prevalence and other anthropometric measures associated with obesity ⁴³).

SRs were excluded if they assessed interventions: aimed at adults only; which did not include any anthropometric outcomes, or where the only outcome of interest was behaviour modification (i.e. increasing physical activity, decreasing sedentariness or improving diet); aimed solely at treatment of existing obesity or expressly targeted weight loss (e.g. pharmacological interventions; bariatric surgery; metabolic or weight loss clinics) or aimed at participants with diagnosed complications linked to obesity; and that solely involved interaction between health professionals and individuals or groups within a clinical setting.

SRs focusing on treatment of existing obesity were excluded as it is typically harder to lose weight than to prevent an initial weight gain. Since our main interest was population-level preventive environmental

interventions, SRs exclusively assessing studies conducted in controlled clinical or laboratory settings were not considered relevant. In the case of reviews that contained a mix of interventions (e.g. aimed at both adults and children; reviewed both treatment and preventive interventions; or reported both behavioural and/or anthropometric outcomes), only data of preventive primary studies aimed at children and which had clearly defined anthropometric outcomes were extracted. Interventions in the included SRs could be multi-component or single-component. Multi-component interventions that combined individual and population-level elements were considered eligible if the population-level component was judged to be more than simple reinforcement of an individual-level intervention.

SRs were included if at least one reviewed primary study described a structural/policy change to the state, community, school, and/or home as a major component of a population-level, obesity-prevention intervention. Policies could be either formal legislative or organizational in scope⁴⁴. Eligible policies within the school setting included changes in school lunch nutrition standards or banning of vending machines. Provision of physical activity (PA) opportunities after school hours was considered to be a population-level environmental intervention, as these can potentially be made available to children living around the school neighbourhood who are not themselves students at the school. However, for the purposes of this overview, modification of school curricula to improve dietary or PA behaviour (e.g. increasing the number, duration or quality of physical education (PE) or nutrition education lessons during school hours through staff training or employing professionally-trained staff; increase in recess time etc.) in isolation of complementary environmental approaches was not considered to be eligible as such interventions have already been adequately described elsewhere³¹⁻³³. Additionally, evidence suggests that studies promoting healthier behaviour rather than focusing on reducing adiposity are less effective at reducing anthropometric outcomes^{45,46}. Eligible community interventions included modifications to the built environment (e.g. creation of walking pathways), whereas acceptable alterations to the home environment could include installation of television monitors. We also included exergaming interventions, as this novel approach to modifying children's leisure-time behaviour can potentially be delivered at population level. Conversely, interventions that focused solely on imparting

information and knowledge (e.g. educational campaigns; nutrition classes) were ineligible because such interventions aim to directly alter individual behaviour, rather than modify children's surroundings.

Systematic review selection and data extraction

DC and KG independently examined titles, abstracts and full-text articles, and extracted data, resolving any disagreement through discussion with a third author (CK). Within each review, interventions having an eligible environmental component that reported on outcomes of interest were identified. The methodological approach of each review (databases searched; language restrictions; inclusion and exclusion criteria; synthesis method; quality evaluation of trials; stated implications for practice and research; limitations; and funding sources) and documented primary review outcome indicators and main findings were recorded. All review-level data can be found in Appendices S1-S7. To avoid biased post hoc decisions, a review protocol based on Cochrane handbook recommendations⁴⁷ was published on PROSPERO⁴⁸ prior to starting the review process.

Quality of included reviews

Two reviewers (DC and KG) independently assessed the methodological quality of the included reviews using the 'Assessment of Multiple Systematic Reviews' (AMSTAR) tool⁴⁹. Reviews were categorised into high (AMSTAR score 8 – 11), medium (5 – 7) or low (0 – 4) based on the Canadian Agency for Drugs and Technologies in Health evaluation criteria⁵⁰. Any disagreement was resolved through consensus.

Identifying relevant primary studies included in the reviews

Most reviews only reported on the direction of effect on outcomes and did not report on the magnitude of effect, making it necessary to review the full text of primary studies in order to identify whether an intervention had a desirable or significant effect. Thus a list of all eligible primary studies (n=76) included in the included reviews was compiled in order to cross-check eligibility, assess clarity of information provided in SRs and to compile a comprehensive, useful list of existing environmental options. We collected information on study design, study setting (school, community, home, state),

characteristics of participants (including sample size, age), type of intervention (including type, intervention duration, length of follow-up, intervention components and source of funding), and outcome data (including difference in change from baseline for intervention vs control, direction and significance of effect). Effects without p-values, confidence intervals or a written statement regarding statistical significance were classified as non-significant. For multiple anthropometric outcomes, one or more statistically-significant benefit in any relevant outcome was counted as an overall beneficial study regardless of the number of non-significant outcomes, and vice-versa. All extracted intervention-level data can be found in Appendix S6. We did not report on quality of primary studies.

To correct for potential discrepancies between reported results for eligible environmental interventions, it was necessary to review the full text of each eligible primary study summarized in SRs. We were interested in the extent to which primary studies with environmental components were correctly identified or described as such in different SRs. In other words, would a policy-maker assessing the evidence on environmental obesity-prevention interventions be able to identify the studies of interest within an SR? We found that in many reviews, primary studies might not necessarily be classified outright as being environmental in nature, yet careful assessment of the information provided in the SR would enable an environmental component to be identified. The converse is also possible. For example, the study by James et al. 2004⁵¹ is classified in at least one SR⁵² as having an environmental component, yet we determined the intervention to be cognitive and behavioural in nature.

Results


Review characteristics

We included 63 systematic reviews (Figure 1) in our study. These searched databases up to 2014, reporting on a wide range of primary studies. Seven reviews focussed specifically on policies or environmental strategies, five explored mainly behavioural interventions, four looked at exergaming or active video games as their intervention of interest, and two concentrated on educational interventions in the school. The remainder (n=45) summarised the effects of a broad mix of interventions. The overall aim of all SRs was broadly to summarise obesity prevention studies – however there was substantial

heterogeneity in terms of setting, population of interest, type of intervention assessed, outcomes considered, review methodology, presentation of results and interpretation of findings (Appendix S3, S4). Eighteen of the reviews were meta-analyses, while the remaining 45 SRs provided a narrative synthesis of results. Twenty-eight SRs restricted their search to articles published in English only, eight SRs did not report on language in their search strategy, while the remainder widened their search to two or more languages. When analysed by setting, 28 assessed school-based studies exclusively⁵³⁻⁸⁰, six summarised community-based interventions^{24,81-85}, two reviews discussed home or family based interventions^{86,87}, and the remainder (n = 27) did not specify study settings in their search strategy^{45,52,88-111}. Most reviews (62%, n=39) assessed quality of the primary studies.

Review quality

Over four-fifths (84%) of the included reviews were assessed to be of low (n = 23) or medium (n = 30) quality according to AMSTAR criteria (Appendix S5). Ten reviews, six of which were meta-analyses, were judged to be of high quality^{52,60,76,81,87,91,92,99,104,111}.



Source: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). *Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement*. PLoS Med 6(6): e1000097. doi:10.1371/journal.pmed1000097

Figure 1: PRISMA flow chart of selection procedure

Summary of potentially effective environmental strategies according to high quality reviews

Fewer than half (n = 26 or 42%) of all SRs specifically discussed built or policy environmental strategies, with most stating that such strategies show promise but require further rigorous assessment and evaluation of social, psychosocial, behavioural, and anthropometric outcomes to prove benefit (Table 1). Overall, reviews were cautious about providing definitive recommendations on which environmental strategies should be implemented. This was due to the often suboptimal methodological quality of primary studies and the challenge of distinguishing which specific components of interventions were necessary to achieve positive outcomes ^{91,111}. Thus, few SRs provided clear statements on the importance of environmental strategies to address childhood obesity, and findings were mixed. However, some environmental interventions were highlighted across high-quality reviews as being particularly promising and likely to be effective in preventing or reducing overweight and obesity in children, especially if part of long-term comprehensive efforts ⁷⁶. These included increased PA sessions ¹¹¹; purchase of PE equipment ⁶⁰; improvements in nutritional quality of the food supply in schools ¹¹¹; ; creation of environments and cultural practices that support consumption of healthier foods and PA at school ^{60,111} and at home ^{76,87,92,111}; and capacity building or professional development for teachers to implement health promotion strategies and activities ¹¹¹.

Table 1: Systematic review comments on environmental strategies to prevent childhood obesity

Environmental interventions....	No. of reviews	References: first author and year of publication
are promising, effective strategies that can support and increase effectiveness of other obesity prevention programmes	10	Avery 2015 ⁸⁸ , Brandt 2010 ⁵³ , De Bourdeaudhuij 2010 ⁵⁸ , Ickes 2014 ⁶² , Katz 2008 ⁶⁵ , Kesten 2011 ¹⁰⁰ , Kropski 2008 ⁶⁷ , Peterson 2007 ⁷¹ , Sharma 2007 ⁷² , Sobol-Goldberg 2013 ⁷⁶
should be prioritised or at least considered for obesity prevention	7	Beauchamp 2014 ⁸⁹ , Budd 2006 ⁵⁵ , Ickes 2014 ⁶² , Kamath 2008 ⁵² , Sharma 2007 ⁷² , Showell 2013, Stice 2006, Waters 2011
minimise health inequalities	1	Beauchamp 2014 ⁸⁹
[specific examples of environmental interventions provided]	15	Avery 2015 ⁸⁸ , Beauchamp 2014 ⁸⁹ , Brandt 2010 ⁵³ , Budd 2006 ⁵⁵ , Chriqui 2014, De Bourdeaudhuij 2010 ⁵⁸ , de Sa 2008, Kamath 2008 ⁵² , Katz 2008 ⁶⁵ , Kropski 2008 ⁶⁷ , Li 2008 ¹¹² , Peterson 2007 ⁷¹ , Sharma 2007 ⁷² , Showell 2013 ⁸⁷ , Waters 2011 ¹¹¹
are of unclear effectiveness or viability	2	Marsh 2014 ⁸⁶ , Towns 2014 ¹⁰⁹
require further rigorous assessment/evaluation of associated policy and environmental, social, psychosocial, behavioural, and biological outcomes	10	Calancie 2015 ²⁴ , Chriqui 2013 ⁹⁴ , de Sa 2008 ⁵⁹ , Ickes 2014 ⁶² , Jaime 2009 ⁶³ , Katz 2008 ⁶⁵ , Peterson 2007 ⁷¹ , Reilly 2003 ¹⁰⁷ , Stice 2006 ⁴⁵ , Waters 2011 ¹¹¹
need for more widespread recognition of environmental influences operating counter to school activities	5	Budd 2006 ⁵⁵ , De Bourdeaudhuij 2010 ⁵⁸ , Katz 2008 ⁶⁵ , Peterson 2007 ⁷¹ , Stice 2006 ⁴⁵
need for research to explore feasibility, cost and effectiveness of environmental strategies	4	Calancie 2015 ²⁴ , Jaime 2009 ⁶³ , Knowlden 2013 ⁶⁶ , Showell 2013 ⁸⁷
need for research on options outside of the school	9	Calancie 2015 ²⁴ , Chriqui 2013 ⁹⁴ , Chriqui 2014 ⁹⁵ , de Sa 2008 ⁵⁹ , Jaime 2009 ⁶³ , Kesten 2011 ¹⁰⁰ , Knowlden 2013 ⁶⁶ , Peterson 2007 ⁷¹ , Reilly 2003 ¹⁰⁷
need for improved longitudinal data with obesity-related outcomes	4	Chriqui 2014 ⁹⁵ , De Bourdeaudhuij 2010 ⁵⁸ , Jaime 2009 ⁶³ , Katz 2008 ⁶⁵

Effect Sizes reported in Meta-Analyses

The meta-analyses identified in our search (Appendix S3, S4) reported BMI/zBMI reduction effect sizes (ES) for interventions aiming to prevent obesity in children. While it is not possible to extract the contribution to overall effect size of specific environmental elements within these interventions, the meta-analyses reported a wide range of effect sizes for different types, settings and duration of interventions. Across all settings, strategies targeting sedentary behaviour emerged as the most consistently successful^{57,65,104,110}, with ES ranging from -0.14 [-0.23, -0.05]¹¹⁰ to -0.35 [-0.63, -0.06]⁶⁵. The impact of physical activity interventions was unclear^{52,65,79,111}, as ES ranged from potential increases in BMI +1.87 [1.31, 2.42]⁶⁵ to reductions of -0.11 [-0.19, -0.02]¹¹¹; whereas dietary interventions^{52,57,65,111} showed a consistently modest beneficial impact, with an ES ranging from -0.02 [-0.07, 0.02]⁷⁹ to -0.39 [-0.56, -0.23]⁶⁵. Williams et al. calculated effect sizes on BMI for provision of school breakfast or lunch to students in isolation⁷⁹. Effect sizes for these interventions varied from a disappointing BMI increase of +0.04 [-0.19, 0.27] following the introduction of school lunches, to a minimal reduction in student BMI of -0.08 [-0.14, -0.02] following the introduction of a healthy school breakfast. Possibly reflecting the wide spread of ES above, meta-analyses assessing combined dietary and PA interventions^{52,65,79,91,104,111} calculated more modest ES ranging from 0.00 [-0.47, 0.47]⁷⁹ (no impact) to a small reduction in children's BMI of -0.18 [-0.27,-0.09]¹¹¹. With regards to the settings where interventions took place, Wolfenden et al.⁸⁵ reported a minor but beneficial (ES: -0.09 [-0.16, 0.02]) impact resulting from community-based interventions of any type, whereas a number of meta-analyses of school-based interventions^{64,65,68,74,76,77,79,111} reported a mix of results, with ES ranging from +0.17 [-0.38, 0.72]⁶⁴ to -0.29 [-0.45, -0.14]⁶⁵. Duration of intervention^{45,52,57,77,111} also seems to positively influence intervention outcome, as interventions lasting more than 12 months showed ES ranging from -0.095⁷⁷ to -0.12 [-0.21, -0.03]¹¹¹. Lastly, specific components of interventions reviewed in meta-analyses included parental involvement^{57,76,77}, which resulted in small but consistently positive ES ranging from -0.094 [p = <0.001]⁷⁷ to -0.151 [-0.334, 0.031]⁷⁶; and substitution of SSB with zero-calorie replacements^{99,105}, where results are less clear (ES: +0.06 [-0.01, 0.13]⁹⁹ to -0.17 [-0.39, 0.05]¹⁰⁵).

Primary studies

Appendix S6 reports on primary study characteristics including the range of environmental strategies undertaken and their effect on anthropometric outcomes, whereas Appendix S7 illustrates the overlap of all eligible primary studies (n = 76) across all SRs. Around half (48%; n = 37) of the primary studies did not result in a significant desirable effect on anthropometric outcomes (i.e. improvement in outcome of intervention compared to control), with some studies showing significant worsening outcomes for the intervention group ^{113–115} or mixed results ^{116–118}. Few RCTs consisted of purely environmental strategies or reported outcome data that can be attributed to environmental change directly. Most studies incorporated modifications to the environment as part of an overall strategy that included nutrition/PE education and curricular changes, and hence it is difficult to disentangle the impact of the environmental component from the overall impact.

Effective environmental strategies discussed in systematic reviews

A total of forty eligible primary studies across all SRs demonstrated a significant beneficial or protective effect on one of the anthropometric measures of childhood overweight and obesity (see Appendix S6 for study outcomes), and were reviewed in further detail to identify potentially effective environmental components. Almost all were primarily school-based, with the exception of two interventions that mainly took place in the home ^{119,120}. The majority contained an educational or information-dissemination component aimed at reinforcing the overall aim of each study. Environmental programme components - with or without additional behavioural and educational components - which appear to have contributed to the beneficial effects observed are outlined below. Anthropometric outcomes (changes from baseline for intervention groups compared to control groups) are provided where the intervention consisted mainly of a single environmental element.

- **Improvement of overall school food environment:** measures included implementation of school nutrition standards/policies; removal of vending machines selling sugar sweetened beverages (SSB) or snacks high in fat, sugar or salt (HFSS); banning sales of HFSS food; reformulation of school lunches to reduce fat content ^{116,117,121–126}

- **Purchase of new PE/sports equipment:** this was made available during recess and at other times throughout the school day, as well as during PE lessons ^{122,124,125,127–131}
- **Daily formal PA session organized after-school:** typically lasting 90 minutes to two hours, involving a substantial proportion of time spent doing MVPA, with or without a healthy snack provided to participants ^{123,132–136}. Three studies assessing impact of two-hour after-school interventions reported improvements in BMI ranging from -0.16 [-0.40, 0.07] ¹³⁵ to -0.45 [-0.79, -0.12] ¹³²; and differences in percentage body fat (%BF) from -0.76 [-1.42, -0.09] ^{135,136} to -2.01 [-2.98, -1.04] ¹³²
- **Provision of free or low-cost fruit:** ensuring that fruit ^{128,131,137} and freshly made fruit juices ¹³⁷ were available at school, and at home
- **Availability of school playgrounds for structured/unstructured PA after regular school hours** ^{124,125,137,138}
- **Provision of free/low cost water in school:** either through installation of water fountains, provision of water bottles or through lowering the cost of bottled water compared to other drinks in the school canteen ^{128,130,131,139–141}. An intervention focusing solely on enhancing water provision in schools reported a small reduction in zBMI (-0.004 [-0.045, 0.036]) and a significant reduction in the risk of overweight (31% reduction, $p = 0.04$) among the intervention group ¹⁴¹.
- **Provision of a healthy breakfast at school** (BMI: -0.11 ($P < 0.05$) in boys; -0.02 ($P < 0.05$ in girls) ¹⁴²
- **Substitution of sweetened beverages:** replacement of SSB with artificially sweetened, zero-calorie substitute (BMI: between -0.13 [-0.21, -0.05; $p = 0.001$] and -0.14 [-0.54 to -0.26] ^{119,143}
- **Reduction in screen time:** at home through the installation of an electronic television time manager device to limit TV watching (BMI: -0.45 [-0.73, -0.17; $p = 0.002$] ¹²⁰

Adjunct elements commonly adopted in the above studies and which are likely to contribute to overall effectiveness, but which our inclusion criteria precluded from further assessment, include: an increase in opportunities for PA during the school day ^{123,124,128–131,138}; increase in number, duration or quality

(e.g. proportion of time spent doing MVPA) of PE lessons ^{116,117,122}; parental involvement ^{126,128,129} and provision of training for food service staff or PE teachers ¹²⁷.

Few identifiable patterns emerged regarding single or multi-component interventions which failed to show any significant beneficial anthropometric outcomes. All active videogame studies, and all after-school PA programmes of less than 90 minutes duration, did not have a significant impact. Multi-level primary studies that had achieved significant positive results in certain contexts and settings (e.g. comprising components such as nutritional changes in schools or increased opportunities for PA, as shown above) failed to show effectiveness elsewhere, suggesting that contextual factors might have an important role in determining intervention success.

Discussion

This study provides a broad and comprehensive overview of environmental strategies to prevent childhood obesity at population level. Most interventions had at best a small to modest impact on childhood anthropometric outcomes (Appendix S6). Single-level interventions that focus on reducing screen time or increasing time spent performing MVPA may also be particularly beneficial. There was no clear link between the number and range of components of an intervention, and effectiveness of outcomes. However, our findings provide some support for consideration of obesity-prevention interventions having one or more environmental components, which may be particularly attractive for increasingly autonomous adolescents who may not respond to conventional nutrition education and behavioural counselling ^{144,145}. It is now important to understand which of these strategies can be combined into an ideal package at population level ¹¹¹. Most SRs did not provide clear recommendations regarding environmental components that should be considered for implementation by policymakers. At primary study level, the main focus was on school-based programmes, with few trials assessing potential environmental influences in the home, community or country setting.

A comparison of SR conclusions suggests that most SRs are cautious about summarising the evidence regarding the relative effectiveness of environmental components, or fail entirely to do so. With few notable exceptions, such as the SR by Waters et al. ¹¹¹, review authors provide limited descriptions of

individual studies without identifying which intervention elements might be most effective. Hence, they are likely to be of limited use to policy makers looking for concrete suggestions regarding which elements are most likely to successfully stop or reverse childhood obesity. This led to our analysis of primary studies within SRs in an attempt to identify those components which achieved the largest magnitude of effect. However, any statement of effectiveness emerging from these studies must be interpreted with caution given the overall methodological challenges and lack of detail provided even in the higher quality reviews and primary studies. In addition, observed changes in weight outcomes in preventive studies having a normal population distribution may be subject to interpretation. For example, in studies where increases in BMI were observed, it is often difficult to discern whether these were due to normal child growth that is within acceptable limits; reflective of normalization of weight outcomes for previously underweight children; or due to an increase in muscle mass. A comparison of intervention effectiveness and cost-effectiveness was beyond the scope of this research, although questions remain regarding whether subjective judgements on the magnitude of a study's impact are useful or acceptable (e.g. is a 10% reduction in risk of population obesity a less or more beneficial outcome than a 0.1 reduction in population BMI?); or whether the ratio of study cost to benefit provided should be considered when evaluating interventions. Furthermore, heterogeneity of data; generic lack of information on costs; the wide range of programme components reviewed; and the difficulties of disentangling the individual contribution of distinct strategies packaged within multi-component programmes to the final magnitude of effect means that it is not possible to distinguish which of these components are the most beneficial. These problems limit our ability to conclude that one strategy or combinations of strategies are more important than others in the prevention of childhood obesity ¹⁴⁶. We sought to identify a shortlist of environmental components having at least some evidence of successful implementation, and identified several strategies that show promise. This is not to say that they will always be effective, or that they can be easily scaled up to population level. Although the 1999 study by Robinson et al. aiming to reduce sedentary behaviour at home by installing a TV locking device was particularly effective, similar studies failed to show a significant impact ¹⁴⁷ or even resulted in higher BMI ^{148,149}. There are also issues around acceptability, long-term sustainability and economic viability of scaling up such interventions ¹⁴⁶.

Gaps in the research

Our results enable us to identify a number of gaps in the existing evidence. We confirm that relatively little review-level evidence is available on the impact of environmental interventions on children and adolescents, and most of what is available concerns school-based interventions²⁸. We purposely selected only reviews which reported anthropometric outcomes – and presented only relevant adiposity-related data for the identified primary studies– because achieving improved nutrition and PA levels does not necessarily translate into improved anthropometric outcomes, and the latter tend to be the main outcomes of interest to policy makers. While this resulted in a good number of reviews for inclusion, preliminary searches showed that there is a dearth of high-quality reviews focusing specifically on upstream environmental approaches to tackling childhood obesity. In addition, few SRs of any quality explicitly addressed macro-level interventions such as the introduction of school food and beverage policies^{59,63,94,95} or implementation of broad community-based strategies^{24,83}. It is unclear whether this is a genuine gap in the literature or a direct consequence of few relevant primary studies having been published in this area. In addition, the search strategy adopted by each review tends to inherently limit the range of primary studies included, and hence the scope of review conclusions and recommendations. In any case, we echo previous calls for further primary and secondary research into this key paradigm of population-level obesity prevention^{65,111}.

Implications for Research and Practice

Policy/environmental approaches to addressing childhood obesity show promise and should be strongly recommended for obesity prevention; however more studies, ideally large, longitudinal natural experiments conducted outside of the school environment with sufficient sampling power to ascertain effect sizes with some confidence, should be carried out^{98,111,150}. The fact that almost a third of reviews focused on school-based studies indicates that schools are considered to be key sites for childhood and adolescent obesity prevention interventions, presumably because children spend a substantial portion of their day there, and the relative ease with which interventions can be trialled. However, interventions

in schools that were successful in reducing obesity at first glance have been shown to lose their effect during the summer ¹¹³ and are often difficult to sustain among the same population in the longer term. Thus, engagement of the community and environmental support within the home are essential aspects of successful interventions. Studies should also implement more rigorous analysis and evaluation of associated social, psychosocial, behavioural, and biological outcomes, particularly adiposity outcomes. Undertaking formal economic evaluations would also add substantially to the utility of a study for policymakers ¹¹¹, yet these are rarely performed, and there is little review-level economic data in this regard.

Limitations

Publication bias may be a potential limitation of this overview. Other SRs might exist but have not been submitted or accepted for publication and therefore could not be identified during our search. We are also aware that since the searches for this overview were carried out in March 2015, other reviews on this topic may have been published. Our criteria with regards to what constitutes an ‘environmental’ intervention meant that a large number of primary studies focusing on improving PE lessons during school hours, which possibly contributes towards achieving desirable anthropometric outcomes, were excluded. Quality appraisal of included reviews using the AMSTAR tool presented a number of issues which have also been raised by other assessors ¹⁵¹. For example, only four reviews (including two Cochrane reviews) achieved a ‘yes’ rating in the ‘conflict of interest included’ criterion, which specifies that sources of funding or support for both the review itself as well as for each of the included studies should be reported. It is debatable whether indicating source of funding for primary studies reflects quality of reporting, rather than methodological quality of the review itself. Additionally, we encountered some difficulty in ascertaining multiple publications evaluating the same study, particularly in the case of long term studies. It was rarely immediately clear that publications were contiguous unless the project name was used in the title of the article, as in the case of the Medical College of Georgia FitKid Project ^{113,135} or APPLE project ^{130,131} publications. In other cases, including ICAPS ¹³⁸ or the Dutch Obesity Prevention Intervention in Teenagers ^{116,117}, references to previous publications were buried in the full text. Our exclusion of non-anthropometric outcomes means that

impact of interventions on other potentially valid outcomes such as improvements in PA or nutrition behaviour, VO2 max, blood pressure and blood cholesterol levels were not assessed. However, there is evidence to suggest that these do not necessarily translate into reduced BMI ^{46,152}.

Like Woodman et al., we had expected to find greater overlap of primary studies between reviews, and agree with their view that this can be attributed to differences in inclusion criteria and outcome assessments of SRs rather than erroneous search strategies ¹⁵³. Where unique primary studies were included in multiple reviews, their intervention design and findings were not necessarily consistently reported, potentially leading to type II errors during this overview process. Furthermore, inconsistencies in the definition of what constitutes an ‘environmental’ component of an intervention were seen. We suggest that the reporting structure of SRs could be improved by specifying the types of strategies (e.g. cognitive, educational, behavioural, parental, environmental) that included interventions employ, perhaps in table form as done by Sobol-Goldberg et al. ⁷⁶ or Kamath et al. ⁵². We found this approach particularly useful because it enabled us to directly assess the primary studies of interest, rather than go through the description of all included studies to judge whether one of its components was environmental in nature. On the other hand, this experience highlights the challenge of understanding what is meant by an ‘environmental intervention’: how is the term operationalized when conducting the review? Few SRs provided a clear, explicit definition, with most opting for, at most, a brief illustrative example ²⁴. More detailed and pragmatic frameworks for describing primary studies would be valuable ²⁸.

Our methods led to the exclusion of a number of high quality SRs which implicitly or explicitly addressed environmental interventions, such as those by Matson-Koffman et al. ²³ and Wang et al. ¹⁵⁰. These failed to provide sufficient descriptive or outcome data for us to clearly identify which included interventions were environmental in nature, or whether they had any impact on obesity-related outcomes in school-aged children. However, to ‘test’ the comprehensiveness of our approach to identifying primary studies using review-level data, we compared the primary studies included in the Wang et al. review ¹⁵⁰ with our final list of 76 eligible primary studies having an environmental

component (Table S5). Wang et al. identify ten studies as having an environmental component in the main text of their article. Five of these were included in our list because they overlapped with SRs that we had included in our overview. Three primary studies were not eligible according to our inclusion criteria. However, our approach missed two relevant primary studies with clearly reported environmental components and anthropometric outcomes^{154,155} that were reported in Wang et al.'s article. The intervention by Pettman et al. (2014)¹⁵⁴ had not been reported upon in any of the SRs included in our study. However, the study by Chang et al. (2010)¹⁵⁵ had been cited the review by Bleich et al. (2010)⁸³ that we had included in our overview, but which was described by the review authors as having “social marketing, strategic partnerships, knowledge mobilization, strategies in multiple sectors” elements. There was no indication of any environmental component for this intervention. Based solely on the study description provided by Bleich et al.'s reporting, and taking into consideration that the SR had clearly described several other primary interventions as being ‘environmental’ in nature (or described a study component in sufficient detail to enable its classification as ‘environmental’ overview authors), we did not look at the full text of the Change et al. (2010) study and thus missed the opportunity to include it in our list of environmental interventions. This highlights one of the disadvantages of using review-level evidence to inform policy: the evidence obtained is only as useful as the quality of presentation of data in the reviews.

Strengths

There are several strengths of this study. As many decisions as possible were made a priori to limit potential bias throughout the overview. All stages of the overview (i.e., inclusion criteria, exclusion criteria, data extraction, AMSTAR tables) were conducted in duplicate to minimize error. Our search strategy was sensitive and inclusive of several databases and grey literature, and our definition of ‘environmental interventions’ was broad. No language restrictions were applied. Consequently, the reviews analysed here, and the list of primary obesity prevention studies having an environmental component derived from this overview, are likely to represent the majority of relevant reviews and interventions available at the time of our search.

Conclusions

Environmental interventions may be modestly effective in addressing childhood obesity. However, there is a dearth of research into the feasibility and effectiveness of implementing environmental strategies in non-school settings, and information on programme cost that may be of use to policymakers seeking to translate evidence into practice is lacking. Most reviews suffered from poor methodology and presentation, making it difficult to assess the true effects of interventions on adiposity outcomes and necessitating retrieval of full texts of primary studies to ensure comprehensiveness.

Author contributions

DC performed the systematic review search, reviewed articles for inclusion, extracted data, generated tables (including appendices) and drafted the manuscript. KG independently reviewed articles for inclusion, extracted data, and reviewed and edited text. CK helped develop the research question, designed and piloted the extraction form for a portion of included articles, and reviewed and edited text. MP helped develop the research question and reviewed and edited text.

Supporting information

Appendix S1. Ovid-Medline search strategy

Appendix S2. Excluded Systematic reviews

Appendix S3. Characteristics of included systematic reviews (part 1)

Appendix S4. Characteristics of included systematic reviews (part 2)

Appendix S5. Quality assessment of systematic reviews (AMSTAR)

Appendix S6. Characteristics of primary studies having an eligible environmental component

Appendix S7. Overlap of primary studies across systematic reviews

References

- 1 Ahluwalia N, Dalmasso P, Rasmussen M *et al.* Trends in overweight prevalence among 11-, 13- and 15-year-olds in 25 countries in Europe, Canada and USA from 2002 to 2010. *Eur J Public Health* 2015; **25 Suppl 2**: 28–32.
- 2 Verschuuren M, Fietje N, Greenwell F, Raj T, Stein C, World Health Organization. The European Health Report 2015: Targets and beyond – reaching new frontiers in evidence. Copenhagen: UN City, 2015 <http://www.euro.who.int/en/data-and-evidence/european-health-report/european-health-report-2015/ehr2015>.
- 3 Dehghan M, Akhtar-Danesh N, Merchant AT. Childhood obesity, prevalence and prevention. *Nutr J* 2005; **4**: 24.
- 4 Parsons TJ, Power C, Logan S, Summerbell CD. Childhood predictors of adult obesity: a systematic review. *Int J Obes Relat Metab Disord* 1999; **23 Suppl 8**: S1–107.
- 5 Serdula MK, Ivery D, Coates RJ, Freedman DS, Williamson DF, Byers T. Do obese children become obese adults? A review of the literature. *Prev Med (Baltim)* 1993; **22**: 167–177.
- 6 World Health Organization. Global status report on noncommunicable diseases 2014. World Health Organization: Geneva, 2014 <http://www.who.int/nmh/publications/ncd-status-report-2014/en/> (accessed 28 Jan 2016).
- 7 Owen CG, Whincup PH, Orfei L *et al.* Is body mass index before middle age related to coronary heart disease risk in later life? Evidence from observational studies. *Int J Obes (Lond)* 2009; **33**: 866–877.
- 8 Gortmaker SL, Swinburn BA, Levy D *et al.* Changing the future of obesity: science, policy, and action. *Lancet* 2011; **378**: 838–847.
- 9 Egger G, Swinburn B. An ‘ecological’ approach to the obesity pandemic. *BMJ* 1997; **315**: 477–480.
- 10 McLeroy KR, Bibeau D, Steckler A, Glanz K. An ecological perspective on health promotion programs. *Health Educ Q* 1988; **15**: 351–377.
- 11 Summerbell CD, Waters E, Edmunds LD, Kelly S, Brown T, Campbell KJ. Interventions for preventing obesity in children. *Cochrane database Syst Rev* 2005; : CD001871.
- 12 Flynn MAT, McNeil DA, Maloff B *et al.* Reducing obesity and related chronic disease risk in children and youth: a synthesis of evidence with ‘best practice’ recommendations. *Obes Rev* 2006; **7 Suppl 1**: 7–66.
- 13 Swinburn B, Egger G, Raza F. Dissecting obesogenic environments: the development and application of a framework for identifying and prioritizing environmental interventions for obesity. *Prev Med (Baltim)* 1999; **29**: 563–570.
- 14 NICE. Clinical Guideline 43: Obesity - Guidance on the prevention, identification, assessment and management of overweight and obesity in adults and children. National Institute for Health and Clinical Excellence, 2006.
- 15 Schwartz MB. Environmental and policy strategies to improve eating, physical activity behaviors, and weight among adolescents. *Adolesc Med State Art Rev* 2012; **23**: 589–609.

- 16 Bero LA, Jadad AR. How consumers and policymakers can use systematic reviews for decision making. *Ann Intern Med* 1997; **127**: 37–42.
- 17 Graham ID, Logan J, Harrison MB *et al*. Lost in knowledge translation: time for a map? *J Contin Educ Health Prof* 2006; **26**: 13–24.
- 18 Wolfenden L, Wiggers J, Tursan D’Espaignet E, Bell a. C. How useful are systematic reviews of child obesity interventions? *Obes Rev* 2010; **11**: 159–165.
- 19 Francis D, Turley R, Thomson H, Weightman A, Waters E, Moore L. Supporting the needs of public health decision-makers and review authors in the UK. *J Public Health (Oxf)* 2015; **37**: 172–4.
- 20 Faith MS, Fontaine KR, Baskin ML, Allison DB. Toward the reduction of population obesity: macrolevel environmental approaches to the problems of food, eating, and obesity. *Psychol Bull* 2007; **133**: 205–226.
- 21 French SA, Story M, Jeffery RW. Environmental influences on eating and physical activity. *Annu Rev Public Health* 2001; **22**: 309–335.
- 22 Hider P. Environmental interventions to reduce energy intake or density: a critical appraisal of the literature. *New Zeal Heal Technol Assessment NZHTA* 2001; **4**.<http://nzhta.chmeds.ac.nz/publications/nzhtav4n2.pdf>.
- 23 Matson-Koffman DM, Brownstein JN, Neiner JA, Greaney ML. A site-specific literature review of policy and environmental interventions that promote physical activity and nutrition for cardiovascular health: what works? *Am J Heal Promot* 2005; **19**: 167–193.
- 24 Calancie L, Leeman J, Jilcott Pitts SB *et al*. Nutrition-related policy and environmental strategies to prevent obesity in rural communities: a systematic review of the literature, 2002-2013. *Prev Chronic Dis* 2015; **12**: E57.
- 25 Mayne SL, Auchincloss AH, Michael YL. Impact of policy and built environment changes on obesity-related outcomes: a systematic review of naturally occurring experiments. *Obes Rev* 2015. doi:10.1111/obr.12269.
- 26 Osei-Assibey G, Dick S, Macdiarmid J *et al*. The influence of the food environment on overweight and obesity in young children: a systematic review. *BMJ Open* 2012; **2**. doi:10.1136/bmjopen-2012-001538.
- 27 Williams J, Scarborough P, Matthews A *et al*. A systematic review of the influence of the retail food environment around schools on obesity-related outcomes. *Obes Rev* 2014; **15**: 359–374.
- 28 Woodman J, Lorenc T, Harden A, Oakley A. Social and environmental interventions to reduce childhood obesity: a systematic map of reviews. EPPI-Centre, Social Science Research Unit, Institute of Education, University of London: London, 2008<http://eppi.ioe.ac.uk/cms/Default.aspx?tabid=2395>.
- 29 Becker L, Oxman AD. Chapter 22: Overviews of reviews. *Cochrane Handb. Syst. Rev. Interv.* Version 5.1.0 (updated March 2011). 2011.
- 30 Smith V, Devane D, Begley CM, Clarke M. Methodology in conducting a systematic review of systematic reviews of healthcare interventions. *BMC Med Res Methodol* 2011; **11**: 15.
- 31 Amini M, Djazayeri A, Majdzadeh R *et al*. Effect of School-based Interventions to Control

- Childhood Obesity: A Review of Reviews. *Int J Prev Med* 2015; **6**: 68.
- 32 Khambalia AZ, Dickinson S, Hardy LL, Gill T, Baur LA. A synthesis of existing systematic reviews and meta-analyses of school-based behavioural interventions for controlling and preventing obesity. *Obes Rev* 2012; **13**: 214–233.
- 33 Safron M, Cislak A, Gaspar T, Luszczynska A. Effects of school-based interventions targeting obesity-related behaviors and body weight change: a systematic umbrella review. *Behav Med* 2011; **37**: 15–25.
- 34 Gebel K, Bauman AE, Petticrew M. The physical environment and physical activity: a critical appraisal of review articles. *Am J Prev Med* 2007; **32**: 361–369.
- 35 Ding D, Gebel K. Built environment, physical activity, and obesity: What have we learned from reviewing the literature? *Health Place* 2012; **18**: 100–105.
- 36 Centre for Evidence Based Medicine. Asking Focused Questions - The PICO model. <http://www.cebm.net/asking-focused-questions/> (accessed 19 Nov2014).
- 37 Community Preventive Services Task Force. Obesity Prevention and Control: Interventions in Community Settings. 2015.<http://www.thecommunityguide.org/obesity/communitysettings.html>.
- 38 McMaster University. Health Systems Evidence. 2015; **2015**.<https://www.healthsystemsevidence.org/>.
- 39 WHO. Health Evidence Network. 2015.<http://data.euro.who.int/HEN/>.
- 40 U.S. Department of Health and Human Services. Agency for Healthcare Research and Quality . 2015.<http://www.ahrq.gov/>.
- 41 Centre for Reviews and Dissemination. CRD Database. 2015.<http://www.crd.york.ac.uk/CRDWeb/>.
- 42 National Institute for Health and Care Excellence. Evidence Search. 2015; **2015**.<http://www.evidence.nhs.uk/>.
- 43 Mei Z, Grummer-Strawn LM, Pietrobelli A, Goulding A, Goran MI, Dietz WH. Validity of body mass index compared with other body-composition screening indexes for the assessment of body fatness in children and adolescents. *Am J Clin Nutr* 2002; **75**: 978–985.
- 44 Schmid TL, Pratt M, Howze E. Policy as intervention: environmental and policy approaches to the prevention of cardiovascular disease. *Am J Public Health* 1995; **85**: 1207–11.
- 45 Stice E, Shaw H, Marti CN. A meta-analytic review of obesity prevention programs for children and adolescents: the skinny on interventions that work. *Psychol. Bull.* 2006; **132**: 667–691.
- 46 Harris KC, Kuramoto LK, Schulzer M, Retallack JE. Effect of school-based physical activity interventions on body mass index in children: a meta-analysis. *C Can Med Assoc J* 2009; **180**: 719–726.
- 47 Higgins JPT, Green S. Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]. 2011.Available from www.cochrane-handbook.org.
- 48 Cauchi D, Glonti K, Knai C, Petticrew MP. Modifying the obesogenic environment to prevent obesity in children: an overview of systematic reviews (CRD42013005768).

- 2015.<http://www.crd.york.ac.uk/PROSPERO/>.
- 49 Shea BJ, Hamel C, Wells GA *et al*. AMSTAR is a reliable and valid measurement tool to assess the methodological quality of systematic reviews. *J Clin Epidemiol* 2009; **62**: 1013–1020.
- 50 Grimshaw J, Hill S, Lowe D *et al*. Canadian Agency for Drugs and Technologies in Health, Methods for Development. 2015; **2015**.<https://www.cadth.ca/fr/methods-development>.
- 51 James J, Thomas P, Cavan D, Kerr D. Preventing childhood obesity by reducing consumption of carbonated drinks: cluster randomised controlled trial. *Bmj* 2004; **328**: 1237.
- 52 Kamath CC, Vickers KS, Ehrlich A *et al*. Behavioral interventions to prevent childhood obesity: a systematic review and metaanalyses of randomized trials. *J Clin Endocrinol Metab* 2008; **93**: 4606–4615.
- 53 Brandt S, Moss A, Berg S, Wabitsch M. School-based obesity prevention. How can it be realized?. [German]Schulbasierte Praventioen der Adipositas: Wie sollte sie aussehen? *Bundesgesundheitsblatt - Gesundheitsforsch - Gesundheitsschutz* 2010; **53**: 207–220.
- 54 Brown T, Summerbell C. Systematic review of school-based interventions that focus on changing dietary intake and physical activity levels to prevent childhood obesity: An update to the obesity guidance produced by the National Institute for Health and Clinical Excellence. *Obes Rev* 2009; **10**: 110–141.
- 55 Budd GM, Volpe SL. School-based obesity prevention: Research, challenges, and recommendations. *J Sch Health* 2006; **76**: 485–495.
- 56 Cole K, Waldrop J, D’Auria J, Garner H. An integrative research review: effective school-based childhood overweight interventions. *J Spec Pediatr Nurs* 2006; **11**: 166–177.
- 57 Cook-Cottone C, Casey Carolyn M, Feeley Thomas H, Baran J. A Meta-Analytic Review of Obesity Prevention in the Schools: 1997-2008. *Psychol Sch* 2009; **46**: p695–719.
- 58 De Bourdeaudhuij I, Van Cauwenberghe E, Spittaels H *et al*. School-based interventions promoting both physical activity and healthy eating in Europe: a systematic review within the HOPE project. *Obes Rev*; **12**: 205–216.
- 59 De Sa J, Lock K. Will European agricultural policy for school fruit and vegetables improve public health? A review of school fruit and vegetable programmes. *Eur J Public Health* 2008; **18**: 558–568.
- 60 Dobbins M, DeCorby K, Robeson P, Husson H, Tirilis D. School-based physical activity programs for promoting physical activity and fitness in children and adolescents aged 6-18. *Cochrane Database Syst Rev* 2013; **(2)**. doi:<http://dx.doi.org/10.1002/14651858.CD007651>.
- 61 Holub CK, Lobelo F, Mehta SM, Sánchez Romero LM, Arredondo EM, Elder JP. School-Wide Programs Aimed at Obesity Among Latino Youth in the United States: A Review of the Evidence. *J Sch Health* 2014; **84**: 239–246.
- 62 Ickes MJ, McMullen J, Haider T, Sharma M. Global school-based childhood obesity interventions: A review. *Int J Environ Res Public Health* 2014; **11**: 8940–8961.
- 63 Jaime PC, Lock K. Do school based food and nutrition policies improve diet and reduce obesity? *Prev Med (Baltim)* 2009; **48**: 45–53.
- 64 Kanekar A, Sharma M. Meta-analysis of school-based childhood obesity interventions in the

- UK and US. *Int. Q. Community Health Educ.* 2008; **29**: 241–256.
- 65 Katz DL, O’Connell M, Njike VY, Yeh MC, Nawaz H. Strategies for the prevention and control of obesity in the school setting: Systematic review and meta-analysis. *Int J Obes* 2008; **32**: 1780–1789.
- 66 Knowlden AE, Sharma M. Systematic Review of School-based Obesity Interventions Targeting African American and Hispanic Children. *J Heal Care Poor Underserved* 2013; **24**: 1194–1214.
- 67 Kropiski JA, Keckley PH, Jensen GL. School-based obesity prevention programs: An evidence-based review. *Obesity* 2008; **16**: 1009–1018.
- 68 Lavelle H V, Mackay DF, Pell JP. Systematic review and meta-analysis of school-based interventions to reduce body mass index. *J Public Health (Bangkok)* 2012; **34**: 360–369.
- 69 Lissau I. Prevention of overweight in the school arena. *Acta Paediatr Suppl* 2007; **96**: 12–18.
- 70 Lobelo F, Garcia de Quevedo IHCKNBJAEM. School-based programs aimed at the prevention and treatment of obesity: evidence-based interventions for youth in Latin America. *J Sch Health* 2013; **83(9)**: 668–677.
- 71 Peterson KE, Fox MK. Addressing the epidemic of childhood obesity through school-based interventions: What has been done and where do we go from here? *J Law, Med Ethics* 2007; **35**: 113–130.
- 72 Sharma M. International school-based interventions for preventing obesity in children. *Obes Rev* 2007; **8**: 155–167.
- 73 Shirley K, Rutfield R, Hall N, Fedor N, McCaughey VK, Zajac K. Combinations of Obesity Prevention Strategies in US Elementary Schools: A Critical Review. *J Prim Prev* 2014; **36**: 1–20.
- 74 Silveira JACD, Taddei JADAC, Guerra PH, Nobre MRC. The effect of participation in school-based nutrition education interventions on body mass index: A meta-analysis of randomized controlled community trials. *Prev Med (Baltim)* 2013; **56**: 237–243.
- 75 Silveira JAC, Taddei JAAC, Guerra PH, Nobre MRC. Effectiveness of school-based nutrition education interventions to prevent and reduce excessive weight gain in children and adolescents: A systematic review. *J Pediatr (Rio J)* 2011; **87**: 382–392.
- 76 Sobol-Goldberg S, Rabinowitz J, Gross R. School-based obesity prevention programs: A meta-analysis of randomized controlled trials. *Obesity* 2013; **21**: 2422–2428.
- 77 Vasques C, Magalhaes P, Cortinhas A, Mota P, Leitao J, Lopes VP. Effects of intervention programs on child and adolescent BMI: A meta-analysis study. *J Phys Act Health* 2014; **11**: 426–444.
- 78 Verstraeten R, Roberfroid D, Lachat C *et al.* Effectiveness of preventive school-based obesity interventions in low- and middle-income countries: a systematic review. *Am J Clin Nutr* 2012; **96**: 415–438.
- 79 Williams AJ, Henley WE, Williams CA, Hurst AJ, Logan S, Wyatt KM. Systematic review and meta-analysis of the association between childhood overweight and obesity and primary school diet and physical activity policies. *Int J Behav Nutr Phys Act* 2013; **10**: 101.
- 80 Zenzen W, Kridli S. Integrative review of school-based childhood obesity prevention programs. *J Pediatr Heal Care* 2009; **23**: 242–258.

- 81 Baker PR, Francis DP, Soares J, Weightman AL, Foster C. Community wide interventions for increasing physical activity. *Cochrane database Syst Rev* 2011; : CD008366.
- 82 Barr-Anderson DJ, Singleton C, Cotwright CJ, Floyd MF, Affuso O. Outside-of-school time obesity prevention and treatment interventions in African American youth. *Obes Rev* 2014; **15**: 26–45.
- 83 Bleich SN, Segal J, Wu Y, Wilson R, Wang Y. Systematic review of community-based childhood obesity prevention studies. *Pediatrics* 2013; **132**: e201–e210.
- 84 Small L, Anderson D, Melnyk BM. Prevention and early treatment of overweight and obesity in young children: a critical review and appraisal of the evidence. *Pediatr. Nurs.* 2007; **33**: 149–152.
- 85 Wolfenden L, Wyse R, Nichols M, Allender S, Millar L, McElduff P. A systematic review and meta-analysis of whole of community interventions to prevent excessive population weight gain. *Prev Med (Baltim)* 2014; **62**: 193–200.
- 86 Marsh S, Foley LS, Wilks DC, Maddison R. Family-based interventions for reducing sedentary time in youth: A systematic review of randomized controlled trials. *Obes Rev* 2014; **15**: 117–133.
- 87 Showell NN, Fawole O, Segal J *et al.* A systematic review of home-based childhood obesity prevention studies. *Pediatrics* 2013; **132**: e193–e200.
- 88 Avery a, Bostock L, McCullough F. A systematic review investigating interventions that can help reduce consumption of sugar-sweetened beverages in children leading to changes in body fatness. *J Hum Nutr Diet* 2015; **28 Suppl 1**: 52–64.
- 89 Beauchamp A, Backholer K, Magliano D, Peeters A. The effect of obesity prevention interventions according to socioeconomic position: a systematic review. *Obes Rev* 2014; **15**: 514–554.
- 90 Branscum P, Sharma M. A systematic analysis of childhood obesity prevention interventions targeting Hispanic children: lessons learned from the previous decade. *Obes Rev* 2011; **12**: e151–8.
- 91 Brown T, Smith S, Bhopal R, Kasim A, Summerbell C. Diet and physical activity interventions to prevent or treat obesity in south asian children and adults: A systematic review and meta-analysis. *Int J Environ Res Public Health* 2015; **12**: 566–594.
- 92 Campbell K, Waters E, O’Meara S, Kelly S, Summerbell C. Interventions for preventing obesity in children. *Cochrane Database Syst Rev* 2002; : CD001871.
- 93 Chen JL, Wilkosz ME. Efficacy of technology-based interventions for obesity prevention in adolescents: A systematic review. *Adolesc Health Med Ther* 2014; **5**: 159–170.
- 94 Chriqui JF. Obesity Prevention Policies in U.S. States and Localities: Lessons from the Field. *Curr Obes Rep* 2013; **2**: 200–210.
- 95 Chriqui JF, Pickel M, Story M. Influence of school competitive food and beverage policies on obesity, consumption, and availability: A systematic review. *JAMA Pediatr* 2014; **168**: 279–286.
- 96 Flodmark C, Marcus C, Britton M. Interventions to prevent obesity in children and

- adolescents: A systematic literature review. *Int J Obes* 2006; **30**: 579–589.
- 97 Gao Z, Chen S. Are field-based exergames useful in preventing childhood obesity? A systematic review. *Obes Rev* 2014; **15**: 676–691.
- 98 Haynos AF, O’Donohue WT. Universal childhood and adolescent obesity prevention programs: review and critical analysis. *Clin Psychol Rev* 2012; **32**: 383–399.
- 99 Kaiser KA, Shikany JM, Keating KD, Allison DB. Will reducing sugar-sweetened beverage consumption reduce obesity? Evidence supporting conjecture is strong, but evidence when testing effect is weak. *Obes Rev* 2013; **14**: 620–633.
- 100 Kesten JM, Griffiths PL, Cameron N. A systematic review to determine the effectiveness of interventions designed to prevent overweight and obesity in pre-adolescent girls. *Obes Rev* 2011; **12**: 997–1021.
- 101 Lamboglia CMGF, Silva VTBLD, Vasconcelos Filho JED *et al.* Exergaming as a Strategic Tool in the Fight against Childhood Obesity: A Systematic Review. *J Obes* 2013; **2013**. doi:Export Date 15 February 2015.
- 102 LeBlanc AG, Chaput JP, McFarlane A *et al.* Active Video Games and Health Indicators in Children and Youth: A Systematic Review. *PLoS One* 2013; **8**. doi:10.1371/journal.pone.0065351.
- 103 Leung MM, Agaronov A, Grytsenko K, Yeh MC. Intervening to reduce sedentary behaviors and childhood obesity among school-age youth: A systematic review of randomized trials. *J Obes* 2012; **2012**. doi:http://dx.doi.org/10.1155/2012/685430.
- 104 Liao Y, Liao J, Durand CP, *et al.* Which type of sedentary behaviour intervention is more effective at reducing body mass index in children? A meta-analytic review. *Obes Rev* 2014; **15**: 159–168.
- 105 Malik VS, An P, Willett WC, Hu FB. Sugar-sweetened beverages and weight gain in children and adults: a systematic review and meta-analysis. *Am J Clin Nutr* 2013; **98**: 1084–1102.
- 106 Pérez-Morales ME, Bacardí-Gascón M, Jiménez-Cruz a. Childhood overweight and obesity prevention interventions among Hispanic children in the United States: systematic review. *Nutr Hosp* 2012; **27**: 1415–21.
- 107 Reilly JJ, McDowell ZC. Physical activity interventions in the prevention and treatment of paediatric obesity: systematic review and critical appraisal. *Proc Nutr Soc* 2003; **62**: 611–619.
- 108 Sbruzzi G, Eibel B, Barbiero SM *et al.* Educational interventions in childhood obesity: a systematic review with meta-analysis of randomized clinical trials. *Database Abstr. Rev. Eff.* 2013; : 254–264.
- 109 Towns C, Cooke M, Rysdale L, Wilk P. Healthy Weights Interventions in Aboriginal Children and Youth: A Review of the Literature. *Can J Diet Pract Res* 2014; **75**: 125–131.
- 110 van Grieken A, Ezendam NP, Paulis WD, van der Wouden JC, Raat H. Primary prevention of overweight in children and adolescents: a meta-analysis of the effectiveness of interventions aiming to decrease sedentary behaviour. *Int J Behav Nutr Phys Act* 2012; **9**: 61.
- 111 Waters E, de Silva-Sanigorski A, Burford BJ *et al.* Interventions for preventing obesity in children. *Cochrane database Syst Rev* 2011; : CD001871.

- 112 Li M, Li S, Baur LA, Huxley RR. A systematic review of school-based intervention studies for the prevention or reduction of excess weight among Chinese children and adolescents. *Obes Rev* 2008; **9**: 548–559.
- 113 Gutin B, Yin Z, Johnson M, Barbeau P. Preliminary findings of the effect of a 3-year after-school physical activity intervention on fitness and body fat: the Medical College of Georgia Fitkid Project. *Int J Pediatr Obes IJPO* 2008; **3 Suppl 1**: 3–9.
- 114 Saksvig BI, Gittelsohn J, Harris SB, Hanley AJ, Valente TW, Zinman B. A pilot school-based healthy eating and physical activity intervention improves diet, food knowledge, and self-efficacy for native Canadian children. *J Nutr* 2005; **135**: 2392–2398.
- 115 Story M, Sherwood NE, Himes JH *et al*. An after-school obesity prevention program for African-American girls: the Minnesota GEMS pilot study. *Ethn Dis* 2003; **13**: S54–64.
- 116 Singh AS, Chin APMJ, Brug J, van Mechelen W. Short-term effects of school-based weight gain prevention among adolescents. *Arch Pediatr Adolesc Med* 2007; **161**: 565–571.
- 117 Singh AS, Chin APMJ, Brug J, van Mechelen W. Dutch obesity intervention in teenagers: effectiveness of a school-based program on body composition and behavior. *Arch Pediatr Adolesc Med* 2009; **163**: 309–317.
- 118 Ask AS, Hernes S, Aarek I, Vik F, Brodahl C, Haugen M. Serving of free school lunch to secondary-school pupils - a pilot study with health implications. *Public Heal Nutr* 2010; **13**: 238–244.
- 119 Ebbeling CB, Feldman HA, Osganian SK, Chomitz VR, Ellenbogen SJ, Ludwig DS. Effects of decreasing sugar-sweetened beverage consumption on body weight in adolescents: a randomized, controlled pilot study. *Pediatrics* 2006; **117**: 673–680.
- 120 Robinson TN. Reducing children's television viewing to prevent obesity: a randomized controlled trial. *JAMA* 1999; **282**: 1561–7.
- 121 Foster GD, Sherman S, Borradaile KE *et al*. A policy-based school intervention to prevent overweight and obesity. *Pediatrics* 2008; **121**: e794–802.
- 122 Foster GD, Linder B, Baranowski T *et al*. A school-based intervention for diabetes risk reduction. *N Engl J Med* 2010; **363**: 443–453.
- 123 Marcus C, Nyberg G, Nordenfelt A, Karpmyr M, Kowalski J, Ekelund U. A 4-year, cluster-randomized, controlled childhood obesity prevention study: STOPP. *Int J Obes* 2009; **33**: 408–417.
- 124 Sallis JF, McKenzie TL, Conway TL *et al*. Environmental interventions for eating and physical activity: a randomized controlled trial in middle schools. *Am J Prev Med* 2003; **24**: 209–217.
- 125 Sanigorski AM, Bell AC, Kremer PJ, Cuttler R, Swinburn BA. Reducing unhealthy weight gain in children through community capacity-building: results of a quasi-experimental intervention program, Be Active Eat Well. *Int J Obes* 2008; **32**: 1060–1067.
- 126 Singhal N, Misra A, Shah P, Gulati S. Effects of controlled school-based multi-component model of nutrition and lifestyle interventions on behavior modification, anthropometry and metabolic risk profile of urban Asian Indian adolescents in North India. *Eur J Clin Nutr* 2010; **64**: 364–373.

- 127 Coleman KJ, Tiller CL, Sanchez J *et al.* Prevention of the epidemic increase in child risk of overweight in low-income schools: the El Paso coordinated approach to child health. *Arch Pediatr Adolesc Med* 2005; **159**: 217–224.
- 128 Haerens L, Deforche B, Maes L, Stevens V, Cardon G, De Bourdeaudhuij I. Body mass effects of a physical activity and healthy food intervention in middle schools. *Obes (Silver Spring)* 2006; **14**: 847–854.
- 129 Kain J, Uauy R, Albala, Vio F, Cerda R, Leyton B. School-based obesity prevention in Chilean primary school children: methodology and evaluation of a controlled study. *Int J Obes Relat Metab Disord J Int Assoc Study Obes* 2004; **28**: 483–493.
- 130 Taylor RW, McAuley KA, Barbezat W, Strong A, Williams SM, Mann JI. APPLE Project: 2-y findings of a community-based obesity prevention program in primary school age children. *Am J Clin Nutr* 2007; **86**: 735–742.
- 131 Taylor RW, McAuley KA, Barbezat W *et al.* Two-year follow-up of an obesity prevention initiative in children: the APPLE project. *Am J Clin Nutr* 2008; **88**: 1371–1377.
- 132 Barbeau P, Johnson MH, Howe CA *et al.* Ten months of exercise improves general and visceral adiposity, bone, and fitness in black girls. *Obes (Silver Spring)* 2007; **15**: 2077–2085.
- 133 Jansen W, Borsboom G, Meima A *et al.* Effectiveness of a primary school-based intervention to reduce overweight. *Int J Pediatr Obes IJPO* 2011; **6**: e70–7.
- 134 Martinez Vizcaino V, Salcedo Aguilar F, Franquelo Gutierrez R *et al.* Assessment of an after-school physical activity program to prevent obesity among 9- to 10-year-old children: a cluster randomized trial. *Int J Obes* 2008; **32**: 12–22.
- 135 Yin Z, Gutin B, Johnson MH *et al.* An environmental approach to obesity prevention in children: Medical College of Georgia FitKid Project year 1 results. *Obes Res* 2005; **13**: 2153–2161.
- 136 Wang LY, Gutin B, Barbeau P *et al.* Cost-effectiveness of a school-based obesity prevention program. *J Sch Heal* 2008; **78**: 619–624.
- 137 Angelopoulos PD, Milionis HJ, Grammatikaki E, Moschonis G, Manios Y. Changes in BMI and blood pressure after a school based intervention: the CHILDREN study. *Eur J Public Health* 2009; **19**: 319–325.
- 138 Simon C, Schweitzer B, Oujaa M *et al.* Successful overweight prevention in adolescents by increasing physical activity: a 4-year randomized controlled intervention. *Int J Obes* 2008; **32**: 1489–1498.
- 139 Millar L, Kremer P, de Silva-Sanigorski A *et al.* Reduction in overweight and obesity from a 3-year community-based intervention in Australia: the ‘It’s Your Move!’ project. *Obes Rev* 2011; **12 Suppl 2**: 20–28.
- 140 Kremer P, Waqa G, Vanualailai N *et al.* Reducing unhealthy weight gain in Fijian adolescents: results of the Healthy Youth Healthy Communities study. *Obes Rev* 2011; **12 Suppl 2**: 29–40.
- 141 Muckelbauer R, Libuda L, Clausen K, Toschke AM, Reinehr T, Kersting M. Promotion and provision of drinking water in schools for overweight prevention: randomized, controlled cluster trial. *Pediatrics* 2009; **123**: e661–7.

- 142 Ask AS, Hernes S, Aarek I, Johannessen G, Haugen M. Changes in dietary pattern in 15 year old adolescents following a 4 month dietary intervention with school breakfast--a pilot study. *Nutr J* 2006; **5**: 33.
- 143 de Ruyter JC, Olthof MR, Seidell JC, Katan MB. A trial of sugar-free or sugar-sweetened beverages and body weight in children. *N Engl J Med* 2012; **367**: 1397–1406.
- 144 Story M, Neumark-Sztainer D, Ireland M, Evans T. Adolescent health and nutrition: a survey of perceived knowledge and skill competencies and training interests among dietitians working with youth. *J Am Diet Assoc* 2000; **100**: 362–4.
- 145 Story MT, Neumark-Sztainer D, French SA. Individual and Environmental Influences on Adolescent Eating Behaviors. *J Am Diet Assoc* 2002; **102**: S40–S51.
- 146 Campbell K, Waters E, O'Meara S, Summerbell C. Interventions for preventing obesity in childhood. A systematic review. *Obes Rev* 2001; **2**: 149–157.
- 147 Todd MK, Reis-Bergan MJ, Sidman CL *et al*. Effect of a family-based intervention on electronic media use and body composition among boys aged 8–11 years: a pilot study. *J Child Heal Care* 2008; **12**: 344–358.
- 148 Ni Mhurchu C, Roberts V, Maddison R *et al*. Effect of electronic time monitors on children's television watching: pilot trial of a home-based intervention. *Prev Med* 2009; **49**: 413–417.
- 149 French SA, Gerlach AF, Mitchell NR, Hannan PJ, Welsh EM. Household obesity prevention: Take Action--a group-randomized trial. *Obesity* 2011; **19**: 2082–2088.
- 150 Wang Y, Cai L, Wu Y *et al*. What childhood obesity prevention programmes work? A systematic review and meta-analysis. *Obes Rev* 2015. doi:10.1111/obr.12277.
- 151 Faggion Jr. CM. Critical appraisal of AMSTAR: challenges, limitations, and potential solutions from the perspective of an assessor. *BMC Med Res Methodol* 2015; **15**: 63.
- 152 Gonzalez-Suarez C, Worley A, Grimmer-Somers K, Dones V. School-based interventions on childhood obesity: a meta-analysis. *Am J Prev Med* 2009; **37**: 418–427.
- 153 Woodman J, Thomas J, Dickson K. How explicable are differences between reviews that appear to address a similar research question? A review of reviews of physical activity interventions. *Syst Rev* 2012; **1**: 37.
- 154 Pettman T, Magarey A, Mastersson N, Wilson A, Dollman J. Improving weight status in childhood: results from the eat well be active community programs. *Int J Public Health* 2014; **59**: 43–50.
- 155 Chang DI, Gertel-Rosenberg A, Drayton VL, Schmidt S, Angalet GB. A statewide strategy to battle child obesity in Delaware. *Health Aff* 2010; **29**: 481–490.