

UNIVERSITY OF BIRMINGHAM

University of Birmingham
Research at Birmingham

Excision of metastatic breast cancer from sternum and reconstruction in two patients with solitary metastatic spread

Mohamed, Saifullah; Mazhar, Khurum; Osman, Ahmed; Patel, Akshay; Srinivasan, Lakshmi; Ghosh, Shilajit

DOI:

[10.1093/jscr/rjaa272](https://doi.org/10.1093/jscr/rjaa272)

License:

Creative Commons: Attribution-NonCommercial (CC BY-NC)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Mohamed, S, Mazhar, K, Osman, A, Patel, A, Srinivasan, L & Ghosh, S 2020, 'Excision of metastatic breast cancer from sternum and reconstruction in two patients with solitary metastatic spread', *Journal of Surgical Case Reports*, vol. 2020, no. 8, rjaa272. <https://doi.org/10.1093/jscr/rjaa272>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

CASE REPORT

Excision of metastatic breast cancer from sternum and reconstruction in two patients with solitary metastatic spread

Saifullah Mohamed¹, Khurum Mazhar¹, Ahmed Osman¹, Akshay Patel^{1,2,*}, Lakshmi Srinivasan¹, and Shilajit Ghosh¹

¹Department of Cardiothoracic Surgery, Royal Stoke University Hospital, Stoke On Trent, UK and ²Institute of Immunology and Immunotherapy, University of Birmingham, Edgbaston, Birmingham, UK

*Correspondence address. Institute of Immunology and Immunotherapy, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK.
Tel: 0121 4143344; E-mail: ajp.788@gmail.com

Abstract

Metastatic breast carcinoma is a relatively common clinical entity. However, the prognosis of oligometastatic and polygometastatic disease differs considerably pertaining to five-year survival. Metastatic breast carcinoma to the sternum has been described as early as 1988. We describe two cases in our institution who presented with solitary sternal metastases with a previous history of treated breast cancer. In both cases, there had been a history of previous left breast cancer treated aggressively with surgical resection and adjuvant oncological therapy and maintenance hormonal therapy. Partial sternectomy or total sternectomy for solitary metastatic sternal deposits is well established with relatively low morbidity and mortality and improvement in quality of life and possible improvement in long-term survival. Furthermore, reconstructive options are inherently dependent on extent of resection performed. These techniques can incorporate the use of sternal plates in order to approximate defects and reinforce the sternum in the setting of partial sternectomy.

INTRODUCTION

Metastatic breast carcinoma is a relatively common clinical entity with 20–30% of patients with early-stage disease developing metastases and 3–6% with *de novo* Stage IV disease at time of initial diagnosis [1]. However, the prognosis of oligometastatic and polygometastatic disease differs considerably pertaining to five-year survival. Furthermore, the survival and procedure-related morbidity and mortality data for any surgical resection are largely based on retrospective data analysis of multiple cases performed at a single-centre or multicentre studies rather

than from any randomized controlled trial. Therefore, there is considerable debate and contention about the efficacy of metastasectomy and its widespread application in clinical practise. Utilizing equivalent strategy and principles of pulmonary metastasectomy, any surgery for metastatic disease is founded on the core principles outlined by Thomford *et al.* [2] in 1965. These include control or eradication of primary site of cancer and exclusion of local recurrence, relative disease-free interval with a longer disease-free interval indicating a better prognosis, surgically resectable disease and availability of adjuvant oncological therapy. There have been discussions regarding the cytoreductive aspect of metastasectomy, but this is

Received: May 13, 2020. Accepted: June 29, 2020

Published by Oxford University Press and JSCR Publishing Ltd. All rights reserved. © The Author(s) 2020.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

well established due to the concept of micrometastatic disease in cases where complete resection is not possible and it is generally accepted that incomplete metastasectomy probably offers no significant advantages but potentially carries a considerable morbidity and mortality risk. Metastatic breast carcinoma to the sternum has been described as early as 1988 [3]. Furthermore, one of the earliest case series in 1988 described a series of nine patients with solitary sternal metastasis who underwent partial or total sternectomy. In this study, patients with solitary metastatic sternal deposits without evidence of other regional spread favoured considerably compared with those with sternal deposit oligometastatic disease that had spread to other sites. The authors of the study concluded that resection of solitary sternal metastasis should be advocated and that in addition to improving quality of life, survival may also be improved [3]. Similarly, the efficacy of partial or total sternectomy in patients with solitary metastatic sternal spread has been demonstrated subsequently [4]. There have been numerous cases discussing sternal resection and reconstruction often with the use of methyl methacrylate and acrylic cemented prostheses [5–6], Marlex mesh and muscle flaps to cover defects [4]. We describe two cases in our institution who presented with solitary sternal metastases with a previous history of treated breast cancer.

CASE REPORT

The first case at our institution was a 39-year-old woman who had a history of left breast carcinoma treated with left mastectomy, axillary node clearance and immediate reconstruction under the care of the breast team 3 years prior to presentation. She had no significant co-morbidities. Her past surgical history consisted of the aforementioned oncological operation, appendectomy and cholecystectomy. She had completed adjuvant oncological therapy in the form of chemoradiotherapy and maintenance hormonal therapy. Her post-operative histology was consistent with a fully excised left breast ductal carcinoma and spread to three out of nine lymph nodes. Due to positive oestrogen receptor-2 status, she had been placed on hormonal therapy. She re-presented with chest wall and sternal pain and subsequent computer tomography (CT) imaging demonstrated a suspicious looking lytic lesion in the mid-sternum (Fig. 1). She underwent a biopsy of this lesion, which confirmed the diagnosis. She had been taking opioid medication for the pain. She was seen in clinic and counselled for partial sternectomy and reconstruction. She underwent partial sternectomy and underwent reconstruction with vertical and horizontal Synthes® plates. Her post-operative recovery was unremarkable. Post-operative histology was consistent with metastatic breast carcinoma completely excised. The recovery was uncomplicated and the patient was seen in clinic with a satisfactory chest radiograph demonstrating the sternal Synthes® plates (Fig. 2).

The second case at our institution was a 52-year-old woman with a history of left breast carcinoma treated for which she had undergone left wide local excision and sentinel node biopsy. Due to uncertainty regarding margins, she required a re-resection of her left breast tissue. She completed adjuvant radiotherapy and received maintenance hormonal therapy. She presented 8 months following her radiotherapy with sternal pain and positron emission tomography-computed tomography (PET-CT) imaging demonstrated localized uptake within the sternum (Fig. 3). Biopsy demonstrated proven recurrence of metastatic breast cancer. She was seen in clinic and counselled for




Figure 1: Lytic mid-sternal lesion on CT (arrow).



Figure 2: Follow-up chest radiograph demonstrating the fixation plates.

partial sternectomy and reconstruction. She underwent partial sternectomy and underwent reconstruction with Synthes® plates. Her post-operative recovery was unremarkable. Post-operative histology was consistent with metastatic breast carcinoma, which was completely excised. The patient made an uncomplicated recovery and was seen in the outpatient clinic with a satisfactory post-operative chest radiograph (Fig. 4).

DISCUSSION

Partial or total sternectomy for solitary metastatic sternal deposits is well established with relatively low morbidity, mortality and improvement in quality of life and long-term survival [7]. Interestingly, the aspect of chest wall or sternal pain is one aspect that considerably improves following sternal metastasectomy. Chest wall resection and reconstruction for metastatic breast cancer has been demonstrated to be effective in certain cohort of breast cancer patients with low mortality and morbidity [8]. In addition to surgery, cryoablation has also been purported as an effective alternative to radical surgical resection to achieve loco-regional control [9]. Upon review of the relevant literature, the application of surgery seems universal in that localized disease and recurrence in the sternum should be resected [4]. However, the type of resection partial or total sternectomy is dependent on the size of the tumour and the principle of achieving complete excision. Reconstructive options




Figure 3: CT-PET demonstrating increased metabolic activity within the sternal lesion.



Figure 4: Follow-up chest radiograph of second patient.

have been performed via a multitude of methods and involve using cemented prostheses, Marlex mesh and muscle flaps [4] and in some cases of total sternectomy, customized 3D-printed titanium prostheses [10]. Our experience involves partial sternectomy for localized sternal recurrence and reconstruction of the defect using Synthes® plates to reinforce and stabilize the sternum. In summary, metastasectomy, either as partial or total sternectomy, should be performed for localized sternal recurrence for metastatic breast cancer as long as the core principles of metastasectomy are observed. Furthermore, reconstructive options are inherently dependent on extent of resection performed. These techniques can incorporate the use of sternal plates in order to approximate defects and reinforce the sternum in the setting of partial sternectomy.

CONFLICT OF INTEREST STATEMENT

None declared.

FUNDING

None.

REFERENCES

1. Tosello G, Torloni MR, Mota BS, Neeman T, Riera R. Breast surgery for metastatic breast cancer. *Cochrane Database Syst Rev* 2018;3.
2. Thomford NR, Woolner LB, Clagett OT. The surgical treatment of metastatic tumors in the lungs. *J Thorac Cardiovasc Surg* 1965;49:357–63.
3. Noguchi S, Miyauchi K, Nishizawa Y, Imaoka S, Koyama H, Iwanaga T. Results of surgical treatment for sternal metastasis of breast cancer. *Cancer* 1988;62:1397–401.
4. Incarbone M, Nava M, Lequaglie C, Ravasi G, Pastorino U. Sternal resection for primary or secondary tumors. *J Thorac Cardiovasc Surg* 1997;114:93–9.
5. Sanna S, Brandolini J, Pardolesi A, Argnani D, Mengozzi M, Dell'Amore A, et al. Materials and techniques in chest wall reconstruction: a review. *J Vis Surg* 2017;3:95.
6. Demetrian AD, Olteanu M, Mîndrîlă I, Macovei A, Râmboiu DS, Enache ME, et al. Long disease-free survival following total sternal resection and reconstruction of the sternum with acrylic cement for unique massive sternal metastasis after operated breast cancer. *Rom J Morphol Embryol* 2018;59:1225–32.
7. Bongiolatti S, Voltolini L, Borgianni S, Borrelli R, Innocenti M, Menichini G, et al. Short and long-term results of sternectomy for sternal tumours. *J Thorac Dis* 2017;9:4336–46.
8. Santillan AA, Kiluk JV, Cox JM, Meade TL, Allred N, Ramos D, et al. Outcomes of locoregional recurrence after surgical chest wall resection and reconstruction for breast cancer. *Ann Surg Oncol* 2008;15:1322–9.
9. Hegg RM, Kurup AN, Schmit GD, Weisbrod AJ, Atwell TD, Olivier KR, et al. Cryoablation of sternal metastases for pain palliation and local tumor control. *J Vasc Interv Radiol* 2014;25:1665–70.
10. Tran MD, Varzaly JA, Chan JCY, Caplash Y, Worthington MG. Novel sternal reconstruction with custom three-dimensional-printed titanium PoreStar prosthesis. *Innovations* 2018;13:309–11.