

# Gold(I)-catalyzed synthesis of 3-sulfenyl pyrroles and indoles by a regioselective annulation of alkynyl thioethers

Simm, Peter; Sekar, Prakash; Richardson, Jeffery; Davies, Paul

DOI:

[10.1021/acscatal.1c01457](https://doi.org/10.1021/acscatal.1c01457)

License:

Creative Commons: Attribution (CC BY)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Simm, P, Sekar, P, Richardson, J & Davies, P 2021, 'Gold(I)-catalyzed synthesis of 3-sulfenyl pyrroles and indoles by a regioselective annulation of alkynyl thioethers', *ACS Catalysis*, vol. 11, no. 11, pp. 6357–6362. <https://doi.org/10.1021/acscatal.1c01457>

[Link to publication on Research at Birmingham portal](#)

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

## Gold(I)-Catalyzed Synthesis of 3-Sulfenyl Pyrroles and Indoles by a Regioselective Annulation of Alkynyl Thioethers

Peter E. Simm, Prakash Sekar, Jeffery Richardson, and Paul W. Davies\*

Cite This: *ACS Catal.* 2021, 11, 6357–6362

Read Online

ACCESS |



Metrics &amp; More




Article Recommendations



Supporting Information

**ABSTRACT:** The combination of nucleophilic nitrenoids and  $\pi$ -acid catalysis has emerged as a powerful tool in heterocycle synthesis. Accessing more varied heterocycle-substitution patterns by maintaining the same reaction pathways across different alkynes remains a challenge. Here we show that Au(I) catalysis of isoxazole-based nitrenoids with alkynyl thioethers provides controlled access to (3 + 2) annulation by a regioselective addition  $\beta$  to the sulfenyl group. The reaction with isoxazole-containing nitrenoids delivers sulfenylated pyrroles and indoles as single regioisomers bearing useful functional groups and structural variety.

**KEYWORDS:** gold, pyrroles, nitrenoid, annulation, thioether, indole



N-nucleophilic nitrenoids have proved to be versatile surrogates for nitrene-containing 1,3-dipoles, providing modular and expedient access to a variety of densely functionalized N-heterocycles by a formal (3 + 2) cycloaddition to gold-activated alkynes.<sup>1</sup> Most of these annulations were first realized with ynamides,<sup>2</sup> and to date there are only a few specific instances where those same transformations have been achieved with other types of alkynes.<sup>2j,3</sup> As the alkyne substitution pattern is directly translated into the product, retaining the same transformation across different alkyne types would greatly expand the potential of these convergent annulation methods for complex molecule synthesis. A challenge lies in finding alkynes that are sufficiently reactive and do not change how the reaction pathway evolves while providing useful substitution patterns.

The use of isoxazole-based N-nucleophilic nitrenoids is illustrative: The groups of Ye<sup>2f</sup> and Hashmi<sup>2j</sup> demonstrated the potential of using isoxazoles and [2,1]benzoxazoles, respectively, to assemble pyrroles and indoles through a formal (3 + 2) cycloaddition across C-aryl ynamides (Figure 1b). Several groups have subsequently established that a tremendous variety of heterocyclic structures can be formed when these nitrenoids are combined with differently substituted alkynes.<sup>4</sup> Because the alkyne structure affects the nature of the intermediate  $\alpha$ -imino gold carbene **B** and the pathways through which it can evolve (Figure 1a), even relatively small changes can divert the pathway away from formal (3 + 2) cycloaddition (Figure 1c).

Few reports have addressed gold-catalyzed intermolecular reactions with alkynyl thioethers, and the majority have


reported nucleophilic addition  $\alpha$  to the sulfur,<sup>5</sup> invoking a dominant gold ketenethionium character. However, a recent study from our group using N-acylpyridinium aminides showed that, in the presence of a Au(III) precatalyst, the outcomes matched selective attack at the  $\beta$ -position of alkynyl thioethers.<sup>3b</sup> We reasoned that if this latter route was generally accessible from other types of nitrenoids, especially those accessed using Au(I) catalysis, then a more generalized approach for (3 + 2)-type annulations with nitrenoids could be realized. Replicating this reactivity would ensure that the auroated carbon is substituted by the sulfenyl group throughout the reaction manifold, potentially facilitating a more consistent reactivity profile that tolerates modifications elsewhere. Sulfur-substituted heterocycles are desirable, not least in medicinal chemistry.<sup>6,7</sup> The increasing number of C–S functionalization methods renders them potentially useful substrates for further elaboration.<sup>8</sup> The prospective utility of sulfenylated pyrrole and indole products combined with the diversity of reaction pathways that can be accessed from (benz)isoxazoles identified them as ideal systems for probing the wider utility of alkynyl thioethers in nitrenoid-based annulations.

During our studies Ye and co-workers reported the Zn(II)-catalyzed reaction of alkynyl thioethers with isoxazoles, where

Received: March 30, 2021

Revised: May 3, 2021






**Figure 1.** Divergent outcomes for the reaction of alkynes and isoxazoles under  $\pi$ -acid catalysis.

selective  $\alpha$ -addition led to sulfur-substituted  $\beta$ -keto enamides with 1,2-S migration (Figure 1d).<sup>9</sup> Here we show that the Au(I)-catalyzed reaction of alkynyl thioethers with isoxazoles and [2,1]benzoxazoles proceeds regioselectively by  $\beta$ -addition to access the formal (3 + 2) cycloaddition pathway.

We found that using alkynyl thioether **1a** and 3,5-dimethylisoxazole **2a** gave rise to regioselective formation of the 3-sulfonylated pyrrole **3a** under gold catalysis. The finalized conditions employ a 1:2 stoichiometry of **1a** to **2a** in 1,1,1-trifluorotoluene at 80 °C in the presence of [IPrAu(CH<sub>3</sub>CN)]-SbF<sub>6</sub> in a capped vial and no measures taken to exclude air and moisture (Scheme 1; see the Supporting Information for an optimization survey). No reaction was seen in the absence of catalyst or on replacing it with a strong Brønsted acid, while gold(I) and gold(III) catalysts with other ligand combinations were less effective. The noncoordinating counterions tetrakis-[3,5-bis(trifluoromethyl)phenyl]borate and hexafluoroantimonate were optimal in comparison to weakly coordinating alternatives. Only a small reduction in yield is seen at lower temperatures or with a lower stoichiometry of **2a**.

The effects of different S-substituents on the annulation were tested (Scheme 1). While an S-phenyl substituent stalled the process, presumably due to its rigid steric bulk, S-benzylated systems were reactive (**3c,d**). 1,5-Enynyl thioether **1e** gave solely the pyrrole (**3e**), despite the potential for

## Scheme 1. Pyrrole Formation from Alkynyl Thioethers: S-Substituent




intramolecular cyclopropanation at an intermediate  $\alpha$ -imino gold carbene.<sup>4k</sup> The S-alkynyl thioester **1f** degraded under the reaction conditions, but the 2-(ethynylthio)acetate derivative **1g** afforded pyrrole **3g**.

The wider scope of the annulation was then explored (Scheme 2). The reaction was largely invariant to steric bulk at the C-terminus of the alkynyl thioether, allowing ortho and diortho substitution (**3h–j**). A single-crystal X-ray diffraction analysis confirmed the structure of **3j**. Electron-donating substituents on the aryl group are beneficial but not required (**3h–q**). An aryl fluoride and an aniline derivative worked well, as did heteroaromatic groups such as indole and thiophene (**3m–p**). Alkyl-substituted alkynyl thioethers reacted sluggishly but did give the desired 3-sulfonylated pyrroles **3r,s** in serviceable yields.

An unsymmetrical acetylene disulfide reacted to give the 2,3-bis-sulfonyl pyrroles as a mixture of regioisomers **3t,t'**. The regioselective preference correlates to the relative reactivity seen on changing the sulfonyl group in alkynyl thioethers (cf. Scheme 1, **1a,b**), the major product arising from C–N bond formation  $\beta$  to the smaller sulfonyl group, although the regioisomer **3t'** was formed despite the recalcitrance of **1b**. Neither triisopropylsilyl nor terminal alkynyl thioethers gave productive reactions (**1u,v**, see Supporting Information).

A variety of differently substituted isoxazoles proved to be compatible with the reaction. 3-Sulfonylated pyrroles can be prepared with alkyl, aryl, and vinyl groups at the 5-position (**3u–3af**), and with aldehydes (**3x**), amides (**3y,z**), and aryl- or alkyl-substituted carbonyl groups at the 4-position (**3u–w,ac–af**). No reaction was seen with the electronically deactivated 3-(perfluorophenyl)-5-phenylisoxazole and 3-phenyl-5-(trifluoromethyl)isoxazole (not shown). The use of isoxazol-5-amines had not previously been reported under gold catalysis,<sup>10</sup> while Liu reported non-pyrrole-forming pathways when unsubstituted isoxazoles were reacted with propiolate derivatives.<sup>4b,c</sup> The formation of pyrroles (**3x–z**) from both of these systems illustrates the consistent reaction outcomes obtained in gold-catalyzed reactions of alkynyl thioethers with nitrenoids. The reaction tolerates a range of functionalities, including 1° and 3° aromatic amines (**3n,y,z**), a free hydroxyl group (**3ac**), an ester (**3ad**), and an aryl halide (**3af**).

The wider generality of alkynyl thioethers was established using anthranil **4** as the nitrenoid to deliver 3-sulfonyl-7-acyl indole motifs **5a–f** by the formal (3 + 2) cycloaddition pathway. 3-Sulfonyl indoles have shown some medicinal potential with activity against HIV<sup>11</sup> and inhibition of tubulin polymerization.<sup>12</sup> As anthranil and the products coeluted, the alkynyl thioether was used in excess (see the Supporting


Scheme 2. Au(I)-Catalyzed Synthesis of Pyrroles and Indoles from Alkynyl Thioethers<sup>a</sup>

<sup>a</sup>Reactions unless stated otherwise were run in capped vials with no precautions against air or moisture with alkynyl thioethers (1.0 equiv), 2a–l (2.0 equiv), [IPrAu(CH<sub>3</sub>CN)]SbF<sub>6</sub> (5 mol %), and PhCF<sub>3</sub> (0.1 M) at 80 °C. The crystal structure of 3j is shown with ellipsoids drawn at the 50% probability level. <sup>b</sup>Reaction at 50 °C; 53% of 3n at 80 °C. <sup>c</sup><sup>1</sup>H NMR spectroscopic analysis of the reaction mixture elucidated a 43% yield of 3t and 19% of the putative 3t'. The regioisomerism of 3t was confirmed by <sup>1</sup>H NOESY. <sup>d</sup>4 equiv of isoxazole. <sup>e</sup>0.4 mmol of 1a with 0.2 mmol of 2i. <sup>f</sup>4 was used as the limiting reagent to 1.

Information). A competition experiment showed that anthranil 4 is less reactive than isoxazole 2a (see the Supporting Information).

Reactions with 3,4,5-trisubstituted isoxazoles 6a,b were investigated (Scheme 3). Reactions with isoxazoles 6a,b saw formation of the N-acylated pyrrole 7 and deacylative annulation product 9, respectively, as has been preceded in ynamide reactions.<sup>13</sup> Reactivity unique to alkynyl thioethers was also elucidated with the formation of desulfenylated 3-acylated pyrroles 8a,b.


The practicality of the annulation protocol is demonstrated by its ready upscaling, which proceeded smoothly under noninert conditions to give 3a on a gram scale (Scheme 4). Selective oxidations of the resulting pyrrole 3a into its sulfonyl and sulfinyl forms (10 and 11) were effective.<sup>14</sup> Treating 11 with triflic anhydride in the presence of base led to protodesulfenylation and the formation of trisubstituted pyrrole 12. The direct use of triflic acid<sup>15</sup> gave a dirtier reaction and only traces of 12. However, Brønsted acid mediated desulfenylation of the sulfide 3d also gave 12. Such acid-

Scheme 3. Au(I)-Catalyzed Reactions of Alkynyl Thioethers with 3,4,5-Trisubstituted Isoxazoles<sup>a</sup>

<sup>a</sup>All reactions were run in capped vials with no precautions against air or moisture, using 1a (1.0 equiv), 6a or 6b (2.0 equiv), [IPrAu(CH<sub>3</sub>CN)]SbF<sub>6</sub> (5 mol %), and PhCF<sub>3</sub> (0.1 M), at 80 °C. The crystal structure of 8a is shown with ellipsoids drawn at the 50% probability level.

mediated desulfuration reactions permit chemoselective reduction of the sulfonyl group in the presence of the ketone

Scheme 4. Scale-up Annulation and Selective Oxidation of a 3-Sulfenyl Pyrrole




and enable the nitrenoid chemistry to be used to access non-heteroatom-substituted pyrroles. Initial attempts to achieve metal-catalyzed C–S activation for cross-coupling have so far been unsuccessful with these elaborated pyrroles.

The isotopically labeled substrate  $^{13}\text{C}$ -II was prepared in order to determine whether there was any skeletal rearrangement of the alkynyl thioether backbone during or after annulation. 1,2-Sulfenyl migrations have been reported in  $\pi$ -acid mediated reactions that invoke carbenoid character,<sup>5a–d</sup> while methylthio-substituted pyrroles have been shown to undergo Brønsted acid mediated isomerization.<sup>16</sup> The single pyrrole isotopomer  $^{13}\text{C}$ -3I was formed under the standard conditions. A combination of HMBC and NOESY experiments was used to confirm the regiochemical outcome on 3I/ $^{13}\text{C}$ -3I and that the connectivity of the alkynyl thioether was maintained (Scheme 5). A small amount of the enriched  $\beta$ -keto enamide  $^{13}\text{C}$ -13 was also isolated. This outcome is in concordance with the pathway reported by Ye under Zn(II) catalysis<sup>9</sup> and provides labeling evidence for the 1,2-S-migration step.

A general mechanism is proposed on the basis of these observations and literature precedent (Scheme 5). The  $^{13}\text{C}$ -labeling experiment is consistent with slippage to either end of the triple bond being energetically accessible:  $\alpha$ -addition, matching a gold ketenethionium type activation  $\text{A}^1$  invoked in the majority of Au(I)-catalyzed reactions of alkynyl thioethers, is the minor pathway and is not viable for the formation of pyrrole. This pathway leads to  $\beta$ -ketoenamide D. The  $\beta$ -addition pathway is more productive ( $\text{A} \rightarrow \text{E}$ ), and we postulate that a stabilizing S–Au interaction<sup>17</sup> promotes irreversible gold carbene formation. This pathway is aided by, but not dependent upon, the presence of more electron rich alkyne substituents (Scheme 2), which would further distort the  $\pi$ -complex toward a nascent Au–S interaction  $\text{A}^2$ .<sup>18</sup> The resulting  $\alpha$ -imino  $\alpha'$ -sulfenyl gold carbene F undergoes cyclization and aromatization to pyrrole I. When C-4-substituted isoxazoles are used, in addition to established deacylative mechanisms ( $\text{H} \rightarrow \text{J/K}$ ),<sup>2h,3c</sup> an unprecedented desulfenylation can follow 1,5-acyl migration,<sup>4b</sup> leading to all-carbon substituted pyrroles ( $\text{H} \rightarrow \text{M}$ ).

In conclusion, alkynyl thioethers react with isoxazoles and anthranils under gold catalysis and provide selective access to the formal (3 + 2) cycloaddition pathway. This pathway is maintained across reactants with broad structural and func-

Scheme 5.  $^{13}\text{C}$ -Labeling Experiment and Outline Mechanism

tional group changes. As a result, these readily accessed substrates can be used to deliver convergent and modular access to sulfenylated pyrroles and indoles. The practical and straightforward protocol has been demonstrated on a multi-millimole scale. Initial investigations into the annulation with trisubstituted isoxazoles also reveal reactivity unique to alkynyl thioethers. One-step access to all-carbon-substituted pyrroles establishes the intriguing potential of using a sulfenyl moiety as a traceless directing and alkyne-activating group for heterocycle synthesis. The observed outcomes and an isotopic labeling study match a  $\beta$ -selective addition to the alkynyl thioether. In contrast to  $\alpha$ -selective approaches, such as from ubiquitously employed ynamides, the gold carbene character is developed adjacent to the heteroatom group and not the other, variable, alkyne substituent. If gold-catalyzed reactions of alkynyl thioethers and nitrenoids consistently proceed through the putative  $\alpha$ -imino  $\alpha'$ -sulfenyl gold carbene, then this can provide the basis of a unifying (3 + 2) cycloaddition strategy for heterocycle synthesis.

## ■ ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acscatal.1c01457>.

Experimental details, compound characterization data, crystallographic data, and NMR spectra (PDF)  
Crystallographic data for 3j (CIF)

Crystallographic data for **8a** (CIF)

## AUTHOR INFORMATION

### Corresponding Author

Paul W. Davies – School of Chemistry, University of Birmingham, Birmingham B15 2TT, U.K.; [orcid.org/0000-0002-0340-2414](https://orcid.org/0000-0002-0340-2414); Email: [p.w.davies@bham.ac.uk](mailto:p.w.davies@bham.ac.uk)

### Authors

Peter E. Simm – School of Chemistry, University of Birmingham, Birmingham B15 2TT, U.K.  
Prakash Sekar – School of Chemistry, University of Birmingham, Birmingham B15 2TT, U.K.  
Jeffery Richardson – Lilly U.K., Windlesham GU20 6PH, U.K.

Complete contact information is available at:  
<https://pubs.acs.org/10.1021/acscatal.1c01457>

### Author Contributions

All authors contributed to the preparation of the manuscript and have given approval to the final version of the manuscript.

### Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENTS

We thank the EPSRC and Eli Lilly for funding (NPIF studentship EP/RS12436/1, P.E.S.). This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 839037 (P.S., SEAROX). P.W.D. is grateful to the Royal Society and Leverhulme Trust for the award of a Senior Research Fellowship (SRF\R1\191033). The authors gratefully acknowledge support from Dr. Louise Male (XRD) and the Centre for Chemical and Materials Analysis in the School of Chemistry (UoB).

## REFERENCES

(1) (a) Davies, P. W.; Garzón, M. Nucleophilic Nitrenoids Through  $\pi$ -Acid Catalysis: Providing a Common Basis for Rapid Access into Diverse Nitrogen Heterocycles. *Asian J. Org. Chem.* **2015**, *4*, 694–708. (b) Aguilar, E.; Santamaría, J. Gold-catalyzed heterocyclic syntheses through  $\alpha$ -imino gold carbene complexes as intermediates. *Org. Chem. Front.* **2019**, *6*, 1513–1540. (c) Ye, L.-W.; Zhu, X.-Q.; Sahani, R. L.; Xu, Y.; Qian, P.-C.; Liu, R.-S. Nitrene Transfer and Carbene Transfer in Gold Catalysis. *Chem. Rev.* **2020**, DOI: [10.1021/acs.chemrev.0c00348](https://doi.org/10.1021/acs.chemrev.0c00348). (2) (a) Davies, P. W.; Cremonesi, A.; Dumitrescu, L. Intermolecular and Selective Synthesis of 2,4,5-Trisubstituted Oxazoles by a Gold-Catalyzed Formal [3 + 2] Cycloaddition. *Angew. Chem., Int. Ed.* **2011**, *50*, 8931–8935. (b) Garzón, M.; Davies, P. W. A Direct Route into Fused Imidazo-diazines and Imidazo-pyridines Using Nucleophilic Nitrenoids in a Gold-Catalyzed Formal [3 + 2]-Dipolar Cycloaddition. *Org. Lett.* **2014**, *16*, 4850–4853. (c) Pawar, S. K.; Sahani, R. L.; Liu, R.-S. Diversity in Gold-Catalyzed Formal Cycloadditions of Ynamides with Azidoalkenes or 2H-Azirines: [3 + 2] versus [4 + 3] Cycloadditions. *Chem. - Eur. J.* **2015**, *21*, 10843–10850. (d) Shu, C.; Wang, Y.-H.; Zhou, B.; Li, X.-L.; Ping, Y.-F.; Lu, X.; Ye, L.-W. Generation of  $\alpha$ -Imino Gold Carbenes through Gold-Catalyzed Intermolecular Reaction of Azides with Ynamides. *J. Am. Chem. Soc.* **2015**, *137*, 9567–9570. (e) Wu, Y.; Zhu, L.; Yu, Y.; Luo, X.; Huang, X. Polysubstituted 2-Aminopyrrole Synthesis via Gold-Catalyzed Intermolecular Nitrene Transfer from Vinyl Azide to Ynamide: Reaction Scope and Mechanistic Insights. *J. Org. Chem.* **2015**, *80*, 11407–11416. (f) Zhou, A.-H.; He, Q.; Shu, C.; Yu, Y.-F.; Liu, S.; Zhao, T.; Zhang, W.; Lu, X.; Ye, L.-W. Atom-economic generation of

gold carbenes: gold-catalyzed formal [3 + 2] cycloaddition between ynamides and isoxazoles. *Chem. Sci.* **2015**, *6*, 1265–1271. (g) Zhu, L.; Yu, Y.; Mao, Z.; Huang, X. Gold-Catalyzed Intermolecular Nitrene Transfer from 2H-Azirines to Ynamides: A Direct Approach to Polysubstituted Pyrroles. *Org. Lett.* **2015**, *17*, 30–33. (h) Xiao, X.-Y.; Zhou, A.-H.; Shu, C.; Pan, F.; Li, T.; Ye, L.-W. Atom-Economic Synthesis of Fully Substituted 2-Aminopyrroles via Gold-Catalyzed Formal [3 + 2] Cycloaddition between Ynamides and Isoxazoles. *Chem. - Asian J.* **2015**, *10*, 1854–1858. (i) Chen, M.; Sun, N.; Chen, H.; Liu, Y. Dioxazoles, a New Mild Nitrene Transfer Reagent in Gold Catalysis: Highly Efficient Synthesis of Functionalized Oxazoles. *Chem. Commun.* **2016**, *52*, 6324–6327. (j) Jin, H.; Huang, L.; Xie, J.; Rudolph, M.; Rominger, F.; Hashmi, A. S. K. Gold-Catalyzed C–H Annulation of Anthranils with Alkynes: A Facile, Flexible, and Atom-Economical Synthesis of Unprotected 7-Acyl Indoles. *Angew. Chem., Int. Ed.* **2016**, *55*, 794–797. (k) Yu, Y.; Chen, G.; Zhu, L.; Liao, Y.; Wu, Y.; Huang, X. Gold-Catalyzed  $\beta$ -Regioselective Formal [3 + 2] Cycloaddition of Ynamides with Pyrido[1,2-b]indazoles: Reaction Development and Mechanistic Insights. *J. Org. Chem.* **2016**, *81*, 8142–8154. (l) Xu, W.; Wang, G.; Sun, N.; Liu, Y. Gold-Catalyzed Formal [3 + 2] Cycloaddition of Ynamides with 4,5-Dihydro-1,2,4-oxadiazoles: Synthesis of Functionalized 4-Aminoimidazoles. *Org. Lett.* **2017**, *19*, 3307–3310. (m) Zeng, Z.; Jin, H.; Xie, J.; Tian, B.; Rudolph, M.; Rominger, F.; Hashmi, A. S. K.  $\alpha$ -Imino Gold Carbenes from 1,2,4-Oxadiazoles: Atom-Economical Access to Fully Substituted 4-Aminoimidazoles. *Org. Lett.* **2017**, *19*, 1020–1023. (n) Allegue, D.; González, J.; Fernández, S.; Santamaría, J.; Ballesteros, A. Regiodivergent Control in the Gold(I) Catalyzed Synthesis of 7-Pyrzolyloindoles from 1-Propargyl-1H-benzotriazoles and Ynamides through  $\alpha$ -Imino Gold(I) Carbene Complexes. *Adv. Synth. Catal.* **2019**, *361*, 758–768. (o) Tian, X.; Song, L.; Han, C.; Zhang, C.; Wu, Y.; Rudolph, M.; Rominger, F.; Hashmi, A. S. K. Gold(III)-Catalyzed Formal [3 + 2] Cycloaddition of N-Acyl Sulfilimines with Ynamides for the Synthesis of 4-Aminooxazoles. *Org. Lett.* **2019**, *21*, 2937–2940. (3) (a) Chatzopoulou, E.; Davies, P. W. Highly regioselective synthesis of 2,4,5-(hetero)aryl substituted oxazoles by intermolecular [3 + 2]-cycloaddition of unsymmetrical internal alkynes. *Chem. Commun.* **2013**, *49*, 8617–8619. (b) Reddy, R. J.; Ball-Jones, M. P.; Davies, P. W. Alkynyl Thioethers in Gold-Catalyzed Annulations To Form Oxazoles. *Angew. Chem., Int. Ed.* **2017**, *56*, 13310–13313. (c) Zhu, X.-Q.; Yuan, H.; Sun, Q.; Zhou, B.; Han, X.-Q.; Zhang, Z.-X.; Lu, X.; Ye, L.-W. Benign catalysis with zinc: atom-economical and divergent synthesis of nitrogen heterocycles by formal [3 + 2] annulation of isoxazoles with ynol ethers. *Green Chem.* **2018**, *20*, 4287–4291. (d) Davies, P. W.; Cremonesi, A.; Dumitrescu, L. Intermolecular and Selective Synthesis of 2,4,5-Trisubstituted Oxazoles by a Gold-Catalyzed Formal [3 + 2] Cycloaddition. *Angew. Chem., Int. Ed.* **2011**, *50*, 8931–8935. (4) (a) Jin, M. S. H.; Tian, M. S. B.; Song, M. S. X.; Xie, J.; Rudolph, M.; Rominger, F.; Hashmi, A. S. K. Gold-Catalyzed Synthesis of Quinolines from Propargyl Silyl Ethers and Anthranils through the Umpolung of a Gold Carbene Carbon. *Angew. Chem., Int. Ed.* **2016**, *55*, 12688–12692. (b) Sahani, R. L.; Liu, R.-S. Development of Gold-catalyzed [4 + 1] and [2 + 2 + 1]/[4 + 2] Annulations between Propiolate Derivatives and Isoxazoles. *Angew. Chem., Int. Ed.* **2017**, *56*, 1026–1030. (c) Sahani, R. L.; Liu, R. S. Gold-Catalyzed [4 + 2] Annulation/Cyclization Cascades of Benzisoxazoles with Propiolate Derivatives to Access Highly Oxygenated Tetrahydroquinolines. *Angew. Chem., Int. Ed.* **2017**, *56*, 12736–12740. (d) Shen, W.-B.; Xiao, X.-Y.; Sun, Q.; Zhou, B.; Zhu, X.-Q.; Yan, J.-Z.; Lu, X.; Ye, L.-W. Highly Site Selective Formal [5 + 2] and [4 + 2] Annulations of Isoxazoles with Heterosubstituted Alkynes by Platinum Catalysis: Rapid Access to Functionalized 1,3-Oxazepines and 2,5-Dihydropyridines. *Angew. Chem., Int. Ed.* **2017**, *56*, 605–609. (e) Kardile, R. D.; Kale, B. S.; Sharma, P.; Liu, R. S. Gold-Catalyzed [4 + 1]-Annulation Reactions between 1,4-Diyn-3-ols and Isoxazoles To Construct a Pyrrole Core. *Org. Lett.* **2018**, *20*, 3806–3809. (f) Giri, S. S.; Liu, R.-S. Gold-catalyzed [4 + 3]- and [4 + 2]-annulations of 3-en-1-ynamides with isoxazoles via novel  $6\pi$ -electrocyclizations of 3-azahepta trienyl

- cations. *Chem. Sci.* **2018**, *9*, 2991–2995. (g) Zhao, Y.; Wang, C.; Hu, Y.; Wan, B. Brønsted acid-catalyzed formal [5 + 2 + 1] cycloaddition of ynamides and isoxazoles with water: access to oxygen-bridged tetrahydro-1,4-oxazepines. *Chem. Commun.* **2018**, *54*, 3963–3966. (h) Kulandai Raj, A. S.; Tan, K. C.; Chen, L. Y.; Cheng, M. J.; Liu, R. S. Gold-Catalyzed Bicyclic Annulations of 4-Methoxy-1,2-dienyl-5-ynes with Isoxazoles to Form Indolizine Derivatives via an Au- $\pi$ -allene Intermediate. *Chem. Sci.* **2019**, *10*, 6437–6442. (i) Singh, R. R.; Skaria, M.; Chen, L.-Y.; Cheng, M.-J.; Liu, R.-S. Gold-Catalyzed [4 + 3]-Annulations of 2-Alkenyl-1-Alkynylbenzenes with Anthranils with Alkyne-Dependent Chemoselectivity: Skeletal Rearrangement Versus Non-Rearrangement. *Chem. Sci.* **2019**, *10*, 1201–1206. (j) Ogunlana, A. A.; Bao, X. Mechanistic insights into the gold(i)-catalyzed annulation of propiolates with isoxazoles: a DFT study. *Chem. Commun.* **2019**, *55*, 11127–11130. (k) Song, L.; Tian, X.; Rudolph, M.; Rominger, F.; Hashmi, A. S. K. Gold(III)-catalyzed chemo-selective annulations of anthranils with N-allylynamides for the synthesis of 3-azabicyclo[3.1.0]hexan-2-imines. *Chem. Commun.* **2019**, *55*, 9007–9010. (l) Zhu, X.-Q.; Wang, Z.-S.; Hou, B.-S.; Zhang, H.-W.; Deng, C.; Ye, L.-W. Zinc-Catalyzed Asymmetric Formal [4 + 3] Annulation of Isoxazoles with Enynol Ethers by  $6\pi$  Electrocyclization: Stereoselective Access to 2H-Azepines. *Angew. Chem., Int. Ed.* **2020**, *59*, 1666–1673. (m) Skaria, M.; More, S. A.; Kuo, T.-C.; Cheng, M.-J.; Liu, R.-S. Gold-Catalyzed Iminations of Terminal Propargyl Alcohols with Anthranils with Atypical Chemoselectivity for C(1)-Additions and 1,2-Carbon Migration. *Chem. - Eur. J.* **2020**, *26*, 3600–3608. (n) Arce, E. M.; Lamont, S. G.; Davies, P. W. Sulfenyl Ynamides in Gold Catalysis: Synthesis of Oxo-functionalised 4-aminoimidazolyl Fused Compounds by Intermolecular Annulation Reactions. *Adv. Synth. Catal.* **2020**, *362*, 2503–2509. (o) Vanjari, R.; Dutta, S.; Prabagar, B.; Gandon, V.; Sahoo, A. K. Ring Expansion and 1,2-Migration Cascade of Benzisoxazoles with Ynamides: Experimental and Theoretical Studies. *Chem. - Asian J.* **2019**, *14*, 4828–4836.
- (5) (a) Ye, X.; Wang, J.; Ding, S.; Hosseini, S.; Wojtas, L.; Akhmedov, N. G.; Shi, X. Investigations on Gold-Catalyzed Thioalkyne Activation Toward Facile Synthesis of Ketene Dithioacetals. *Chem. - Eur. J.* **2017**, *23*, 10506–10510. (b) Bai, Y.-B.; Luo, Z.; Wang, Y.; Gao, J.-M.; Zhang, L. Au-Catalyzed Intermolecular [2 + 2] Cycloadditions between Chloroalkynes and Unactivated Alkenes. *J. Am. Chem. Soc.* **2018**, *140*, 5860–5865. (c) Wang, J.; Zhang, S.; Xu, C.; Wojtas, L.; Akhmedov, N. G.; Chen, H.; Shi, X. Highly Efficient and Stereoselective Thioallylation of Alkynes: Possible Gold Redox Catalysis with No Need for a Strong Oxidant. *Angew. Chem., Int. Ed.* **2018**, *57*, 6915–6920. (d) Sharma, P.; Singh, R. R.; Giri, S. S.; Chen, L.-Y.; Cheng, M.-J.; Liu, R.-S. Gold-Catalyzed Oxidation of Thioalkynes To Form Phenylthio Ketene Derivatives via a Non-carbene Route. *Org. Lett.* **2019**, *21*, 5475–5479.
- (6) (a) Iardi, E. A.; Vitaku, E.; Njardarson, J. T. Data-Mining for Sulfur and Fluorine: An Evaluation of Pharmaceuticals To Reveal Opportunities for Drug Design and Discovery. *J. Med. Chem.* **2014**, *57*, 2832–2842. (b) Scott, K. A.; Njardarson, J. T. Analysis of US FDA-Approved Drugs Containing Sulfur Atoms. *Top. Curr. Chem.* **2018**, *376*, 5.
- (7) Bhardwaj, V.; Gumber, D.; Abbot, V.; Dhiman, S.; Sharma, P. Pyrrole: A Resourceful Small Molecule in Key Medicinal Hetero-Aromatics. *RSC Adv.* **2015**, *5*, 15233–15266.
- (8) Lou, J.; Wang, Q.; Wu, P.; Wang, H.; Zhou, Y.-G.; Yu, Z. Transition-metal mediated carbon–sulfur bond activation and transformations: an update. *Chem. Soc. Rev.* **2020**, *49*, 4307–4359.
- (9) Zhu, X.-Q.; Sun, Q.; Zhang, Z.-X.; Zhou, B.; Xie, P.-X.; Shen, W.-B.; Lu, X.; Zhou, J.-M.; Ye, L.-W. Zinc-catalyzed reaction of isoxazoles with thioynol ethers involving an unprecedented 1,2-sulfur migration. *Chem. Commun.* **2018**, *54*, 7435–7438.
- (10) (a) Cao, Z.; Zhu, J.; Liu, L.; Pang, Y.; Tian, L.; Sun, X.; Meng, X. AgNTf<sub>2</sub>-Catalyzed Formal [3 + 2] Cycloaddition of Ynamides with Unprotected Isoxazol-5-amines: Efficient Access to Functionalized 5-Amino-1H-pyrrole-3-carboxamide Derivatives. *Beilstein J. Org. Chem.* **2019**, *15*, 2623–2630. (b) Chen, C.; Cui, S. Iterative Assembly of Nitrile Oxides and Ynamides: Synthesis of Isoxazoles and Pyrroles. *J. Org. Chem.* **2019**, *84*, 12157–12164.
- (11) Ragno, R.; Coluccia, A.; La Regina, G.; De Martino, G.; Piscitelli, F.; Lavecchia, A.; Novellino, E.; Bergamini, A.; Ciapri, C.; Sinistro, A.; Maga, G.; Crespan, E.; Artico, M.; Silvestri, R. Design, Molecular Modeling, Synthesis, and Anti-HIV-1 Activity of New Indolyl Aryl Sulfones. Novel Derivatives of the Indole-2-carboxamide. *J. Med. Chem.* **2006**, *49*, 3172–3184.
- (12) La Regina, G.; Edler, M. C.; Brancale, A.; Kandil, S.; Coluccia, A.; Piscitelli, F.; Hamel, E.; De Martino, G.; Matesanz, R.; Díaz, J. F.; Scovassi, A. I.; Prospero, E.; Lavecchia, A.; Novellino, E.; Artico, M.; Silvestri, R. Arylthioindole Inhibitors of Tubulin Polymerization. 3. Biological Evaluation, Structure–Activity Relationships and Molecular Modeling Studies. *J. Med. Chem.* **2007**, *50*, 2865–2874.
- (13) Zhu, X.-Q.; Yuan, H.; Sun, Q.; Zhou, B.; Han, X.-Q.; Zhang, Z.-X.; Lu, X.; Ye, L.-W. Benign Catalysis with Zinc: Atom-Economical and Divergent Synthesis of Nitrogen Heterocycles by Formal [3 + 2] Annulation of Isoxazoles with Ynol Ethers. *Green Chem.* **2018**, *20*, 4287–4291.
- (14) For some recent examples assembling sulfonylated and sulfinylated pyrroles, see: (a) Prabagar, B.; Mallick, R. K.; Prasad, R.; Gandon, V.; Sahoo, A. K. Umpolung Reactivity of Ynamides: An Unconventional [1,3]-Sulfonyl and [1,5]-Sulfinyl Migration Cascade. *Angew. Chem., Int. Ed.* **2019**, *58*, 2365–2370. (b) Dutta, S.; Mallick, R. K.; Prasad, R.; Gandon, V.; Sahoo, A. K. Alkyne Versus Ynamide Reactivity: Regioselective Radical Cyclization of Yne-Ynamides. *Angew. Chem., Int. Ed.* **2019**, *58*, 2289–2294.
- (15) Brutiu, B. R.; Klose, I.; Maulide, N. Facile C–S Bond Cleavage of Aryl Sulfoxides Promoted by Brønsted Acid. *Synlett* **2021**, *32*, 488–490.
- (16) Gilow, H. M.; Brown, C. S.; Copeland, J. N.; Kelly, K. E. Sulfonylation of some pyrroles and indoles. *J. Heterocycl. Chem.* **1991**, *28*, 1025–1034.
- (17) (a) Luo, Q.; Jia, G.; Sun, J.; Lin, Z. Theoretical Studies on the Regioselectivity of Iridium-Catalyzed 1,3-Dipolar Azide–Alkyne Cycloaddition Reactions. *J. Org. Chem.* **2014**, *79*, 11970–11980. (b) Beno, B. R.; Yeung, K.-S.; Bartberger, M. D.; Pennington, L. D.; Meanwell, N. A. A Survey of the Role of Noncovalent Sulfur Interactions in Drug Design. *J. Med. Chem.* **2015**, *58*, 4383–4438. (c) Song, W.; Zheng, N.; Li, M.; He, J.; Li, J.; Dong, K.; Ullah, K.; Zheng, Y. Rhodium(I)-Catalyzed Regioselective Azide-internal Alkynyl Trifluoromethyl Sulfide Cycloaddition and Azide-internal Thioalkyne Cycloaddition under Mild Conditions. *Adv. Synth. Catal.* **2019**, *361*, 469–475.
- (18) Even without an aryl substituent on the alkyne, the  $\beta$ -keto enamide product from the  $\alpha$ -addition pathway was only observed in small amounts (<10%) in the <sup>1</sup>H NMR spectra of the crude reaction mixture for the formation of **3r**.