



University of Dundee

Editorial

Gissot, Mathieu; Pawlowic, Mattie C.; Modrzynska, Katarzyna K.; Francia, Maria E.

Published in:
Frontiers in Cellular and Infection Microbiology

DOI:
[10.3389/fcimb.2021.680184](https://doi.org/10.3389/fcimb.2021.680184)

Publication date:
2021

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):
Gissot, M., Pawlowic, M. C., Modrzynska, K. K., & Francia, M. E. (2021). Editorial: Molecular Basis of Stage Conversion in Apicomplexan Parasites. *Frontiers in Cellular and Infection Microbiology*, 11, [680184]. <https://doi.org/10.3389/fcimb.2021.680184>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Editorial: Molecular Basis of Stage Conversion in Apicomplexan Parasites

Mathieu Gissot^{1*}, Mattie C. Pawlowic^{2*}, Katarzyna K. Modrzynska^{3*} and Maria E. Francia^{4,5*}

¹ Univ. Lille, CNRS, Inserm, CHU Lille, Institut Pasteur de Lille, U1019 - UMR 9017 - CIL - Center for Infection and Immunity of Lille, Lille, France, ² Wellcome Centre for Anti-Infectives Research, School of Life Sciences, University of Dundee, Dundee, United Kingdom, ³ Wellcome Centre for Integrative Parasitology, University of Glasgow, Glasgow, United Kingdom, ⁴ Laboratory of Apicomplexan Biology, Institut Pasteur de Montevideo, Montevideo, Uruguay, ⁵ Department of Parasitology and Mycology, School of Medicine, Universidad de la República, Montevideo, Uruguay

Keywords: Apicomplexa, *Toxoplasma*, *Plasmodium*, *Cryptosporidium* spp., Eimeria, parasite, differentiation

Editorial on the Research Topic

Molecular Basis of Stage Conversion in Apicomplexan Parasites

OPEN ACCESS

Edited and reviewed by:

Tania F. De Koning-Ward,
Deakin University, Australia

*Correspondence:

Mathieu Gissot
mathieu.gissot@pasteur-lille.fr
Katarzyna K. Modrzynska
katarzyna.modrzynska@glasgow.ac.uk
Mattie C. Pawlowic
m.c.pawlowic@dundee.ac.uk
Maria E. Francia
mfrancia@pasteur.edu.uy

Specialty section:

This article was submitted to
Parasite and Host,
a section of the journal
Frontiers in Cellular
and Infection Microbiology

Received: 13 March 2021

Accepted: 23 March 2021

Published: 16 April 2021

Citation:

Gissot M, Pawlowic MC,
Modrzynska KK and Francia ME
(2021) Editorial: Molecular Basis
of Stage Conversion in
Apicomplexan Parasites.
Front. Cell. Infect. Microbiol. 11:680184.
doi: 10.3389/fcimb.2021.680184

INTRODUCTION

The Apicomplexa is a large phylum of intracellular parasitic protozoa with more than 6000 species that present a set of invasion organelles organized in an apical complex (Adl et al., 2007). It is estimated that every metazoan can be infected by at least one species from this group (Morrison, 2009), indicating their diversity and adaptation to the parasitic lifestyle. Most of what we know about this phylum is based on the study of a subset of parasites of medical and veterinary importance, such as those causing malaria, cryptosporidiosis, neosporosis, and toxoplasmosis.

All Apicomplexa studied so far are characterized by complex life cycles, during which parasites transition through multiple life forms, often requiring intermediate hosts and vectorial transmission. This process involves multiple cell types and, consequently, differentiation and proliferation steps to adapt to the changing environment. Apicomplexa switch from intra- to extracellular environments; encysted slow-dividing forms, inaccessible to the immune system, to fast-dividing forms exposed to immunity; motile to non-motile. These transitions are all accompanied both by morphological and metabolic changes. The collection of articles in this Topic “Molecular Basis of Stage Conversion in Apicomplexan Parasites” reviews and updates current knowledge of the molecular mechanisms controlling stage conversion in some of the best-described Apicomplexa and presents new findings contributing to this field of study. It also integrates new insight into less-studied species, expanding our breadth of knowledge and understanding of the phylum as a whole.

Plasmodium spp. or *Toxoplasma gondii* parasites benefit from the improvement of a flexible genetic toolbox that permits in-depth dissection of stage transitions. Studies dissecting the regulation of gene expression and, ultimately, of stage transition have shown that a fine-tuned multilayer regulation of gene expression is necessary to control each transition. Understanding the dynamic morphological and metabolic changes at the structural and molecular level is paramount to developing effective prophylactic and treatment strategies as shown in Mévêlec et al. In particular, stage-specific transcriptional regulation (involving epigenetic regulation, transcription factors, and chromatin organization) and translation repression are two mechanisms that shape the life cycle

progression in both *Plasmodium* spp as shown in Briquet et al. and Hollin and Le Roch and *T. gondii* (Sokol-Borelli et al.). In that respect, the recent characterization of the RNA Binding Protein UIS12 provides further evidence of the importance of post-transcriptional mechanisms in the control of *P. berghei* sexual stage development.

In addition to *Plasmodium* spp. and *T. gondii*, a number of other apicomplexan parasites are now amenable to genetic manipulation which has expanded our understanding of stage conversion in this phylum. The molecular basis of stage transition of coccidian parasites such as *Eimeria* spp. are now being investigated, demonstrating how metabolic pathways can be reshaped during transitions and contribute to the parasite's adaptation to the environment as illustrated by Martorelli Di Genova and Knoll. Dissecting the biology of a wider selection of parasites from the phylum, such as *T. gondii*'s closest relatives *Hammondia hammondi* and *Neospora caninum*, is also important as it can also contribute to our understanding of the evolution of adaptations of these parasites to specific biological niches and hosts. For example, Sokol-Borelli et al. show high similarity between parasite genomes and the presence of identified key factors for transition in *T. gondii*, which does not necessarily indicate that these same actors are controlling stage transition in other parasites.

However, despite this progress, some parts of the complex life cycles of Apicomplexa are still not experimentally amenable. The emergence of new *in vitro* models is now allowing for access to large parts of the complex life cycle of these parasites. For example, dissecting the biology of the *T. gondii* sexual cycle may now be within reach, as illustrated by Tomasina and Francia. Such fast progress in the field would not be possible without a number of technological advancements, including in particular improvements in the use of next-generation sequencing technologies. This approach is now powerful enough to investigate Apicomplexa species or life cycle stages that are difficult to access. For example, from an extensive collection of *Cryptosporidium* transcriptomics data, Li et al. produced a new bioinformatic pipeline that identified almost 400 developmentally regulated long non-coding RNAs. These

RNAs cover ~10% of the protein-coding genome and are upregulated in time points corresponding to gametocytogenesis, suggesting a role in sexual development. Next-generation sequencing also provides an in-depth characterization of the *T. gondii* transcriptome. Markus et al. mapped transcriptional start sites (TSS) of *T. gondii* with high resolution and stage-specific alternative TSS were identified for bradyzoite genes. *T. gondii* TSS were also analyzed in reference to recently published nucleosome data and found to be located unusually deep into the nucleosome, suggesting a role of chromatin architecture in stage conversion. It is clear that non-coding elements play important roles in differentiation and require further exploration.

In conclusion, the collection of articles in this Research Topic highlight the contribution of the study of underrepresented species and life stages, as well as the new improved technologies and model systems to current understanding of the molecular basis of stage transitions in Apicomplexa.

AUTHOR CONTRIBUTIONS

MG wrote the manuscript. MEF, KKM, and MCP edited the manuscript. All authors contributed to the article and approved the submitted version.

FUNDING

MG would like to acknowledge the following funding agencies for their support: ANR, CNRS, Inserm, and Institut Pasteur de Lille (CPER-CTRL). MEF is funded by a Banco de Seguros del Estado installment grant, an ACIP grant from the RIIP, and the National Agency for Research and Innovation (ANII). MCP was supported by a Sir Henry Dale Fellowship jointly funded by the Wellcome Trust and the Royal Society (Grant Number 213469/Z/18/Z) and start-up funds from a Wellcome Trust Centre Award (Grant Number 203134/Z/16/Z). KKM is supported by the Wellcome Trust/Royal Society Sir Henry Dale award (202600/Z/16/Z).

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Gissot, Pawlowic, Modrzynska and Francia. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

REFERENCES

- Adl, S. M., Leander, B. S., Simpson, A. G. B., Archibald, J. M., Anderson, O. R., Bass, D., et al. (2007). Diversity, Nomenclature, and Taxonomy of Protists. *Syst. Biol.* 56, 684–689. doi: 10.1080/10635150701494127
- Morrison, D. A. (2009). Evolution of the Apicomplexa: where are we now? *Trends Parasitol.* 25, 375–382. doi: 10.1016/j.pt.2009.05.010