

## **Associations between child mental health, carer worry and help-seeking**

Shaun Liverpool<sup>1,2</sup>

Julian Edbrooke-Childs<sup>1</sup>

<sup>1</sup>Evidence Based Practice Unit, University College London and Anna Freud National Centre  
for Children and Families

<sup>2</sup>Faculty of Health, Social Care & Medicine, Edge Hill University

### Abstract

Carers may not always express child mental health concerns to health professionals. Therefore, identifying factors delaying help-seeking are important. The aim of this study was to examine the relationship between carer affect and help-seeking. In a secondary analysis of data collected from school-aged children ( $N = 1,857$ , mean age = 9.85 years, 51% female), we used logistic regression to examine the associations between carer worry, help-seeking, and child mental health. Regarding worry, higher levels of emotional problems (OR = 1.42, 95% CI = 1.33-1.52), conduct problems (OR = 1.24, 95% CI = 1.12, 1.33), peer problems (OR = 1.17, 95% CI = 1.05-1.27), or functional impairment (OR = 1.37, 95% CI = 1.2-.56) were associated with higher levels of carer worry. Regarding help-seeking, higher levels of functional impairment were associated with higher levels of help-seeking (OR = 1.51, 95% CI = 1.09-2.11). After controlling for mental health problems, carers who reported being worried about their child's mental health were less likely than other carers to seek help (OR = 0.13, 95% CI = 0.05-0.35). Knowledge of these factors may inform early interventions. Alongside implications for future research and practice, limitations of the study are discussed.

*Keywords:* child; adolescent; parents and carers; help-seeking behaviour; mental health

### **Associations between child mental health, carer worry and help-seeking**

A high prevalence of young people experience mental health problems at some point during childhood and adolescence (Oh et al., 2015; Rickwood et al., 2005). However, only a proportion of young people experiencing mental health problems will receive mental health support (Mitchell et al., 2017; Oh & Bayer, 2015). In the extant literature, several theoretical explanations have been proposed for help-seeking, albeit these are predominantly for adults (Gulliver et al., 2012). For example, the theory of planned behaviour has been used to demonstrate the mediating effect of attitudes on psychological help-seeking intentions (Ajzen, 2011). Similarly, the health belief model posits that help-seeking is determined by the individual's appraisal of the perceived threat of the illness, its severity, and barriers and benefits of help-seeking (Rosenstock, 1977). A meta-analysis has found that perceived barriers and perceived benefits are strong predictors of help-seeking behaviour, although perceived threats and severity were not consistent predictors (Carpenter, 2010).

Parents and carers are key to identifying when their child is experiencing mental health problems and critical to the help-seeking process (Haine-Schlagel & Walsh, 2015). The gateway provider model posits that carers' decision to seek help may be influenced by a range of internal and external factors, such as perceived need, demographic characteristics, and awareness and availability of resources (Stiffman, Pescosolido & Cabassa, 2004). In particular, limited awareness of formal help-seeking processes, carers' stress levels and carers' perception of the severity of the child's mental health problem are barriers to help-seeking (Bonis, 2016; O'Brien et al., 2016; Reardon et al., 2017; Reardon et al., 2019). Previous studies also highlight that carers experience additional barriers due to negative attitudes towards seeking help, concerns about trust, and fear of being stigmatized or judged (Andershed et al., 2017; Bradby et al., 2007; Coogle & Hanline, 2016).

Quantitative studies thus far have mainly focused on families already accessing specialist mental health care (Oh et al., 2015; Rickwood et al., 2005; Skylstad et al., 2019). Further, studies in the United States (US) have shown that higher levels of carer worry were

associated with accessing formal help-seeking (Ellingson et al., 2004; Godoy et al., 2014). This is not mirrored in findings from qualitative studies in the United Kingdom (UK), where higher levels of carer worry, and perceiving a child's mental health problems as being more severe, were reported as barriers to help-seeking (Bradby et al., 2007; Sayal et al., 2010). This pattern of findings can potentially be explained using the affective appraisal approach to decision making in child mental health settings (Liverpool et al., 2021). This evidence-informed model posits that carer child mental health decision making is a cyclical process, underpinned by affective states. Hence, high levels of worry and concern that a child is experiencing severe problems may contribute to negative affective states, which act as a barrier to formal help-seeking. Notwithstanding, carers may be more likely to access informal, rather than formal, support when their child is experiencing mental health difficulties (Reardon et al., 2019). These authors also highlighted that carers may feel a sense of blame and concern that they will be perceived to have failed as a carer, placing further barriers to help-seeking. Despite the authors' focus on anxiety, these findings highlight the importance of exploring carer worry, specific to the child's mental health, which can be a potential underlying factor influencing barriers to recognising a problem and subsequently seeking help. This knowledge could further evidence the role of affective states in carer help-seeking and child mental health.

To improve the quality of care and early interventions for carers and their children, it would be important to further investigate the factors influencing help-seeking behaviours and identify possible predictors. In light of the current waiting periods to access child and adolescent mental health services (CAMHS) (NHS Digital, 2019), and only a subset of parents accessing and engaging with CAMHS, it is just as important to investigate the gap between service need and service access in a non-clinical population. It is important to extend the literature further to understand how these same factors affect help-seeking from both informal and formal sources. Additionally, by exploring help-seeking experiences of parents in a non-clinical population (i.e. pre-CAMHS) we can identify; 1) areas suitable for

early interventions; 2) families who are less likely to access early intervention or carer support; and 3) areas to target that will require further investigation.

## **Aims**

Consequently, the aim of the present research was to examine characteristics associated with carer help-seeking and child mental health, using a secondary analysis of data collected from school-aged children in the UK. In particular, we examined the associations between carer worry, help-seeking, and their child's mental health. We expected that carers would have higher levels of worry when their child's mental health problems were more severe. We also expected that carers would be less likely to seek help when they had higher levels of worry and when their child's mental health problems were more severe.

## **Methods**

### **Data source**

A secondary analysis was conducted on data collected from carers of children 8-12 years from primary and secondary schools in the UK who participated in the 2008-2011 Targeted Mental Health in Schools programme (TaMHS). All families were eligible to participate in the study. Further details of the TaMHS programme have been reported elsewhere (Department for Education, 2011). Taking into account the length of time it generally takes for data to be released for further analysis (Johnston, 2014), the authors recently received permission to analyse and publish the findings from the baseline data. Nonetheless, owing to the limited knowledge on carer help-seeking for child mental health, the dataset was deemed valid for exploratory purposes. The original TaMHS study received ethical approval for collection and analysis of the primary data. However, this secondary data analysis included no identifiable data and permission was granted for further use and analysis (Tripathy, 2013).

## **Measures**

### ***Demographic characteristics***

The demographic profile of the sample included relationship to the child, the child's age and sex, and the families first language. Relationship to the child, as reported by the carer, was categorised as mother, father, guardian, or other. The child's age was calculated based on the child's date of birth, and the child's sex, based on available data, was categorised as male or female. The family's first language was categorised as "English" or "Other". Self-report ethnicity was recorded using the superordinate categories of the 2011 Census: Asian, Black, mixed-race, "other", or White (Office for National Statistics, 2012).

### ***Child mental health problems***

To capture child mental health problems, carers completed the Strengths and Difficulties Questionnaire (SDQ), which consists of 25 items divided into 5 sub-scales (emotional problems, conduct problems, hyperactivity, peer relationship problems, and prosocial behaviour), in addition to the impact supplement (Goodman et al., 2003). Higher scores indicate higher levels of mental health difficulties, except for the prosocial sub-scale where lower scores indicate higher levels of prosocial behaviour difficulties (Goodman et al., 2003). Items required parents to respond on a Likert scale (not true, somewhat true, or certainly true) based on the child's behaviour (eg. Constantly fidgeting or squirming) or the child's mood (eg. Many worries or Often seems worried) within the previous six months. A total difficulties score, including emotional problems, conduct problems, hyperactivity, and peer relationships problems, can also be calculated ranging from 0 to 40, with an increased score corresponding to an "increase in the risk" of developing a mental health disorder. The SDQ displayed adequate internal consistency (Cronbach's alpha = 0.74) for the current sample.

### ***Carer worry about their child's mental health***

Worry has been defined as a state of feeling anxious or troubled about a person or situation (Hirsch & Mathews, 2012). Carer worry was assessed using a binary item based on

a modified version of the General Help-Seeking Questionnaire (Wilson et al., 2005). Carers were asked “Have you ever been worried because your child seemed to be unhappy or disruptive?” to which they responded “Yes/No”. Similar to the Health Anxiety Questionnaire (Lucock & Morley, 1996) and the Health Anxiety Inventory (Salkovskis et al., 2002), which are used to report on an individual’s worry about their own physical or mental health, the current measure aimed at capturing carer worry about their child’s mental health.

### ***Carer help-seeking for child mental health problems***

Based on the modified version of the General Help-Seeking Questionnaire, a brief measure was used to capture carer help-seeking behaviours. Experts encourage that where possible the use of such measures are necessary to reduce burden and ease of interpretation (Bowling, 2005). Carers were asked “Did you try to get help from any of the following: A family member/ A friend/ A form/class teacher/ A family doctor?” to which they responded “No” or “Yes, but not helpful” or “Yes, a little helpful” or “Yes, and helped a lot”. To address our research questions, the responses were coded as “Yes” or “No” responses. It was important to note that only a subset of our sample ( $n = 529$ ) responded to this question. Differences between completers and non-completers were statistically significant only in relation to prosocial behaviour:  $\chi^2(17) = 1998.19, p < .05$ . Carers reporting higher prosocial scores for their children were less likely to complete the help-seeking item.

### **Design and statistical analysis**

First, an exploration of the variation among carer help-seeking between schools was conducted to investigate if carers at the same schools displayed similar help-seeking behaviours. The intraclass correlation coefficient was  $<.05$ , suggesting that the school-level accounted for less than 5% of the variation in carer help-seeking behaviour. Therefore, logistic regressions were acceptable (LeBreton & Senter, 2008).

Once all assumptions were tested and confirmed, binary logistic regression analysis was conducted to explore the associations between child mental health with carer worry, controlling for demographic variables. Similarly, binary logistic regression analysis was conducted to explore the associations between child mental health and carer worry with

carer help-seeking behaviour, controlling for demographic variables. Odds ratios (ORs) and corresponding 95% confidence intervals (CI) were used as estimates of the associations. Statistical analyses were conducted using SPSS (version 24), and p-values <0.05 were considered significant (Du Prel et al., 2009).

## Results

### Participants

The included sample comprised  $N = 1,857$  children (MeanAge=9.85, SD=1.56; 950 or 51.2% girls). The sample was predominantly English-speaking mothers of White ethnic origins. The overall sample had “close to average” scores on all subscales of the SDQ (Goodman et al., 2003). The majority of carers (1,352, 73%) reported feeling worried about their child’s mental health, of whom less than half (529, 39%) had complete data on seeking help from formal (teachers or general practitioners) and/or informal sources (family members or friends). See Table 1 for a descriptive summary of the sample.

**Table 1**

*Demographic and clinical characteristics of the total sample.*

Characteristics	Frequencies
Child’s age, M ( <i>SD</i> )	9.85 (1.56)
Child’s sex, n (%)	
Boy	907 (48.8%)
Girl	950 (51.2%)
Family’s first language	
English	1683 (90.6%)
Other	174 (9.4%)
Ethnicity	
Asian	134 (7.2%)
Black	44 (2.4%)
Mixed-race	45 (2.4%)
Other	11 (0.6%)
White	1623 (87.4%)
Relationship to child	
Father	275 (14.8%)



Mother	1560 (84%)
Guardian	18 (1%)
Other	4 (0.2%)
Child mental health problems, Mdn ( <i>IQR</i> )	
Hyperactivity and inattention	3 (4)
Emotional problem	2 (3)
Conduct problems	1 (2)
Peer problems	1 (2)
Prosocial behaviours	9 (3)
Impact score	0 (1)
Total difficulties score	7 (8)
Carer worry	
Yes	1352 (72.8%)
No	505 (27.2%)

*Note.*  $N = 1,857$ . Continuous variables were described by the mean (M) and standard deviation (SD) or median (Mdn) and interquartile range (IQR). Categorical variables were described by count (n) and percentages (%).

### Associations between carer worry and child mental health

The logistic regression model was statistically significant:  $\chi^2 (16) = 576.54, p < .05$ . Asian carers were about 3.9 times less likely to report being worried about child mental health than White parents (OR=.258, 95% CI [0.107, 0.624]). Guardians were about 11 times less likely to report being worried than mothers (OR = .092, 95% CI [0.011, 0.941]). As the carer ratings of their child's emotional problems (OR=1.418, 95% CI [1.326, 1.516]), conduct problems (OR=1.237, 95% CI [1.123, 1.363]), and peer problems (OR=1.172, 95% CI [1.084, 1.268]) increased, they were more likely to indicate that they were worried about their child's mental health. As functional impairment increased (i.e., impact score) carers were about 1.4 times more likely to indicate that they were worried (OR=1.371, 95% CI [1.202, 1.563]). The child's age, gender, first language, hyperactivity and inattention problems, and prosocial behaviours were not independently significant in predicting carer worry (see Table 2).

**Table 2***Logistic regression with demographic and clinical characteristics predicting carer worry.*

Variable	OR	95% Confidence Interval
Child's age	0.994	[0.918, -1.076]
Child's sex (Ref: Girl)		
Boy	1.279	[0.987, 1.657]
Language (Ref: English)		
Other	0.564	[0.264, 1.207]
Ethnicity (Ref: White)		
Black	0.817	[0.332, 2.012]
Asian	0.258	[0.107, .624]*
Mixed	1.584	[0.719, 3.490]
Other	0.340	[0.058, 1.999]
Relationship to child (Ref: Mother)		
Father	0.690	[0.472, 1.007]
Guardian	0.092	[0.011, .941]*
Other	0.594	[0.053, 6.694]
Psychosocial difficulties		
Hyperactivity	1.020	[0.960, 1.085]
Emotional problems	1.418	[1.326, 1.516]**
Conduct problems	1.237	[1.123, 1.363]**
Peer problems	1.172	[1.084, 1.268]**
Prosocial behaviours	0.932	[0.858, 1.013]
Impact score	1.371	[1.202, 1.563]**

*Note.*  $N=1,852$ . OR = odds ratio; CI = confidence interval; Ref = Reference group was selected as per the larger percentage of the sample

\* =  $p < .05$ , \*\* =  $p < .001$ .

### **Associations between child mental health and carer worry with help-seeking**

Analyses were conducted for the sample of carers who completed the help-seeking measure ( $n = 529$ ). The logistic regression model was statistically significant,  $\chi^2 (17) = 45.94$ ,  $p < .05$ . Fathers were about 2.6 times less likely to report seeking help than mothers (OR=0.386, 95% CI [0.163, 0.915]). Carers who reported being worried about their child's mental health were about 7.5 times less likely to seek help than carers who did not report

being worried (OR=.134, 95% CI [.051, .349]). However, as functional impairment increased (i.e., impact score), carers were about 1.5 times more likely to seek help (OR=1.514, 95% CI [1.085, 2.114]). None of the other variables were independently significant in predicting carer help-seeking (see Table 3).

**Table 3**

*Logistic regression with demographic, clinical characteristics and carer worry predicting help-seeking.*

Variable	OR	95% CI
Child's age	0.973	[0.792, 1.197]
Child's sex (Ref: Girl)		
Boy	0.787	[0.404, 1.535]
Language (Ref: English)		
Other	0.601	[0.088, 4.092]
Relationship to child (Ref: Mother)		
Father	0.386	[0.163, .915]*
Psychosocial difficulties		
Hyperactivity	1.045	[0.899, 1.227]
Emotional problems	0.891	[0.768, 1.033]
Conduct problems	0.885	[0.716, 1.095]
Peer problems	1.124	[0.921, 1.372]
Prosocial behaviours	1.016	[0.832, 1.240]
Impact score	1.514	[1.085, 2.114]*
Parental worry (Ref: No)		
Yes	0.134	[.051, .349]**

*Note.*  $N = 529$  OR = odds ratio; CI = confidence interval; Ref = Reference group was selected as per the larger percentage of the sample

\* $p < .05$ , \*\* $p < .001$

## Discussion

The aim of the present research was to examine characteristics associated with carer help-seeking and child mental health, using a secondary analysis of data collected from school-aged children in the UK. In particular, we examined the associations between carer worry, help-seeking, and their child's mental health. The findings from the logistic

regressions suggest that carer perceptions of their child's mental health difficulties and carer worry play important roles in help-seeking.

Regarding carer worry, higher levels of emotional problems, conduct problems, peer problems, and functional impairment were associated with increased likelihood of worry. These findings are in line with previous studies suggesting that carers' appraisal of their child's mental health is associated with their level of worry (Godoy et al., 2014). The findings of the present research also add to the evidence-base indicating that carer worry may go beyond the activities involved in help-seeking (e.g. concerns about stigma and lack of resources) and be specific to the child's condition or its impact.

Regarding help-seeking, higher levels of functional impairment were associated with increased likelihood of help-seeking. This is in line with previous research suggesting that higher levels of functional impairment for children were associated with higher levels of help-seeking for carers (Ellingson et al., 2004; Godoy et al., 2014). This finding is also in line with previous evidence suggesting that individuals are more likely to seek help when mental health difficulties impair functioning in education, work, training, and interpersonal relationships (DosReis et al., 2010; Patel et al., 2007). Nevertheless, previous research suggests that problem severity in addition to functional impairment is associated with help-seeking, which was not the case in the present study as problem severity was not associated with help-seeking (Reardon et al., 2017; 2019). The impact of the mental health problem was significant in our model, but the severity of the individual problems were not, therefore, implicating a need for further investigations to differentiate between presence and impact of the child's mental health on the carers' decision to seek help.

This study further adds to the literature highlighting carer worry as a potential influencing factor that may be a barrier to the help-seeking process. This finding may be partly explained by the perceived level of burden the mental health problems may have on the family (Angold et al., 1998) or the level of blame parents place on themselves (Reardon et al., 2019). However, this finding is contrary to similar US based studies that reported a positive association between parental worry and help-seeking (Ellingson et al., 2004; Godoy

et al.,2014). The contrary findings may be explained by differences in the studies' samples. Previous studies targeted parents of younger children (<5 years), parents seeking formal support, and included smaller samples (<300). Nonetheless, based on raw scores, previous studies highlighted that less than one third of parents actually spoke to a healthcare provider about the problem (Ellingson et al., 2004; Oh & Baker, 2015; Sayal & Taylor, 2004). This raises a further question about which problems parents consider as atypical in preteens and therefore warrants seeking help. Taken together, this finding underscores the importance of the parents' affective states as an influencing factor to help-seeking and the need for further investigations (Liverpool et al., 2021). The future findings can further inform existing help-seeking models (eg. Gateway Providers Model) by explicitly acknowledging and accounting for parents' emotional states.

The current findings also reflect potential demographic disparities. Asian parents and guardians were less likely to worry about their child's mental health, and fathers were less likely to seek help. The results partly support previous qualitative findings suggesting British-Asian parents are worried about being 'gossiped' about or stigmatized which prevented them from seeking help (Bradby et al., 2007). Similarly, although studies show that fathers do worry about their child's mental health (Skylstad et al., 2019), the current findings added that they are less likely to seek help. Although these findings may help to identify socially defined groups for targeted interventions, these groups were represented in less than 15% of the total sample, and therefore need to be interpreted with caution.

### **Limitations**

The strength of this exploratory study lies in its ability to provide some insight into carer worry about child mental health problems and their decision to seek help. However, potential limitations should be considered when interpreting the findings of the present research. This was a study about parents' decision to seek help and not a formal preference trial or evaluation of the depth of services received, so no conclusions about the effectiveness of the support or which support parents prefer can or should be drawn from

these findings. The authors only accessed available data from 2008 of the TaMHS dataset, with majority White English-speaking mothers, to represent a cross-section of the population, and as a result, causality and generalizability cannot be inferred. In acknowledging the time of data collection, this study acts as an exploratory study to support findings from more recent studies and inform future research. The authors also take caution when comparing the current findings to studies involving data more recently collected as there have been many advancements in the field.

Owing to the limitations of secondary data analysis (Johnston, 2014), the available dataset only provided a single item measure to assess parental worry. Due to parents' interpretation of worry, and the subjective nature of the question, participants may have selected to express their first or most recent recall of their child being unhappy or disruptive, therefore neglecting other instances or lack thereof. Similarly, parents may not have reported being worried when their children experienced other mental health challenges.

Another limitation is that the data collected was based on self-report data from parents. Due to self-report bias, some parents may have reported underestimated or inflated scores on psychosocial difficulties (Cheng et al., 2018). Further, less than half of the total sample (n=529) responded to the help-seeking measure. Moreover, help-seeking from online resources, traditional mental health services (e.g. psychologist) or other sources (eg. charities) were not reflected in the measure. Further, investigations into the differences in completers and non-completers of the help-seeking measure showed that increased prosocial scores was associated with completion. Therefore, the current findings may not be generalisable to families reporting higher prosocial scores.

### **Implications for Future Research**

Further studies on effectiveness and preference on various sources of support in child mental health is warranted as satisfaction has an important influence on help-seeking and whether parents go on to seek further help (Gulliver et al., 2010), or to accept the support offered. Recognising the need for help can be challenging as carers' perceptions of

their child's mental health differ from that of the children, teachers and health professionals (Cleridou et al., 2017; Hawley & Weisz, 2003). These disagreements may develop into a cycle of deciding to seek help then deciding to not seek further help. More rigorous and up to date studies in this area, controlling for the views of others and any previous experience parents have had with accessing support would be beneficial to update the theories outlined in the help-seeking literature.

### **Implications for Practice**

Results from this study are useful in identifying possible areas of shortcomings. Carers of school aged children may have regular contact with schools and therefore continuing to establish stronger links between mental health services and education providers may lead to more children receiving appropriate support. In addition, parents do access informal support and therefore, these support systems should be provided with the knowledge base to offer the necessary support. Consequently, information about identifying child mental health disorders, ways to access help and treatment, and early-stage interventions should be widely disseminated and accessible to the public. Additionally, embedded in practice, should be the necessary emotional support guidelines for working with parents of children accessing child mental health support. It may also be important to consider lowering carers' worry through routine screening and education about child mental health problems.

### **Conclusion**

Although these findings could be viewed as exploratory, they suggest that carers' perception of child mental health problems and carer worry may influence their decision to seek help. Knowledge of these factors may improve development and access to early carer-targeted interventions and further inform theories of pathway to CAMHS. Involving informal support systems in the development and dissemination of information resources and interventions may help reach those parents who are less likely to access CAMHS.

**Acknowledgements**

The authors would like to thank Dr. Mohammed Moinuddin and Thomas Bardsley for guidance and advice on data analysis and interpretation. We would also like to thank Professor Miranda Wolpert, Professor Jessica Deighton and Professor Peter Fonagy for guidance and support at various points throughout this study. Lastly, we would like to thank Miles Weekes for his support at the time of writing the manuscript.

**Contributorship**

All authors contributed to the conception and design of the study. Analysis and interpretation of the data was conducted by the first author. All authors contributed to the drafting and refinement of the article before submission.

**Conflict of interest**

The authors have declared that they have no competing or potential conflicts of interest.



**Abbreviations**

CAMHS- Child and adolescent mental health services

CI- Confidence interval

OR- Odds ratio

SDQ- Strengths and Difficulties Questionnaire

TaMHS- Targeted Mental Health in Schools

### References

- Andershed, B., Ewertzon, M., & Johansson, A. (2017). An isolated involvement in mental health care – experiences of parents of young adults. *Journal of Clinical Nursing* 26(7–8): 1053–1065. <https://doi.org/10.1111/jocn.13560>
- Angold, A., Messer, S. C., Stangl, D., Farmer, E. M., Costello, E. J., & Burns, B. J. (1998). Perceived parental burden and service use for child and adolescent psychiatric disorders. *American journal of public health* 88(1): 75–80. <https://doi.org/10.2105/ajph.88.1.75>
- Ajzen, I. (2011). The theory of planned behaviour: Reactions and reflections. *Psychology & Health*, 26(9): 1113-1127.
- Bøe, T., Hysing, M., Skogen, J. C., & Breivik, K. (2016). The Strengths and Difficulties Questionnaire (SDQ): Factor Structure and Gender Equivalence in Norwegian Adolescents. *PloS One* 11(5): e0152202–e0152202. <https://doi.org/10.1371/journal.pone.0152202>
- Bonis, S. (2016). Stress and Parents of Children with Autism: A Review of Literature. *Issues in Mental Health Nursing* 37(3): 153-163 <https://doi.org/10.3109/01612840.2015.1116030>
- Bowling, A (2005). Just one question: If one question works, why ask several? *Journal of Epidemiology & Community Health* 59(5): 342-345.
- Bradby, H., Varyani, M., Oglethorpe, R., Raine, W., White, I., & Helen, M. (2007). British Asian families and the use of child and adolescent mental health services: A qualitative study of a hard to reach group. *Social Science and Medicine* 65(12): 2413–2424.
- Butler, A. M., Weller, B., & Titus, C. (2015). Relationships of Shared Decision Making with Parental Perceptions of Child Mental Health Functioning and Care. *Administration and Policy in Mental Health* 42(6): 767–774. <https://doi.org/10.1007/s10488-014-0612-y>
- Carpenter C. J. (2010). A meta-analysis of the effectiveness of health belief model variables in predicting behavior. *Health communication* 25(8): 661–669. <https://doi.org/10.1080/10410236.2010.521906>

- Cheng, S., Keyes, K. M., Bitfoi, A., Carta, M. G., Koç, C., Goelitz, D., Otten, R., Lesinskiene, S., Mihova, Z., Pez, O., & Kovess-Masfety, V. (2018). Understanding parent–teacher agreement of the Strengths and Difficulties Questionnaire (SDQ): Comparison across seven European countries. *International Journal of Methods in Psychiatric Research* 27(1): e1589. <https://doi.org/10.1002/mpr.1589>
- Choudhry, F. R., Mani, V., Ming, L. C., & Khan, T. M. (2016). Beliefs and perception about mental health issues: a meta-synthesis. *Neuropsychiatric Disease and Treatment* 12: 2807–2818. <https://doi.org/10.2147/NDT.S111543>
- Cleridou, K., Patalay, P., & Martin, P. (2017). Does parent–child agreement vary based on presenting problems? Results from a UK clinical sample. *Child and Adolescent Psychiatry and Mental Health* 11(1): 22. <https://doi.org/10.1186/s13034-017-0159-2>
- Coogle, C. G., & Hanline, M. F. (2016). An exploratory study of family-centred help-giving practices in early intervention: Families of young children with autism spectrum disorder. *Child & Family Social Work* 21(2): 249–260.
- Department for Education. (2011). *Me and My School: Findings from the National Evaluation of Targeted Mental Health in Schools 2008-2011*. Available at: [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/184060/DFE-RR177.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/184060/DFE-RR177.pdf) (accessed 25 January 2021).
- DosReis, S., Barksdale, C. L., Sherman, A., Maloney, K., & Charach, A. (2010). Stigmatizing Experiences of Parents of Children With a New Diagnosis of ADHD. *Psychiatric Services* 61(8): 811–816.
- Du Prel, J.-B., Hommel, G., Röhrig, B., & Blettner, M. (2009). Confidence interval or p-value?: part 4 of a series on evaluation of scientific publications. *Deutsches Arzteblatt International* 106(19): 335–339. <https://doi.org/10.3238/arztebl.2009.0335>
- Edbrooke-Childs, J., Jacob, J., Argent, R., Patalay, P., Deighton, J., & Wolpert, M. (2015). The relationship between child- and parent-reported shared decision making and child-, parent-, and clinician-reported treatment outcome in routinely collected child mental

- health services data. *Clinical Child Psychology and Psychiatry* 21(2): 324–338.  
<https://doi.org/10.1177/1359104515591226>
- Ellingson, K. D., Briggs-Gowan, M. J., Carter, A. S., & Horwitz, S. M. (2004). Parent Identification of Early Emerging Child Behavior Problems: Predictors of Sharing Parental Concern With Health Providers. *JAMA Pediatrics* 158(8): 766–772.  
<https://doi.org/10.1001/archpedi.158.8.766>
- Godoy, L., Mian, N. D., Eisenhower, A. S., & Carter, A. S. (2014). Pathways to service receipt: modeling parent help-seeking for childhood mental health problems. *Administration and Policy in Mental Health* 41(4): 469–479.  
<https://doi.org/10.1007/s10488-013-0484-6>
- Goodman, R., Meltzer, H., & Bailey, V. (2003). The Strengths and Difficulties Questionnaire: a pilot study on the validity of the self-report version. *International Review of Psychiatry* 15(1–2): 173–177. <https://doi.org/10.1080/0954026021000046137>
- Gopalan, G., Goldstein, L., Klingenstein, K., Carolyn Sicher Psy, D., Blake, C., & McKay, M. M. (2010). Engaging families into child mental health treatment: Updates and special considerations. *Journal of the Canadian Academy of Child and Adolescent Psychiatry* 19(3): 182-196.
- Gulliver, A., Griffiths, K. M., & Christensen, H. (2010). *Perceived barriers and facilitators to mental health help-seeking in young people: A systematic review* 10(1): 113.  
<https://doi.org/10.1186/1471-244X-10-113>
- Haine-Schlagel, R., & Walsh, N. E. (2015). A Review of Parent Participation Engagement in Child and Family Mental Health Treatment. In *Clinical Child and Family Psychology Review* 18(2): 133-150. <https://doi.org/10.1007/s10567-015-0182-x>
- Hawley, K. M., & Weisz, J. R. (2003). Child, parent and therapist (dis)agreement on target problems in outpatient therapy: The therapist's dilemma and its implications. *Journal of Consulting and Clinical Psychology*, 71(1), 62–70.  
<http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=psyc4&AN=2003-01069-009>

- Hirsch, C. R., & Mathews, A. (2012). A cognitive model of pathological worry. *Behaviour Research and Therapy* 50(10): 636-646.. <https://doi.org/10.1016/j.brat.2012.06.007>
- Jackson, C., Cheater, F. M., & Reid, I. (2008). A systematic review of decision support needs of parents making child health decisions. *Health Expectations* 11(3): 232-251  
<https://doi.org/10.1111/j.1369-7625.2008.00496.x>
- Johnston, M. (2017). Secondary Data Analysis: A Method of which the Time Has Come. *Qualitative And Quantitative Methods In Libraries* 3(3): 619-626
- Karver, M. S., Handelsman, J. B., Fields, S., & Bickman, L. (2006). Meta-analysis of therapeutic relationship variables in youth and family therapy: The evidence for different relationship variables in the child and adolescent treatment outcome literature. *Clinical Psychology Review* 26(1): 50–65.  
<https://doi.org/https://doi.org/10.1016/j.cpr.2005.09.001>
- LeBreton, J. M., & Senter, J. L. (2008). Answers to 20 questions about interrater reliability and interrater agreement. *Organizational Research Methods* 11(4): 815-852.  
<https://doi.org/10.1177/1094428106296642>
- Lerner, J. S., Li, Y., Valdesolo, P., & Kassam, K. S. (2015). Emotion and Decision Making. *Annual Review of Psychology* 66(1): 799–823. <https://doi.org/10.1146/annurev-psych-010213-115043>
- Lucock, M. P., & Morley, S. (1996). The Health Anxiety Questionnaire. *British Journal of Health Psychology* 1(2): 137–150. <https://doi.org/10.1111/j.2044-8287.1996.tb00498.x>
- Macdonald, E., Lee, E., Geraghty, K., McCann, K., Mohay, H., & O'Brien, T. (2007). Towards a developmental framework of consumer and carer participation in child and adolescent mental health services. *Australasian Psychiatry* 15(6): 504-508.  
<https://doi.org/10.1080/10398560701458228>
- Mitchell, C., McMillan, B., & Hagan, T. (2017). Mental health help-seeking behaviours in young adults. *British Journal of General Practice* 67(654): 8-9.  
<https://doi.org/10.3399/bjgp17X688453>

- NHS Digital. (2019). Waiting times for children and young people's mental health services, 2018 - 2019 additional statistics. Available at: <https://digital.nhs.uk/data-and-information/find-data-and-publications/supplementary-information/2019-supplementary-information-files/waiting-times-for-children-and-young-peoples-mental-health-services-2018---2019-additional-statistics> (accessed 25 January 2021).
- O'Brien, D., Harvey, K., Howse, J., Reardon, T., & Creswell, C. (2016). Barriers to managing child and adolescent mental health problems: A systematic review of primary care practitioners' perceptions. *British Journal of General Practice* 66(651): E693-E707.. <https://doi.org/10.3399/bjgp16X687061>
- Office for National Statistics (2012). 2011 Census Key Statistics for England and Wales, UK Data Service Census Support. Available at: <https://wicid.ukdataservice.ac.uk> (accessed 25 January 2021).
- Oh, E., & Bayer, J. K. (2015). Parents' help-seeking processes for early childhood mental health problems. *Child and Adolescent Mental Health* 20(3): 149-154. <https://doi.org/10.1111/camh.12081>
- Oh, E., Mathers, M., Hiscock, H., Wake, M., & Bayer, J. (2015). Professional help seeking for young children with mental health problems. *Australian Journal of Psychology* 67(3): 187-195. <https://doi.org/10.1111/ajpy.12072>
- Ozbek, A., Gencer, O., & Mustan, A. T. (2019). Which parents dropout from an evidence-based parenting programme (Triple-P) at CAMHS? Comparison of programme-completing and dropout parents. *Clinical Child Psychology and Psychiatry* 24(1): 144-157. <https://doi.org/10.1177/1359104518792294>
- Patel, V., Flisher, A. J., Hetrick, S., & McGorry, P. (2007). Mental health of young people: a global public-health challenge. *The Lancet* 369(9569), 1302–1313. [https://doi.org/10.1016/S0140-6736\(07\)60368-7](https://doi.org/10.1016/S0140-6736(07)60368-7)
- Reardon, T., Harvey, K., Baranowska, M., O'Brien, D., Smith, L., & Creswell, C. (2017). What do parents perceive are the barriers and facilitators to accessing psychological treatment for mental health problems in children and adolescents? A systematic review

- of qualitative and quantitative studies. *European Child & Adolescent Psychiatry* 26(6): 623–647. <https://doi.org/10.1007/s00787-016-0930-6>
- Reardon, T., Harvey, K., & Creswell, C. (2019). Seeking and accessing professional support for child anxiety in a community sample. *European Child & Adolescent Psychiatry* 29(5): 649-664.
- Rickwood, D., Deane, F. P., Wilson, C. J., & Ciarrochi, J. (2005). Young people's help-seeking for mental health problems. *Australian E-Journal for the Advancement of Mental Health* 4(3): 218-251. <https://doi.org/10.5172/jamh.4.3.218>
- Rosenstock, I. M. (1977). The Health Belief Model and Preventive Health Behavior. *Health Education & Behavior* 2(4): 354-386. <https://doi.org/10.1177/109019817400200405>
- Salkovskis, P. M., Rimes, K. A., Warwick, H. M. C., & Clark, D. M. (2002). The Health Anxiety Inventory: development and validation of scales for the measurement of health anxiety and hypochondriasis. *Psychological Medicine* 32(5): 843–853. <https://doi.org/10.1017/S0033291702005822>
- Sayal, K., & Taylor, E. (2004). Detection of child mental health disorders by general practitioners. *British Journal of General Practice* 54(502): 348-352. [https://doi.org/10.1016/s0084-3954\(07\)70218-x](https://doi.org/10.1016/s0084-3954(07)70218-x)
- Skylstad, V., Akol, A., Ndeezi, G., Nalugya, J., Moland, K. M., Tumwine, J. K., & Engebretsen, I. M. S. S. (2019). Child mental illness and the help-seeking process: a qualitative study among parents in a Ugandan community. *Child and Adolescent Psychiatry and Mental Health* 13(1): 3. <https://doi.org/10.1186/s13034-019-0262-7>
- Starcke, K., & Brand, M. (2012). Decision making under stress: A selective review. *Neuroscience & Biobehavioral Reviews* 36(4): 1228–1248. <https://doi.org/https://doi.org/10.1016/j.neubiorev.2012.02.003>
- Stiffman, A. R., Pescosolido, B., & Cabassa, L. J. (2004). Building a model to understand youth service access: the gateway provider model. *Mental health services research* 6(4): 189–198. <https://doi.org/10.1023/b:mhsr.0000044745.09952.33>

Tripathy, J. P. (2013). Secondary Data Analysis: Ethical Issues and Challenges. *Iranian Journal of Public Health* 42(12): 1478–1479.

<https://www.ncbi.nlm.nih.gov/pubmed/26060652>

Wilson, C. J., Deane, F. P., Ciarrochi, J., & Rickwood, D. (2005). Measuring Help-Seeking Intentions: Properties of the General Help-Seeking Questionnaire. *Canadian Journal of Counselling* 39(1): 15–28.