

Accepted Manuscript

BJGP OPEN

Online and telephone access to general practice: a cross sectional patient survey

Bryce, Carol; O'Connell, Matthew; Dale, Jeremy; Underwood, Martin; Atherton, Helen

DOI: <https://doi.org/10.3399/BJGPO.2020.0179>

To access the most recent version of this article, please click the DOI URL in the line above.

Received 26 November 2020

Revised 04 March 2021

Accepted 08 March 2021

© 2021 The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0/>). Published by BJGP Open. For editorial process and policies, see: <https://bjgpopen.org/authors/bjgp-open-editorial-process-and-policies>

When citing this article please include the DOI provided above.

Author Accepted Manuscript

This is an 'author accepted manuscript': a manuscript that has been accepted for publication in BJGP Open, but which has not yet undergone subediting, typesetting, or correction. Errors discovered and corrected during this process may materially alter the content of this manuscript, and the latest published version (the Version of Record) should be used in preference to any preceding versions

Online and telephone access to general practice: a cross sectional patient survey

Carol Bryce MA (Hons), MA, PhD¹, Matthew DL O'Connell, PhD², Jeremy Dale MB, BS, PhD, FRCGP¹, Martin Underwood, FRCGP MD³, Helen Atherton, BSc (Hons), MSc, MPH, PhD¹

¹ Unit of Academic Primary Care, Warwick Medical School, University of Warwick, Coventry, UK

² Department of Population Health and Environmental Sciences, King's College London, London, UK

³ Warwick Clinical Trials Unit, Warwick Medical School, University of Warwick, Coventry, UK

For correspondence: Carol Bryce, C. Bryce.1@warwick.ac.uk

Accepted Manuscript - BJGP Open - BJGP2020.0179

Abstract

Background

Improving access to primary healthcare in the United Kingdom has focused on the use of telephone and online access but little is known about how awareness of and use varies between different patient groups.

Aim

To determine how patients are interacting with telephone and online channels for accessing general practice services and information, and to analyse how this varies according to patient characteristics and health status.

Design and setting

A cross sectional self-administered survey of adult patients in general practices across the West Midlands, UK.

Method

Descriptive statistics were used to show participants' awareness of and interaction with online information sources and remote access. Multivariable logistic regression was used to model the relationships between demographic and health characteristics and awareness and use of online services and alternatives to face to face consultations (e.g. telephone).

Results

2789 patients (response rate 19.0%) from 43 general practices participated. 60.8% (1651/2715) of participants were aware of online services and 30.3% (811/2674) reported having used one. Daily internet usage and frequently visiting the GP showed the strongest associations with knowledge and use of online services.

Conclusion

We have shown that there is the potential for inequitable awareness and use of telephone and online services in general practice populations. Given that their use has greatly increased due to the COVID-19 pandemic future service design will need to ensure this is taken into account.

Keywords

Digital health, primary care, inequalities

How this fits in

- Evidence on how patients interact with telephone and online channels for accessing general practice services and information is lacking.

- We have shown that prior to the pandemic, awareness and use of telephone and online channels of access in the UK was higher in certain groups within the population. Less frequent internet use and not attending the general practice were associated with lower awareness and use of online services. When accounting for this we have also shown differences according to factors including age, education level and deprivation.
- The onset of the pandemic led to rapid introduction of telephone and video consultation and there remains an urgent need for strategies to avoid exacerbating existing inequalities as we move forward.

Introduction

Communications technology is seen as a tool for improving access to primary healthcare. [1-4] In the United Kingdom (UK) there has been a recent policy drive for increased use of digital services. [4, 5] However, before the COVID-19 pandemic adoption by general practices was slow. [6-8] At the onset of the pandemic the UK National Health Service (NHS) rapidly implemented telephone triage and remote consultation (telephone, online, video) in response, as a way to reduce the number of face-to-face consultations. [9] GP face-to-face consultations dropped during March 2020 from 80.5% to 51%. [10] As health services plan for the future, it is important to consider benefits [11] and consequences that may be associated with rapid implementation of remote access.

From 2019 it was mandatory for GPs in England to provide patients with online appointment booking, online repeat prescriptions, and access to their medical records online. [12] Data from early 2020 show fewer than a third of patients (29.6%) were registered to use any of these online services [13] and uptake was low (ordering prescription online (18.8%), booking appointment online (18.1%) and requesting access to patient record online (5.8%)). [14] In the UK most individuals are connected to the internet [15] but frequency of use and ability to use it varies greatly. Some groups may be disadvantaged by a move to online access e.g. older adults, those with a disability or long term health condition, lower socioeconomic groups, migrants and ethnic minorities. [16]

We examined online and telephone access to general practice services by conducting a cross-sectional survey of patients registered at general practices in the West Midlands, UK. We determined how patients were interacting with telephone and online channels for accessing general practice services and information, and analysed how this varied according to patient characteristics and health status. The survey now also provides baseline data before the COVID-19 pandemic.

Methods

Study Design

We conducted a cross-sectional, self-administered survey to adults registered at general practices across the West Midlands, UK. The survey was administered between February and June 2019

Survey instrument

The survey had 32 questions. It was designed to collect data on demographic and health factors known to influence whether and how patients access online services and/or general practice as well as on awareness and use of online services.[17-20] We used validated questions from the NHS general practice patient survey [21] (on internet usage, caring responsibilities, respondent's experience of using their GP surgery, its website and getting an appointment and repeat prescription) and the Office for National Statistics [22] (on participant characteristics including age, gender, ethnicity, education level, health status).

Questions relating to access, including knowledge and use of alternatives to a face-to-face consultation and private online providers (outside of NHS), were devised by the study team, drawing on surveys conducted in this field and other related work. [23,24]

The survey was piloted by conducting five cognitive interviews with members of the public to check understanding, particularly in relation to the questions devised by the study team. [25]

Sampling and recruitment

We aimed to sample approximately 15,000 people. Based on estimates from previous surveys conducted in general practice we expected to see a 20% response rate. [26,27] Patients were sampled at the general practice level. We aimed to recruit 40 practices selected to ensure the inclusion of a wide range of patients, sampling general practices purposively to ensure variation in rural/urban location, list size, deprivation score and proportion of patients in ethnic minority groups. Sampling of patients was proportional to general practice list size, achieved by randomly sampling 5% of eligible patients registered at each practice. Patients were excluded if they were under 18 years, at the end of life or lacked capacity to consent to participate in the survey. An index of multiple deprivation score was assigned to participants within each practice based on the area score

for location of that practice. [28] This scores practices from 1-10 with a score of 1 indicating the most deprivation. [28]

Data collection

Patients were sent a paper copy of the survey with a postage paid return envelope. Participants were given the option to complete the survey online using a web link or a QR code on the front of the paper survey. The online survey was administered using the software package Qualtrics. [29] We sent a reminder letter to all sampled patients two weeks after the initial survey this thanked them for participating (where they had) and prompted a response where they had not.

Data Analysis

Descriptive statistics were used to show participants' awareness of and interaction with online information sources and remote access. Ethnicity was split into two categories, White, and Black And other Minority Ethnic Groups (BAME): it was not possible to split the categories further owing to small numbers of respondents from BAME groups. Multivariable logistic regression models were used to model the relationships between demographic characteristics (age, sex, ethnicity, education level, employment status), health characteristics (number of long term conditions, number of visits to GP in last year), other personal characteristics (level of internet use, caring responsibilities) and awareness and use of services. The predictor variables were selected a priori based on the factors known to influence whether and how patients access online services and/or general practice that comprised the questions in the survey. Univariable models for each variable were also constructed for comparative purposes (Supplementary tables S1-S4). These models included clustered standard errors to account for the survey design (individuals nested within practices). Participating general practices were contacted to determine what online and telephone services they offered. General practices not offering any particular online or telephone service featured on the survey were excluded from analysis for that service. Results are expressed as odds ratios (OR) and 95% confidence intervals (95% CI). The statistical software package Stata 15 was used. [30]

Results

In total, 43 general practices (7% of those in the West Midlands) participated; see Table 1. Of 14,694 patients sampled across the practices, 2789 (19.0%) responded, with 2413 (86.5%) returning the survey via post and 376 (13.5%) completing it online. All participating practices had a website, offered online booking of appointments and of repeat prescriptions. Telephone appointments or call back systems were available in 40/43 practices including six practices which operated a telephone

triage system after all on the day appointments were booked. No practices offered email or video consultation.

Respondents' characteristics are in Table 2. Around half of respondents (51.4%; 1360/2647) reported they had a long term physical or mental health condition, disability, or illness. Of these, 26.6% (680/2556) reported a physical health problem, 7.9% (201/2556) a sensory condition, 13.4% (343/2556) a mobility problem and 5.6% (143/2556) a mental health problem. The majority of participants (89.0%; 2447/2748) reported their experience of general practice as good and most had visited their practice at least once in the last 12 months (60.1%; 1652/2748). Patient awareness of their general practice website was high (85.0%; 2266/2666) with 42.1% (1147/2725) having accessed the website on at least one occasion. Awareness of alternatives to face-to-face consultations as a concept was low with 0.5% (12/2643) aware of email consultation and 0.5% (14/2643) of video consultation.

Awareness and use of online services

Table 3 outlines the percentage of patients aware of and using the different online services. In the multivariable model, we observed an association between awareness of online booking for appointments and daily internet usage (OR 4.39 [CI 2.98,6.46]), higher frequency of GP visits (OR 2.13 [CI 1.41,3.24]), having higher education qualifications (1.83 [CI 1.35,2.48]), having at least one long term condition (OR 1.35 [CI 1.10,1.66]), being retired (OR 1.51 [CI 1.09,2.08]), and being female (OR 1.5 [CI 1.24,1.80]) (Supplementary table 5). Usage patterns showed the same associations with daily internet usage (OR 14.92 [6.84,32.57]), higher frequency of GP visits (OR 10.64 [5.01,22.59]), having further education qualifications (OR 1.90 [1.33,2.73]), having at least one long term condition (OR 1.47 [1.20,1.81]), being female (OR 1.43 [1.14,1.79]). There was also a weak association between practice deprivation score and awareness of online appointment booking with higher deprivation associated with lower awareness (OR 1.09 [CI 1.02,1.15]), this was less clear for use (OR 1.05 [CI 0.99,1.12]).

We observed an association between awareness of online repeat prescriptions being offered and daily internet use (OR 5.14 [CI 3.74,7.07]), visiting the GP more than 5 times (OR 2.22 [CI 1.56,3.14]), having a long term condition (OR 1.81 [CI 1.50,2.18]), being retired (OR 1.54 [CI 1.17,2.03]), having education qualifications (OR 1.53 [CI 1.15,2.03]), being female (OR 1.34 [CI 1.12,1.61]), aged 55-64 years (OR 1.47 [CI 1.03,2.10]) and lower deprivation (OR 1.10 [CI 1.03,1.17]) (Supplementary table 6). Reported use of the repeat prescription service online showed the

association was strongest for those who visited the GP five or more times a year (OR 2.22 [CI 1.56,3.14]) and those who used the internet daily (OR [11.86 [6.57,21.42])). It was also significant for those with education qualifications that were not further or higher education (OR 1.79 [1.10,2.90]), caring 10+ hours a week (OR 1.51 [1.08,2.11]), being retired (OR 1.55 [1.11,2.16]) and lower deprivation (OR 1.11 [1.02,1.21]).

Awareness of online access to medical records was associated with using the internet everyday (OR 4.20 [CI 2.90,6.09]), visiting the GP more than 5 times a year (OR 3.29 [CI 1.80,6.03]), being older (aged 65-74 OR 1.77 [CI 1.24,2.52], having a long term condition (OR 1.38 [CI 1.12,1.70], being retired (OR 1.40 [CI 1.06,1.85], and lower deprivation (OR 1.18 [CI 1.07,1.30]). Being from a non-white ethnic group was also associated with awareness (OR 2.05 [CI 1.35,3.11]) (Supplementary table 7). Access was more strongly associated with use of the internet (OR 8.00 [CI 2.63,24.30]), frequently visiting the GP (OR 19.88 [CI 2.50,157.87]) and having a long term condition (OR 2.21 [CI 1.34,3.63]).

Awareness and use of telephone consultation

Telephone consultations were widely offered as an alternative to a face-to-face consultation. Overall, 55.7% of respondents (1471/2643) were aware of telephone consultation and 36.7% (987/2691) had used the service. Awareness of telephone consultation was associated with being female (OR 1.82 [CI 1.48,2.23]), frequent internet use (OR 1.81 [CI 1.27,2.59]), visiting the GP more than 5 times a year (OR 2.74 [CI 1.76,4.24]), having a long term condition (OR 1.40 [CI 1.12,1.75]), being a parent of a child under the age of 16 (OR 1.44 [CI 1.08,1.93]) and having an informal caring role of ten hours or more per week (OR 1.54 [CI 1.08,2.20]) (Supplementary table 8). Reported use of telephone consultations followed the same pattern, with higher education (OR 1.51 [CI 1.04, 2.20]) also being associated with use of telephone consultation.

Discussion

Summary

We show that, prior to the pandemic, in our study population, awareness and use of online primary care services was higher in individuals who use the internet daily and those who attend the general practice frequently. Having a long term condition, being female, and registered at a practice in an area of low deprivation were all associated with greater awareness and use of telephone and online services. We also demonstrated awareness and use of telephone and online access being associated with being retired and with higher education levels. Parents and carers were most likely to use

telephone access. We observed that awareness of the practice website and online services was higher than usage.

Strengths and limitations

The survey included a diverse range of practices. A 19.0% patient response rate is consistent with unsolicited community postal surveys. [31] However, it is likely that those who responded were more interested in the topic and may have differed in other ways to the non-responders therefore their responses cannot necessarily be extrapolated to the entire community.

Respondents were older than the resident population in the West Midlands (age 65+ survey 45.1%; West Midlands 18.6%) [32]. However, apart from high consultation rates in infants, consultation rates increase with increasing age from the lowest levels in the 15-24 age group.[8] Consultation rates are also higher in areas of high deprivation [33] and in this study deprivation score was only available at the practice level which lacked granularity. This means we cannot be confident that we engaged a wide enough range of patients across the deprivation gradient and that in controlling for deprivation in our models we used only an approximation for deprivation level. The views of ethnic minority groups are also under-represented in our findings, with the proportion of respondents from these groups lower than in the general population (White ethnicity, this survey 91.5%; West Midlands region 82.7% [34]). The findings of this study should be interpreted in light of these limitations.

The survey tool used validated questions from both the ONS and the GPPS, alongside questions developed to address the topic of access to general practice. Although we did not conduct additional validation of the survey as a whole, we used cognitive interviews with public contributors to check understanding of the survey instrument before we used it in our study population.

We sampled practices purposively and consequently we did not include practice level variables other than deprivation (as a proxy for patient deprivation level) in our models. We included clustered standard errors to account for the nesting of participants within practices. Practice level factors such as location (rural vs urban) often influence patient behaviour in general practice and so any future surveys should consider these practice level variables.

Comparison with existing literature

Our data found similar results to the national general practice patient survey by NHS England [17], with 89% of our respondents and 83% of the national survey rating overall experience of their general practice as good and 71.7% of our sample compared to 67% of the national sample rating their experience of making an appointment as good. This suggests our participants are broadly similar in views despite our sample being smaller and skewed towards older people.

We found that online services, including online booking, repeat prescription ordering and access to patient records, were available at the majority of the general practices at the time of our survey but uptake was limited. This is in line with national data which shows that in January 2020 less than a third of patients in England (29.6%) were registered for at least one online service. [35,36] By 31st December 2020, the number had risen to 32.42%, [35] a modest increase given that the majority of contacts with UK general practices became remote during the COVID-19 pandemic.[37] Previous research shows that uptake of online or digital services remain low due to a lack of awareness or engagement within practice populations alongside staff reservations and understanding about appropriate use. [38,39] It is also known that deprivation has an impact on service quality, satisfaction, and usage. [40,41]

Implications for research and practice

The COVID-19 pandemic markedly increased levels of use of telephone consultation and online consultation and it looks likely that the rapid move to using online methods of contact and access will be retained after the pandemic to a certain degree. [9, 11] Our sample was comprised of mostly older participants with white ethnicity, who were retired. It is likely that awareness and use of online and telephone services is even lower in groups known to be disadvantaged in relation to accessing services; people who use the internet less frequently, those who visit the general practice infrequently, older people, those with lower levels of education and those who live in areas of high deprivation. Practices should raise awareness of services available, being cautious to avoid assumptions over how patients get their information. Awareness of the general practice website was high but less than half of our sample had used it, so this is unlikely to provide a primary information source for patients about what services are available, nor does information provision alone lead to use. Patients may require support to both learn about and use services that provide alternative routes of access.

Patients who regularly attend the practice differ in relation to awareness and use of online services relative to those who attend less frequently if at all, and so different approaches are likely to be needed depending on the patient.

The burden of COVID-19 ill health and economic disadvantage is not equally distributed across communities [42-44] and may be hitting hardest where awareness and use of online and telephone access is lowest, risking inequality. Future research studies should explore the role of the practice, as well as patient characteristics, in determining awareness and use of services. They should also aim to produce practical recommendations about what can be done in everyday practice to ensure patients have a suitable route of access available to them.

Ethical approval

Ethical approval was granted by North East -York Research Ethics Committee (reference number 18/NE/0333)

References

1. Danish Government Den digitale vej til fremtidens velfærd 2011 Available from: <https://horsensbibliotek.dk/om-bibliotekerne/projekter/den-digitale-vej-til-fremtidens-velfaerd>. 8 Feb 2017
2. Danish Medical Association. aktivitet-og-oekonomi-i-almen-praksis [Activity and Economy in General Practice] 2017 <https://www.laeger.dk/nyhed/aktivitet-og-oekonomi-i-almen-praksis>. 16 Jan 2018
3. Pearl R. Kaiser Permanente Northern California: current experiences with internet, mobile, and video technologies. Health Aff (Millwood) 2014 33(2):251-7.
4. Royal College of General Practitioners. The 2022 GP: a vision for general practice in the future NHS. (2013) London: RCGP
5. NHS England. Investment and evolution: A five-year framework for GP contract reform to implement The NHS Long term Plan. 2019 <https://www.england.nhs.uk/wp-content/uploads/2019/01/gp-contract-2019.pdf>
6. Whitfield L. Netscape navigators 2013 <https://www.digitalhealth.net/2013/02/netscape-navigators/> 8 Feb 2017
7. British Medical Association. The future of general practice - survey results 2015 <http://bma.org.uk/working-for-change/negotiating-for-the-profession/bma-general-practitioners-committee/surveys/future-of-general-practice> 8 Feb 2017.
8. Hobbs FDR, Bankhead C, Mukhtar T, et al. Clinical workload in UK primary care: a retrospective analysis of 100 million consultations in England, 2007-14. Lancet. 2016 387(10035):2323-30.
9. Thornton, J., COVID-19-19: how coronavirus will change the face of general practice forever. BMJ. 2020 368.
10. NHS Digital Appointments in General Practice March 2020 <https://digital.nhs.uk/data-and-information/publications/statistical/appointments-in-general-practice/march-2020> 1 Oct 2020
11. Duckett, S. What should primary care look like after the COVID-19-19 pandemic? Aust J Prim Health. 2020 May 27.

12. England, N.H.S. and British Medical Association. Investment and evolution: a five-year framework for GP contract reform to implement the NHS long term plan. 2019 London: NHS England. <https://www.england.nhs.uk/wp-content/uploads/2019/01/gp-contract-2019.pdf>
13. NHS Digital Appointments in General Practice January 2020, 2020
<https://digital.nhs.uk/data-and-information/publications/statistical/appointments-in-general-practice/january-2020> 27 Aug 2020
14. Department of Health. NHS. The GP Patient Survey. 2020
file:///C:/Users/u1472402/Downloads/GPPS_2020_National_report_PUBLIC.pdf 1 Oct 2020
15. Office for National Statistics (ONS) 2019
<https://www.ons.gov.uk/businessindustryandtrade/itandinternetindustry/bulletins/internetusers/2019> 27 Aug 2020
16. Public Health England. Local action on health inequalities: Improving health literacy to reduce health inequalities. 2015
https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/460709/4a_Health_Literacy-Full.pdf 27 Aug 2020
17. Ford JA, Turley R, Porter T, Shakespeare T, Wong G, Jones AP, et al. Access to primary care for socio-economically disadvantaged older people in rural areas: A qualitative study. *PLoS One*. 2018;13(3):e0193952.
18. Burt, J., Lloyd, C., Campbell, J., Roland, M. and Abel, G., 2016. Variations in GP-patient communication by ethnicity, age, and gender: evidence from a national primary care patient survey. *British Journal of General Practice*, 66(642):e47-e52.
19. Huntley AL, Johnson R, Purdy S, Valderas JM, Salisbury C. Measures of multimorbidity and morbidity burden for use in primary care and community settings: a systematic review and guide. *Ann Fam Med*. 2012;10(2):134-41
20. Stormacq, C., Van den Broucke, S. and Wosinski, J., 2019. Does health literacy mediate the relationship between socioeconomic status and health disparities? Integrative review. *Health promotion international*, 34(5): e1-e17.
21. Ipsos Mori. *GP Patient Survey 2017*, <https://gp-patient.co.uk/>.
22. UK Data Service Opinions and Lifestyle Survey, Internet Access Module, January, February and April, 2017. 2017
<https://beta.ukdataservice.ac.uk/datacatalogue/studies/study?id=8299&type=Data%20catalogue> 27 Aug 2020
23. Brant H, Atherton H, Ziebland S, et al. Using alternatives to face-to-face consultations: a survey of prevalence and attitudes in general practice. *Br J Gen Pract*. 2016 66(648):e460-6.
24. van Velthoven MH, Atherton H, Powell J. A cross sectional survey of the UK public to understand use of online ratings and reviews of health services. *Patient Educ Couns*. 2018 1;101(9):1690-6.
25. Drennan J. Cognitive interviewing: verbal data in the design and pretesting of questionnaires. *J Adv Nurs*. 2003 42(1):57-63.
26. Peeters MJ, Krijgsman WJ, Brabers EA, et al. Use and Uptake of eHealth in General Practice: A Cross-Sectional Survey and Focus Group Study Among Health Care Users and General Practitioners. *JMIR Med Inform*. 2016 4(2):e11.
27. Newbould J, Abel G, Ball S, et al. Evaluation of telephone first approach to demand management in English general practice: observational study. *BMJ*. 2017 358.
28. Ministry of Housing, Communities & Local Government The English Indices of Deprivation (IoD2019) 2019
https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/835115/IoD2019_Statistical_Release.pdf 27 Aug 2020
29. Warwick IT services. Software: Qualtrics 2018. Available from:
<https://warwick.ac.uk/services/its/servicessupport/software/list/qualtrics>.
30. StataCorp. *Stata Statistical Software: Release 15*. 2017 College Station, TX: StataCorp LLC.

31. Sinclair M, O'Toole J, Malawaraarachchi M and Leder K. Comparison of response rates and cost-effectiveness for a community-based survey: postal, internet and telephone modes with generic or personalised recruitment approaches. *BMC Med Res Methodol.* 2012 12(1):132
32. Office for National Statistics
<https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationestimates/bulletins/annualmidyearpopulationestimates/mid2019estimates> 27 Aug 2020
33. Mukhtar, T.K., Bankhead, C., Stevens, S., Perera, R., Holt, T.A., Salisbury, C. and Hobbs, F.R., 2018. Factors associated with consultation rates in general practice in England, 2013–2014: a cross-sectional study. *Br J Gen Pract*, 68(670): e370-e377.
34. Office for National Statistics
<https://www.ons.gov.uk/peoplepopulationandcommunity/culturalidentity/ethnicity/articles/ethnicityandnationalidentityinenglandandwales/2012-12-> 1 Oct 2020
[11#:~:text=The%20West%20Midlands%20was%20the,group%20at%2079.2%20per%20cent.&text=The%20West%20Midlands%20had%20a,Caribbean%20at%201.5%20per%20cent.](https://www.ons.gov.uk/peoplepopulationandcommunity/culturalidentity/ethnicity/articles/ethnicityandnationalidentityinenglandandwales/2012-12-1#:~:text=The%20West%20Midlands%20was%20the,group%20at%2079.2%20per%20cent.&text=The%20West%20Midlands%20had%20a,Caribbean%20at%201.5%20per%20cent.)
35. Digital N. Patient online management information (POMI) 2019. <https://digital.nhs.uk/data-and-information/data-tools-and-services/data-services/general-practice-data-hub/patient-online-management-information-pomi> 27 Aug 2020
36. Edwards HB, Marques E, Hollingworth W, et al Use of a primary care online consultation system, by whom, when and why: evaluation of a pilot observational study in 36 general practices in South West England. 2017 *BMJ Open*.7:e016901.
37. Murphy, M., Scott, L.J., Salisbury, C., Turner, A., Scott, A., Denholm, R., Iyer, G., Lewis, R., MacLeod, J. and Horwood, J., 2021. The implementation of remote consulting in UK primary care following the COVID-19 pandemic: a mixed-methods longitudinal study. *Br J Gen Pract BJGP*.2020.0948
38. Atherton H, Brant H, Ziebland S, Bikker A, Campbell J, Gibson A, McKinstry B, Porqueddu T, Salisbury C. Alternatives to the face-to-face consultation in general practice: focused ethnographic case study. *British Journal of General Practice*. 2018 Apr 1;68(669):e293-300.
39. Rodgers M, Raine GA, Thomas S, Harden M, Eastwood AJ. Informing NHS policy in 'digital-first primary care': a rapid evidence synthesis. *Health Services and Delivery Research*. 2019:1-54.41.
40. Fisher R, Dunn P, Asaria M, and Thorlby R. Level or not? 2020. (<https://doi.org/10.37829/HF-2020-RC13>)
41. Fisher, R and C. Fraser [Who gets in? | The Health Foundation](https://www.health.org.uk/news-and-comment/charts-and-infographics/who-gets-in) 2020
<https://www.health.org.uk/news-and-comment/charts-and-infographics/who-gets-in>
42. Bamba, C., Riordan, R., Ford, J., Matthews, F. The COVID-19 pandemic and health inequalities. *J Epidemiol Community Health*. 2020 June 12
43. Douglas, M., Katikireddi, S.V., Taulbut, M., et al. Mitigating the wider health effects of COVID-19-19 pandemic response. *BMJ*. 2020 369.
44. Khilnani, A., Schulz, J. and Robinson, L. The COVID-19 pandemic: new concerns and connections between eHealth and digital inequalities. *J. Inf. Commun. Ethics Soc.* 2020 June 25.

Table 1 Practice characteristics

Location	Practices N=43
Urban	32
Rural	11
List size (number of patients)	
Small <6000	16
Medium 6000-12000	20
Large > 12000	7
Deprivation (IMD 2019) <small>1=more deprived</small> <small>https://www.gov.uk/government/statistics/english-indices-of-deprivation-2019</small>	
Score 1-3	14
Score 4-7	17
Score 8-10	12
Ethnicity % BAME <small>West Midlands BAME 17.4% 2011 Census</small>	
Highest 82% Lowest 1%	
<5%	16
5-10%	7
11-20%	5
21-50%	10
>50%	5

Accepted Manuscript - BJGP Open - BJGPO.2020.0179

Table 2: Characteristics of respondents

		N	%
Age N=2756	18-34	248	9.0
	35-54	680	24.6
	55-64	589	21.4
	65-74	723	26.2
	75+	516	18.9
Gender N=2748	Male	1181	42.8
	Female	1567	56.8
	Prefer not to say	9	0.3
Ethnic group N=2728	White	2496	91.5
	Black and other Minority Ethnic Groups	232	8.5
Education N=2657	No formal qualifications	419	15.8
	Secondary level	386	14.5
	Further	588	22.1
	Higher	1115	42.0
	Still Studying	18	0.7
	Other	131	4.9
Caring responsibilities Parent or guardian for any young person aged under 16 years living at home N=2706	Yes	402	14.9
	No	2304	85.1
Caring responsibilities for family or friends N=2795	No	2112	78.4
	Yes (Up to 9 hours/week)	334	12.4
	Yes (10+ hours/week)	249	9.2
Health status: long term physical or mental health conditions, disabilities or illnesses. N=2647	Yes	1360	51.4
	No	1287	48.6
Frequency of using general practice services in last 12 months N=2748	0	255	9.3
	1-5 times	1652	60.1
	5+	841	30.6
Internet use in the last 3 months N=2646	Everyday	1788	67.6
	<Everyday	420	15.9
	Never	438	16.6

Table 3: Awareness and use of online services

	Awareness N=2715		Use N=2674	
	N	%	N	%
Any online service	1651	60.8	811	30.3
Booking appointments	1479	54.5	552	20.7
Ordering repeat prescriptions	1362	50.2	584	21.9
Accessing medical records	629	23.2	172*	7.8
	Awareness N=2628		Use N=2697	
Private online general practice services	657	25.0	126	4.7
Private online GP consultations	451	17.2	32	1.2

* For this variable N=2207

Accepted Manuscript - BJGP Open - BJGPO-2020.0179