

Análise dos acidentes de trabalho na equipe de enfermagem: uma revisão integrativa

Analysis of the accidents at work in the nursing staff: an integrative review

Análisis de los accidentes de trabajo en el equipo de enfermería: una revisión integrativa

Patrick Leonardo Nogueira da Silva¹, Larissa Mendes Carvalho², José Márcio Girardi de Mendonça³, Renata Patrícia Fonseca Gonçalves⁴, Simone Guimarães Teixeira Souto⁵, Marcelo Rocha Torres⁶

How to quote this article:

Silva PLN; Carvalho LM; Mendonça JMG, et al. Analysis of the accidents at work in the nursing staff: an integrative review. Rev Fund Care Online. 2016 out/dez; 8(4):5163-5176. DOI: <http://dx.doi.org/10.9789/2175-5361.2016.v8i4.5163-5176>

ABSTRACT

Objective: To analyze work-related injuries in nursing staff. **Method:** Integrative review in which the sample was obtained by articles of national and international journals that were available on the SciELO and LILACS databases, between 2007-2011. **Results:** We found 212 publications, of which 18 were selected after applying the inclusion criteria. A large number of publications were found in SciELO (61%) and there was equality in the number of publications in the years 2007-2009 and 2011 (22.2%). After examination, it was noticed that the nursing staff is very struck by accidents, and that the sharps are the main villains in this sense, and biological material is also very present. **Conclusion:** It is concluded that not only the workers should have care knowledge in their workplace, but also the health institutions have to meet their obligations in order to protect the workers.

Descriptors: work accidents, nursing crew, occupational risks.

¹ Nurse, Occupational Health Nursing Specialist, University of Montes Claros / UNIMONTES, Montes Claros (MG), Brazil. E-mail: Patrick_mocesp70@hotmail.com.

² Nurse, University of Montes Claros / UNIMONTES, Montes Claros (MG), Brazil. E-mail: laramc18@hotmail.com.

³ Nurse, Master nursing professor of the State University of Montes Claros / UNIMONTES, Montes Claros (MG), Brazil. E-mail: girardidemendonca@terra.com.br.

⁴ Nurse, Master nursing professor of the State University of Montes Claros / UNIMONTES, Montes Claros (MG), Brazil. E-mail: renatapfonseca@yahoo.com.br.

⁵ Nurse, Master nursing professor of the State University of Montes Claros / UNIMONTES, Montes Claros (MG), Brazil. E-mail: simonegts28@yahoo.com.br.

⁶ Nurse, Specialist in Emergency, Trauma and Critical Care, University of Montes Claros / UNIMONTES, Montes Claros (MG), Brazil. E-mail: mrtorres26@yahoo.com.br.

RESUMO

Objetivo: Analisar os acidentes de trabalho na equipe de enfermagem.

Métodos: Revisão integrativa na qual a amostra de artigos captados estava disponível em periódicos nacionais e internacionais e se encontravam nas bases do SCIELO e LILACS, durante os anos de 2007-2011. **Resultados:** Encontrou-se 212 publicações, sendo que 18 foram selecionados após aplicação dos critérios de inclusão. Encontrou-se um maior número de publicações na SCIELO (61%) e houve igualdade no número de publicações nos anos de 2007-2009 e 2011 (22,2%). Após a análise, percebeu-se que a equipe de enfermagem é bastante atingida pelos acidentes de trabalho, e que os materiais perfurocortantes são os principais vilões nesse sentido, e o material biológico se faz bastante presente. **Conclusão:** Conclui-se que não somente os trabalhadores devem ter a ciência dos cuidados em seu local de trabalho, mas também as instituições de saúde em cumprir suas obrigações visando à proteção do trabalhador.

Descritores: Acidentes de trabalho, Equipe de enfermagem, Riscos ocupacionais.

RESUMEN

Objetivo: Analizar las lesiones relacionadas con el trabajo del personal de enfermería. **Método:** Revisión integradora en la que se obtuvo la muestra de artículos disponibles en revistas nacionales e internacionales en las bases SciELO y LILACS, durante los años 2007-2011. **Resultados:** Se encontraron 212 publicaciones, de las cuales 18 fueron seleccionadas después de aplicar los criterios de inclusión. Se reunió un gran número de publicaciones en SciELO (61%) y hubo igualdad en el número de publicaciones en los años 2007-2009 y 2011 (22,2%). Después del examen, se observó que el personal de enfermería es muy atingido por los accidentes, y que los objetos punzantes son los principales villanos en este sentido, el material biológico está muy presente. **Conclusión:** Se concluye que no sólo los trabajadores deben tener la ciencia del cuidado en su lugar de trabajo, sino también las instituciones de salud para cumplir con sus obligaciones con el fin de proteger al trabajador.

Descriptor: Accidentes de Trabajo; Equipo de Enfermería; Riesgos Ocupacionales.

INTRODUCTION

For some theorists of the twentieth century and early twenty-first century, the work has a great value in all the writings of the philosopher Karl Marx, being the life-affirming activity that forms the existence of individuals and establishes them as social characters. It is the work that expresses human superiority against other living beings. It would be the realization of man himself, the source of all wealth and material goods.¹ Work in health is essential for human life. It is understood work as a transforming action, and health specific to the identity of nature between subjects receiving care and caregivers, in addition to the inseparability of the production process and the product of labor.² When performing any type of work, people are exposed to occupational hazards and possible work accidents (AT).

Occupational hazards are defined as: Health risks or the worker's life due to their occupational activities.³ Concern about the AT is old. There are reports of its existence since

the time before Christ, Greek and Roman empires, in which diseases at work already occurred with slaves and servants. In Brazil, this concern is more recent, being translated by the law promulgated on 15 January 1919. The issue of AT nursing gained greater concern from the 1980s.⁴

AT is a sudden event that occurs in the course of labor activity, which causes damage to health, potential or immediate, and causes injury or functional disorder, which can cause death or permanent loss or temporary reduced ability to work. This includes even the accident in any situation where the employee is representing the interests of the company or acting in defense of its assets, in addition to occurring during the journey between the workplace and the home or otherwise.⁵

The Regulatory Standard (NR) 32, the Ministry of Labor and Employment (MTE) of 11 November 2005, created by Ordinance N° 485, establishing measures to protect the health and safety of health workers in any health service. This standard calls for hand hygiene, vaccination of professionals for hepatitis B (HB), tetanus and diphtheria. Determine also some situations in regard to clothing and changing facilities, waste, continuous and ongoing training in the specific area of expertise, among other provisions. In addition to this standard, the MTE produced the Ordinance N° 939 of 18 November 2008, which led employers to promote the replacement of sharps by other safety devices within 24 months from the date of publication of this Regulation. In 2011, the MTE repealing Ordinance N° 939 with then Ordinance N° 1748 of 30 August 2011, the employer shall devise and program plan Accident Risk Prevention with sharps, and the companies that produce or market sharps shall make available to the workers of health services, training on proper use of the safety device.⁶

In the last two decades, occupational accidents involving biological materials with the workers in the health sector have been gaining ground in the global survey, since exposure to pathogens that are linked to blood can lead professionals to contract infections, and the consequent harm to Cheers. As a historical example, one can refer to the first case occupational acquisition of *Human Immunodeficiency Virus* (HIV) occurred in England in 1984, after a nurse have been contaminated by a needle containing infected patient blood.⁷ Nurses play a direct relief work and continued to the patient, making it susceptible to contamination by biological material, especially regarding accidents that occur by percutaneous inoculation mediated needles or sharp instruments, which are the most responsible for occupational transmission of blood-borne infections.⁸

This study aims to analyze the scientific publications on AT in the Nursing Team in national and international journals available in the Virtual Health Library (VHL), on the basis of the *Scientific Electronic Library Online* (SciELO) and the Latin American and Caribbean Health Sciences (LILACS), from 2007 to 2011.

Thus, the identification of studies related to AT can contribute to nursing professionals use measures in order to prevent/avoid these occupational accidents that occur in the workplace. These measures are also highlighted by other authors⁹ suggesting the importance of programming more effective measures to diminish the AT and updates courses, educational programs, clinical follow-up of exposure, vaccination and use of safety devices to perform invasive procedures.

Theoretical framework

The OA can be classified as typical or path. Typical AT occurs during work performance; Path AT happens during displacement from the residence to the workplace. There is also the occupational disease, defined as the one which is produced or triggered by the exercise inherent to the profession and occupational disease that is acquired or triggered by special conditions where the work is performed and with which it relates. It is noteworthy also the duty that every AT must be registered with the competent social security body, using for this purpose the Communication of Work Accident (CAT).¹⁰

From the 1980s, the AT in the nursing team have been gaining ground in the global research.⁷ Regarding legislation, the first way to protect the workers in relation to AT was through AT insurance institution in 1919.¹¹ From these constant and growing health concerns of the health worker, laws have emerged, ordinances NR to professionals be protected for accidents at work.

Historically, health professionals were not considered as a category that had high risk for AT. Concern about biological risks arose from the aggravation of finding health professionals working in the laboratory, which was handling both microorganisms as clinical material, and this since the 1940s.¹² Among the various types of accidents, the sharps are not only more frequent but also more serious, as they allow the development of life-threatening diseases for workers.¹³ This is also highlighted in another study in which the accident sharps is a major accident occurs with the team of nursing. It was noted in the study that needlestick injuries are in the range of 79.87%; accidents with blood, organic material involved 86.58% and 59.06% lumen needles.¹⁴

In São Paulo, for example, of the accidents notifications with biological material in health professionals, 86% involved percutaneous exposure and 82% blood.¹⁵ In relation to the most affected sites at the time of the accident, there are the hands, followed eyes, legs, trunk and other parts.¹⁶

Possible causes of these accidents with the nursing staff may be related to the unavailability/inappropriateness of personal protective equipment (PPE), work overload, lack of training on the correct use of biosecurity measures exist to be performed, as well as own. Feeling of invulnerability and wrong habit of recapping needles contaminated by some workers.¹⁷ The main causes attributed to the occurrence

of AT by these sharps are: disposal in inappropriate places or in overcrowded containers, transport or handling of unprotected needles and needle off the syringe, but the main associated factor is the recapping of needles.¹⁸

With the enactment of the Federal Constitution (FC) 1988, it was established the current Unified Health System (SUS), which separated the health of the pension plan, and this would be responsible for concessions and management of retirement and pension AT insurance. This role of Social Security appears in Section III, Article 201 of the Constitution of 1988.¹⁹ On 19 setembro 1990 was drafted Law N° 8080, which provides for the conditions for the promotion, protection and recovery of health, the organization and operation of corresponding services and other measures. In its Article 5 establishes as SUS playing field executions of health surveillance, epidemiological surveillance, worker health. It also aims at the recovery and rehabilitation of health workers victims of risks and hazards from working conditions.²⁰

Through Ordinance N° 1679 of 19 September 2002, the National Network of Care for Occupational Health (RENAST).²¹ Ordinance N° GM was created. 2,437, of December 7, 2005, provides for the expansion and strengthening of RENAST and other measures, such as supporting the organization and structuring of average attendance and high complexity, local and regional, which meets the AT and injuries that are on the list of diseases and disorders related to constant work in GM Ordinance N° 1339 of 18 November 1999, and also to the notifiable diseases contained in Ordinance N° GM. 777 of 28 April 2004.²²⁻²⁴

On 11 November 2005, the MTE, through Ordinance N° 485, creates the NR-32 aimed at establishing basic guidelines to plan measures to protect the safety and health of the health care workers and those who carry out promotion and health care activities in general. Calls, therefore, the use of PPE; hand hygiene; HB vaccination, tetanus and diphtheria. For this standard complement the purpose of establishing the Ordinance N° 939 of 19 November 2008 establishing maximum period of 24 months from the date of publication of this ordinance, for companies to do the replacement of sharps by others with safety device. In 2011 the MTE repealing Ordinance N° 939 with then Ordinance N° 1748 of 30 August 2011, the employer shall devise and program plan Accident Risk Prevention with sharps, and the companies that produce or market sharps shall make available to the workers of health services, training on proper use of the safety device.²⁵⁻²⁷

According to law N° 8213, 24 July 1991 all AT must be registered in the relevant social security instance, and must use the CAT for this purpose.²⁸ One way to avoid contamination by pathogens via the blood is to use EPI, forming protective barriers reducing contact with organic materials. The basic PPE are gloves, masks, gowns, cap and eye protection. Besides the use of such equipment, the care with immunization should be followed by the vaccine,

care for the environment. In addition, other approaches are essential for individual protection, as well as hand washing.²⁹ The basic objective of a precaution system is to prevent the transmission of a micro-organism from one patient to another, or to a health professional. This prevention encompasses measures concerning the transmission of the agents involved.³⁰

METHODS

This is an integrative literature review which theme was AT the Nursing Team, presents itself also as a study of descriptive and exploratory.

The exploratory research aims to exhibit characteristics of a given population or particular phenomenon. In turn, can also establish correlations between variables and define its nature.³¹ The descriptive research aims at the description of characteristics of a given population or phenomenon or to establish relationships between variables.³² The integrative literature review consists of building an extensive literature review, contributing to discussions on methods and results of research, as well as reflections on the future studies. The initial objective of the study is to deeply understand a certain phenomenon based on previous studies.³³ The term “integrative” comes from the integration of opinions, concepts or ideas from the surveys used in the method.³⁴

The literature integrative review is divided into six phases. In the first phase is to identify the theme or question the integrative review, which is prepared to issue the theme delimited search for integrative review, and then the keywords are determined for the studies search strategy. The second phase is the sampling or literature search, since the problem or theme is set, the search begins in the literature. The key element for the proper conduct is a comprehensive literature search. After finishing reading titles and abstracts, should be items selected. Aiming at a complete work, it is important that the studies are analyzed in full and that the search is as complete as possible. The third phase is the categorization of studies involving the preparation or use of a data collection tool that has been validated, which aims to extract the key information for each item that was selected. The fourth phase is the evaluation of the studies included in the integrative review. At this stage the items that were selected are reviewed in relation to the authenticity criteria, methodological quality, importance of information and representation.³⁵

The fifth stage is to discuss the results at this stage after the interpretation and synthesis of the results, the data disclosed are compared in the analysis of articles to the theoretical framework.³⁶ The sixth phase, and final, is the presentation of the review/synthesis knowledge, it is said that the review should allow the replication of the study. In this last step should be drawn up a document that should include a description of all the stages covered by the researcher, judiciously, and should present the main results obtained.³⁴

This last step includes the data visualization. The display mode can be expressed in tables, graphs or charts, in which the comparison between all the selected studies is possible, and therefore, the identification of patterns, differences and subletting these topics as part of the general discussion.³⁶

To survey the articles to be used, it was carried out a search in the Virtual Library database in Health (BVS), and initially without selection of titles or sources, and thus the articles that were selected for the study belonged to the database *Scientific Electronic Library online* (SCIELO) and the Latin American and Caribbean Health Sciences (LILACS). To carry out the research the following key words were used: “AT and Nursing Team” and “Occupational Risks and Nursing Team.”

The study population was found in publications contained such data in the banks, and was selected the years from 2007 to 2011, being the most recent past years. A total of 212 articles were found, and of these, 103 were with the keywords “Occupational Risks and Nursing Team” and 109 with “AT and Nursing Team.” They were first read all the titles and abstracts of articles. The following criteria for inclusion in the study were used: articles published between 2007-2011 that were in Portuguese, English and/or Spanish and articles that have the full text.

93 were related articles with both keywords and full text. Of these, 18 articles were selected for meeting all inclusion criteria, including the years referred to preparation of the final work, and found seven in LILACS database and eleven found in SCIELO. In the search it was found two articles in English, but we managed to open them in Portuguese; an article was in Spanish, Portuguese and English and the rest in Portuguese.

The search for articles was conducted in October 2012, to collect data. For data analysis it was constructed a table with eight items, which included: title, author(s), year, periodic, objective, methods, results and conclusion. When the data were collected, the articles were passed to the tables, which provided opportunity to analyze and relate these articles so that the display of which was published in the last five years related to TA in Nursing Team were visualized and understood the best possible.

A table in order to analyze all the selected items was built, these were transcribed and organized in that table. They also followed the steps of an integrative literature review: identification of theme or questioning of the integrative review; sampling or literature search; categorization of studies; evaluation of studies included in the integrative review; discussion of the results and final phase consists of presentation of the review/synthesis of knowledge.

RESULTS

This integrative literature review was made from articles selected by a previous survey conducted in the Virtual Health Library (VHL) without selection of titles or sources.

The articles chosen were indexed in the Scientific Electronic Library Online (SCIELO) and the Latin American and Caribbean Health Sciences (LILACS). The keywords chosen to carry out this research were: "AT/Nursing Team" and "Occupational Risk/Nursing Team." It was found 212 articles, and after applying the inclusion criteria: "full text", left over 93 articles, and finally selected a total of 18 articles for research. The articles are distributed in the table below.

Table 1 - Total distribution of articles with descriptors in the Virtual Library database in Health (BVS). Montes Claros, 2012

| Data base | Keywords | articles found | articles selected |
|--------------|--|----------------|-------------------|
| VHL | Occupational hazards; Nursing team. | 103 | 02 |
| | AT; Nursing team | 109 | 16 |
| Total | | 212 | 18 |

Source: Data collection. Integrative review 2012.

Table 2 - Total distribution of articles with descriptors in the LILACS and SciELO database. Montes Claros, 2012

| Base accessed | n | % |
|----------------------------|-----------|------------|
| SCIELO | 11 | 61 |
| LILACS | 07 | 39 |
| Language | n | % |
| Portuguese | 15 | 83.5 |
| English | 02 | 11 |
| Portuguese English Spanish | 01 | 5.5 |
| Total | 18 | 100 |

Source: Data collection. Integrative review 2012.

Conducting the survey in the VHL was possible to highlight articles 18 to carry out this research, and of these, 39% were found at the base of LILACS and 61% in the SCIELO database. Of the sample, 83.5% were available in Portuguese and 11% in English, but we managed to open them in Portuguese, and 5.5% was available in three languages.

Table 3 - Transcribed articles according to the requirements chosen for evaluation. Montes Claros, 2012

| Article | Title | Author | Year | Periodic | Goal | Methodology | Results | Conclusion |
|---------|--|--|------|------------------------|---|--|---|--|
| I | Accidents with biological material among public health workers units | Chioldi, Marziale, Robazzi ¹⁸ | 2007 | Rev Latino-am Enferm | To investigate the occurrence of AT with exposure to biological material among health workers, who work in Public Health Units. | Descriptive quantitative approach. | It was recorded via CAT, AT 155 in 2004, and 40% involved exposure to biological material susceptible to infection by Hepatitis and <i>Acquired Immunodeficiency Syndrome</i> (AIDS). Needles were responsible for 80.6% of the injuries and the blood was the material involved in most of the exhibits. | This subject needs greater attention so that preventive measures can be implemented, considering the peculiarities of the activities performed in the different professional categories. |
| II | Accidents with needlestick materials: knowing the feelings and emotions of nursing professionals | Lima, Pine, Vieira ⁸ | 2007 | Esc Anna Nery R Enferm | Identify the knowledge of the feelings and emotions of nursing professionals who had accidents with needlestick injuries in a hospital in the public schools. | This is a descriptive research with a qualitative approach. | It was found that the occurrence of AT Sharps may be favored by the realization of hard work carried out quickly, in more than one health facility, but also by inattention and distraction. Fear before the change of lifestyle was expressed by respondents after the accident. | Points to the in - service training, technical improvement and professional development developed by the continued education sector as important to minimize the risk AT. |
| III | Accidents at work in the nursing team of Paraná University Hospital - Brazil | Secco, Robazzi ¹⁶ | 2007 | Ciênc Enferm | Analyze the typical AT recorded by nursing staff according to the variables related to time, space and person and estimate risk indicators. | Descriptive epidemiological study of cross-sectional design. | The most affected were with AT Technical/Nursing Assistants. Accidents involving exposure to biological materials were the most common. And the hands were the hardest hit in the handling of sharps. | The findings showed that the typical AT presented to hegemonic every year. It emphasizes the importance of preventive measures through educational strategies and review of work processes to prevent accidents. |

(To be continued...)

(Continuation)

| Article | Title | Author | Year | Periodic | Goal | Methodology | Results | Conclusion |
|---------|---|---|------|----------------------|---|---|--|---|
| IV | Work accidents involving nursing workers | Ribeiro, Shimizu ⁴ | 2007 | Rev Bras Enferm | The aim of this study was to identify and analyze accidents and workloads they are exposed to the nursing staff in the development of their activities. | This is a case study, descriptive and exploratory. | It was found that the workers suffered 76 AT, among which, 83.95% were caused by sharps, 8.64% by falls, 6.17% by exposure to biological fluids and 1,24% by injuries. | To achieve adequate and safe working conditions, workers in nursing need to be technically skilled to perform functions and ultimately participate in the institutional development processes of labor policies affecting them. |
| V | Factors associated with needlestick injuries in the nursing team of a university tertiary referral hospital | Canini, Moraes, Gir, Freitas ⁷ | 2008 | Rev Latino-am Enferm | To identify factors associated with needlestick injuries in the nursing team of a tertiary hospital. | This is a case-control with a quantitative approach. Used multivariate logistic regression. | Six predictors of percutaneous injuries: "recapping needles"; "Workweek ≥ 50 hours"; "Nursing experience ≤ 5 years"; "Working on night journey"; "Self-assess how low the risk of accidents" and "previous needlestick injuries." | It is recommended that institutions directed to tertiary care, despite the limits imposed by conventional measures of promotion and prevention in health, take into account the implementation of effective and reasonable measures to prevent needlestick injuries informed knowledge management and its predictors. |
| VI | Occupational health: analyzing the risks related to the nursing team in an intensive care unit | Leitão, Fernandes, Ramos ³⁷ | 2008 | Ciênc Cuid Health | To analyze the occupational hazards they are exposed to the nurses working in the ICU. | Descriptive, exploratory study with a qualitative approach. | The main occupational hazards found were excess noise in the unit, inadequate temperature of the environment, failure to observe the control of gases and vapors, inadequate use of PPE during and radiation exposure. Recorded daily exposure to biological agents, psychosocial factors and ergonomic nature. | It is emphasized that there must be a concentration of efforts and resources to promote changes in the working environment, with the implementation of prevention programs and awareness of safe practices and the provision of continuous and uniform, the safety equipment on all the professionals. |
| VII | Biosecurity and work sharps injuries among university hospital nursing professionals in Fortaleza-CE | Paulino Lopes, Rolim ¹¹ | 2008 | Cogitare Enferm | Characterize the AT occurred among health professionals; analyze the frequency with which it occurs AT sharps devices between reported nursing staff in Accident Notification Form and identify the factors contributing to its occurrence among nursing workers. | This is a documentary, exploratory study with a quantitative approach. | The object of the accident was visibly contaminated with blood or other organic substances in 61.5% of cases. In 28.2% of the accidents, the cause of the accident instrument had been used in the patient, but had no visible contamination. Syringe needles were the objects that caused more accidents, with 51.3% of the total. Then contains the peripheral access catheters (25.6%). | The education service is essential for the worker to see the importance of standard precautions, among which is included the prevention of sharps injuries and adoption of safe practices. |
| VIII | Experience report about the direction of conduct against Occupational accidents with sharps and body fluids | Sousa Campos ¹⁵ | 2008 | Cogitare Enferm | Report the lived experience of the orientation to the nursing staff about the conduct in case of accidents with sharps and body fluids. | This is a descriptive study, supported in a report of academic experience on the course for nursing staff about the conduct in case of accidents with sharps and body fluids. | The empirical observation of practice identified the significant manipulation of sharps and body fluids by the staged unit nursing professionals, exposing them constantly accident risks. The consequences of possible accidents damage the professional, patient and institution. | It is emphasized that it is for the nurse to seek information and develop actions aimed at education methodology in service, promoting the involvement of the nursing team. |

(To be continued...)

(Continuation)

| Article | Title | Author | Year | Periodic | Goal | Methodology | Results | Conclusion |
|---------|---|---|------|------------------------|--|---|---|--|
| IX | Repercussions of accident with needlestick for nursing: a construction from the focus group | Castro Farias ³⁸ | 2009 | Esc Anna Nery R Enferm | Knowing the repercussions of the accident Sharps for nursing worker and discuss these repercussions. | Descriptive study with a qualitative approach. | Be surmised that the consequences of the accident have two facets: unfavorable impact on workers' health, involving feelings of fear, despair, worry, shame, anxiety and insecurity, and positive impact on the professional conduct of this work involving the need for further care and attention when handling sharps. | It was evident that this kind of accident affects the worker's subjectivity, the adoption of safety practices making it necessary to manipulate sharps in view of its high potential for health problems both in the physical and emotional nursing workers. |
| X | Accidents at work, occupational risks and absenteeism among hospital nursing staff. | Giomo, Freitas Alves, Robazzi ¹⁰ | 2009 | Rev Nurse UERJ | The objectives of this study were to identify AT, absenteeism and occupational risks relate with absenteeism related to two hospitals nursing workers in the city of Ribeirao Preto. | Descriptive study with a quantitative approach. | They identified 140 AT and of these, 85% for women, 81% among nursing assistants and 92% typical accidents. Among the route accidents happened, most occurred before the start of the work shift. All these accidents have generated 117 days of absence. | It is considered important to conduct further research to further study this issue so traumatic for the worker and for better planning of care and prevention of accidents. |
| XI | Occupational accidents involving biological material and the nursing staff of a teaching hospital | Gomes, Agy, Malaguti, Canini Cross Gir ⁹ | 2009 | Rev Nurse UERJ | To evaluate the occurrence of accidents with biological material among nursing professionals of a large teaching hospital, São Paulo. | Descriptive study, document, and quantitative approach. | Most accidents were percutaneous (85.7%) and 67.8% of exposures hollow needle was the most involved causing object. The most frequent occurrences were vascular punctures (26.8%) and drug administration (13.3%). | There is a need to program a permanent educational program and new strategies to enable the review of the work process. |
| XII | Occupational risks in a hemodialysis unit from the perspective of workers of the nursing staff | Silva, Zeitoune ³⁹ | 2009 | Esc Anna Nery R Enferm | Describe the occupational risks in the context of workers of the nursing team in a hemodialysis unit, analyze the knowledge of the team about the protection and safety measures to the unit and discuss the knowledge of these workers on occupational hazards. | Descriptive exploratory study of qualitative approach. | Workers have the knowledge about occupational hazards and on the protection and safety measures, although not always apply them in their professional practice. They were cited as the main health implications respiratory problems, column and contagious diseases. | Despite the knowledge of the risks and safety and security measures, there is no practical application of these measures for the reduction of exposure to risk and even of occupational diseases. The aim is to review awareness strategies health care professional to risk prevention and occupational diseases. |
| XIII | Needlestick accidents among nursing professionals at a university hospital | Silva, Rock, Ayres, Juliani ¹⁴ | 2010 | Rev Gaucha Enferm | Analyze needlestick injuries in the period from 2002 to 2006, involving the nursing staff of a university hospital, to understand the context in which they occur. The use of this information can be a prevention tool. | Descriptive, retrospective study quantitative. | There were 149 accidents during the study period, with sharps, of which seven were recurrences involving the nursing staff in the categories: nurse, technician, auxiliary and nursing attendant. | The findings showed that there is still a significant degree of unawareness or trivialization of accidents among health professionals. |

(To be continued...)

(Continuation)

| Article | Title | Author | Year | Periodic | Goal | Methodology | Results | Conclusion |
|---------|---|---|------|----------------------|--|---|---|---|
| XIV | Work accidents involving sharps unit nursing professionals hospital emergency | Simon Smith, Souza, Borges, Cortez ² | 2010 | Rev Nurse UERJ | The aim of this study was to identify and analyze the occurrence of AT with needlestick materials among the nursing staff. | Descriptive, exploratory study with a quantitative approach. | It was 44 (43.6%) professionals involved in this type of accident, and the hollow needle most often associated object (68.2%) and the needle recapping responsible for 38.6% of accidents. | The analysis of the results shows that a significant part of the hospital's emergency unit nursing framework has been AT victim involving sharps. |
| XV | Investigation of biological accidents among professionals of the multidisciplinary team of a hospital | Camera, Lyre, Junior, Vilella, Hinrichsen ⁴⁰ | 2011 | Rev Nurse UERJ | Analyze the occurrence of accidents with biological material among professionals in a hospital general care. | Descriptive study documentary with a quantitative approach. | The most exposed professional category was nursing technicians (50%), followed by nurses (25.5%), and the leading cause of occurrence was due to improper disposal of sharps (43.6%). | No need to program a continuing education program, monitoring of procedures performed and implementation of biosecurity protocols. |
| XVI | Workloads and nursing working conditions: integrative review | Schmöller, Trinity, Neis, Gelbcke Saucer ² | 2011 | Rev Gaucha Enferm | Knowing the theoretical production on workloads and working conditions of nurses. | This is a descriptive study, supported in an integrative review of scientific papers, theses and dissertations of the last ten years. | The results indicated workloads as responsible for the drain of professionals, influencing the occurrence of accidents and health problems. | Studies suggest some strategies, such as adequacy of staffing quantitative, continuing education and better working conditions. |
| XVII | Evaluation of occupational exposure to biological material in health services | Valim, Marziale ¹³ | 2011 | Text Context Enferm | Identify the occurrence and characteristics of AT with biological material in health institutions of a city in the interior of São Paulo. | Descriptive transversal study, document, and quantitative approach. | The disposal of sharps in inappropriate places accounted for 18.8% of the 85 AT; 80% had contact with blood, and 20% was used chemoprophylaxis. | There is need to improve the quality of records and research possible underreporting. |
| XVIII | Analysis of accidents with biological material among health workers | Vieira, Padilha, Pine Tree ¹⁷ | 2011 | Rev Latino-am Enferm | Knowing the AT with exposure to biological material and workers profile, from the reporting forms the Regional Reference Center for Health Florianopolis Macroregion worker. | Descriptive, retrospective documentary with a quantitative approach. | Predominated nursing technicians, female, mean age of 34.5 years. Accidents, 73% involved percutaneous exposure, 78% had contact with blood and / or fluid with blood and 44.91% resulted from invasive procedures. | The strategies to prevent the occurrence of AT with biological material should include joint actions between workers and management of services should be directed to the improvement of the conditions and organization of work. |

Table 4 - Items distributed as the year of publication, type of methodology, journals found, study setting and subject matter. Montes Claros (MG) 2012

| Year of Publication | n | % |
|--|----------|----------|
| 2007 | 04 | 22,2 |
| 2008 | 04 | 22,2 |
| 2009 | 04 | 22,2 |
| 2010 | 02 | 11,1 |
| 2011 | 04 | 22,2 |
| Type Method | n | % |
| documentary exploratory | 01 | 5,5 |
| Descriptive, retrospective, quantitative and qualitative | 01 | 5,5 |
| integrative review | 01 | 5,5 |
| if control | 01 | 5,5 |
| descriptive | 01 | 5,5 |
| descriptive exploratory | 01 | 5,5 |
| Descriptive and qualitative | 04 | 22,5 |
| Descriptive and quantitative | 03 | 17 |
| Descriptive, cross | 02 | 11 |
| Descriptive, exploratory and quantitative | 01 | 5,5 |
| Descriptive, retrospective and quantitative | 01 | 5,5 |
| Experience report | 01 | 5,5 |
| Journal | n | % |
| cogitare Nursing | 02 | 11 |
| Nursing Journal of Rio Grande do Sul | 02 | 11 |
| Anna Nery School Journal of Nursing | 03 | 17 |
| Latin American Journal of Nursing | 03 | 17 |
| Brazilian Journal of Nursing | 01 | 5,5 |
| UERJ Nursing Journal | 04 | 22 |
| Science and Nursing | 01 | 5,5 |
| Text & Context Nursing | 01 | 5,5 |
| Science, Care and Health | 01 | 5,5 |
| Study Scenario | n | % |
| Universitary hospital | 06 | 33 |
| Public hospital | 05 | 28 |
| General hospital | 01 | 5,5 |
| Hospital | 02 | 11 |
| Database | 01 | 5,5 |
| Health facilities | 01 | 5,5 |
| Public Health Unit | 01 | 5,5 |
| Worker's Health Reference Center | 01 | 5,5 |
| Subject Addressed | n | % |
| Types of AT | 17 | 94,4 |
| Causes of AT | 16 | 88,8 |
| Legislation | 07 | 38,8 |
| biosecurity | 15 | 83,3 |

Source: Data collection. Integrative review 2012.

¹ Cumulative Percentage

After analyzing the articles chosen by the year of publication, it was noted that there was equality in the years

2007, 2008, 2009 and 2011 - four publications each year, which is equivalent to 22.2% each year, and that number fell in 2010, we were found two publications (11.1%). According to the methodology used to realize a larger share of exploratory descriptive with 22.5% of publications, followed by descriptive quantitative with 17%, and descriptive transversal character with 11% and the other followed a standard 5.5 %. It was realized that each article opts for different methodologies.

Regarding the journals found in the search, it is observed that the magazine received more publications on the subject was the Journal of Nursing of UERJ, with 22% of publications, followed by Anna Nery School and Latin American Journal of Nursing, 17 % each, Cogitare Nursing and Nursing Journal of Rio Grande do Sul with 11% each, and finally the other obtained a percentage of 5.5% of publications. The University Hospitals were those who had a higher number of searches performed, with a total of 33% of the articles, followed by Public Hospitals were second with regard to conducting research, the hospital has averaged 11% followed by other study scenarios with 5.5%.

In the selected articles it was revealed that, in response to the question of the study, "What has been published about AT in nursing team in the last five years?" In 94.4% was approaching "types of TA" in 88.8% possible "causes of these accidents," 38.8% in the "law" with regard to worker safety, the rights and duties of the same, and 83.3% addressed to "biosafety".

DISCUSSION

In this study, it was found 212 articles on the topic searched. Of these, only 18 fulfilled the inclusion criteria of the search. The surveyed items are in the languages English, Portuguese and Spanish. And of these, one was in Portuguese, English and Spanish, two were in English, but it was possible to open them in Portuguese, and the others were exclusively in Portuguese. There was an equal number of publications in the years 2007-2009 and 2011, each with a total of 22.2% of the publications and in 2010 this figure fell to 11.1%. Regarding the methodology most used by the authors of the selected studies highlight the descriptive exploratory with 22.5%, followed by descriptive quantitative with 17%, and descriptive transversal character with 11% and the other followed a pattern of 5.5%. But also realize is that each article opts for different methodologies.

As for the journals that had a higher number of publications are was the UERJ Nursing Journal, with 22% of publications, followed by Latin American Journal of Nursing and School Anna Nery, each with a total of 17% of publications. The study scenario that prevailed more in the research were the University Hospitals that reached a total of 33% of all selected publications, followed by public hospitals with 28%. These were the two leading scenarios. The subjects that had more relevance in pre-selected articles were related

to the types of TA (94.4%); the possible causes of these accidents (88.8%); the legislation governing the protection of workers, as well as guidance on their duties (38.8%); and biosafety (83.3%).

In the articles analyzed stands out an average of 385,000 accidents per year involving sharps to health workers. An investigation by the Center for Disease Control and Prevention (CDC) has identified 57 cases of seroconversion and other 140 possible cases of occupational post exposure AIDS in the period 1981 and 2006. It was also noted that, percutaneous accident appeared in 48 cases and skin mucus were five cases.¹³ In this review, the other selected items not found absolute number of number of accidents, but are specific percentages of the types of accidents.

Sharps injuries are the ones that occur in health institutions.⁹ Other research supports in order to report large number of accidents with sharps, and with hollow needle registers 68.2%, with scalp/Gelco 22.7% and finally the scalpel blade 4.5%.¹² In a study conducted in São Paulo, for notification of accidents with biological material in health professionals, 86% involved percutaneous exposure and 82% blood was involved.¹⁵ Accidents caused by organic materials are the most evident due to exposure to blood and bodily fluids causing infection such as hepatitis B and C and AIDS virus, and can even be lethal. This contamination occurs more frequent way through the skin, as a result of accidents with sharps.¹⁸

Accidents involving biological material should be highlighted and percutaneous exposure appeared in 73% of the total analyzed. Then there are the accidents involving mucous membranes (10%), fair skin exposure (10%) and non-intact skin (7%). In 69.49% of accidents biological material had blood in 9.32% had fluid with blood.¹⁷ The main causes of accidents in the nursing team can be directly related to non-observance of regulations, malpractice, inadequate working conditions, instructions incorrect or insufficient, gaps in supervision and guidance, lack or inadequacy in the use of PPE.¹⁴ The act of "recapping needles" was an important predictor of percutaneous injuries in nursing professionals. In addition, a long day working week corroborates in greater chances of occurrence of such accidents and this may be due to a greater worker time exposure to a hazardous situation and/or the fatigue generated by this long journey.⁷

In emergency departments, for example, the risk of accidents increases further due to work overload. The professionals working in these sectors have to perform a number of tasks in a short time and also are associated with the stress caused by nature craft, and this can cause a decrease in concentration and increase the chances of accidents.¹²

The causes of AT may be linked to low wages, which leads to workers having more than one job, and causes attention at work decreases and workers are susceptible to strong physical and emotional pressures.⁴ Corroborating the subject, another study points out that the work overload and mental suffering make nursing workers more susceptible to occupational

diseases. And these factors accompanied by socio-economic difficulties, since professionals are underpaid, which forces them to adopt double shifts to support, it can be seen how these professionals are subject to high risks of AT.¹⁰

Nursing workers who make use of sharps are responsible for their disposal in order to meet the NR-32 MTE.³⁸ Confirming this assumption, another study says that if they had been complied with the provisions of NR-32, possibly many accidents occurred could have been avoided, which would minimize the suffering of the victims.¹⁰ Other studies also follow the same line of reasoning, where workers should make use of PPE and fulfill precisely the NR-32 requirements.¹³

In one study, noted the importance of reactivating the Internal Commission for Accident Prevention (ICAP) in the studied site, it must be expanded in its scope of action and provide spaces for workers, so that they can negotiate improvements in the workplace with the institution for the safety in carrying out their daily activities.⁴ In terms of biosafety, engineering services specialized in safety, occupational health and worker health institutions should be more committed to remedy the problems in the workplace, which greatly hurt the workers, and even hurt the hospitals themselves.¹⁰

The care must be redoubled when handling or disposing sharps, which in addition to promoting the health care of all professionals who share the work space, promotes care for themselves regarding their health and safety.³⁸ The adoption of hygiene measures hand, proper use of PPE, immunization of professionals, handling and disposal of sharps properly are key measures to minimize the exposure of workers to body fluids and sharps.⁴⁰ It is necessary attention when handling sharps and the adoption of safe practices and use of the principles of biosecurity and use of PPE.³⁸

The care when handling sharp items is evident. However, the institution is responsible for the applicability of the biosecurity measures in the activities of workers of the nursing staff, through human and material resources, providing PPE, encouraging lifelong learning, and further adoption of health and safety measures in the workplace.¹⁴ There is a poor work organization, especially lack of PPE in adequate quantity and quality and next to this is added the lack of investment in continuous training, thus increasing the risk of accidents.⁴ The health of nursing workers is of interest not only of the workers themselves, but of professional associations, trade unions, nursing schools, and above all the service users. Therefore, it is expected that workers seek decent work, guided by the responsibility, scientific expertise and ethics.¹⁶

CONCLUSION

After the analysis, it could be seen that the AT in health sectors occur mostly with nursing professionals. The nursing team is in constant contact with the patient. In addition, the

low wages that make the professional has to look for other jobs to increase monthly income, the workload generated by the long hours that must be met, poor working conditions especially in hospitals, lack the use of PPE; inattention, indifference, confidence in itself, the lack of measures that should from the service to promote updates courses and providing adequate protection materials in sufficient numbers are factors that make the risk of accidents to gain greater proportion and They may be responsible for the large number of accidents.

Studies always emphasize accidents that occurred more, which most often had the biological material involved, on biosafety, and on legislation that gives action on the AT in health workers. It may be noted that accidents more frequently cited refer to those involved with sharps due to needle recapping, for example, that the most affected parts were superior members and more specifically hands, as they are the part of the body most used in industrial activities.

It is concluded that more was published in the last five years on the AT in the nursing team were the types of TA, possible causes, and biosafety legislation. And you can notice that accidents do not occur only by professional errors, but more than that is linked to the inadequacy of service to preventive measures, work overload, poor conditions of service, low wages that involve increasing the number employment and hours worked with a few hours of rest. If the services gave better conditions and workers had greater awareness the number of accidents would decrease significantly.

REFERENCES

- Oliveira RA. A concepção de trabalho na filosofia do jovem Marx e suas implicações antropológicas. *Kinesias* [Internet]. 2010 [Citado em 2012 oct 25];2(3):72-88. Disponível em: http://www.mabilia.unesp.br/Home/RevistasEletronicas/Kinesias/6_RenatoAlmeidadeOliveira.pdf
- Schmoeller R, Trindade LL, Neis MB, Gelbcke FL, Pires DEP. Cargas de trabalho e condições de trabalho da enfermagem: revisão integrativa. *Rev Gaúcha Enferm* [Internet]. 2011 [Citado em 2012 oct 18];32(2):368-77. Disponível em: <http://www.scielo.br/pdf/rgenf/v32n2/a22v32n2.pdf>
- Jakobi HR. Mapa de risco ocupacional no Estado de Rondônia baseado em tecnologia de georeferenciamento [Dissertação]. Porto Velho (RO): Universidade Federal de Rondônia, 2008 [Citado em 2012 oct 25]. Disponível em: http://bvsm.sau.gov.br/bvs/artigos/mapa_risco_occupacional_ro.pdf
- Ribeiro EJJ, Shimizu HE. Acidentes de trabalho com trabalhadores de enfermagem. *Rev Bras Enferm* [Internet]. 2007 [Citado em 2012 oct 18];60(5):535-40. Disponível em: <http://www.scielo.br/pdf/reben/v60n5/v60n5a10.pdf>
- Brasil. Ministério da Saúde. Secretaria de Atenção à Saúde. Departamento de Ações Programáticas e Estratégias. Notificação de acidentes do trabalho fatais, graves e com crianças e adolescentes. Saúde do trabalhador: protocolos de complexidade diferenciada 2. Brasília (DF); 2006 [Citado em 2012 oct 24]. Disponível em: http://bvsm.sau.gov.br/bvs/publicacoes/protocolo_not_acidentes_trab.pdf
- Brasil. Ministério da Saúde. Portaria GM nº. 485, de 11/11/2005, D.O.U. em 16/11/2005. Portaria GM nº. 939, de 18/11/2008, D.O.U. em 19/11/2008. Portaria GM nº. 1.748, de 30/08/2011, D.O.U. em 31/08/2011. Dispõe sobre a Norma Regulamentadora nº. 32 na qual tem por finalidade estabelecer as diretrizes básicas para a programação de medidas de proteção à segurança e à saúde dos trabalhadores dos serviços de saúde, bem como daqueles que exercem atividades de promoção e assistência à saúde em geral. 2006 [Citado em 2012 oct 23]. Disponível em: [http://portal.mte.gov.br/data/files/8A7C812D36A28000138812EAFCE19E1/NR-32%20\(atualizada%202011\).pdf](http://portal.mte.gov.br/data/files/8A7C812D36A28000138812EAFCE19E1/NR-32%20(atualizada%202011).pdf)
- Canini SRMS, Moraes SA, Gir E, Freitas ICM. Percutaneous injuries correlates in the nursing team of a Brazilian tertiary-care university hospital. *Rev Latino-am Enferm* [Internet]. 2008 [Citado em 2012 oct 18];16(5):818-23. Disponível em: <http://www.scielo.br/pdf/rlae/v16n5/04.pdf>
- Lima FA, Pinheiro PNC, Vieira NFC. Acidentes com material perfurocortante: conhecendo os sentimentos e as emoções dos profissionais de enfermagem. *Esc Anna Nery* [Internet]. 2007 [Citado em 2012 oct 18];11(2):205-11. Disponível em: <http://www.scielo.br/pdf/ean/v11n2/v11n2a04.pdf>
- Gomes AC, Agy LL, Malaguti SE, Canini SRMS, Cruz EDA, Gir, E. Acidentes ocupacionais com material biológico e equipe de enfermagem de um hospital-escola. *Rev Enferm UERJ* [Internet]. 2009 [Citado em 2012 oct 18];17(2):220-3. Disponível em: <http://www.facenf.uerj.br/v17n2/v17n2a14.pdf>
- Giommo DB, Freitas FCT, Alves LA, Robazzi MLCC. Acidentes de trabalho, Riscos Ocupacionais e Absenteísmo entre trabalhadores de Enfermagem Hospitalar. *Rev Enferm UERJ* [Internet]. 2009 [Citado em 2012 oct 19];17(1):24-9. Disponível em: <http://files.bvs.br/upload/S/0104-3552/2009/v17n1/a004.pdf>
- Paulino DCR, Lopes MVO, Rolim ILTP. Biossegurança e acidentes de trabalho com pérfuro-cortantes entre os profissionais de enfermagem de Hospital Universitário de Fortaleza-CE. *Cogitare Enferm* [Internet]. 2008 [Citado em 2012 oct 15];13(4):507-13. Disponível em: <http://ojs.c3sl.ufpr.br/ojs2/index.php/cogitare/article/view/13109/8867>
- Simão SAF, Soares CRG, Souza V, Borges RAA, Cortez EA. Acidentes de trabalho com material perfurocortante envolvendo Profissionais de Enfermagem de Unidade de Emergência Hospitalar. *Rev Enferm UERJ* [Internet]. 2010 [Citado em 2012 oct 18];18(3):400-4. Disponível em: <http://www.facenf.uerj.br/v18n3/v18n3a11.pdf>
- Valim MD, Marziale MHP. Avaliação da exposição ocupacional a material biológico em serviços de saúde. *Texto Contexto Enferm* [Internet]. 2011 [Citado em 2012 oct 18];20(esp):138-46. Disponível em: <http://www.scielo.br/pdf/tce/v20nspe/v20nspea18.pdf>
- Silva TR, Rocha SA, Ayres JA, Juliani CMC. Acidente com material perfurocortante entre profissionais de enfermagem de um hospital universitário. *Rev Gaúcha Enferm* [Internet]. 2010 [Citado em 2012 oct 18];31(4):615-22. Disponível em: <http://www.scielo.br/pdf/rgenf/v31n4/a02v31n4.pdf>
- Sousa JV, Campos LF. Relato de experiência quanto à orientação de conduta frente a acidentes de trabalho com perfurocortantes e fluidos orgânicos. *Cogitare Enferm* [Internet]. 2008 [Citado em 2012 oct 30];13(4):602-6. Disponível em: <http://ojs.c3sl.ufpr.br/ojs2/index.php/cogitare/article/view/13124/8883>
- Secco IAO; Robazzi MLCC. Acidentes de trabalho na equipe de enfermagem de um hospital de ensino do Paraná – Brasil. *Ciênc Enferm* [Internet]. 2007 [Citado em 2012 oct 18];13(2):65-78. Disponível em: <http://www.scielo.cl/pdf/cienf/v13n2/art08.pdf>
- Vieira M, Padilha MI, Pinheiro RDC. Analysis of accidents with organic material in health workers. *Rev Latino-am Enferm* [Internet]. 2011 [Citado em 2012 oct 18];19(2):332-9. Disponível em: <http://www.scielo.br/pdf/rlae/v19n2/15.pdf>
- Chioldi MB, Marziale MHP, Robazzi MLCC. Occupational accidents involving biological material among public health workers. *Rev Latino-am Enferm* [Internet]. 2007 [Citado em 2012 oct 18];15(4):632-8. Disponível em: <http://www.scielo.br/pdf/rlae/v15n4/v15n4a17.pdf>
- Brasil. Ministério da Saúde. Senado Federal. Secretaria Especial de Informática. Constituição da República Federativa do Brasil – Texto promulgado em 05 de outubro de 1988. Brasília, 2013. Disponível em: http://www.senado.gov.br/legislacao/const/con1988/CON1988_05.10.1988/CON1988.pdf
- Brasil. Ministério da Saúde. Presidência da República. Lei Nº. 8.080, de 19 de setembro de 1990 [Citado em 2012 oct 24]. Disponível em: http://www.planalto.gov.br/ccivil_03/leis/L8080.htm
- Brasil. Ministério da Saúde. Portaria GM nº. 1679, de 19/09/2002. Dispõe sobre a estruturação da rede nacional de atenção integral à saúde do trabalhador no SUS e dá outras providências. 2002 [Citado em 2012 oct 23]. Disponível em: http://www.sau.gov.br/sites/default/files/portaria_n.1679_de_18.09.2002.pdf
- Brasil. Ministério da Saúde. Portaria GM nº. 2.437, de 7/12/2005. Dispõe sobre a ampliação e o fortalecimento da Rede Nacional de Atenção Integral à Saúde do Trabalhador - RENAST no Sistema Único de Saúde - SUS e dá outras providências. 2005 [Citado em 2012 oct 21]. Disponível em: http://www.registro.sp.gov.br/cerest/arquivos/portarias/PORTARIA_2437.pdf
- Brasil. Ministério da Saúde. Portaria GM nº. 1339, de 18/11/1999. Dispõe sobre a lista de doenças relacionadas ao trabalho, a ser adotada como referência dos agravos originados no processo de trabalho no Sistema Único de Saúde, para uso clínico e epidemiológico. 1999 [Citado em 2012 oct 22]. Disponível em: <http://dtr2001.sau.gov.br/sas/PORTARIAS/Port99/GM/GM-1339.html>
- Brasil. Ministério da Saúde. Portaria GM nº. 777, de 28/04/2004. Dispõe sobre os procedimentos técnicos para a notificação compulsória de agravos à saúde do trabalhador em rede de serviços sentinela específica, no Sistema Único de Saúde – SUS. 2004 [Citado em 2012 oct 21]. Disponível em: <http://portal.sau.gov.br/portal/arquivos/pdf/Portaria777.pdf>
- Brasil. Ministério do Trabalho e Emprego. Portaria MTE nº. 485, de 11/11/2005. 2005 [Citado em 2012 oct 20]. Dispõe sobre as diretrizes básicas para a programação de medidas de proteção à segurança e à saúde dos trabalhadores dos serviços de saúde, bem como daqueles que exercem atividades de promoção e assistência à saúde em geral. Disponível em: http://www.sau.gov.br/images/documentos/Portaria_485.pdf
- Brasil. Ministério do Trabalho e Emprego. Portaria nº. 939, de 18/11/2008. Dispõe sobre os deveres dos empregadores em promover a substituição dos materiais perfurocortantes por outros com dispositivo de segurança no prazo máximo de vinte e quatro meses a partir da data de publicação desta portaria. 2008 [Citado em 2012 oct 21]. Disponível em: http://portal.mte.gov.br/data/files/8A7C812D3226A41101322A9577176D1D/p_20081118_939.pdf
- Brasil. Ministério do Trabalho e Emprego. Portaria nº. 1.748, de 30/08/2011. 2011 [Citado em 2012 oct 20]. Disponível em: http://csa.fau.com.br/site/arquivos/arquivo_20111207110600.pdf
- Brasil. Ministério da Saúde. Lei nº. 8.213, de 24/07/1991. Dispõe sobre os planos de benefícios da Previdência Social e dá outras providências. 1991 [Citado em 2012 oct 22]. Disponível em: http://www.ipsm.gov.br/arquivos/legislacoes/legislacao/leis/lei_8213.pdf

29. Mancini PC, Teixeira LC, Resende LM, Gomes AM, Vicente LCC, Oliveira PM. Medidas de biossegurança em audiologia. Rev CEFAC [Internet]. 2008 [Citado em 2012 nov 7];10(4):603-10. Disponível em: <http://www.scielo.br/pdf/rcefac/v10n4/v10n4a22.pdf>
30. Anvisa. Agência Nacional de Vigilância Sanitária. Infecção relacionada à assistência à saúde. Módulo 05. 2004 [Citado em 2012 oct 25]. Disponível em: <http://www.anvisa.gov.br/servicosaude/manuais/iras/M%F3dulo%205%20-%20Risco%20Ocupacional%20e%20Medidas%20de%20Precau%E7%F5es%20e%20Isolamento.pdf>
31. Moresi E. Metodologia da pesquisa. Universidade Católica de Brasília – UCB. Brasília, 2003 [Citado em 2012 oct 23]. Disponível em: <http://www.inf.ufes.br/~falbo/files/MetodologiaPesquisa-Moresi2003.pdf>
32. Siena O. Metodologia da pesquisa científica: elementos para elaboração e apresentação de trabalhos acadêmicos. Porto Velho, 2007 [Citado em 2012 oct 23]. Disponível em: http://www.mestradoadm.unir.br/site_antigo/doc/manualdetrabalhoacademicoatual.pdf
33. Mendes KDS, Silveira RCCP, Galvão CM. Revisão integrativa: método de pesquisa para a incorporação de evidências na saúde e na enfermagem. Texto Contexto Enferm [Internet]. 2008 [Citado em 2012 oct 30];17(4):758-64. <http://www.scielo.br/pdf/tce/v17n4/18.pdf>
34. Botelho LLR, Cunha CCA, Macedo M. O método da revisão integrativa nos estudos organizacionais. Rev Eletr Gestão Soc [Internet]. 2011 [Citado em 2012 oct 23];5(11):121-36. Disponível em: <http://www.gestoesociedade.org/gestoesociedade/article/view/1220/906>
35. Pompeo DA, Rossi LA, Galvão CM. Integrative literature review: the initial step in the validation process of nursing diagnoses. Acta Paul Enferm [Internet]. 2009 [Citado em 2012 oct 30];22(4):434-8. Disponível em: http://www.scielo.br/pdf/ape/v22n4/en_a14v22n4.pdf
36. Souza MT; Silva MD, Carvalho R. Revisão integrativa: o que é e como fazer. Einstein [Internet]. 2010 [Citado em 2012 oct 30];8(1 Pt 1):102-6. Disponível em: http://apps.einstein.br/revista/arquivos/PDF/1134-Einsteinv8n1_p102-106_port.pdf
37. Leitão IMTA, Fernandes AL, Ramos IC. Saúde ocupacional: analisando os riscos relacionados à equipe de enfermagem numa unidade de terapia intensiva. Ciênc Cuid Saúde [Internet]. 2008 [Citado em 2012 oct 19];7(4):476-84. Disponível em: <http://www.periodicos.uem.br/ojs/index.php/CiencCuidSaude/article/view/6630/3907>
38. Castro MR, Farias SNP. Repercussões do acidente com perfurocortantes para a enfermagem: uma construção a partir do grupo focal. Esc Anna Nery [Internet]. 2009 [Citado em 2012 oct 18];13(3):523-9. Disponível em: <http://www.scielo.br/pdf/ean/v13n3/v13n3a10.pdf>
39. Silva MKD, Zeitoune RCG. Riscos ocupacionais em um setor de hemodiálise na perspectiva dos trabalhadores da equipe de enfermagem. Esc Anna Nery [Internet]. 2009 [Citado em 2012 oct 18];13(2):279-86. Disponível em: <http://www.scielo.br/pdf/ean/v13n2/v13n2a07.pdf>
40. Câmara PF, Lira C, Santos Junior BJ, Vilella TAS, Hinrichsen SL. Investigação de acidentes biológicos entre profissionais da equipe multidisciplinar de um hospital. Rev Enferm UERJ [Internet]. 2011 [Citado em 2012 oct 19];19(4):583-6. Disponível em: <http://www.facenf.uerj.br/v19n4/v19n4a13.pdf>

Received on: 24/02/2014
Reviews required: 24/05/2016
Approved on: 15/06/2016
Published on: 01/10/2016

Mailing address:

Patrick Leonardo Nogueira da Silva
Phone: (038) 991312287
Rua Pedro Álvares Cabral, 530
São Cristóvão, Espinosa, MG, Brazil
ZIP Code: 39.510-000