

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/344380672>

Psychometric Properties of the Inventory of Statements About Self-Injury (ISAS) in Iranian Opioid and Alcohol Abusers

Article in *Iranian Journal of Psychiatry and Behavioral Sciences* · September 2020

DOI: 10.5812/ijpbs.88494

CITATION

1

READS

80

9 authors, including:



Mehran Zarghami

Mazandaran University of Medical Sciences

99 PUBLICATIONS 467 CITATIONS

[SEE PROFILE](#)



Masoudeh Babakhanian

Mazandaran University of Medical Sciences

57 PUBLICATIONS 109 CITATIONS

[SEE PROFILE](#)



Mojtaba Habibi

School of Behavioral Sciences and Mental Health (Tehran Institute of Psychiatry), ...

102 PUBLICATIONS 328 CITATIONS

[SEE PROFILE](#)



Hamid Sharif Nia

Mazandaran University of Medical Sciences

296 PUBLICATIONS 1,091 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Time-series analysis of weather and Acute Myocardial Infarction in Iran [View project](#)



Model of stress of exposure to sharps injury in Iranian nurses: A Path Analysis [View project](#)



Psychometric Properties of the Inventory of Statements About Self-Injury (ISAS) in Iranian Opioid and Alcohol Abusers

Mehran Zarghami^{1,2}, Masoudeh Babakhanian^{3,*}, Mojtaba Habibi Asgarabad⁴, Masumeh Ghazanfanpour⁵, Faezeh Sadat Akrami⁶, Najmodin Nazeri⁷, Yasaman Allameh⁸, Hamid Sharif Nia⁹ and Nader Molavi¹⁰

¹Psychiatry and Behavioral Sciences Research Center, Addiction Institute, Mazandaran University of Medical Sciences, Sari, Iran

²Department of Psychiatry, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

³Student Research Committee, Psychiatry and Behavioral Research Centre, Mazandaran University of Medical Sciences, Sari, Iran

⁴Health Promotion Research Center, Department of Health Psychology, School of Behavioral Sciences and Mental Health (Tehran Institute of Psychiatry), Iran University of Medical Sciences, Tehran, Iran

⁵Department of Midwifery, Razi School of Nursing and Midwifery, Kerman University of Medical Sciences, Kerman, Iran

⁶Department of Clinical Psychology, School of Humanities and Social sciences, Islamic Azad University, Science and Research Branch, Tehran, Iran

⁷Lorestan Department of Education, Poldokhtar, Lorestan, Iran

⁸Psychiatry and Behavioral Research Centre, Mazandaran University of Medical Sciences, Sari, Iran

⁹Department of Medical-Surgical Nursing, School of Nursing and Midwifery Amol, Mazandaran University of Medical Sciences, Sari, Iran

¹⁰Kashan University of Medical Sciences, Kashan, Iran

*Corresponding author: Student Research Committee, Psychiatry and Behavioral Research Centre, Mazandaran University of Medical Sciences, Sari, Iran. Email: babakhanian.m@gmail.com

Received 2019 January 01; Revised 2020 June 15; Accepted 2020 July 11.

Abstract

Background: Non-suicidal self-injury (NSSI) is one of the most dangerous behaviors linked to substance abuse. The Inventory of Statements About Self-Injury (ISAS) has been developed to better understanding the factors associated with the onset and maintenance of NSSI behaviors.

Objective: The ISAS was translated into the Persian language to study its psychometric properties in Iranian population of opioid and alcohol abusers.

Methods: This is a psychometric study investigating opioid and alcohol abusers in Iran, including those residing in addiction rehabilitation camps, prisons, hospitals, and addiction treatment clinics dispensing methadone in 2017. The sample size of this study was 470. The subjects completed the Inventory of Statements About Self-Injury, Barrat Impulsiveness Scale (BIS), and the Distress Tolerance Scale (DTS).

Results: The test-retest results were estimated for two weeks using the Intra-class Correlation Coefficient (ICC). The absolute reliability was determined to be 2.62%. There was a significant correlation between convergent and divergent instruments. The results of the exploratory factor analysis on 235 individuals in the sample showed all subscales of the inventory measure a single factor in the Iranian population. The total Cronbach's alpha coefficient for this subscale was 0.93. Also, the results of confirmatory factor analyses on the rest of the sample (235), after applying the AMOS software suggestions to improve the model, showed this inventory was a good indicator for this population ($\chi^2 = 131.69$; $P < 0.001$; $df = 56$; $\chi^2/df = 2.35$; CFI = 0.97; RMSEA = 0.76; GFI = 0.92).

Conclusions: The Persian version of the Inventory of Statements About Self-Injury has high validity and reliability among the opioid and alcohol-abusing population in Iran.

Keywords: Self-Injurious Behavior, Alcohol, Iran

1. Background

Based on epidemiological studies, a minor, yet significant, part of the society exhibits intentional self-injurious behaviors (1). In its latest classification (DSM-5), the American Psychiatric Association defines non-suicidal self-injury (NSSI) as deliberate, self-inflicted destruction of body tissue without suicidal intent and for purposes not socially sanctioned. These behaviors include cutting, burning, bit-

ing, and scratching the skin to reduce mental discomfort (2, 3).

Understanding pathological behaviors in terms of reason and purpose, such as self-injurious behaviors at a certain time in a particular patient, is often difficult, but it can greatly help manage the patient. Different models conceptualize possible reasons for self-injurious behaviors, considering the effect of different factors in play (4). In or-

der to better understanding the factors associated with the onset and maintenance of NSSI behaviors, Klonsky et al. developed The Inventory of Statements About Self-Injury (ISAS). The four major categories for NSSI are environmental, drive, affect regulation, and interpersonal functions. The instrument includes six specific functional models, namely, environmental, anti-suicide, sexual, affect regulation, dissociation, and boundaries (5, 6).

Self-injury is one of the most dangerous behaviors among substance abusers. They are six to eight times more likely to engage in self-injurious behavior (7). In the Iranian substance abuse population, it seems trait impulsivity is the main psychological index for self-injurious behavior. In order to evaluate risky and impulsive behaviors, Barrett's, Dickman's, and Zakerman's measures of impulsiveness have been localized and used among substance abusers in Iran (8).

2. Objectives

The present study was conducted to study the psychometric properties of ISAS in Iranian population of opioid and alcohol abusers for the identification of intentional self-injury and the associated factors in this population.

3. Methods

This psychometric study investigates the statistical population of substance and alcohol abusers in Iran. Monroe believes a sample size of 200 to 500 is suitable for factor analysis study (9). Using a convenient sampling method, 470 participants were selected. Data were collected from multiple provinces in Iran (Tehran, Mazandaran, Lorestan, Khuzestan, and Semnan). Informed consent was obtained from all participants before the study.

In this descriptive cross-sectional research, the statistical population included abusers of opioids and/or alcohol recruited from addiction rehabilitation camps, prisons, hospitals, and addiction treatment clinics dispensing methadone. The minimum and maximum age of the participants were 16 and 67 years, respectively (mean = 31.9 ± 9.37). Almost all participants (99.5%) were opioid abusers only, and 68 individuals (14.5%) were abusing both opioid and alcohol. The mean age at the time of the first and the last self-injury events were 19.1 years and 24.7 years, respectively.

The inclusion criteria were: a history of intentional self-injurious behaviors; a history of opioid abuse documented by urine dipstick and/or alcohol abuse; no addiction to stimulants; clinical diagnosis of opioid abuse by a psychiatrist or a clinical psychologist based on DSM 5 criteria; the patient's informed consent for participation in the

research; a history of induced injury or intentional damage to self (based on self-declaration). The exclusion criteria were: psychotic symptoms; neurological disorders; serious suicidal thoughts based on a psychiatric interview; and history of stimulant abuse.

3.1. Measures

3.1.1. The Inventory of Statements of Self-Injury

Klonsky and colleagues developed ISAS in 2008 and evaluated it on 235 college youth with a history of at least one occasion of non-suicidal self-injury (5, 6). The first part of the questionnaire examines deliberate behaviors without suicidal intentions. These include cutting, biting, burning, pinching, hair plucking, intense scratching, hitting oneself against a surface (such as hitting one's fist or head against the wall), preventing wound healing (such as scaling the scalp), rubbing the skin on a rough surface, needling the body, swallowing harmful objects, etc. Participants were asked to estimate the number of times that each behavior was performed.

There were five other questions in the descriptive section including age at the time of onset, the experience of pain while performing self-injurious behaviors, frequency of these behaviors in the presence or absence of others, time interval between feeling the motive for self-injury to do it, and whether the individual wanted to stop the behavior. The second part deals with assessing 13 self-injurious behaviors: affect regulation, interpersonal boundaries, self-punishment, self-care, anti-dissociation/feeling-generation, anti-suicide, sensation-seeking, peer-bonding, interpersonal influence, toughness, marking distress, revenge, and autonomy (5, 6).

Individual's experiences in each function were evaluated by one of the following three scores: 0-not relevant (two points), 1-somewhat relevant (four points), 2-very relevant (six points). Therefore, the score of each of the 13 self-harming behaviors could range from two to six (5, 6). This version of the questionnaire has been localized in other countries such as Turkey, Sweden, and Australia (10-12).

3.1.2. Barrat Impulsiveness Scale

Ernest Barrett developed Barrat Impulsiveness Scale (BIS) in 1977 and its eleventh edition in 2004. This questionnaire consists of 30 items, evaluating cognitive impulsivity, motor impulsivity, and disorganization as three distinctive factors. Each item has four choices, and the highest score is 120. Two independent evaluators categorized 58 impulsivity actions with a kappa coefficient of 0.83 ($P < 0.0001$). The calculated Cronbach's alpha coefficient for reliability of Barrat's impulsivity scale was 0.845 with the following values for the subscales: 0.781 for cognitive impulsivity, 0.741 for motor impulsivity, and 0.437 for disorgani-

zation factor (13). Ekhtiari et al. observed that the Persian translation of Barrett's impulsivity scale has desirable validity and reliability in Iran with a reliability level of 0.83 (7).

3.1.3. The Distress Tolerance Scale

Simons and Gaher developed the Distress Tolerance Scale (DTS) in 2005 as a self-assessment for distress tolerance and studied its reliability for six months. Their results showed the questionnaire had a high internal consistency ($\alpha = 0.89$). In addition, distress tolerance was stable in six months ($r = 0.61$). The scale includes 15 items and 4 subscales called emotional distress tolerance (items 1, 3, and 5), being attracted by negative emotions (items 2, 4, and 15), mental distress estimation (items 6, 7, 9, 10, 11, and 12), and adjustment attempts to alleviate distress (items 4, 8, and 13) (14). Azizi estimated the alpha coefficient for each of the subscales of tolerance, attraction, evaluation, and adjustment to be 0.75, 0.77, 0.70, and 0.75, respectively. The re-test reliability coefficient for the whole scale was 0.81, and for the subscales of tolerance, attraction, evaluation, and adjustment was 0.71, 0.69, 0.77, and 0.73, respectively (9).

3.2. Procedure

First, two psychologists who were experts in both Persian and English languages translated the scale into Persian. Next, a native translator expert in the field back-translated the scale into English. Then, the translation team compared the English back-translation with the original text to ensure the accuracy of the concepts and to identify any differences. After agreement on the final English back-translation, the Persian version was adjusted accordingly. To assess the content validity, two methods of (CVR [Content validity ratio]) and (CVI [Content validity index]) were used. Rubio suggested experts' opinions are the best way to evaluate the content of an inventory (15). No questions were omitted or changed. The obtained values indicated the adequacy of the content validity of the Persian version of the statements about self-harm. After verifying the accuracy of the translation, the questionnaire was tested and re-tested within 20 days in a small sample (30 qualified participants) in a preliminary study. Afterward, authorization to distribute the questionnaire among the target population was obtained. Then we distributed the questionnaire to the sample population in prisons, addiction rehabilitation camps, hospitals, addiction treatment clinics dispensing methadone, and NA centers.

4. Results

In this study, we used both descriptive, to indicate the central and distributive tendencies, and inferential statis-

tical methods. Confirmatory factor analysis was used to investigate and verify the subscales of the scale, and in other words, to confirm the construct validity. To conduct the aforementioned statistical calculations, we used SPSS.24 and Amos.21 software.

Table 1 shows the demographic and descriptive characteristics of the participants, including age and the type of drug they used.

Table 1. Frequency of Self-Injurious Behaviors Without Suicidal Intention

Self-Injurious Behaviors	N	Mean of Repeat
Swallowing chemicals	469	2.96
Needle-sticking	466	20.29
Rubbing skin against rough surfaces	467	15.01
Wound picking	468	0.33
Punching	469	3.39
Severe scratching	470	2202.56
Hair pulling	470	8.9
Pinching	470	1.48
Carving	469	8.44
Burning	470	2.46
Biting	467	4.21
Cutting	468	19.46

The main self-injurious methods used were cutting, severe scratching, punching, needle sticking, wound picking, burning, biting, hair pulling, and self-pinching. Among participants, 39% reported pain during self-injury, and 18.1% of them felt no pain. The interval between having self-injurious motivation and carrying it out was less than one hour in 62.3% and between one to three hours in 30.8% of the cases.

First, using the exploratory factor analysis, we confirmed the subscales and verified the validity of the construct in an Iranian population. Before conducting exploratory analysis, we used Kaiser-Myer-Olkin (KMO) test and Bartlett's test to examine the sample size. Kaiser-Myer-Olkin value of 0.909 was highly significant (approximate. chi-square 9225.388, $df = 741$, $P < 0.001$), which exceeds the recommended value of 0.6 for Bartlett's test of sphericity.

4.1. Intra-class Correlation Coefficient

After estimating Intra-class Correlation Coefficient (ICC), we calculated it using the standard error of measurement and the SD formula. After verifying the accuracy of the translation, the questionnaire was tested and re-tested in a preliminary study within 20 days in a small sample (30 qualified participants) with an ICC result of 0.761. The final ICC was ± 2.62 , which showed desirable reliability for the questionnaire.

4.2. Exploratory Factor Analysis

We randomly divided the data into two parts and conducted exploratory factor analysis (EFA) with principal components analysis (PCA) method on the first half and confirmatory factor analyses (CFA) on the other hand in order to confirm the factor structure obtained in EFA of the first part as recommended by Harrington (16). Principal component analysis indicated a single factor with 13 items could explain 68% of the variance (Eigen value = 8.85). Factors Loadings ranged from 0.80 to 0.87 (Table 2)

As shown in Table 2, all sub-tests are under one sub-factor. Hence, this questionnaire is a one-factor instrument in Iranian population.

4.3. Consistency

The internal consistency of all questions was high. In Table 3, the mean, standard deviation, and Cronbach's alpha of all subscales are calculated. The total Cronbach's alpha coefficient was 0.93.

Cronbach's alpha of 0.7 indicates a high internal consistency (17); therefore, the internal consistency of this subscale is desirable.

4.4. Convergent and Divergent Questionnaire

The correlation between this instrument with a convergent instrument (Barrat Impulsiveness Scale) was positive and significant (P = 0.01, r = 0.82).

In addition, its correlation with the distress tolerance scale, a divergent instrument, was negative and significant (r = -0.179, P = 0.000).

Table 2. Exploratory Factor Analysis with Varimax Rotation (N = 235)

Function Scale	Factor Loading
Affect regulation	0.77
Interpersonal boundaries	0.81
Self-punishment	0.80
Self-care	0.85
Anti-dissociation/feeling-generation	0.83
Anti-suicide	0.84
Sensation-seeking	0.83
Peer-bonding	0.85
Interpersonal influence	0.71
Toughness	0.82
Marking distress	0.77
Revenge	0.87
Autonomy	0.81




Figure 1. Factor confirmatory factor analysis for "ISAS"

4.5. Confirmatory Factor Analyses

Confirmatory factor analyses were performed with AMOS 21 on the second part of the data (N = 235) and concluded the primary model was unsuccessful in fit with data ($\chi^2 = 287.70$; $P < 0.001$; $df = 56$; $\chi^2/df = 4.42$, CFI = 0.91; RMSEA = 1.12; GFI = 0.83) (Figure 1). Based on the modification indices provided by AMOS, many error covariances were uncorrelated, and observed variables were related. After correlating five error terms according to the largest modification indices, satisfactory fit indices were obtained ($\chi^2 = 158.09$; $P < 0.001$; $df = 58$; $\chi^2/df = 2.7$; CFI = 0.96; RMSEA = 0.08; GFI = 0.90) (Figure 2).

In addition, question S4 and question S8 had the heaviest and weakest factor loadings, respectively.

5. Discussion

The present study provided evidence for psychometric properties of the Inventory of Statements About Self-Injury in a sample of opioid and alcohol abusers in Iran. The results showed ISAS is a useful inventory for the Iranian population. Findings have noetic and conceptual implications for understanding the measurement of non-suicidal self-injury in opioid and alcohol abusers. The results show the frequency of NSSI and its functions are relatively stable over time. The behavior with the greatest stability was severe scratching. Burning, pinching, and pulling hair were three of the most common and clinically severe NSSI behaviors.

The results of the confirmatory factor analysis after the exploratory factor analysis showed the factor loadings of all sub-inventories were more than 0.6, indicating that the factor rotation caused an increase in the power of factor

Table 3. Cronbach's Alpha for the Hidden Factor of "ISAS" in Iranian Population^a

Subscales	Questions: When I Self-Harm, I Am...	Mean \pm SD
Affect regulation	Calming myself down	1.06 \pm 0.7
	Releasing emotional pressure that has built up inside of me	1.11 \pm 0.73
	Reducing anxiety, frustration, anger, or other overwhelming emotions	0.47 \pm 0.66
Interpersonal boundaries	Creating a boundary between myself and others	0.95 \pm 0.72
	Demonstrating that I am separate from other people	0.84 \pm 0.74
	Establishing a barrier between myself and others	0.80 \pm 0.75
Self-punishment	Punishing myself	1.07 \pm 0.75
	Expressing anger towards myself for being worthless or stupid	0.83 \pm 0.79
	Reacting to feeling unhappy with myself or disgusted with myself	0.99 \pm 0.75
Self-care	Giving myself away to care for myself (by attending to the wound)	0.70 \pm 0.78
	Creating a physical injury that is easier to care for than my emotional distress	0.81 \pm 0.74
	Allowing myself to focus on treating the injury, which can be gratifying or satisfying	0.80 \pm 0.78
Anti-dissociation/ feeling generation	Causing pain so I will stop feeling numb	0.73 \pm 0.72
	Trying to feel something (as opposed to nothing) even if it is physical pain	1.00 \pm 1.18
	Making sure I am still alive when I do not feel real	0.76 \pm 0.77
Anti-suicide	Avoiding the impulse to attempt suicide	0.76 \pm 0.77
	Responding to suicidal thoughts without actually attempting suicide	0.81 \pm 0.70
	Putting a stop to suicidal thoughts	0.85 \pm 0.77
Sensation-seeking	Doing something to generate excitement or exhilaration	0.57 \pm 0.70
	Entertaining myself or others by doing something extremely	0.63 \pm 0.70
	Pushing my limits in a manner akin to skydiving or other extreme activities	0.75 \pm 0.70
Peer-bonding	Bonding with peers	0.61 \pm 0.71
	Fitting in with others	0.63 \pm 0.73
	Creating a sign of friendship or kinship with friends or loved ones	0.65 \pm 0.74
Interpersonal influence	Letting others know the extent of my emotional pain	1.02 \pm 0.74
	Seeking care or help from others	0.72 \pm 0.74
	Keeping a loved one from leaving or abandoning me	0.97 \pm 0.74
Toughness	Seeing if I can stand the pain	0.86 \pm 0.75
	Demonstrating I am tough or strong	0.72 \pm 0.70
	Proving I can take the physical pain	0.77 \pm 0.74
Marking distress	Creating a physical sign that I feel awful	0.77 \pm 0.74
	Proving to myself that my emotional pain is real	1.01 \pm 0.73
	Signifying the emotional distress I experience	0.81 \pm 0.75
Revenge	Getting back at someone	0.84 \pm 0.74
	Getting revenge against others	0.74 \pm 0.77
	Trying to hurt someone close to me	0.67 \pm 0.72
Autonomy	Ensuring that I am self-sufficient	0.40 \pm 0.63
	Demonstrating that I do not need to rely on others for help	0.84 \pm 0.79
	Establishing that I am autonomous/independent	0.71 \pm 0.70

^aThe Cronbach alpha (α) for all items was 0.93.

loadings sub-inventories. Cronbach's alpha for the total score and both hidden factors were similar to the original version of the inventory in a study by Klonsky et al. (5, 6), which was higher than 0.7, indicating this inventory had a good fit for the Iranian population of opioid and alcohol abusers.

The results of the study by Klonsky et al. have suggested separate categories (interpersonal and intraper-

sonal) for NSSI functions (5, 6). However, some studies emphasize NSSI functions overlap and are interrelated (18). According to a meta-analysis, intrapersonal functions are less common causes of NSSI compared to interpersonal functions such as emotional regulation, which is often difficult for substance abusers. Nevertheless, the above-mentioned study, as well as the current study, do not suggest that focus should only be on one function in an individual (19).




Figure 2. Factor confirmatory factor analysis for "ISAS"

In factor analysis, "Self-care" and "peer-bonding" had the highest and lowest factor loadings, respectively. The theory of social learning supports these findings. Sometimes adolescents, by injuring themselves, reinforce self-care behavior, seek attention from their family, and gain social status among their peers.

Substance abuse is associated with suicidal tendencies. Non-lethal suicide attempts can be a risk factor for future suicide attempts and one of the main reasons for hospitalization in psychiatric wards (20). This study examined self-cutting and slit wrestling, the most important types of self-injurious behaviors among drug abusers (21, 22). Limited access to dangerous means of suicide may play an important role in reducing self-injury attempts in the studied sample. The most important groups at risk are youths, prisoners, and all those committed intentional self-injurious behaviors during their lifetime (23).

Given the convergent and divergent validity, there is a positive and significant correlation between both distress tolerance and impulsivity with intentional self-injurious behaviors, consistent with previous studies. In addition, alcohol abuse and self-injurious behaviors are associated with low degrees of distress tolerance (19). Distress tolerance, an individual difference variable, points to the capacity of experience and resistance to emotional distress.

Distress tolerance has increasingly been seen as an important concept in development, a new insight into the onset and maintenance of mental health, as well as prevention and treatment of mental disorders. People with low distress tolerance engage in behavioral deviances such as drug abuse in an erroneous attempt to deal with their negative emotions and to relieve their psychological pain (10). Impulsivity is also a key factor in the incidence of NSSI behaviors (23). Addiction is a powerful predictor of the persistence of drug abuse, and understanding the relationships between these factors can help reduce the recurrence of drug abuse (24).

Moreover, ISAS is the first inventory localized among opioid and alcohol abusers in Iran, which can safely and reliably examine NSSI behavior in this population. Therapists can now evaluate patients' responses to the treatment and follow up on their progress over time. In addition to interview, therapists can use this instrument to create treatment plans. This tool can also be used to monitor treatment progress and reduction in self-injurious behaviors. Therefore, it is important to adequately train the staff to use this instrument.

A limitation of this study was respondents' low educational level; the researcher resolved this limitation by personally reading all the questions for the participants. Also, the fact that very few study participants (1.06%) were females makes it difficult to generalize the results to the female population. In addition, this study only examined individuals, abusing opioids, and alcohol. Manifestations and related functions may be different in abusers of other substances and hence limits the generalizability of the findings. Future studies can examine self-injurious behaviors and the applicability of ISAS, as an instrument, in the female population as well as in abusers of substances other than opioids and alcohol.

The most important achievement of this study is that it provides thorough evidence based on data collected from a sample covering a variety of cultures and ethnicities from multiple regions of the country, including the north, south, east, west, and central parts of Iran. Therefore, this tool can be used as a representative measure of relevant behaviors in the Iranian population.

5.1. Conclusions

The Persian version of the Inventory of Statements About Self-Injury has high validity and reliability among the Iranian population of opioid and alcohol abusers. Further research is needed to investigate the causal relationship between clinical symptoms and suicidal ideation and decreased resilience.

Footnotes

Authors' Contribution: Mehran Zarghami, Masoudeh Babakhanian, and Mojtaba Habibi Asgarabad designed the study; Masoudeh Ghazanfarpour, Mojtaba Habibi Asgarabad, and Masoudeh Babakhanian conducted the statistical analysis; Faezeh Sadat Akrami, Nader Molavi, Yasaman Allameh, and Najmodin Nazeri contributed to data collection; Hamid Sharif Nia and Masoudeh Babakhanian and Mehran Zarghami wrote the first draft of the manuscript. All authors read and approved the contents of the final manuscript.

Conflict of Interests: The authors declare that they have no conflict of interest.

Ethical Approval: The Ethics Committee of the Student Research Committee of Mazandaran University of Medical Sciences approved this study on 14/01/2017 (IR.MAZUMS.REC.1395.371).

Funding/Support: Mazandaran University of Medical Sciences (grant number 371).

Informed Consent: All individual participants were informed about the study and were asked to complete the questionnaires only if they consent to the study.

References

- Sansone RA, Sansone LA. Measuring self-harm behavior with the self-harm inventory. *Psychiatry (Edgmont)*. 2010;7(4):16–20. [PubMed: 20508804]. [PubMed Central: PMC2877617].
- American Psychiatric Pub. *Diagnostic and statistical manual of mental disorders (DSM-5®)*. 2013.
- Zetterqvist M. The DSM-5 diagnosis of nonsuicidal self-injury disorder: a review of the empirical literature. *Child Adolesc Psychiatry Ment Health*. 2015;9:31. doi: 10.1186/s13034-015-0062-7. [PubMed: 26417387]. [PubMed Central: PMC4584484].
- Suyemoto KL. The functions of self-mutilation. *Clin Psychol Rev*. 1998;18(5):531–54. doi: 10.1016/s0272-7358(97)00105-0. [PubMed: 9740977].
- Klonsky ED, Oltmanns TF, Turkheimer E. Deliberate self-harm in a nonclinical population: prevalence and psychological correlates. *Am J Psychiatry*. 2003;160(8):1501–8. doi: 10.1176/appi.ajp.160.8.1501. [PubMed: 12900314]. [PubMed Central: PMC4362719].
- Poorolajal J, Haghtalab T, Farhadi M, Darvishi N. Substance use disorder and risk of suicidal ideation, suicide attempt and suicide death: a meta-analysis. *J Public Health (Oxf)*. 2016;38(3):e282–91. doi: 10.1093/pubmed/fdv148. [PubMed: 26503486].
- Ekhtiari H, Safaei H, Esmaeeli Djavid G, Atefvahid MK, Edalati H, Mokri A. Reliability and validity of Persian versions of Eysenck, Barratt, Dickman and Zuckerman Questionnaires in assessing risky and impulsive behaviors. *Iranian journal of psychiatry and clinical psychology*. 2008;14(3):326–36.
- Munro BH. *Statistical methods for health care research*. Lippincott Williams & Wilkins; 2005.
- Azizi AR. Reliability and validity of the Persian version of distress tolerance scale. *Iran J Psychiatry*. 2010;5(4):154–8. [PubMed: 22952509]. [PubMed Central: PMC3395925].
- Bildik T, Somer O, Kabukcu Basay B, Basay O, Ozbaran B. [The validity and reliability of the Turkish version of the inventory of statements about self-injury]. *Turk Psikiyatri Derg*. 2013;24(1):49–57. Turkish. [PubMed: 23446540].
- Kortge R, Meade T, Tennant A. Interpersonal and Intrapersonal Functions of Deliberate Self-Harm (DSH): A Psychometric Examination of the Inventory of Statements About Self-Injury (ISAS) Scale. *Behaviour Change*. 2013;30(1):24–35. doi: 10.1017/bec.2013.3.
- Lindholm T, Bjarehed J, Lundh LG. Functions of nonsuicidal self-injury among young women in residential care: a pilot study with the Swedish version of the Inventory of Statements about Self-Injury. *Cogn Behav Ther*. 2011;40(3):183–9. doi: 10.1080/16506073.2011.565791. [PubMed: 21877957].
- Barratt ES, Stanford MS, Kent TA, Felthous A. Neuropsychological and cognitive psychophysiological substrates of impulsive aggression. *Biol Psychiatry*. 1997;41(10):1045–61. doi: 10.1016/s0006-3223(96)00175-8. [PubMed: 9129785].
- Simons JS, Gaher RM. The Distress Tolerance Scale: Development and Validation of a Self-Report Measure. *Motivation and Emotion*. 2005;29(2):83–102. doi: 10.1007/s11031-005-7955-3.
- Rubio DM, Berg-Weger M, Tebb SS, Lee ES, Rauch S. Objectifying content validity: Conducting a content validity study in social work research. *Social Work Research*. 2003;27(2):94–104. doi: 10.1093/swr/27.2.94.
- Harrington D. *Confirmatory factor analysis*. Oxford University Press; 2009. doi: 10.1093/acprof:oso/9780195339888.001.0001.
- Hair J. *Multivariate data analysis*. Pearson College Division; 2010.
- Lloyd-Richardson EE, Perrine N, Dierker L, Kelley ML. Characteristics and functions of non-suicidal self-injury in a community sample of adolescents. *Psychol Med*. 2007;37(8):1183–92. doi: 10.1017/S003329170700027X. [PubMed: 17349105]. [PubMed Central: PMC2538378].
- Taylor PJ, Jomar K, Dhingra K, Forrester R, Shahmalak U, Dickson JM. Corrigendum to "A Meta-Analysis of the Prevalence of Different Functions of Non-Suicidal Self-Injury" [Journal of Affective Disorders (2017) 759–769]. *J Affect Disord*. 2019;259:440. doi: 10.1016/j.jad.2019.08.072. [PubMed: 31611001].
- Lee J, Bang YS, Min S, Ahn JS, Kim H, Cha YS, et al. Characteristics of adolescents who visit the emergency department following suicide attempts: comparison study between adolescents and adults. *BMC Psychiatry*. 2019;19(1):231. doi: 10.1186/s12888-019-2213-5. [PubMed: 31349782]. [PubMed Central: PMC6660711].
- Yeum TS, Kim B, Kim EY, Kim SH, Ha K, Ahn YM. Factors Affecting Suicide Method Lethality Among Suicide Attempters in the Korea National Suicide Survey. *J Nerv Ment Dis*. 2018;206(3):202–10. doi: 10.1097/NMD.0000000000000735. [PubMed: 28902068].
- Ersen B, Kahveci R, Saki MC, Tunali O, Aksu I. Analysis of 41 suicide attempts by wrist cutting: a retrospective analysis. *Eur J Trauma Emerg Surg*. 2017;43(1):129–35. doi: 10.1007/s00068-015-0599-4. [PubMed: 26626872].
- Wu N, Hou Y, Chen P, You J. Peer Acceptance and Nonsuicidal Self-Injury among Chinese Adolescents: A Longitudinal Moderated Mediation Model. *J Youth Adolesc*. 2019;48(9):1806–17. doi: 10.1007/s10964-019-01093-0. [PubMed: 31385231].
- Viana AG, Woodward EC, Raines EM, Hanna AE, Zvolensky MJ. The role of emotional clarity and distress tolerance in deliberate self-harm in a sample of trauma-exposed inpatient adolescents at risk for suicide. *Gen Hosp Psychiatry*. 2018;50:119–24. doi: 10.1016/j.genhosppsych.2017.10.009. [PubMed: 29161582].