

Pericarditis in a 56-year-old female with Hashimoto's disease

Zapalenie osierdzia u 56-letniej kobiety z chorobą Hashimoto

Angelika Miazga¹, Anna Wolska-Bułach¹, Beata Begier-Krasińska¹, Maciej Osiński², Andrzej Tykarski¹

¹Department of Hypertensiology, Angiology, and Internal Diseases, Poznan University of Medical Sciences, Poznan, Poland

²Poznan University of Medical Sciences, Poznan, Poland

A 56-year-old woman was hospitalised in the Department of Hypertensiology due to pericardial effusion revealed in routine echocardiographic examination. *First hospitalisation:* The patient reported malaise and exercise intolerance for 18 months. On physical examination regular cardiac function, 80/min with proper accentuation and blood pressure 143/91 mm Hg was seen. *Comorbidities:* Hypothyroidism from 8 years (during the last year TSH varied from 22.93 to 41.42 μ IU/mL) and well-controlled hypertension. *In ECG:* Sinus rhythm, negative T wave in V1. *In echocardiography:* Echo-free space was distinguished behind the lateral wall of the left ventricle (LV) in the amount of 2.0–2.5 cm, and behind the posterior wall: 1.5–1.9 cm (Fig. 1). The sizes of heart cavities and systolic function (LVEF = 65%) were correct. Meagre tricuspid regurgitation was recorded (RVSP \sim 38 mm Hg). The inferior vena cava was undulated with correct respiratory mobility. Significant variability of flows through the atrioventricular valves during respiration was seen. Slight pressure on the right atrium and its collapse during ventricular contraction was observed. Right ventricular (RV) diastolic collapsing and paradoxical movement of the intraventricular septum were not distinguished. Despite significant pericardial effusion, symptoms of tamponade were not observed. At discharge, pericardial effusion decreased slightly (Fig. 2). *Laboratory results:* Elevated levels of anti-thyroid peroxidase and antithyroglobulin antibodies; TSH = 1.7 μ IU/mL within normal range. *Treatment:* Methylprednisolone (16 mg/day) in gradually decreasing dose, colchicine (1 mg/day), levothyroxine (175/200 μ g), and perindopril. The patient was recommended to take these drugs after discharge. *Control hospitalisation.* After 6 weeks the patient did not report any complaints. *In echocardiography:* Echo-free space was distinguished behind the lateral wall of the LV in the amount of 2.1–2.6 cm (Figs. 3, 4), thus significant decrease of effusion did not occur. Pericardial thickening with echo enhancement in the area of tricuspid annulus and RV was found. LV filling profile and inflow from the pulmonary veins were correct. *Laboratory results:* Decreased level of TSH (0.2 μ IU/mL); erythrocyte sedimentation rate, C-reactive protein, and procalcitonin were normal. *Magnetic resonance imaging (MRI) of the heart* revealed a substantial quantity of partially organised fluid in the pericardial sac (Fig. 5). The maximum quantity of fluid occurred behind the LV (26 mm). Over the RV and partially over the LV, pericardial visceral lamina thickening (up to 9 mm) and calcifications were distinguished. Based on MRI, constrictive pericarditis was diagnosed. *Conclusions:* Our patient suffered from long-lasting hypothyroidism. This is the reason why we did not see a rapid decrease in pericardial effusion despite adequate therapy. Proper thyroid replacement therapy causes a slow process of reabsorption of pericardial effusions. According to literature, pericardial effusion incidence in hypothyroidism amounts to 3–6%. Tamponade and obstructive pericarditis occur very rarely in these patients. Echocardiographic assessment of pericardial thickening is troublesome due to the phenomenon of complete ultrasound bouncing from pericardium (between the heart muscle and lung tissue), so MRI is a powerful tool to establish constrictive pericarditis diagnosis.



Figure 1. Echocardiography — first hospitalisation (apical four-chamber projection); LV — left ventricle




Figure 2. Echocardiography at discharge; LV — left ventricle




Figure 3. Control echocardiography (apical four-chamber projection); LV — left ventricle




Figure 4. Control echocardiography (subcostal short-axis projection at the height of the papillary muscles)




Figure 5. Magnetic resonance imaging of the heart; LV — left ventricle; RV — right ventricle

Address for correspondence:

Angelika Miazga, MD, Department of Hypertensiology, Angiology, and Internal Diseases, Poznan University of Medical Sciences, ul. Starowiejska 1B, 61–664 Poznań, Poland, e-mail: angelika.miazga@gmail.com

Conflict of interest: none declared