

## SHORT COMMUNICATION

# Crossover from transradial to ipsilateral transulnar access after sheath insertion into the radial artery

Farshad Roghani-Dehkordi<sup>1,2</sup>, Mohammad Kermani-Alghoraishi<sup>1,2</sup>, Kiyan Heshmat-Gahdarijani<sup>2,3</sup>

1 Interventional Cardiology Research Center, Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran

2 Cardiology Department, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

3 Cardiac Rehabilitation Research Center, Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran

**Introduction** The radial artery has been the preferred access site for percutaneous coronary procedures since it has some proven clinical benefits.<sup>1-3</sup> However, in some studies, the failure of transradial access (TRA) was reported in 1% to 7% of cases, mainly due to the inability of puncture, radial artery spasm or dissection, hypoplastic or small radial artery, and failure of catheter passage to the subclavian artery and ascending aorta.<sup>1,4</sup> In these situations, the ipsilateral transulnar access (TUA) has shown to be a safe and feasible alternative to contralateral TRA for coronary interventions.<sup>5,6</sup> In the present report, we describe 11 cases of successful crossover performed from the radial to ipsilateral ulnar artery after sheath insertion into the radial artery.

**Methods** This was a prospective, descriptive, observational study on 11 patients who were candidates for elective coronary angiography due to stable ischemic heart disease with unsuccessful TRA catheterization at 2 distinctive hospitals (Shahid Chamran [7 patients] and Khorshid [4 patients] hospitals, Isfahan, Iran) from January 2012 to September 2018. Our alternative planned approach was to perform a crossover from the radial to ipsilateral ulnar artery after sheath insertion into the radial artery. The crossover from the radial to femoral arteries was not possible due to severe obesity and/or peripheral vascular disease. All procedures were performed by 2 interventional cardiologists with expertise in TRA and ipsilateral TUA approaches.

The right radial access was used for all patients. After sheath insertion into the radial artery, the advancement of the catheter was unsuccessful due to complications; therefore,

the procedure was incomplete. In 6 patients, a Tiger catheter (Terumo, Tokyo, Japan) could not be advanced due to tortuosity or loop of the radial artery (FIGURE 1A), which caused patient discomfort after several attempts to repeat the procedure. In 2 patients, the catheter could not be advanced due to a severe spasm of the radial artery (FIGURE 1B); in 2 other patients, dissection of the radial artery occurred; and in 1 patient, advancement failed because the patient had a high take-off radial artery (FIGURE 1C). Therefore, due to limitation of the femoral access, we performed the crossover from TRA to ipsilateral TUA. Local anesthesia (2 ml of 2% lidocaine) was infiltrated about 1 inch proximal to the flexor crease where the most powerful pulsation of the ulnar artery was sensed. Then the ulnar artery was accessed and the hydrophilic 5F or 6F sheath was introduced over the guide wire (FIGURE 1D and 1E). Intravenous unfractionated heparin was administered (50–70 U/kg, up to 5000 units),<sup>7</sup> and to reduce ulnar artery spasm, intra-arterial injection of diluted verapamil (2.5 mg) was used. Coronary angiography or angioplasty was successful in all patients without any further hindrance and complication.

All patients provided written informed consent to participate in the study. No ethics committee approval was required in this study.

**Statistical analysis** Continuous and categorical variables for the 11 analyzed cases were reported as mean (SD) and frequency, respectively.

**Results and discussion** The mean (SD) age of patients was 57.8 (5.7) years. There were 7 men and 4 women. The mean body mass index of the patients was 32.5 kg/m<sup>2</sup>. Atherosclerotic


**Correspondence to:** Mohammad Kermani-Alghoraishi, MD, Interventional Cardiology Research Center, Isfahan Cardiovascular Research Institute, Salman Farsi, PO Box 8158388994, Isfahan, Iran, phone: +98 31 36115310, email: sm.kermani@med.mui.ac.ir  
Received: October 23, 2019.

Revision accepted: February 11, 2020.

Published online: February 18, 2020.

Kardiol Pol. 2020; 78 (4): 325-327  
doi:10.33963/KP.15193

Copyright by the Author(s), 2020



**FIGURE 1** Angiographic images of the radial artery tortuosity (A), radial artery spasm (B), radial artery with high take-off (C), and radial and ipsilateral ulnar artery sheath insertion (D); E – cannulation of both radial and ulnar arteries

risk factors were as follows: diabetes in 7 patients; hypertension, in 8; dyslipidemia in 6; and current smoking, in 5. Five of the patients underwent percutaneous coronary intervention in addition to coronary angiography. All of the procedures were successful. In the first cases, we achieved hemostasis after sequential

compression, ie, the radial artery compression via TR BAND (Lepu Medical Technology Co., Beijing, China) followed by the ulnar artery compression. In most cases, we used a simultaneous hemostasis method with 2 overlapping balloon-based compression devices on the radial and ulnar arteries. No complications such as hematoma, pain or paresthesia, pseudoaneurysm formation, arterial obstruction, or limb ischemia were recorded during hospitalization or the 1-year follow-up. Radial artery occlusion occurred in 1 and 3 cases in the early (first 24 hours) and late period, respectively.

It has been shown that TUA can be a safe and feasible alternative approach for cardiovascular interventions when the ipsilateral radial artery is inaccessible.<sup>6</sup> Recent studies have reported a high success rate and an extremely low incidence of puncture site complications for TUA, which was similar to data reported for TRA.<sup>8,9</sup> On the other hand, the cannulation of the ulnar artery is associated with longer procedural and fluoroscopy times and a higher crossover rate compared with TRA.<sup>10</sup> While the radial artery has a more superficial course, it is readily palpable and compressible, which makes TRA a more preferable approach than TUA.

After TRA failure, the most common alternative approach is transfemoral or contralateral TRA. Despite the limitations of the femoral access, when the mechanism of failure is the radial artery itself, ipsilateral TUA may be considered.<sup>11</sup> In our patients, tortuosity, dissection or perforation, severe spasm, and the radial artery with a high take-off were the reasons for discontinuation of the procedure and the crossover from TRA to TUA; however, the use of distal radial artery access has been reported to be safe and helpful in these cases.<sup>12</sup>

There are 2 major concerns about simultaneous sheath insertion in both the radial and ulnar arteries. The first concern is hand ischemia due to obstruction of the 2 major arteries supplying the hand by 2 sheaths during the procedure. Kedev et al<sup>6</sup> reported no occurrence of hand ischemia in patients with radial artery occlusion undergoing ipsilateral transulnar catheterization procedures. This was most likely due to rapid recruitment of collateral flow from the interosseous arteries. The second concern is the simultaneous hemostasis of both arteries. Manual compression is feasible but using 2 overlapping balloon-based compression devices on the radial and ulnar arteries is also helpful. A Pulsera hemostatic device (Accumed Radial Systems LLC., Ann Arbor, Michigan, United States) also could be safe and practical for achieving simultaneous hemostasis in the radial and ulnar arteries.

Regarding complications, it should be noted that there are several causes of a relatively high incidence of radial artery occlusion (approximately 35%), including multiple radial artery

puncture and manipulation, prolonged sheath removal time, atretic or small-diameter radial artery, severe tortuosity and loop, high take-off, and limited experience in hemostatic strategies.

This report demonstrated that ipsilateral TUA could be a secure and viable alternative approach for cardiovascular interventions in case of inaccessible radial and femoral arteries.

## ARTICLE INFORMATION

**CONFLICT OF INTEREST** None declared.

**OPEN ACCESS** This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND 4.0), allowing third parties to download articles and share them with others, provided the original work is properly cited, not changed in any way, distributed under the same license, and used for non-commercial purposes only. For commercial use, please contact the journal office at [kardiologiapolska@ptkardio.pl](mailto:kardiologiapolska@ptkardio.pl).

**HOW TO CITE** Roghani-Dehkordi F, Kermani-Alghoraishi M, Heshmat-Ghahdarjani K. Crossover from transradial to ipsilateral transulnar access after sheath insertion into the radial artery. *Kardiol Pol.* 2020; 78: 325-327. doi:10.33963/KP.15193

## REFERENCES

- 1 Jolly SS, Yusuf S, Cairns J, et al. Radial versus femoral access for coronary angiography and intervention in patients with acute coronary syndromes (RIVAL): a randomized, parallel group, multicenter trial. *Lancet.* 2011; 377: 1409-1420.
- 2 Wagener JF, Rao SV. A comparison of radial and femoral access for cardiac catheterization. *Trends Cardiovasc Med.* 2015; 25: 707-713.
- 3 Mason PJ, Shah B, Tamis-Holland JE, et al. An update on radial artery access and best practices for transradial coronary angiography and intervention in acute coronary syndrome: a scientific statement from the American Heart Association. *Circ Cardiovasc Interv.* 2018; 11: e000035.
- 4 Roberts JS, Niu J. A novel technique for simultaneous hemostasis of ipsilateral radial and ulnar artery access sites. *Catheter Cardiovasc Interv.* 2018; 9: 901-904.
- 5 de Andrade PB, Tebet MA, Nogueira EF, et al. Transulnar approach as an alternative access site for coronary invasive procedures after transradial approach failure. *Am Heart J.* 2012; 164: 462-467.
- 6 Kedev S, Zafirovska B, Dharma S, Petkoska D. Safety and feasibility of transulnar catheterization when ipsilateral radial access is not available. *Catheter Cardiovasc Interv.* 2014; 83: E51-E60.
- 7 Roghani F, Shirani B, Hashemifard O. The effect of low dose versus standard dose of arterial heparin on vascular complications following transradial coronary angiography: Randomized controlled clinical trial. *ARYA Atheroscler.* 2016; 12: 10-17.
- 8 Aptekar E, Pernes JM, Chabane-Chaouch M, et al. Transulnar versus transradial artery approach for coronary angioplasty: the PCVI-CUBA study. *Catheter Cardiovasc Interv.* 2006; 67: 711-720.
- 9 Roghani-Dehkordi F, Mansouri R, Khosravi A, et al. Transulnar versus transradial approach for coronary angiography and angioplasty: Considering their complications. *ARYA Atheroscler.* 2018; 14: 128-131.
- 10 Hahalis G, Tsigkas G, Xanthopoulos I, et al. Transulnar compared with transradial artery approach as a default strategy for coronary procedures: a randomized trial. The Transulnar or Transradial Instead of Coronary Transfemoral Angiographies Study (the AURA of ARTEMIS Study). *Circ Cardiovasc Interv.* 2013; 6: 252-261.
- 11 Singh V, Cohen MG. Crossover from radial to ipsilateral ulnar access: an additional strategy in the armamentarium of the "radialist." *Cath Lab Digest.* 2015; 23: 10-11.
- 12 Kontopodis E, Rigatou A, Tsiafoutis I, et al. Snuffbox to the rescue: distal transradial approach for cardiac catheterization after failed ipsilateral radial puncture. *Kardiol Pol.* 2018; 76: 1491.