



University of Dundee

Abreast of Health

Sinclair, Julia M. A.; Dutey-Magni, Peter F; Anderson, Annie; Baird, Janis; Barker, Mary E; Cutress, Ramsey I

Publication date:
2020

Document Version
Early version, also known as pre-print

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Sinclair, J. M. A., Dutey-Magni, P. F., Anderson, A., Baird, J., Barker, M. E., Cutress, R. I., Kaner, E. F. S., McCann, M., Priest, C. K., & Copson, E. R. (2020, Jan 24). Abreast of Health: Development of a contextspecific digital alcohol brief intervention (ABI) in symptomatic breast clinics . (1 ed.)

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Original Paper

Abreast of Health: Development of a context-specific digital alcohol brief intervention (ABI) in symptomatic breast clinics

Abstract

Background: Potentially modifiable risk factors account for approximately 23% of breast cancer cases. In the UK, alcohol consumption alone is held responsible for 8-10% of cases diagnosed every year. Symptomatic breast clinics focus on early detection and treatment, but also offer scope for delivery of low-cost lifestyle interventions to encourage a cancer prevention culture within the cancer care system. Careful development work is required to effectively translate such interventions to novel settings.

Objective: To develop a theory of change and delivery mechanism for a context-specific alcohol and lifestyle brief intervention aimed at women attending screening and symptomatic breast clinics.

Methods: A formative study combined evidence reviews, analysis of mixed-method data, and user experience research to develop an intervention model, following the Six Steps in Quality Intervention Development (6SQulD) framework.

Results: A web application focused on: improving awareness, encouraging self-monitoring, and reframing alcohol reduction as a positive choice to improve health was found to be acceptable to women. Accessing this in the clinic waiting area on a tablet computer was shown to be feasible. An important facilitator for change may be the heightened readiness to learn associated with a salient health visit (a 'teachable moment'). Women may have increased motivation to change if they can develop a belief in their capability to monitor and, if necessary, reduce their alcohol consumption.

Conclusions: Using the 6SQulD framework supported the prototyping and maximized acceptability and feasibility of an alcohol brief intervention for women attending symptomatic breast clinics, regardless of their level of alcohol consumption.

Keywords: Cancer; Lifestyle determinants; Intervention theory; Health promotion; Alcohol brief intervention; Digital health intervention

Introduction

Breast cancer is the most common type of cancer worldwide, and its incidence is rising [1]. The World Health Organisation considers that sufficient knowledge is available to prevent 30-50% of cancer cases globally and that 'prevention offers the most cost-effective long-term strategy for the control of cancer' [2]. In the UK, the proportion of breast cancer cases attributable to lifestyle factors is: insufficient physical activity: 2%; overweight/obesity: 8%; and alcohol consumption: between 8% [3] and 10% [4]. Alcohol increases the risk of breast cancer in a dose-dependent fashion even from low levels, with an estimated relative risk of 1.09 for 10g/day [5]. Observational evidence shows that alcohol consumption may also increase the risk of recurrence of breast cancer in survivors [6,7]. New UK clinical guidelines advise this group to observe an upper limit of 5 units per week [8].

Systematic reviews of alcohol interventions indicate that, outside of regulatory interventions, alcohol brief interventions (ABIs) demonstrate the greatest effectiveness and cost-effectiveness [9-11], with small reductions in alcohol consumption (20g/week) which can be sustained for at least a year [12,13]. Despite this, ABIs remain relatively underutilised across health care systems. In England, fewer than 7% of 'increased-risk' drinkers recall receiving advice from their GP on their alcohol consumption in the past year, compared with 50% of smokers who recalled receiving tobacco cessation advice [14].

The use of 'teachable moments' is increasingly advocated to encourage modification of lifestyle determinants of cancers [15-17], but more research is required as to how best to situate health prevention interventions into current health systems. In England, over 540,000 women annually attend UK National Health Service (NHS) symptomatic breast clinics [18] as part of a rapid referral (two-week wait) system to prevent delay in diagnosis. However, as fewer than 8% of women attending are found to have breast cancer [19], but health promotion information is not offered to those without a diagnosis, in prevention terms the majority do not currently benefit from attending the clinic.

Previous research has criticised the premature trialling of ABIs in new environments, with recommendations that 'applications of brief intervention to novel settings should begin with foundational research and developmental studies' [20]. This paper describes the development of a context-specific ABI aimed at women attending symptomatic breast clinics guided by the Six Steps in Quality Intervention Development (6SQUID) [21], a framework commonly employed in the development of public health interventions.

Methods

Framework

The 6SQuID framework [21] is intended to improve the design of public health interventions and, consequently, their effectiveness. The present study synthesized information from four sources of data (reviews, empirical data from the target population, theory and concept mapping, and iterative content appraisal and design) to complete these steps in the breast health setting (Table 1).

Table 1. Six Steps in Quality Intervention Development as applied in the development of Abreast of Health (adapted from [21])

Step	Data provenance ^a	Methods
1. Define and understand the problem and its causes.	Attitudes literature (E)	Scoping review
	Scoping study [22] (E)	
2. Clarify which causal or contextual factors are malleable and have greatest scope for change.	Risk attitude literature (E)	Scoping review; theory mapping
	Scoping study [22] (E)	
	Review of existing apps (N)	
3. Identify how to bring about change: the change mechanism.	Behaviour change technique review (E)	Theory and concept mapping
4. Identify how to deliver the change mechanism.	Behaviour change technique review (E)	Concept mapping
	User testing (N)	Agile prototyping
5. Test and refine on small scale.	User testing (N)	'Think aloud' and 'teach me back' cognitive interviewing
6. Collect sufficient evidence of effectiveness to justify rigorous evaluation/implementation	To be addressed in future publication (N)	

ion.	
------	--

^a(E): existing data from the public domain; (N): new data generated during this study.

Reviews

The academic and grey literature were reviewed iteratively in three different areas relevant to the intervention (Multimedia Appendix 1).

1. Knowledge and social attitudes to alcohol amongst women (particularly in the UK), and amongst health care staff. This included information on knowledge of alcohol volumes, effect of alcohol on health, and confidence in managing alcohol related health risks.
2. Knowledge and social attitudes in relation to modifiable risk factors for cancer. Particular attention was paid to interaction with social determinants of health, including health literacy, socio-economic status and social deprivation.
3. Findings from existing reviews on behaviour change mechanisms and techniques for reducing alcohol consumption. In addition to reviews from the Cochrane library, we focused on systematic and narrative reviews of features of digitally-delivered ABIs [23–27].

Mixed Method Study with the Target Population

A mixed-method study was undertaken to complement evidence from the literature reviews with data from the target environment: symptomatic breast clinics and an NHS Breast Screening Programme unit in Southampton, UK. A total of 205 women attending appointments were recruited to take part in (a) a survey of knowledge of risk factors for breast cancer, and alcohol beverage content, and (b) five focus groups. Thirty-three health professionals took part in a similar survey, of whom eight also participated in semi-structured interviews. The full detail is reported separately [22], but will be referred to here as part of the intervention development process.

Theory and Concept Mapping

As part of 6SQulD steps 3-4, relevant theories and behaviour change constructs were reviewed, and mapped onto harmonised constructs from two systematic collations of health psychology theories commonly used in meta-analyses. These were (a) the 26 Mechanisms of Action [28] consolidating and extending the pre-existing Theoretical Domains Framework [29]; and (b) the 93 Behaviour Change Techniques (BCTs) from the BCT Taxonomy v1.1 [30].

Iterative Content Appraisal and Design

The structure and content (both textual and visual) of the intervention prototype were designed by JMAS, PDM, CKP and SEC in an 'Agile' approach [31] between December 2016 and April 2017. This method relied on rapid prototyping and testing of small components using short cycles:

- (a) The research team scoped, reviewed and appraised existing alcohol information leaflets designed by health organisations and charities, as well as alcohol web and smartphone apps. This involved mapping BCTs, appraising the language, tone and focus of different approaches to consolidate a view of the most adapted content. A particular focus was placed on identifying features that were deemed difficult to understand, insufficiently relevant, or that could be perceived by some women as 'scary' and/or judgemental. Similarly, features that appeared most helpful at implementing target mechanisms of change were also noted.
- (b) Ten women recruited from symptomatic breast clinics were invited to test and comment on a range of existing health apps in one focus group, adding to findings from the team's own analysis (Abreast of Health Phase 2; REC reference 17/LO/0953; IRAS 227642).
- (c) The research team sketched the visual layout of small components of the intervention.
- (d) Immediate comments and reactions on early versions of wording and visual features of these components were invited from 161 women recruited from symptomatic breast clinics (Abreast of Health Phase 3; REC reference 18/SC/0120; IRAS 241069). Participants took part in face-to-face cognitive interviews, which invited them to 'think aloud' and 'teach back' information gathered while testing the prototype to the researcher [32,33].

New findings were discussed by the research team on a weekly basis, setting objectives for the next data collection cycle the following week. Conclusions from these activities were mapped to a particular component of the emerging prototype intervention and recorded on a Kanban board (using the Trello software [34]) together with lists of actions, to incorporate them in the design work at every iteration of the weekly cycles.

All participants were recruited from women attending the symptomatic breast clinics at Southampton General Hospital on referral from their primary care physician. All participants were approached in the waiting room and, having given consent, either participated at that time, and/or agreed to take part in a focus group/testing session at a later date.

Results

Steps 1-2: Causal and Contextual Factors of the Target Problem

Having identified alcohol consumption as a potentially modifiable lifestyle cause of breast cancer, we undertook a broad review of underpinning factors (Multimedia Appendix 1). Table 2 gives a thematic summary of dominant themes of social and psychological determinants of knowledge, attitudes and behaviour around alcohol consumption.

Key findings were that, although 60-72% of women attending breast screening appointments or symptomatic breast clinics drink alcohol, only 20% were aware it was a risk factor for breast cancer [22,35-37]. Despite efforts from public campaigns informing the population of the effects of alcohol on long-term health, recent studies still demonstrate that the UK population recognize these far less than the social harms of alcohol. This focus on risks associated with 'binge' drinking (high-intensity, single-occasion alcohol use) can dim awareness of the effects of consuming alcohol in lesser quantities across a sustained period. A recent UK-based qualitative study by Khadjesari *et al.* [38] examined attitudes to alcohol and UK low-risk drinking UK guidelines (recommendation not to drink more than 14 units a week on a regular basis, keeping several drink-free days per week [39]) amongst adults attending primary care facilities. Authors argue that this incomplete public understanding of risks reduces the perceived relevance of low-risk drinking guidelines, and explains the participants' belief that the 14-unit threshold is unnecessarily low.

From steps 1-2, we concluded that the greatest scope for change resides in increasing awareness of alcohol's role in promoting chronic conditions such as cancer, even at low levels. This interacts with other behavioural predictors listed in Table 2, some of which are situated in the cancer context. For instance, attitudes and beliefs such as cancer pre-determinism and fatalism affect engagement with prevention behaviours [40,41] and the perceived relevance of information of lifestyle risk factors.

Table 2. Thematic summary of social and psychological determinants of knowledge, attitudes and behaviour around alcohol consumption

Domain	Evidence
Knowledge: low alcohol literacy	Only 20% of women in breast clinics [22] identified alcohol consumption as a risk factor for breast cancer, a similar proportion as in the general population [35,36]. This lack of awareness is singled out as an obstacle in promoting low-risk drinking by the UK Chief Medical Officers [39].

	<p>Some common beliefs about alcohol and cancer are incorrect, for instance that alcohol only becomes a health risk in ‘problem drinkers’ or people who are alcohol-dependent; red wine being the only type of alcohol which causes cancer, and conversely, red wine/moderate alcohol intake being good for health; physical exercise mitigating the effects of heavy drinking [38].</p>
<p>Knowledge: low alcohol numeracy</p>	<p>Individuals do not always accurately recall the frequency, volume, and concentration of alcohol they drink [42,43]. Improving numeracy, and encouraging monitoring of alcohol intake within primary care has been proposed by some [44,45] as a population prevention strategy.</p>
<p>Social role and identity of health professionals</p>	<p>In addition to lacking time and relevant training on lifestyle interventions, health care staff may not believe it is part of their clinical role to discuss lifestyle factors in relation to modifiable risk factors for cancer [46–49]. Evidence also points to health professionals lacking awareness of the causes of cancer, relevant lifestyle guidance and the appropriate advice to give [46,50,51], and lacking confidence that information will motivate women to change behaviours [46,50–52], sometimes hindered by the health care professionals’ own lifestyle choices [53]. Clinicians perceive a lack of patient interest in the subject [51] and tend to underestimate evidence that changing behaviours affects breast cancer risk [46,50–52].</p>
<p>Beliefs about capability and readiness to learn</p>	<p>Patients are more concerned by genetic determinants, rather than modifiable risk factors for breast cancer [54].</p> <p>Previous research has found some scepticism and defensiveness in relation to health promotion messages in relation to alcohol [55,56].</p> <p>In some individuals, health literacy levels may be an obstacle to processing and making decisions based on the information given [57]. Many lack skill or confidence in taking practical steps to reduce alcohol consumption [57,58].</p>
<p>Health beliefs: cancer pre-determinism and fatalism</p>	<p>A proportion of the population believes that incidence of cancer is purely down to ‘fate’, or known genetic causes. ‘Cancer fatalism’ is thought to have a negative impact on health behaviours, including screening uptake. Evidence suggests it is more prevalent amongst women from Black and minority</p>

	ethnic backgrounds, and that beliefs that cancer is pre-determined are strongest amongst women (a) born outside the UK, (b) whose main language is not English, or (c) exhibiting lower levels of health literacy [59]. Fatalistic beliefs are correlated with lifestyle [40] and mediate the relationship between health literacy and information seeking [41].
Exposure to fear appeal messages	Alcohol and cancer are health themes in which public health campaigns have traditionally appealed to fear processes, seeking impact by evoking a strong emotional response. Alcohol harm reduction video advertisements, in particular, tend to have a negative emotional tone (74%) and focus on short-term risks (53%), with only 18% focusing on how to adapt lifestyle to improve long-term health [60]. This contributes to a subtext which may trigger fear by association, even when unintended.
Perceived relevance of alcohol prevention	General ABIs are frequently poorly tailored to individuals. We found many leaflets contain messages and recommendations aimed at higher-risk drinkers and therefore not relevant to many recipients' level of alcohol consumption or lifestyles. These messages may therefore be easily dismissed by the majority of readers as irrelevant [38].

Step 3: Mechanisms of Action

Beyond the sole need to increase knowledge of the long-term health effects of alcohol (commonly invoked as a necessary mechanism of action to promote behaviour change [28]), we note the role of attitudes towards the behaviour as well as perceived susceptibility/vulnerability. We explored the role of emotions and perceived susceptibility/vulnerability in mediating or moderating alcohol behaviour change from existing reviews of behaviour change mechanisms and techniques.

The 'teachable moment' model [61] posits that some health events can facilitate behaviour change by affecting the subject's perception of personal risks, due to affective responses (such as a worry), and challenge their health-related beliefs to the point of promoting behaviour change. However, this effect could be moderated by other processes in situations perceived as threats to life (for example, a potential cancer diagnosis). Under the assumption that a symptomatic breast referral raises the level of fear or perceived vulnerability, the Extended Parallel Process Model by Witte et al. [62] anticipates one of two main responses: participants either accept related health

messages (Danger Control processes) or reject them (Fear Control processes).

Danger Control processes predict an enhanced 'readiness to learn', which we define as the propensity to absorb information on health risks, reflect on its meaning, and use it in relation to everyday lifestyle choices. An ABI could capitalise on Danger Control processes by establishing an association between alcohol and the risk of breast cancer, and redirecting the individual's attention towards achievable methods of reducing alcohol consumption.

Conversely, an ABI could fail by triggering 'Fear Control' processes, by exacerbating fatalistic thoughts in women attending clinic who believe that cancer risk is largely predetermined and beyond their control. Such beliefs are known to be more prevalent in populations with limited health literacy [59]. If Fear Control processes dominate, recipients of the ABI may be inclined to discard lifestyle advice in an effort to manage or control their fear of cancer.

Data from our focus groups indicated that while Fear Control processes occur amongst women attending breast clinics (e.g. 'information overload', avoidance of health literature), the desire to learn about modifiable risk factors is also present [61,63,64]. Studies by Anderson et al. [65] have shown that the anxiety generated by a breast mammogram, far from constituting an obstacle to health promotion, can be used for opportunistic large-scale lifestyle interventions. Adapting the content of the intervention so as to minimise Fear Control processes is thus the main avenue to activating the potential efficacy of a teachable moment.

In addition to the findings from our reviews, qualitative evidence we collected [22,64] suggested that an intervention would need to enhance the perception that, out of all cancer risk factors, alcohol is one of the most easily modified, and it is necessary to emphasise the health and wellbeing **gains** of adopting and/or maintaining a lower level of alcohol consumption. Framing low-risk alcohol consumption levels in terms of 'health gains' [66], using positive language, may be particularly important in the areas of cancer and alcohol use, where health promotion has been dominated by fear appeal techniques (e.g. campaigns on missing the early signs of cancer or against drink-driving). As individuals targeted by the proposed intervention will be influenced by their previous exposure to primarily fear-based messages, we specifically monitored the meaning early testers gave to health promotion messages embedded in the prototype intervention.

From this step 3, we concluded that the intervention is most likely to succeed if it provides reassurance that alcohol is a controllable determinant of cancer, and promotes positive benefits of limiting alcohol use to long-term health and wellbeing.

Step 4: How to Deliver the Change Mechanism

Our previous work identified that the most feasible and scalable mode of delivering a lifestyle intervention in clinics was a web application accessed by women in the clinic waiting area on a tablet computer [22]. In addition to circumventing the health care professional's lack of time and confidence in delivering lifestyle brief interventions, preliminary user testing confirmed that electronic delivery was acceptable and brought advantages in terms of privacy.

Within the constraints set by a web application, and with the help of the third review, we identified candidate BCTs to deliver the following mechanisms of action (see Table 3):

- improving knowledge of the health benefits of low-risk drinking
- increasing skills in relation to estimating the alcohol content of beverages
- changing attitudes to, and beliefs about consequences of, alcohol consumption
- capitalising on perceived susceptibility/vulnerability heightened by the symptomatic breast clinic attendance to increase motivation while emphasising personal control and belief in capability to reduce cancer risk.

The four BCTs employed with the highest degree of fidelity across the prototype were: provision of information on health consequences of alcohol; feedback on behaviour; discrepancy between current behaviour and goals; and social comparison. Other techniques, for example self-monitoring or instructions on how to perform the behaviour, informed the design of prompts or suggestions deeper in the application interface available to those who were interested in exploring them rather than being delivered procedurally by the interface to all users.

Table 3. Behaviour change techniques (BCTs) and features identified for prototyping

BCTs (taxonomy number)	Prototype features
Information about health consequences (5.1)	Information on alcohol's dose-response association with breast cancer and the absence of safe threshold.
	Information of the proportion of breast cancer cases attributable to alcohol in the UK.
	Information about benefits of low-risk drinking beyond lower risks of breast cancer (other types of

	cancers, mental health, dementia, liver, etc.).
	'Myth busting' quiz on risk factors for breast cancer.
Feedback on behaviour (2.2); Discrepancy between current behaviour and goal (1.6)	<p>Assessment of current alcohol consumption in units per week.</p> <p>Personalised feedback based on the UK Chief Medical Officers' low-risk drinking guidance [39].</p> <p>Automated suggestion of one of four goals in line with the same guidance [39], as a function of the current pattern of alcohol consumption measured by the app:</p> <ul style="list-style-type: none"> • maintain current low-risk drinking • have no more than 5 units of alcohol in any one day (low frequency and high intensity drinkers) • reduce alcohol consumption by a specified number of units per week (difference to 14 units within a maximum of 10 units) with equivalent amount presented in number of wine glasses • reduce alcohol consumption (amount unspecified for users in the region of 14 units per week or ≥ 24 units per week).
Social comparison (6.2)	Personalised feedback of current alcohol consumption compared to (a) other women in England and (b) other women in the clinic.
Framing/reframing (13.2)	<p>Frame alcohol as an easily controllable risk factor for breast cancer.</p> <p>Focus messages on risk reduction by changing behaviour rather than risk being increased by current behaviour (gain framing).</p> <p>Frame alcohol as any another health-related lifestyle by embedding alcohol information within information content covering other key lifestyle determinants: physical activity, diet, weight.</p> <p>Offer ways to reduce alcohol consumption, promote them as simple and easy steps.</p> <p>Emphasise choice, presenting change as an easy</p>

	option, with advice on how to cut down.
Self-monitoring of behaviour (2.3)	'Top tips': recommend keeping a diary of alcohol intake with a smartphone app (hyperlink to NHS drinks tracker app) or a paper diary (hyperlink to a diary template).
Credible source (9.1)	'Myth busting' quiz challenging common misunderstandings on risk factors believed to promote breast cancer. Breast Cancer Now charity logo and endorsement. NHS branding of the app (requested by women, to be implemented subject to relevant authorisations). Delivery of the intervention within the clinic waiting room, endorsement by health care staff.
Instruction on how to perform a behaviour (4.1); Behaviour substitution; Problem solving (1.2)	'Top tips': examples of techniques to reduce alcohol consumption on social occasions, by setting goals, self-monitoring, involving relatives. 'Top tips': advice on choosing beverages with lower alcohol content and/or smaller volume; alternating drinks with glasses of water. Drink calculator: information on beverage sizes and alcohol content in UK units. Hyperlinks to further resources: drinking diary template, Public Health England drink tracker application, 'Soberistas', 'Club Soda'.

Information about Health Consequences (BCT 5.1)

Information related to consequences for the risk of breast cancer was designed to convey the dose-dependent nature of the association between alcohol and breast carcinogenesis, emphasising that no 'safe' threshold exists for alcohol consumption in relation to breast cancer risk. The material designed by the team is adapted from an existing information leaflet [67] developed by a partner charity (Breast Cancer Now) on the basis of extensive qualitative research.

Feedback on Behaviour (BCT 2.2); Discrepancy Between Current Behaviour and Goal (BCT 1.6); Social Comparison (BCT 6.2)

As women are often unsure about their alcohol risk levels (Table 2), study participants indicated that personalised feedback needed to be the first step of the intervention. Therefore, we assessed a range of questionnaires to assess current alcohol consumption or risk level. Testing of existing alcohol smartphone and web applications in the focus group confirmed women wished to position themselves on a risk gradient to identify the scale of change they needed to undertake. We found that stratification tools that included items measuring social risks of alcohol were off-putting (e.g. the items on injuries or feelings of guilt in the Alcohol Use Disorders Identification Test (AUDIT) [68]). Such items triggered perceptions associated with substance 'abuse' which diverted attention from dose-dependent processes putting them at risk of chronic medical conditions. We therefore chose a short consumption-focused three-item questionnaire, the Extended AUDIT-C, and are currently validating an algorithm that estimates average weekly alcohol consumption based on these three items.

Framing/Reframing of Alcohol (BCT 13.2)

The content of the intervention sought to reframe alcohol as one of the more controllable lifestyle risk factors for chronic illness (Table 3). We aimed to:

- offer a new perspective on low-risk drinking as a positive choice (gain framing) made to improve future health prospects;
- challenge binary stereotyping of alcohol use opposing 'safe drinkers' and 'alcoholics'/'boozers'; instead, represent the risks of drinking as a continuum. The language describing alcohol risks was kept as neutral as possible to adapt to a wide audience, and excluded references to addiction or social harms of alcohol [38].

Some BCTs were potentially unhelpful in the context of the teachable moment within our target health settings because of their potential to trigger fear control reactions. In particular, we did not wish to enhance the *salience of health consequences of alcohol drinking* (BCT 5.2) or evoke *anticipated regret* (BCT 5.5) as the situational context of the breast clinic already made potential consequences of breast cancer tangible and memorable.

Finally, we identified other features likely to mediate the efficacy of the intervention, which required consideration as part of the iterative design and testing stage. As the usability of an electronic intervention is a predictor of engagement [69], we paid attention to women's evaluation of its quality and discoverability (the extent to which women were able to find content on the app without being told it existed). We allowed the user to assess the alcohol content of their

own preferred alcoholic drinks, sought to make 'top tips' easy to navigate to facilitate participants to focus on specific information of interest to them.

Step 5: Iterative Design, Testing and Refining of a Prototype Intervention

Following a phase of testing, with cycles of refinement of the prototype with 161 women in clinics, the final prototype consisted of:

- (a) **An initial assessment of alcohol consumption, smoking, height and weight.**
- (b) **Personalised feedback on alcohol intake integrated with other risk factors:** A feedback page presents the estimated number of units per week, and drinking risk level, assisted by a graphic visualising alcohol risk levels based on the UK Chief Medical Officers' guidance [39] (Multimedia Appendix 2). Personal drinking risk level is compared to the general population of women, as well as other women attending the same clinic, and the Department of Health low-risk drinking guidelines. To reduce stigma, this feedback is integrated with more succinct personalized feedback on benefits of not smoking, success rates of quit attempts; and ranges of healthy weights corresponding to the person's height with a button linking to health promoting physical activity and diet. Study participants improved the wording of the personalised feedback where it proved confusing or off-putting (e.g. feedback aimed at low-frequency but high-intensity alcohol consumption was rephrased from 'drinking large quantities' to 'having no more than 5 units' on any single day).
- (c) **An overview page linking to other health promotion information including:**
 - (i) A myth-busting quiz testing knowledge on modifiable risk factors for breast cancer, including alcohol.
 - (ii) Information on the dose-response association between breast cancer and alcohol.
 - (iii) An interactive drink calculator providing alcohol units and calories of standard drinks as well as larger volumes (e.g. bottles). This was refined to help participants add up, over any time period, how many units of alcohol they may be consuming, how many kilocalories these drinks contain, as well as food equivalents (in hamburgers and biscuits) and

metabolic equivalents in minutes of a tasks such as running, swimming or housework.

- (iv) Example goals for maintaining low-risk drinking or reducing alcohol consumption.
- (v) Specific information pages on: weight management; physical activity; diet; and smoking. A section on breast symptoms initially designed and tested was removed to refocus content on lifestyle promotion.

Discussion

Principal Results

This study applied a rigorous intervention development framework, drawing on a suite of reviews of the risk factor literature, attitudes towards modifiable risk factors for cancer, and digital health interventions. We involved women attending breast clinics in the design, prototyping and testing of a context-specific digital ABI in breast health settings with a potential to reach over 540,000 women per year in England alone, at very low costs, and where little information is currently provided in relation to modifiable risk factors for breast cancer. Coined as ‘teachable moments’ in the cancer prevention literature [15], breast appointments constitute a privileged opportunity to raise awareness of potentially preventable causes of breast cancer. This assumes the provision of relevant, acceptable and effective health promotion messages delivered with the highest level of fidelity.

The mechanisms of actions identified in this paper, and our reviews of their evidence base, suggest potential to achieve small reductions in alcohol consumption. Several moderators of the mechanisms of change for this intervention have been identified: acceptability to women, particularly those whose anxiety makes them potentially averse to health-related information; usability of the web application delivering the intervention; and engagement with sub-components of the digital interface. The next phase of research will evaluate the feasibility, acceptability and usability of the intervention in clinics with the target population and produce the necessary evidence on how to optimise the effect of such moderators.

Comparison with Prior Work

The design of the proposed intervention differs to that of other digital ABIs, which focus either on student populations, or longer-term engagement with smartphone or web apps [13,70]. In a clinical setting characterised by a high throughput, and a narrow window for engagement, our development has focused on designing content that

engages with the user as quickly as possible, and is relevant to the widest range of women attending. This is a marked difference from other UK-based precedents such as Down Your Drink [71], which enrolled participants from primary care into a 6-week programme through an online account. Our intervention is designed to promote the take-up of other resources for longer-term engagement, where required. Effective engagement with such resources (for instance a smartphone drink tracker) is likely to constitute a key mediator of the intervention's effect.

Limitations

This prototype intervention was developed in a single site in Southampton, UK. Feasibility and acceptability remain to be demonstrated in other sites, with different population demographics. The proposed intervention is also designed around the characteristics of the UK cancer detection model and may require adaptation to other health systems.

Conclusions

Breast cancer is the most common type of cancer in women, and alcohol one of the most feasible risk factors to moderate for the prevention of breast cancer [3]. Symptomatic breast clinics constitute a context in which targeted health improvement interventions could take place. Unlike other ABIs, the proposed intervention aims to be acceptable and feasible to deliver to all women who attend symptomatic breast clinics, irrespective of their level of alcohol consumption. In spite of extensive research on ABIs, current evidence is predominantly restricted to increased-risk drinkers. It also provides little data on the maintenance of effects of digitally-delivered ABIs beyond 12 months [13]. The effectiveness of the proposed intervention thus requires further research.

Acknowledgements

This work was funded by the Medical Research Council [grant number MR/P016960/1] and an internal grant from the supported by the NIHR Southampton Biomedical Research Centre. Professor Kaner is a member of Fuse, the Centre for Translational Research in Public Health (www.fuse.ac.uk), the NIHR School of Public Health Research and NIHR School of Primary Care Research. MMcC holds a Medical Research Council/University fellowship supported by an MRC partnership grant (MC/PC/13027) and is supported by the Medical Research Council and Chief Scientist Office through the Complexity (MC_UU_12017/14/SPHSU14) in Health Improvement programme. We gratefully acknowledge the advice and comments throughout the

project from our project partners Eluned Hughes of Breast Cancer Now, and Lucy Rocca of Soberistas.

Conflicts of Interest

EC declares honoraria, advisory board meetings and support to attend educational meetings from: Roche, Astra-Zeneca, Lilly, Nanostring and Pfizer.

Abbreviations

6SQulD: six steps in quality intervention development

ABI: alcohol brief intervention

AUDIT-C: alcohol use disorders identification tool, consumption items (short form)

BCT: behaviour change technique

NHS: National Health Service

References

1. Fitzmaurice C, Allen C, Barber RM, Barregard L, Bhutta ZA, Brenner H, Dicker DJ, Chimed-Orchir O, Dandona R, Dandona L, Fleming T, Forouzanfar MH, Hancock J, Hay RJ, Hunter-Merrill R, Huynh C, Hosgood HD, Johnson CO, Jonas JB, Khubchandani J, Kumar GA, Kutz M, Lan Q, Larson HJ, Liang X, Lim SS, Lopez AD, MacIntyre MF, Marczak L, Marquez N, Mokdad AH, Pinho C, Pourmalek F, Salomon JA, Sanabria JR, Sandar L, Sartorius B, Schwartz SM, Shackelford KA, Shibuya K, Stanaway J, Steiner C, Sun J, Takahashi K, Vollset SE, Vos T, Wagner JA, Wang H, Westerman R, Zeeb H, Zoeckler L, Abd-Allah F, Ahmed MB, Alabed S, Alam NK, Aldhahri SF, Alem G, Alemayohu MA, Ali R, Al-Raddadi R, Amare A, Amoako Y, Artaman A, Asayesh H, Atnafu N, Awasthi A, Saleem HB, Barac A, Bedi N, Bensenor I, Berhane A, Bernabé E, Betsu B, Binagwaho A, Boneya D, Campos-Nonato I, Castañeda-Orjuela C, Catalá-López F, Chiang P, Chibueze C, Chittheer A, Choi J-Y, Cowie B, Damtew S, das Neves J, Dey S, Dharmaratne S, Dhillon P, Ding E, Driscoll T, Ekwueme D, Endries AY, Farvid M, Farzadfar F, Fernandes J, Fischer F, G/hiwot TT, Gebru A, Gopalani S, Hailu A, Horino M, Horita N, Hussein A, Huybrechts I, Inoue M, Islami F, Jakovljevic M, James S, Javanbakht M, Jee SH, Kasaeian A, Kedir MS, Khader YS, Khang Y-H, Kim D, Leigh J, Linn S, Lunevicius R, El Razek HMA, Malekzadeh R, Malta DC, Marcenes W, Markos D, Melaku YA, Meles KG, Mendoza W, Mengiste DT, Meretoja TJ, Miller TR, Mohammad KA, Mohammadi A, Mohammed S, Moradi-Lakeh M, Nagel G, Nand D, Le Nguyen Q, Nolte S, Ogbo FA, Oladimeji KE, Oren E, Pa M, Park E-K, Pereira DM, Plass D, Qorbani M, Radfar A, Rafay A, Rahman M, Rana SM, Søreide K, Satpathy M, Sawhney M,

- Sepanlou SG, Shaikh MA, She J, Shiue I, Shore HR, Shrimme MG, So S, Soneji S, Stathopoulou V, Stroumpoulis K, Sufiyan MB, Sykes BL, Tabarés-Seisdedos R, Tadese F, Tedla BA, Tessema GA, Thakur JS, Tran BX, Ukwaja KN, Uzochukwu BSC, Vlassov VV, Weiderpass E, Wubshet Terefe M, Yebyo HG, Yimam HH, Yonemoto N, Younis MZ, Yu C, Zaidi Z, Zaki MES, Zenebe ZM, Murray CJL, Naghavi M. Global, Regional, and National Cancer Incidence, Mortality, Years of Life Lost, Years Lived With Disability, and Disability-Adjusted Life-years for 32 Cancer Groups, 1990 to 2015. *JAMA Oncol* [Internet] 2017 Apr 1;3(4):524. PMID:27918777
2. World Health Organisation. Cancer prevention [Internet]. 2014 [cited 2018 Jun 22]. Available from: <http://www.who.int/cancer/prevention/en/>
 3. Brown KF, Rungay H, Dunlop C, Ryan M, Quartly F, Cox A, Deas A, Elliss-Brookes L, Gavin A, Hounscome L, Huws D, Ormiston-Smith N, Shelton J, White C, Parkin DM. The fraction of cancer attributable to modifiable risk factors in England, Wales, Scotland, Northern Ireland, and the United Kingdom in 2015. *Br J Cancer* 2018;118(8):1130–1141. PMID:29567982
 4. Jones L, Bellis MA. Updating England-Specific Alcohol-Attributable Fractions [Internet]. 2013. Available from: <http://www.cph.org.uk/wp-content/uploads/2014/03/24892-ALCOHOL-FRACTIONS-REPORT-A4-singles-24.3.14.pdf>
 5. World Cancer Research Fund/American Institute for Cancer Research. Continuous Update Project Expert Report 2018. Alcoholic drinks and the risk of cancer [Internet]. 2018. Available from: www.dietandcancerreport.org
 6. Kwan ML, Kushi LH, Weltzien E, Tam EK, Castillo A, Sweeney C, Caan BJ. Alcohol Consumption and Breast Cancer Recurrence and Survival Among Women With Early-Stage Breast Cancer: The Life After Cancer Epidemiology Study. *J Clin Oncol* [Internet] 2010;28(29):4410–4416. PMID:20805458
 7. Simapivapan P, Boltong A, Hodge A. To what extent is alcohol consumption associated with breast cancer recurrence and second primary breast cancer?: A systematic review. *Cancer Treat Rev* [Internet] Elsevier Ltd; 2016;50:155–167. PMID:27693930
 8. National Institute of Health and Clinical Excellence. Early and locally advanced breast cancer: diagnosis and management. NICE guideline NG101. July 2018 [Internet]. 2018. Available from: <https://www.nice.org.uk/guidance/ng101>
 9. Angus C, Li J, Romero-Rodriguez E, Anderson P, Parrott S, Brennan A. Cost-effectiveness of strategies to improve delivery of brief interventions for heavy drinking in primary care: results from the ODHIN trial. *Eur J Public Health* [Internet] 2019 Apr 1;29(2):219–225. PMID:30239676
 10. Anderson P, Chisholm D, Fuhr DC. Effectiveness and cost-

effectiveness of policies and programmes to reduce the harm caused by alcohol. *Lancet* 2009;373(9682):2234–2246. PMID:19560605

11. Martineau F, Tyner E, Lorenc T, Petticrew M, Lock K. Population-level interventions to reduce alcohol-related harm: An overview of systematic reviews. *Prev Med (Baltim)* [Internet] 2013 Oct;57(4):278–296. PMID:23811528
12. Kaner EF, Beyer FR, Muirhead C, Campbell F, Pienaar ED, Bertholet N, Daepfen JB, Saunders JB, Burnand B. Effectiveness of brief alcohol interventions in primary care populations. *Cochrane Database Syst Rev* [Internet] 2018 Feb 24;(12). PMID:29476653
13. Kaner EF, Beyer FR, Garnett C, Crane D, Brown J, Muirhead C, Redmore J, O'Donnell A, Newham JJ, de Vocht F, Hickman M, Brown H, Maniatopoulos G, Michie S. Personalised digital interventions for reducing hazardous and harmful alcohol consumption in community-dwelling populations. *Cochrane Database Syst Rev* [Internet] 2017 Sep 25; PMID:28944453
14. Brown J, West R, Angus C, Beard E, Brennan A, Drummond C, Hickman M, Holmes J, Kaner E, Michie S. Comparison of brief interventions in primary care on smoking and excessive alcohol consumption: a population survey in England. *Br J Gen Pract* [Internet] 2016 Jan 30;66(642):e1–e9. PMID:26719481
15. Demark-Wahnefried W, Aziz NM, Rowland JH, Pinto BM. Riding the Crest of the Teachable Moment: Promoting Long-Term Health After the Diagnosis of Cancer. *J Clin Oncol* [Internet] 2005 Aug 20;23(24):5814–5830. PMID:16043830
16. Senore C, Giordano L, Bellisario C, Di Stefano F, Segnan N. Population Based Cancer Screening Programmes as a Teachable Moment for Primary Prevention Interventions. A Review of the Literature. *Front Oncol* [Internet] 2012;2. PMID:22649789
17. Anderson AS, Mackison D, Boath C, Steele R. Promoting Changes in Diet and Physical Activity in Breast and Colorectal Cancer Screening Settings: An Unexplored Opportunity for Endorsing Healthy Behaviors. *Cancer Prev Res* [Internet] 2013 Mar 1;6(3):165–172. PMID:23324132
18. NHS England. Waiting Times for Suspected and Diagnosed Cancer Patients. 2017-18 Annual Report [Internet]. 2018. Available from: <https://www.england.nhs.uk/statistics/statistical-work-areas/cancer-waiting-times/>
19. Public Health England. NHS England Cancer Waiting Times Database: Two-week referrals resulting in a diagnosis of cancer (Conversion rate: as % of all TWW referrals) [Internet]. National Cancer Registration and Analysis Service; 2019 [cited 2019 Aug 5]. Available from: https://fingertipsws.phe.org.uk/api/all_data/csv/by_indicator_id?parent_area_code=E92000001&parent_area_type_id=152&child_

area_type_id=7&profile_id=92&indicator_ids=91845&category_area_code=&sex_id=4&age_id=1

20. Heather N. The efficacy-effectiveness distinction in trials of alcohol brief intervention. *Addict Sci Clin Pract* [Internet] 2014;9(1):13. PMID:25127717
21. Wight D, Wimbush E, Jepson R, Doi L. Six steps in quality intervention development (6SQuID). *J Epidemiol Community Health* 2016 May;70(5):520–525. PMID:26573236
22. Sinclair J, McCann M, Sheldon E, Gordon I, Brierley-Jones L, Copson E. The acceptability of addressing alcohol consumption as a modifiable risk factor for breast cancer: a mixed method study within breast screening services and symptomatic breast clinics. *BMJ Open* [Internet] 2019 May 17;9(6):e027371. [doi: 10.1136/bmjopen-2018-027371]
23. Crane D, Garnett C, Brown J, West R, Michie S. Behavior Change Techniques in Popular Alcohol Reduction Apps. *J Med Internet Res* [Internet] 2015 May 14;17(5):e118. PMID:25977135
24. Michie S, Whittington C, Hamoudi Z, Zarnani F, Tober G, West R. Identification of behaviour change techniques to reduce excessive alcohol consumption. *Addiction* [Internet] 2012 Aug;107(8):1431–1440. PMID:22340523
25. Black N, Mullan B, Sharpe L. Computer-Delivered Interventions for Reducing Alcohol Consumption: Meta-Analysis and Meta-Regression using Behaviour Change Techniques and Theory. *Health Psychol Rev* [Internet] Taylor & Francis; 2016;7199(April):1–33. PMID:26999311
26. Beyer F, Lynch E, Kaner E. Brief Interventions in Primary Care: an Evidence Overview of Practitioner and Digital Intervention Programmes. *Curr Addict Reports* [Internet] *Current Addiction Reports*; 2018 Jun 3;5(2):265–273. PMID:29963364
27. Johnson M, Jackson R, Guillaume L, Meier P, Goyder E. Barriers and facilitators to implementing screening and brief intervention for alcohol misuse: a systematic review of qualitative evidence. *J Public Health (Oxf)* 2011 Sep 1;33(3):412–421. PMID:21169370
28. Carey RN, Connell LE, Johnston M, Rothman AJ, de Bruin M, Kelly MP, Michie S. Behavior Change Techniques and Their Mechanisms of Action: A Synthesis of Links Described in Published Intervention Literature. *Ann Behav Med* [Internet] 2018 Oct 10; PMID:30304386
29. Cane J, O'Connor D, Michie S. Validation of the theoretical domains framework for use in behaviour change and implementation research. *Implement Sci* [Internet] 2012 Dec 24;7(1):37. PMID:22530986
30. Michie S, Richardson M, Johnston M, Abraham C, Francis J, Hardeman W, Eccles MP, Cane J, Wood CE. The Behavior Change Technique Taxonomy (v1) of 93 Hierarchically Clustered

- Techniques: Building an International Consensus for the Reporting of Behavior Change Interventions. *Ann Behav Med* [Internet] 2013 Aug 20;46(1):81–95. PMID:23512568
31. Hekler EB, Klasnja P, Riley WT, Buman MP, Huberty J, Rivera DE, Martin CA. Agile science: creating useful products for behavior change in the real world. *Transl Behav Med* [Internet] 2016 Jun 26;6(2):317–328. PMID:27357001
 32. Jaspers MWM, Steen T, Bos C van den, Geenen M. The think aloud method: a guide to user interface design. *Int J Med Inform* 2004;73(11):781–795. PMID:15491929
 33. Health Education England. Health Literacy ‘How To’ Guide [Internet]. 2017. Available from: <https://www.hee.nhs.uk/our-work/health-literacy>
 34. Trello [Internet]. 2018 [cited 2019 Mar 24]. Available from: <https://trello.com/>
 35. Buykx P, Li J, Gavens L, Lovatt M, Gomes De Matos E, Holmes J, Hooper L, Meier P. An investigation of public knowledge of the link between alcohol and cancer [Internet]. 2015. Available from: http://www.cancerresearchuk.org/sites/default/files/an_investigation_of_public_knowledge_of_the_link_between_alcohol_and_cancer_buykx_et_al.pdf
 36. Shahab L, McGowan JA, Waller J, Smith SG. Prevalence of beliefs about actual and mythical causes of cancer and their association with socio-demographic and health-related characteristics: Findings from a cross-sectional survey in England. *Eur J Cancer* [Internet] 2018 Nov;103:308–316. PMID:29705530
 37. Bates S, Holmes J, Gavens L, de Matos EG, Li J, Ward B, Hooper L, Dixon S, Buykx P. Awareness of alcohol as a risk factor for cancer is associated with public support for alcohol policies. *BMC Public Health* [Internet] 2018 Dec 4;18(1):688. [doi: 10.1186/s12889-018-5581-8]
 38. Khadjesari Z, Stevenson F, Toner P, Linke S, Milward J, Murray E. ‘I’m not a real boozer’: a qualitative study of primary care patients’ views on drinking and its consequences. *J Public Health (Oxf)* [Internet] 2018 Apr 18;(April):1–7. PMID:29912419
 39. UK Department of Health. UK Chief Medical Officers’ Low Risk Drinking Guidelines. August 2016 [Internet]. 2016. Available from: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/545937/UK_CMOs__report.pdf
 40. Niederdeppe J, Levy AG. Fatalistic Beliefs about Cancer Prevention and Three Prevention Behaviors. *Cancer Epidemiol Biomarkers Prev* [Internet] 2007 May 1;16(5):998–1003. [doi: 10.1158/1055-9965.EPI-06-0608]
 41. Kobayashi LC, Smith SG. Cancer Fatalism, Literacy, and Cancer Information Seeking in the American Public. *Heal Educ Behav* [Internet] 2016 Aug 16;43(4):461–470. [doi:

- 10.1177/1090198115604616]
42. Boniface S, Kneale J, Shelton N. Drinking pattern is more strongly associated with under-reporting of alcohol consumption than socio-demographic factors: evidence from a mixed-methods study. *BMC Public Health* [Internet] 2014;14(1):1297. PMID:25519144
 43. Boniface S, Kneale J, Shelton N. Actual and Perceived Units of Alcohol in a Self-Defined "Usual Glass" of Alcoholic Drinks in England. *Alcohol Clin Exp Res* [Internet] 2013 Jun;37(6):978-983. PMID:23278164
 44. Nutt DJ, Rehm J. Doing it by numbers: A simple approach to reducing the harms of alcohol. *J Psychopharmacol* [Internet] 2014 Jan 7;28(1):3-7. PMID:24399337
 45. Rehm J, Anderson P, Manthey J, Shield KD, Struzzo P, Wojnar M, Gual A. Alcohol Use Disorders in Primary Health Care: What Do We Know and Where Do We Go? *Alcohol Alcohol* [Internet] 2016 Jul;51(4):422-427. PMID:26574600
 46. Rees G, Young M-A, Gaff C, Martin PR. A Qualitative Study of Health Professionals' Views Regarding Provision of Information About Health-Protective Behaviors During Genetic Consultation for Breast Cancer. *J Genet Couns* [Internet] 2006 Apr 16;15(2):95-104. PMID:16541332
 47. Daley AJ, Bowden SJ, Rea DW, Billingham L, Carmicheal AR. What advice are oncologists and surgeons in the United Kingdom giving to breast cancer patients about physical activity? *Int J Behav Nutr Phys Act* [Internet] 2008;5(1):46. PMID:18803812
 48. Anderson AS, Caswell S, Wells M, Steele RJC. Obesity and lifestyle advice in colorectal cancer survivors - how well are clinicians prepared? *Color Dis* [Internet] 2013 Aug;15(8):949-957. PMID:23480570
 49. Spellman C, Craike M, Livingston P. Knowledge, attitudes and practices of clinicians in promoting physical activity to prostate cancer survivors. *Health Educ J* [Internet] 2014 Sep 13;73(5):566-575. [doi: 10.1177/0017896913508395]
 50. Miles A, Simon A, Wardle J. Answering Patient Questions about the Role Lifestyle Factors Play in Cancer Onset and Recurrence. *J Health Psychol* [Internet] 2010 Mar 5;15(2):291-298. [doi: 10.1177/1359105309351245]
 51. Williams K, Beeken RJ, Fisher A, Wardle J. Health professionals' provision of lifestyle advice in the oncology context in the United Kingdom. *Eur J Cancer Care (Engl)* [Internet] 2015 Jul;24(4):522-530. [doi: 10.1111/ecc.12305]
 52. James-Martin G, Koczwara B, Smith EL, Miller MD. Information needs of cancer patients and survivors regarding diet, exercise and weight management: a qualitative study. *Eur J Cancer Care (Engl)* [Internet] 2014 May;23(3):340-348. [doi:

- 10.1111/ecc.12159]
53. Bakhshi S, While A. Health Professionals' Alcohol-Related Professional Practices and the Relationship between Their Personal Alcohol Attitudes and Behavior and Professional Practices: A Systematic Review. *Int J Environ Res Public Health* [Internet] 2013 Dec 23;11(1):218–248. [doi: 10.3390/ijerph110100218]
 54. Dumalaon-Canaria JA, Hutchinson AD, Prichard I, Wilson C. What causes breast cancer? A systematic review of causal attributions among breast cancer survivors and how these compare to expert-endorsed risk factors. *Cancer Causes Control* [Internet] 2014 Jul 27;25(7):771–785. [doi: 10.1007/s10552-014-0377-3]
 55. Conway E, Wyke S, Sugden J, Mutrie N, Anderson AS. Can a lifestyle intervention be offered through NHS breast cancer screening? Challenges and opportunities identified in a qualitative study of women attending screening. *BMC Public Health* [Internet] 2016 Dec 11;16(1):758. [doi: 10.1186/s12889-016-3445-7]
 56. Scheideler JK, Klein WMP. Awareness of the Link between Alcohol Consumption and Cancer across the World: A Review. *Cancer Epidemiol Biomarkers Prev* [Internet] 2018 Apr 2;27(4):429–437. [doi: 10.1158/1055-9965.EPI-17-0645]
 57. Berkman ND, Sheridan SL, Donahue KE, Halpern DJ, Crotty K. Low Health Literacy and Health Outcomes: An Updated Systematic Review. *Ann Intern Med* [Internet] 2011 Jul 19;155(2):97. [doi: 10.7326/0003-4819-155-2-201107190-00005]
 58. Sinclair JMA, E. Chambers S, C. Manson C. Internet Support for Dealing with Problematic Alcohol Use: A Survey of the Soberistas Online Community. *Alcohol Alcohol* [Internet] 2016 Oct 25; [doi: 10.1093/alcalc/agw078]
 59. Vrinten C, Wardle J, Marlow LA. Cancer fear and fatalism among ethnic minority women in the United Kingdom. *Br J Cancer* [Internet] 2016 Mar 11;114(5):597–604. [doi: 10.1038/bjc.2016.15]
 60. Dunstone K, Brennan E, Slater MD, Dixon HG, Durkin SJ, Pettigrew S, Wakefield MA. Alcohol harm reduction advertisements: a content analysis of topic, objective, emotional tone, execution and target audience. *BMC Public Health* [Internet] *BMC Public Health*; 2017;17(1):312. PMID:28399829
 61. McBride CM, Puleo E, Pollak KI, Clipp EC, Woolford S, Emmons KM. Understanding the role of cancer worry in creating a “teachable moment” for multiple risk factor reduction. *Soc Sci Med* 2008;66(3):790–800. PMID:18037204
 62. Witte K, Cameron KA, McKeon JK, Berkowitz JM. Predicting risk behaviors: Development and validation of a diagnostic scale. *J Health Commun* 1996;1(4):317–341. PMID:7611301
 63. Johnson SB, Bradshaw CP, Wright JL, Haynie DL, Simons-Morton

- BG, Cheng TL. Characterizing the Teachable Moment: Is an Emergency Department Visit a Teachable Moment for Intervention Among Assault-Injured Youth and Their Parents? *Pediatr Emerg Care* [Internet] 2007 Aug [cited 2014 Nov 26];23(8):553-559. [doi: 10.1097/PEC.0b013e31812c6687]
64. Chambers SE, Copson ER, Dutey-Magni PF, Priest C, Anderson AS, Sinclair JMA. Alcohol use and breast cancer risk: A qualitative study of women's perspectives to inform the development of a preventative intervention in breast clinics. *Eur J Cancer Care (Engl)* [Internet] 2019 Apr 30;(e13075):e13075. [doi: 10.1111/ecc.13075]
 65. Anderson AS, Macleod M, Mutrie N, Sugden J, Dobson H, Treweek S, O'Carroll RE, Thompson A, Kirk A, Brennan G, Wyke S. Breast cancer risk reduction - is it feasible to initiate a randomised controlled trial of a lifestyle intervention programme (ActWell) within a national breast screening programme? *Int J Behav Nutr Phys Act* [Internet] 2014 Dec 17;11(1):156. [doi: 10.1186/s12966-014-0156-2]
 66. Gallagher KM, Updegraff JA. Health Message Framing Effects on Attitudes, Intentions, and Behavior: A Meta-analytic Review. *Ann Behav Med* [Internet] 2012 Feb 13;43(1):101-116. [doi: 10.1007/s12160-011-9308-7]
 67. Breast Cancer Now. Alcohol and breast cancer risk: How much does drinking alcohol affect your risk? [health information leaflet]. Reviewed and updated May 2014. Breast Cancer Now; 2014.
 68. Saunders JB, Aasland OG, Babor TF, de la Fuente JR, Grant M. Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO Collaborative Project on Early Detection of Persons with Harmful Alcohol Consumption--II. *Addiction* [Internet] 1993 Jun;88(6):791-804. PMID:8329970
 69. Merolli M, Gray K, Martin-Sanchez F. Health outcomes and related effects of using social media in chronic disease management: A literature review and analysis of affordances. *J Biomed Inform* [Internet] Elsevier Inc.; 2013 Dec;46(6):957-969. PMID:23702104
 70. Garnett C, Crane D, West R, Brown J, Michie S. The development of Drink Less : an alcohol reduction smartphone app for excessive drinkers. *Transl Behav Med* [Internet] 2019 Mar 1;9(2):296-307. [doi: 10.1093/tbm/iby043]
 71. Linke S, McCambridge J, Khadjesari Z, Wallace P, Murray E. Development of a Psychologically Enhanced Interactive Online Intervention for Hazardous Drinking. *Alcohol Alcohol* [Internet] 2008;43(6):669-674. PMID:18693217