

Analysis of putative quadruplex-forming sequences in fungal genomes: novel antifungal targets?

Emily F. Warner^{1†}, Natálie Bohálová^{2,3}, Václav Brázda², Zoë A. E. Waller⁴ and Stefan Bidula^{5,*}

Abstract

Fungal infections cause >1 million deaths annually and the emergence of antifungal resistance has prompted the exploration for novel antifungal targets. Quadruplexes are four-stranded nucleic acid secondary structures, which can regulate processes such as transcription, translation, replication and recombination. They are also found in genes linked to virulence in microbes, and ligands that bind to quadruplexes can eliminate drug-resistant pathogens. Using a computational approach, we quantified putative quadruplex-forming sequences (PQS) in 1359 genomes across the fungal kingdom and explored their presence in genes related to virulence, drug resistance and biological processes associated with pathogenicity in *Aspergillus fumigatus*. Here we present the largest analysis of PQS in fungi and identify significant heterogeneity of these sequences throughout phyla, genera and species. PQS were genetically conserved in *Aspergillus* spp. and frequently pathogenic species appeared to contain fewer PQS than their lesser/non-pathogenic counterparts. GO-term analysis identified that PQS-containing genes were involved in processes linked with virulence such as zinc ion binding, the biosynthesis of secondary metabolites and regulation of transcription in *A. fumigatus*. Although the genome frequency of PQS was lower in *A. fumigatus*, PQS could be found enriched in genes involved in virulence, and genes upregulated during germination and hypoxia. Moreover, PQS were found in genes involved in drug resistance. Quadruplexes could have important roles within fungal biology and virulence, but their roles require further elucidation.

DATA SUMMARY

The authors confirm that links to access all supporting data, genomes, code and protocols have been provided within the article or are accessible through the supplementary data files. The genomes used were obtained from public repositories (<https://doi.org/10.5281/zenodo.3783970>; <https://mycocosm.jgi.doe.gov/mycocosm/home>; <ftp://ftp.ncbi.nlm.nih.gov/genomes/>).

INTRODUCTION

Compared to viruses and bacteria, fungi are underappreciated [1]. They are key contributors to the food and drink, biotechnology and textile industries, whilst also being an important source of novel antimicrobial compounds [2–5].

However, >1.5 million deaths per year are attributed to fungi in humans globally, more than malaria and on par with tuberculosis [6, 7]. Fungi can also cause blindness, serious skin conditions, promote allergic responses and can cause secondary infections in cystic fibrosis, tuberculosis, human immunodeficiency virus (HIV) and SARS-CoV-2 patients [8, 9]. Fungal plant pathogens and oomycetes also destroy around a third of crops annually; enough to feed 600 million people [1]. Whilst fungal infections of amphibians, bats, bees, animals and trees have a huge impact on biodiversity [7]. Thus, fungi are an important asset, but they also pose a devastating global burden, and a deeper understanding of their biology is therefore essential.

Received 23 December 2020; Accepted 26 March 2021; Published 06 May 2021

Author affiliations: ¹School of Clinical Medicine, University of Cambridge, Cambridge, UK; ²Institute of Biophysics of the Czech Academy of Sciences, Brno, Czechia; ³Department of Experimental Biology, Faculty of Science, Masaryk University, Brno, Czechia; ⁴School of Pharmacy, University College London, London, UK; ⁵School of Biological Sciences, University of East Anglia, Norwich, UK.

*Correspondence: Stefan Bidula, s.bidula@uea.ac.uk

Keywords: Fungi; *in-silico*; virulence; drug resistance; *Aspergillus fumigatus*; G-quadruplexes; i-motifs.

Abbreviations: BUSCO, Benchmarking Universal Single-Copy Orthologs; G4, G-quadruplex; GO, gene ontology; HIV, human immunodeficiency virus; iM, i-motif; LTR, long terminal repeat; PQS, putative quadruplex-forming sequence; WHO, World Health Organisation.

†Present address: Ikarovec Limited, Norwich Research Park Innovation Centre, Norwich, UK.

Data statement: All supporting data, code and protocols have been provided within the article or through supplementary data files. Six supplementary tables and six supplementary figures are available with the online version of this article.

000570 © 2021 The Authors



This is an open-access article distributed under the terms of the Creative Commons Attribution License. This article was made open access via a Publish and Read agreement between the Microbiology Society and the corresponding author's institution.

The negative effects of fungi are exacerbated by a lack of antifungals and the emergence of multidrug-resistant pathogens with intrinsic resistance such as *Candida auris* and *Lomentospora prolificans* [7, 10]. Current classes of antifungals include azoles (e.g. fluconazole), echinocandins (e.g. caspofungin) and polyenes (e.g. amphotericin B), but resistance to these antifungals is ever more prevalent and there are no vaccines [11]. Indeed, a recent meeting of the World Health Organization (WHO) Expert Group on Identifying Priority Fungal Pathogens highlighted azole-resistant *Aspergillus fumigatus* as one such priority pathogen [12]. Therefore, we have an urgent requirement to identify potential novel antifungal targets.

G-quadruplexes (G4s) and i-motifs (iMs) are intriguing four-stranded (quadruplex) secondary structures in nucleic acids that are enriched in regulatory regions, particularly the promoters and telomeric regions of prokaryotic and eukaryotic genomes [13, 14]. G4s can also be found in other segments of DNA and RNA, such as untranslated regions, exons and introns [15]. Here, four guanine bases associate through Hoogsteen hydrogen bonding to form the basic unit of the G4, the G-tetrad [16]. These can then stack on top of each other to form the G4 structure itself (Fig. 1a–e). These stacks of G-tetrads are connected by loops of mixed-sequence nucleotides and can form intramolecular or intermolecular associations [17, 18]. This structure is further stabilized by the presence of monovalent cations, especially potassium [19]. Moreover, the 5'- to 3'-directionality of the strands, glycosidic bonding in the G-tetrads, the cation present, and number of stacked G-tetrads contribute to the wide variation of observed G4 structures and topologies [13]. Conversely, iMs form within cytosine-rich regions of DNA and can typically be found on the complementary strand opposite a G4 [20]. Like G4s, they are four-stranded structures but are composed of two intercalated hairpins, which are stabilized by hemi protonated cytosine-cytosine⁺ (C-C⁺) base pairs (Fig. 1d–f) [21]. Studies into iMs have been limited compared to G4s as it was thought they were not physiologically relevant based on them being most stable in slightly acidic conditions. However, they have since been shown to form under physiological conditions, including neutral pH and molecular crowding, and have recently been identified within the nuclei of human cells [21–24].

Once deemed structural curiosities, G4s and iMs have been highlighted to participate within host–pathogen interactions, display important roles within gene regulation, and trigger phase separation of RNA in cells [25–27]. However, the regulation of biological functions by quadruplexes is complex and is influenced by their position within DNA/RNA, the surrounding topology, and the environmental factors within the cell. In recent years, there has been increased interest in the therapeutic potential of targeting quadruplexes following the implication of these secondary structures in disease, especially cancer, due to their prevalence in oncogene promoters [26].

There is a growing number of pathogens in which G4s have been shown to contribute to virulence phenotypes, including

Impact Statement

Fungal infection results in more than 1 million deaths annually and can cause life-threatening illness in the immune compromised; such as those suffering from cancer, tuberculosis, HIV, cystic fibrosis and SARS-CoV-2. Moreover, antifungal drug resistance has emerged as a global threat and we have a critical requirement for new drug targets. Quadruplexes are intriguing four-stranded DNA/RNA structures, which have been implicated in key biological functions and have emerged as potential drug targets to kill drug-resistant micro-organisms. A comprehensive analysis of sequences with the potential to form these structures in fungi has not been conducted to date and their roles within fungi are practically unknown. This article provides the largest analysis of predicted quadruplex-forming sequences in the fungal kingdom and the first in-depth analysis of the location and potential functions of these sequences in a pathogenic fungus. This preliminary screen also unveils quadruplex-forming sequences within virulence and drug-associated genes in the clinically relevant pathogen, *Aspergillus fumigatus*, which could represent novel antifungal targets to explore. We hope this initial study will prompt the generation of interesting ideas within the field and initiate studies which promote the exploration of quadruplexes as potential antifungal targets.

viruses (human papilloma virus; Epstein–Barr virus, HIV, Adenovirus, Zika virus, Nipah virus and SARS-CoV-2), prokaryotic bacterial pathogens (*Staphylococcus aureus*, *Streptococcus pneumoniae*, *Klebsiella pneumoniae*, *Vibrio cholerae*, *Salmonella enterica*, *Enterococcus* spp., *Borrelia* spp., *Neisseria meningitidis* and *N. gonorrhoeae*), and eukaryotic pathogens (*Trypanosoma brucei*, *Plasmodium falciparum*) [25, 28–34]. Notably, G4 DNA-binding agents have been shown to be active against methicillin-resistant *S. aureus* and vancomycin-resistant *Enterococcus* spp.; potentially providing a novel target to overcome drug resistance [35, 36]. In contrast, iMs have so far only been identified in the long terminal repeat promoter of the HIV-1 pro-viral genome where they modulate the transcription of viral genes [37]. Although, one could assume that this is only the tip of the iceberg, considering the ubiquitous nature of quadruplexes within the genomes of almost every organism.

Practically all organisms possess quadruplexes and due to the advances in the development of drugs that bind and stabilize these structures, the ability to target specific quadruplexes is starting to become a reality [38–40]. A thorough analysis of putative quadruplex-forming sequences (PQS) in fungi has not been conducted to date and could unveil potential drug targets. Here, we identified PQS in publicly available genomes across the fungal kingdom with a focus on the Ascomycota, which contains many known clinically and agriculturally




Fig. 1. G-quadruplex (G4) and i-motif (iM) structures. Representative examples of G4s and iMs. (a) Side (b) and top-down view of the human telomere DNA quadruplex in K^+ solution hybrid-1 form (PDB:2HY9). (c) The basic structure of the G-tetrad. (d) Side (e) and top-down view of an intramolecular iM DNA structure with C-C⁺ base pairing (PDB 1A83). (f) C-C⁺ base pairing found in iM structures. Images were generated using the Protein Imager software.

important fungal pathogens. Moreover, we explored and discussed the potential roles of quadruplexes in the human pathogen *A. fumigatus*, and their possible contribution to the virulence of this microorganism.

METHODS

Selection of genome sequences

The 1107 genome sequences from fungi in the Ascomycota that were analysed in this study were obtained from a publicly available repository (<https://doi.org/10.5281/zenodo.3783970>). These genomes encompassed the Saccharomycotina (332 genomes from 12 major clades), Pezizomycotina (761 genomes representing 9 of 16 classes) and Taphrinomycotina (14 genomes representing 4 of 5 classes). The completeness of these genomes has been assessed previously [41]. Analysis of the genome completeness based on 1315 full-length Benchmarking Universal Single-Copy Orthologs (BUSCO) genes revealed that 1021/1107 (~92%) genomes contained >90% of the 1315 BUSCO genes.

A further 252 published genomes from the Basidiomycota, Mucoromycota, Zoopagomycota, Chytridiomycota, Microsporidia and Cryptomycota were also obtained from the Joint Genome Institute MycoCosm portal (<https://mycocosm.jgi.doe.gov/mycocosm/home>) and GenBank via FTP access number (<ftp://ftp.ncbi.nlm.nih.gov/genomes/>). The completeness of these genomes was not assessed, and

thus the analysis of these genomes can be found in the Supplementary Material for reference. Information for all the genomes used in this study can be found in Table S1 (available in the online version of this article).

The principle of the G4Hunter algorithm and process of analysis

The G4Hunter web application was used to identify PQS within fungal genomes [42]. The algorithm used in G4Hunter considers the G-richness and G-skewness of a genome. Each position within a sequence is given a score between -4 and 4, with G's giving a positive score and C's giving a negative score. A's and T's are neutral and have a score of 0. An increasing G4Hunter score (either positive or negative) correlates with increased propensity to form quadruplexes. A near-zero average score is indicative of a sequence that is most likely to form stable duplexes. The G4Hunter score is the arithmetic mean value of the sequence.

When analysing the genomes, the scored nucleic acid sequence was computed for a sliding window of 30 nucleotides and a threshold above 1.5 for stringent analyses, or a window of 25 nucleotides and thresholds between 1–2 for complete analyses. Regions in which the value of the mean score was above a threshold (either positive or negative) were extracted. The algorithm used specifically recognizes




Fig. 2. Heterogeneity of PQS across fungal phylum. The total number and frequency of PQS within 1359 fungal genomes were analysed using G4Hunter with a threshold of 1.5 and a window size of 30. The average number of PQS (a, e), PQS/kbp (b, f), PQS/GC% (c, g) and GC content (d, h) in fungi from the Ascomycota ($n=1107$), Basidiomycota ($n=186$), Mucoromycota ($n=31$), Zoopagomycota ($n=12$), Chytridiomycota ($n=12$), Microsporidia ($n=9$) and Cryptomycota ($n=2$). (e–h) contain boxplots with Tukey whiskers. The outliers are indicated by dots and the line within the boxplot is representative of the median value.

PQS, therefore all sequences obtained from the G4Hunter application can be considered PQS (a mixture of potential G4 and iM sequences).

To identify the location of PQS within annotated genomic features, the file containing the annotations for known genomic features within the genome of the selected fungi were downloaded from the NCBI database. The presence of PQS within a pre-defined genomic feature (e.g. gene, mRNA, mobile element) or within ± 100 bp of these genomic features were analysed. Discrepancies in the frequencies of PQS between mRNA and genes are due to how the feature tables downloaded from the NCBI database present their annotations. mRNA length is slightly shorter than the gene length due to the absence of intronic sequences and alternative splicing (e.g. each alternatively spliced mRNA is considered individually by the script).

Tools used to conduct the analysis

Analysis of the genomes for PQS was conducted using the G4Hunter DNA analyser web application (<http://bioinformatics.ibp.cz:8080/#/>) [42]. The location of PQS in known genomic features were identified using a publicly available script found at <https://pypi.org/project/dna-analyser-ibp/>. The protein classes of genes containing PQS were determined via PANTHER Protein Class v.15.0 [43]. Gene ontology (GO)-term analysis of PQS-containing genes was conducted using the FungiFun web tool V2.2.8 (<https://elbe.hki-jena.de/fungifun/>) and significance was determined via Fisher's exact test [44]. The PQS-containing genes identified in G4Hunter were verified via another PQS-predictive

analysis tool called QGRS Mapper (<http://bioinformatics.ramapo.edu/QGRS/analyze.php>) [45]. Sequences were analysed using the default settings of max length (30), min G-group (2) and loop size (0–36). The highest scoring sequences with the shortest loop length were selected in each case.

ChemDraw Ultra v12.0 was used to draw structures and The Protein Imager (<https://3dproteinimaging.com/protein-imager/>) [46] was used to generate 3D images from PDBs. These data were processed, and graphs were generated using GraphPad Prism software v6.01.

Analysis of transcriptomes

Upregulated genes in germinating conidia and hyphae [47], during hypoxia [48], and during iron limitation or oxidative stress [49] were identified using publicly available transcriptome datasets. These were analysed using the FungiDB transcriptomic resources tool (<https://fungidb.org/>) [50]. Upregulated genes were identified by comparing a reference sample (control) with a comparison sample (test). In each case, the difference between the minimum expression value of each gene in the control sample and maximum expression value in the test sample was quantified. All genes upregulated >twofold were noted. The identity of genes upregulated >twofold in *A. fumigatus* biofilms [51] were obtained from the Supplementary Material. PQS frequencies in the top 20 most upregulated genes in each condition were identified using G4Hunter and QGRS Mapper using the default search settings. To determine whether the mean frequencies observed for each independent condition were




Fig. 3. PQS within the sub-divisions of the phylum Ascomycota. The total number and frequency of PQS within fungal sub-divisions of the phylum Ascomycota were quantified using G4Hunter with a threshold of 1.5 and window size of 30. The total number of PQS (a), PQS/kbp (b), PQS/GC% (c) and GC content (d) in fungi from the Taphrinomycotina ($n=14$), Saccharomycotina ($n=332$) and Pezizomycotina ($n=761$). (a–d) contain boxplots with Tukey whiskers. The outliers are indicated by dots and the line within the boxplot is representative of the median value. Significance was determined by one-way ANOVA with multiple comparisons.

significantly different from the mean frequency observed for the entire genome, outliers were removed using the ROUT method ($Q=1.0\%$) and normality was tested with Shapiro–Wilk, prior to analysis via one-sample T test. A P -value < 0.05 was considered statistically significant.

Statistical analysis

Statistical analyses were conducted via parametric one-way ANOVA with Tukey’s multiple comparisons, Student’s T test, one-sample T test or Pearson’s correlation coefficient; or non-parametric Kruskal–Wallis tests with post-hoc Dunn’s test and Bonferroni corrections. Outliers were removed when necessary using the ROUT method ($Q=1.0\%$) and normality was tested using Shapiro–Wilk. All analyses were performed using GraphPad Prism software v6.01. $P < 0.05$ were considered statistically significant.

RESULTS

There is large heterogeneity in the frequency and number of PQS throughout the fungal kingdom

A thorough analysis of putative quadruplex-forming sequences in fungi has not been conducted to date. Using G4Hunter, the number of PQS were quantified in 1359 genomes across the fungal kingdom. Due to the high variability in genome size and chromosome number between fungal species, to normalize the data, the total number of PQS in addition to the frequency of PQS/kbp and number of PQS relative to the GC content (PQS/GC%) were noted.

Across the divisions, the Chytridiomycota had the largest average number of PQS (24734 PQS) and highest PQS frequencies (0.555 PQS/kbp and 473 PQS/GC%, respectively; Fig. 2). Large numbers of PQS were also found in




Fig. 4. Higher genome GC-content is positively correlated with the frequency of PQS. The frequency of PQS relative to the GC content of fungi was plotted for (a) all fungal genomes in the Ascomycota, (b) Taphrinomycotina, (c) Saccharomycotina, (d) and the Pezizomycotina. The Pearson correlation coefficient was used to determine the association between PQS and GC content. $P < 0.05$ was considered statistically significant.

fungi from the Mucoromycota and Basidiomycota (19575 and 16474 PQS, respectively; Fig. 2a, e). The Basidiomycota and Zoopagomycota had the highest PQS frequencies relative to genome size (0.434 and 0.373 PQS/kbp, respectively; Fig. 2b, f). However, the Mucoromycota and Basidiomycota displayed high PQS frequencies relative to GC content (459 and 321 PQS/GC%, respectively; Fig. 2c, g). Fungi within the Basidiomycota had the highest average GC content (52.2%; Fig. 2d, h). The Microsporidia and Cryptomycota scored lowest for total number of PQS (300 and 372, respectively), PQS/kbp (0.091 and 0.029, respectively) and PQS/GC% (8 and 11, respectively; Fig. 2). Moreover, they also had low GC content (39.6 and 35.0%, respectively).

Fungi from the Pezizomycotina have the highest average PQS frequencies in the Ascomycota

More than 1000 of the genomes investigated were from the Ascomycota, and because this phylum contains many of the most prevalent fungal pathogens and ~92% of the ascomycete genomes contained >90% of the 1315 BUSCO genes, this phylum was investigated in greater detail. Further analysis identified that there was significant heterogeneity in the total numbers and frequency of PQS between the sub-divisions Taphrinomycotina, Saccharomycotina and

Pezizomycotina (Fig. 3). Fungi from the Saccharomycotina had the fewest total PQS on average (1146 PQS), whereas fungi from the Pezizomycotina contained significantly more (17021 PQS). Moreover, this pattern was observed for PQS frequency (0.078/kbp vs. 0.436/kbp), GC content (40.6 vs. 49.6%), and PQS frequency relative to the GC content (27/GC% vs. 344/GC%) for the Saccharomycotina and Pezizomycotina, respectively (Fig. 3). A complete breakdown of the G4Hunter results for all fungi can be found in Table S2.

Considering G4s and iMs form in guanine or cytosine-rich regions, respectively, one would expect fungi with a higher genome GC content to have a higher PQS frequency by chance. To investigate this further, the frequency of PQS/kbp relative to the GC content in all fungi were plotted. As expected, there was a positive correlation between GC content and PQS frequency amongst all the ascomycete genomes analysed ($r=0.5620$; $P < 0.0001$; Fig. 4a). Moreover, this positive correlation was observed for the Saccharomycotina and Pezizomycotina ($r=0.5068$ and 0.2883 , respectively; both $P < 0.0001$; Fig. 4b, c). However, the genomes with the highest GC contents did not contain the greatest frequency of PQS, so this association cannot be taken at face value. This correlation was not seen in the




Fig. 5. PQS in *Aspergillus* spp. are genetically conserved. (a) *Aspergillus* spp. were categorized into sections based upon a phylogenetic tree generated by Steenwyk et al. [46]. The frequency of PQS was shown to be closely associated with the genetic relatedness of these fungi. (b) A PQS was found in an exonic region of *cyp51A* from *A. fumigatus*. A BLAST search of this sequence identified conservation of this exact sequence or variations of this sequence within the section *Fumigati* (highlighted in red). Alignments were performed using the Clustal Omega web tool (EMBL-EBI). Sequence is reverse complemented.

Taphrinomycotina, but this is likely due to the much lower number of genomes used ($r=0.4052$, $P>0.05$; Fig. 4d). All the analyses above were also conducted in genomes from the Basidiomycota, Mucromycota, Zoopagomycota, Chytridiomycota, Microsporidia and Cryptomycota. However, due to their unknown genome completeness, these analyses can be found in Figs S1 and S2.

PQS are evolutionarily conserved in *Aspergillus* spp.

Evolutionary conservation of genetic motifs within the genome are a hallmark of their fundamental importance to how that organism functions. Therefore, we endeavoured to explore whether there was evolutionary conservation of PQS within fungal genomes. We chose to explore this relationship in *Aspergillus* spp., due to the robustness and accuracy of the phylogenetic tree available [52].

The frequency of PQS/kbp appeared to be intrinsically linked to how closely related species were, with species within the same section displaying similar PQS frequencies (Fig. 5). *Aspergillus* were divided into 13 sections (range of PQS/kbp in brackets), the *Aspergillus* (0.364–0.461), the *Fumigati* (0.204–0.224), the *Candidi* (0.747–0.816), the

Circumdati (0.395–0.467), the *Flavi* (0.225–0.289), the *Ochraceoros* (0.421–0.422), the *Usti* (0.385–0.396), the *Versicolores* (0.282–0.308) and the *Nigri* (0.495–0.782). The *Nigri* displayed the largest variation in PQS, however the lesser related species *A. carbonarius* and *A. aculeatus* skewed this range and if only the closest related species were noted, this range would be from 0.495 to 0.611 (Fig. 5a). As the *Nidulantes*, *Terrei* and *Clavati* only contained one member each, correlations in these sections could not be made. Notably, we found a sequence likely to form a quadruplex within *cyp51A* in *A. fumigatus*, and due to the important implications of *cyp51A* in azole resistance, we chose to see if this sequence was conserved within the section *Fumigati*. Interestingly, *A. fischeri*, *A. novofumigatus* and *A. lentulus* all retained the PQS observed in *A. fumigatus*. Moreover, this sequence was retained in cryptic *Aspergillus* spp. such as *A. felis* and *A. viridinutans*, whilst a variation on this sequence could be found in *A. arcoverdensis* (Fig. 5b). However, no sequence with significant similarity could be found in *A. turcosus* and the sequences in *A. udagawae*, *A. wyomingensis*, *A. siamensis* and *A. aureolus* would be unlikely to form the same quadruplex structure predicted for *A. fumigatus*.




Fig. 6. Pathogenic *Aspergillus* species have lower PQS frequencies compared to their non-pathogenic/inrequently pathogenic counterparts. The frequency of PQS/kbp and PQS/GC% were quantified and compared between species within *Aspergillus* spp. (a) The PQS/kbp in all species of *Aspergillus* in the study. The frequency of PQS/kbp (b) and PQS/GC% (c) in pathogenic and non-pathogenic/inrequently pathogenic species. Dots represent individual species within a genus. Pink bars represent pathogenic species, whilst blue bars represent non-pathogenic/inrequently pathogenic organisms. The error bars represent the sd. Significance was determined by Student's *T* test. Asterisks indicate $P < 0.05$.

Loss of PQS may be associated with pathogenicity in *Aspergillus* spp

The Ascomycota contains many of the most prevalent fungal pathogens of both plants and humans, including the genera *Aspergillus* spp. which contains the important human pathogen, *A. fumigatus*. Although, not all species within *Aspergillus* spp. are frequent causes of infection in humans and we found high variation in the PQS frequencies amongst species (Fig. 6a). Therefore, we compared the PQS frequency between species currently considered to be the most pathogenic and those which are non-pathogenic or infrequently pathogenic to humans, to explore whether there was a link between PQS-frequency and pathogenicity.

Indeed, comparing 72 species of *Aspergillus* (22 pathogenic, 50 non-pathogenic/inrequently pathogenic) pathogenic species had a significantly lower frequency of PQS/kbp (0.364 vs. 0.523) and PQS/GC% (249.9 vs. 380.5) within their genomes on average, compared to non-pathogenic species (Fig. 6b and c). This was an observation that was also noted for *Cryptococcus* spp. (containing the important basidiomycete pathogen *Cryptococcus neoformans*) but not for *Candida* spp. (containing the clinically relevant pathogen *Candida albicans*; Fig. S3). Furthermore, the most pathogenic species, *A. fumigatus*, had a lower than average frequency

of PQS (0.221 PQS/kbp; Fig. 6a). Although PQS frequency within *A. fumigatus* was one of the lowest amongst aspergilli, this did not mean that PQS were not enriched within certain genomic regions or within certain genes. Therefore, we explored *A. fumigatus* in further detail, and endeavoured to identify the genetic location and potential functional roles of quadruplexes in this organism.

PQS are enriched in repeat regions and mRNA, depleted in the tRNA, and are predominantly found in genes involved in metabolism in *A. fumigatus*

All previous wide-scale analyses were conducted using a stringent threshold of 1.5, as this highlighted sequences that were practically guaranteed to form quadruplexes. However, many sequences scoring below this threshold also have the potential to form these secondary structures. Therefore, the default G4Hunter threshold of 1.2 was used for the in-depth analyses of *A. fumigatus* to identify the entire repertoire of fungal PQS, as quadruplex formation for sequences above this threshold have been experimentally validated [53].

Genomic rearrangements have been suggested to contribute to the success of fungal pathogens, their ability to colonize and invade, and their adaptation to host niches. Repetitive genomic regions in *A. fumigatus* (e.g. coding tandem




Fig. 7. PQS in *A. fumigatus* are enriched in the repeat regions and mRNA. The location of PQS found 100 bp before, within, and 100 bp after annotated genomic features in *A. fumigatus* with a G4Hunter score ≥ 1.2 were identified. (a) The total number of PQS in known genomic features and (b) frequency of PQS comparative to the genomic length of the annotated features. (c) PQS-containing genes are primarily associated with metabolism. Associated protein classes were determined using PANTHER Protein Class v.15.0.

repeats) serve as hotspots for genome rearrangement, and mRNA splicing events in around 30% of genes (~3000 genes) contribute to significant functional diversity, which likely contributes to the rapid adaptation to new environments [54, 55]. Quadruplexes have previously been found within several genomic regions in eukaryotes, including mRNA, introns, exons, promoters and telomeric regions. Evidence for their involvement in regulating processes such as mRNA splicing is ever increasing, and it is likely that quadruplexes could regulate these processes in fungi [56]. Therefore, we explored the genomic location of PQS within annotated genomic features in *A. fumigatus* to provide some insight into their potential functional roles.

To evaluate the position of PQS within *A. fumigatus*, the annotation information for the strain Af293 was obtained from the NCBI database and the presence of PQS within defined genomic features, or within 100 bp before or after these features were analysed. Both the total number of PQS per feature and the frequency of PQS/kbp of the described features were noted.

When only total PQS were considered, the largest number of PQS could be found within the genes and mRNA, with few PQS found in other genomic features (Fig. 7a). This was also the same in *C. albicans* and *C. neoformans* (Fig. S4). However, this was not the same when considering the frequency of PQS/kbp of the genomic features. In *A. fumigatus*, the greatest frequency of PQS could be found in the repeat regions and mRNAs, whilst the lowest frequency could be found within the tRNAs (Fig. 7b). It is well established that PQS are enriched in telomeres and the PQS frequency in *A. fumigatus* telomeres was high (14.95 PQS/kbp). However, PQS were also abundant in non-telomeric repeat regions (2.07 PQS/kbp in all repeat regions vs. 1.93 PQS/kbp in repeat regions without telomeres; Fig. 7b). Notably, tandem repeats found

in *A. fumigatus* genes play important roles in host–pathogen interactions and contribute to a diverse range of functional proteins. Leviansky *et al.*, [55] highlighted 292 genes, which contained coding tandem repeats in the *A. fumigatus* genome. We explored these tandem repeat sequences and found that 130 (44.5%) of these genes contained at least one PQS in their tandem repeat regions (Table S3). These observations were suggestive of the potential involvement of quadruplexes in the promotion of genetic diversity and adaptation in *A. fumigatus*.

In *A. fumigatus*, 35.1% of genes contained at least one PQS likely to form quadruplex structures. In *C. neoformans*, this number was almost double, with 59.9% of genes containing PQS. Conversely, PQS were only found in 5.6% of genes in *C. albicans*. Despite the discrepancies in the number of genes where PQS can be found between the organisms, in all cases, PQS were primarily located in genes that encoded proteins involved in metabolism, nucleic acid binding, cell transport and protein modification (Fig. 7c). They were least likely to be found in genes encoding for calcium-binding proteins, extracellular matrix proteins, cell adhesion molecules and defence/immunity proteins.

Quadruplexes may be involved in numerous important biological functions in *A. fumigatus* and genes upregulated in hyphae or during hypoxia are enriched in PQS

As we knew the genomic location of the PQS, we could then identify the number and identity of the genes that contained these sequences. This further enabled us to identify their potential biological implications. GO-term analysis was conducted to identify the potential biological/molecular functions that PQS-containing genes might participate in. In *A. fumigatus*, we observed enrichment of genes involved in numerous functions such as oxidoreductase and hydrolase




Fig. 8. PQS-containing genes may be involved in numerous biological/molecular functions within pathogenic fungi. GO-term enrichment analysis of PQS-containing genes was performed using the FungiFun web tool V2.2.8. White bars indicate terms associated with molecular function, pink bars indicate terms associated with biological functions, and blue bars indicate terms associated with cellular components. The first number in brackets represents the number of genes and the second is the percentage of genes associated represented in the annotated category. Significance was determined via Fisher's exact test. $P < 0.05$ was considered significant.

activity, zinc and molybdenum ion binding, transcription, and the biosynthesis of riboflavin, tryptophan, melanin and secondary metabolites (Fig. 8a). All the enriched terms can be found in Table S4.

Although PQS could be found in genes associated with key fungal processes, we aimed to link the presence of PQS within these organisms with potential biological and pathophysiological functions. To investigate this, transcriptome datasets of *A. fumigatus* during germination [47], hypoxia [48], iron limitation and oxidative stress [49], or in biofilms [51] were analysed. In each instance, the top 20 upregulated genes (compared to dormant/unstressed *A. fumigatus* controls) were investigated for the presence and frequency of PQS. Notably, PQS could be found in 77.5% of the genes investigated (Fig. 9a). This included genes that were upregulated in germinating conidia (16/20; 80.0%), hyphae (14/20; 70.0%), after 12 h of hypoxia (17/20; 85.0%), following iron limitation (17/20; 85.0%), undergoing oxidative stress (14/20; 70.0%), and in biofilms (15/20; 75.0%). These genes were investigated further to assess whether they were enriched with PQS and contained a higher frequency of PQS comparative to the average frequency found in the *A. fumigatus* genome. In all cases, the average PQS frequencies in the upregulated genes were higher than the average PQS observed throughout the entire genome (Fig. 9b, Table S5). However, only the average PQS frequencies in the hyphae and hypoxia groups

were deemed significantly higher compared to the average genome frequency. Moreover, 71.4% of genes upregulated in hyphae and 69.2% upregulated in hypoxia contained PQS frequencies greater than the average in the genome. The genes containing the highest PQS frequencies in each condition were Afu8g01710 in germinating conidia and hyphae (11.90 PQS/kbp), Afu4g09580 in hypoxic fungi (5.59 PQS/kbp), Afu3g03650 during iron limitation (8.50 PQS/kbp), Afu5g10220 during oxidative stress (5.28 PQS/kbp) and Afu8g01980 in biofilms (5.90 PQS/kbp). Interestingly, each of these genes were upregulated in at least three out of the six conditions investigated.

PQS are in genes linked to virulence and drug resistance in *A. fumigatus*

G4s have been characterized in virulence genes from several pathogens and have arisen as a promising target for antimicrobial therapy and overcoming drug resistance [25]. Moreover, an iM in the promoter of the HIV-1 pro-viral genome has also been recently described [37]. Thus, whether PQS could be found in genes associated with virulence/drug resistance in *A. fumigatus* was explored.

Although the functions of many *A. fumigatus* genes are unknown, at least 56 genes have been found to be associated with the virulence and drug resistance of this organism




Fig. 9. PQS can be found in genes that are highly upregulated during fungal germination, in response to environmental stresses, and in biofilms in *A. fumigatus*. Transcriptomes of *A. fumigatus* during germination, hypoxia, iron-limiting conditions, oxidative stress and in biofilms, were analysed and the 20 most upregulated genes were further investigated using QGRS Mapper and G4Hunter. (a) Genes that were highly upregulated (compared to dormant or untreated *A. fumigatus* conidia) contained PQS. Data points represent genes that contain PQS. (b) The average PQS frequency of genes upregulated in hyphae and during hypoxia were significantly higher when compared to the average PQS/kbp for the entire genome (1.55 PQS/kbp, red-dashed line), when analysed with the default G4Hunter settings. Data points represent the PQS frequency of an individual gene. Boxplot represents the median, maximum and minimum values. Whether the mean frequency of PQS within the upregulated genes of each condition was significantly higher compared to the mean of the entire genome was analysed via one-sample *T* test. * indicates a significant difference ($P < 0.05$).

(Table 1). PQS could be found in ~78% of these genes (44/56 genes). This included genes that have been implicated in azole resistance (*cyp51A*, *cyp51B*, *abcC*, *hsp90*, *atrF*, *atrR*, *hmg1*) and echinocandin resistance (*fks1*). Genes implicated in virulence, which were enriched in PQS compared to the average frequency for the entire genome (PQS/kbp in brackets), included *mkk2* (4.78), *zafA* (4.04), *laeA* (2.36), *rasB* (4.97), *medA* (4.25), *glfA* (2.56), *pacC* (3.60), *sidG* (7.06), *hapX* (2.71), *calA* (7.44), *zrfC* (5.23), *aspf13* (8.65), *cpcA* (2.46), *gel2* (2.35) and *mcsA* (2.63). Furthermore, numerous genes also contained high-scoring sequences likely to form quadruplexes, which were potentially composed of three or more G-tetrads; including, *zafA*, *pes1*, *laeA*, *pkac1*, *hasA*, *pacC*, *lysF*, *kre2*, *zrfC*, *hscA*, *mcsA*, *atrR*, *cyp51A*, *cyp51B*, *abcC*, *atrF* and *hmg1* (Table 1).

The highest scoring potential quadruplex-forming sequences for each of these genes were then re-analysed in an alternative PQS predictive algorithm called QGRS Mapper. The scores of sequences known to form quadruplexes were compared to scores of the PQS identified in *A. fumigatus* using this algorithm. This was conducted to provide further insight into whether these sequences were likely to form quadruplex structures. Known quadruplex-forming sequences were obtained from the G4IPDB database ($n=94$ sequences chosen at random) and were shown to have QGRS scores of around 21, 42 or 63 (depending on the number of G-tetrads – 2, 3 or 4, respectively). All the PQS-containing virulence genes

contained at least one sequence scoring higher than 20, with many sequences scoring between 29 and 42 thus, very likely to form quadruplex structures, at least *in vitro* (Fig. S5).

Altered PQS frequencies in orthologous genes in the close genetic relative of *A. fumigatus*, *A. fischeri*

A. fischeri is one of the closest genetic relatives of *A. fumigatus* and around ~90% of the proteome is shared between the two species [57]. However, whilst *A. fumigatus* is regularly isolated from patients, only a few known cases of human infection with *A. fischeri* have ever been reported. Indeed, *A. fischeri* has been shown to be far less virulent compared to *A. fumigatus* in several murine models, but interestingly, *A. fischeri* appears to intrinsically display reduced susceptibility to azoles [57]. Furthermore, orthologues of all the *A. fumigatus* genes could be found in *A. fischeri*. Therefore, the PQS frequencies in orthologous genes in *A. fischeri* were quantified. Only quadruplex frequencies in the coding regions were compared between species, as other genetic regions of the *A. fischeri* orthologues are poorly understood.

Considering the genetic similarity between the two species and high sequence similarities in orthologues of the virulence genes under investigation (average similarities of 96.4% and average sequence coverage of 99%), the distribution of the PQS between the two appeared quite different (Fig. 10a). The PQS frequencies for *sidG*, *aspf13*, *cpcA*, *gliZ*, *gliG*, *gliK*, *gliT*

Table 1. PQS in genes associated with virulence and drug resistance in *A. fumigatus*

Identifier	Gene	Description	Sequence
Afu2g02690	<i>atrR</i>	Fungal specific transcription factor, putative	GGGGGAGGGTGCGGGATGCTCTTCGGG
Afu4g06890	<i>cyp51A</i>	14-alpha sterol demethylase	GGGTCCCAGGGAACGGGCAAGGGG
Afu7g03740	<i>cyp51B</i>	14-alpha sterol demethylase	GGGGTAGGGGGGCCAGGG
Afu1g14330	<i>abcC</i>	Azole transporter	GGGCCTGGGACAGGGTCCGGG
Afu6g04360	<i>atrF</i>	Putative ABC transporter	GGGTATTTGTTTGGGGGATGGG
Afu5g04170	<i>hsp90</i>	Heat shock protein	GGAGGAGGAGG
Afu6g12400	<i>fks1</i>	Putative 1,3-beta-glucan synthase catalytic subunit, major subunit of glucan synthase	GGAAGGGGCTCGGGCATGGG
Afu2g03700	<i>hmg1</i>	Hydroxymethylglutaryl-CoA (HMG-CoA) reductase	GGGCGGCAGGGACAGGGGCAACTTCGGG
Afu1g05800	<i>mkk2</i>	Putative mitogen-activated protein kinase kinase (MAPKK)	GGTCGGTGGGTGG
Afu1g06900	<i>crzA</i>	C2H2-type zinc finger transcription factor involved in calcium ion homeostasis	GGAATGGGAGGAGGG
Afu1g10080	<i>zafA</i>	Putative C2H2 zinc-responsive transcriptional activator	GGGCAGGGCAGAGCAGGGCAGGGG
Afu1g10380	<i>nrps1</i>	Non-ribosomal peptide synthetase (NRPS)	GGGAGCAAGGGGAAATCGCTGAGGGTGGG
Afu1g10880	<i>pmcA</i>	Putative P-type calcium ATPase	GGGCGGGTCAGAAATCAGACTTGGGAAGGG
Afu1g13140	<i>gpaB</i>	G protein-coupled receptor alpha subunit	GGTGGAGGTGG
Afu1g14660	<i>laeA</i>	Involved in regulation of secondary metabolism	GGGATGTGGGACAGGGATTTGGG
Afu1g17200	<i>sidC</i>	Fusarinine C non-ribosomal peptide synthetase (NRPS), putative	GGAGAAGGATCGGGCAAGG
Afu2g07680	<i>sidA</i>	L-ornithine N5-oxygenase	GGCAATGGCCAGGGAAGCGG
Afu2g07770	<i>rasB</i>	Ras family GTPase protein	GGGACGGGGAGGGGCGG
Afu2g12200	<i>pkaC1</i>	cAMP-dependent protein kinase catalytic subunit	GGGCTGGGGGTAAGGGGAGGGGG
Afu2g13260	<i>medA</i>	Putative regulator of adherence, host cell interactions and virulence	GGCGGAGGGATGG
Afu2g17600	<i>pksP</i>	Conidial pigment polyketide synthase alb1	GGTTGGTCGGCGGG
Afu3g12890	<i>hasA</i>	C6 transcription factor hasA	GGGAAGGGAAGGGAAGGG
Afu3g12690	<i>glfA</i>	Putative UDP-galactopyranose mutase, enzyme in the first step of galactofuranose biosynthesis	GGTCTTGGGCTTGGTGTGG
Afu3g11970	<i>pacC</i>	C2H2 finger domain transcription factor	GGGGGGGGGAATGCTTGGG
Afu3g03650	<i>sidG</i>	Putative acetyltransferase with a predicted role in iron metabolism	GGAAGTGGGTCTGGCCGAGG
Afu3g03420	<i>sidD</i>	Nonribosomal peptide synthetase 4	GGAAGGGGACCCGGACGGGG
Afu3g03400	<i>sidF</i>	Siderophore biosynthesis acetylase AceI, putative	GGATCGGATGGCGTGG
Afu3g02270	<i>cat1</i>	Mycelial catalase	GGACGTGGAGTGGGAATGGG
Afu5g03920	<i>hapX</i>	bZIP transcription factor HapX	GGAGATGGAGATGGAGATGG
Afu5g08570	<i>pkaC2</i>	Class II protein kinase A (PKA)	GGGGGTCCGAAGG
Afu5g08890	<i>lysF</i>	Putative homoconitase	GGGACGCGGGGGCGGCATCGGGAGCGGG
Afu3g09690	<i>calA</i>	Invasive calA	GGGCAGGGCAGTGGAGAGGG
Afu5g10760	<i>mnt1</i>	Putative alpha-1,2-mannosyltransferase with a predicted role in protein glycosylation	GGGAGAGGGCACAGGGGGG

Continued

Table 1. Continued

Identifier	Gene	Description	Sequence
Afu5g11230	<i>rasA</i>	Ras family GTPase protein	GGTGGTGGTGG
Afu4g09560	<i>zrfC</i>	Zinc transporter that functions in neutral or alkaline environments	GGGATTGGTGGGGTCCAGGGGAGGG
Afu4g10460	<i>hcsA</i>	Homocitrate synthase, essential enzyme of the alpha-aminoadipate pathway of lysine biosynthesis	GGGTGGGGTTTATGAGGGCGGG
Afu2g12630	<i>asf13</i>	Allergen Asp f 13	GGATGGTGGGGG
Afu4g12470	<i>cpcA</i>	Transcriptional activator of the cross-pathway control system of amino acid biosynthesis	GGTGGCGGGGG
Afu6g09660	<i>gliP</i>	Nonribosomal peptide synthetase gliP	GGGATGGCCGGGGATGG
Afu6g11390	<i>gel2</i>	GPI-anchored 1,3-beta-glucanosyltransferase	GGTACGGAGAGGATGAGG
Afu8g02750	<i>cgrA</i>	Nucleolar rRNA processing protein	GGAACGGGAGGGGAGG
Afu8g01670	<i>cat2</i>	Putative bifunctional catalase-peroxidase	GGTCGGGGTCTTGGTCCAGG
Afu6g04820	<i>pabA</i>	Para-aminobenzoic acid synthetase, an enzyme catalysing a late step in the biosynthesis of folate	GGTGGCGGGGG
Afu6g03590	<i>mcsA</i>	Methylcitrate synthase	GGGGTTCGGGTATTCTGGGTCTGGGG

and *gliH* were higher in *A. fumigatus* compared to *A. fischeri*. Conversely, the PQS frequencies for *laeA*, *sidA*, *medA*, *glfA*, *sidD*, *cat1*, *lysF*, *rodA*, *kre2*, *rasA*, *atrF*, *hmg1* and *gliI* were higher in *A. fischeri* compared to *A. fumigatus* (Fig. 10b, Table S6). Furthermore, the average PQS frequencies for the PQS-containing upregulated genes during hypoxia and in hyphae in *A. fumigatus* were higher in *A. fumigatus* when compared to *A. fischeri* orthologues (3.19 vs. 2.59 PQS/kbp and 3.65 vs. 3.00 PQS/kbp, for genes upregulated during hypoxia and in hyphae, respectively; Fig. S6). Moreover, *A. fischeri* orthologues of *hmg1*, *pkaC1*, *hasA*, *glfA*, *kre2* and *hcsA* also do not share the high-scoring PQS that were found in *A. fumigatus*. The implication of these differences to the functions of these organisms, and whether the altered PQS frequencies are associated with virulence, are currently unknown. However, future studies investigating these differences might unveil how these species differ in their pathogenicity.

DISCUSSION

In this study, we computationally predicted PQS within the genomes of fungi. We made several important observations. This was the first study to investigate the frequency of PQS amongst numerous genetically distinct fungal species. Moreover, we highlighted the genomic location of PQS in *A. fumigatus*, their potential involvement in key biological processes, and discussed their presence in genes associated with virulence and drug resistance.

The frequency of PQS throughout genomes is highly variable; for example, human genomes were shown to contain around 0.228 PQS/kbp, whereas the genomes of *Escherichia coli* contain around 0.028 PQS/kbp [15]. In this study we also found significant differences in the number and frequency of PQS throughout fungal genomes. For example,

fungi from the Saccharomycotina (primarily composed of yeasts) contained a low frequency of PQS. Conversely, filamentous fungi from the Pezizomycotina (containing many important pathogens of both plants and humans) had much higher PQS frequencies and >15 fold more PQS than fungi from the Saccharomycotina on average. It has previously been shown that G4s contribute to genetic instability in yeasts (*Saccharomyces* spp.) and as suggested for bacteria, it is possible that G4s may also have been deselected through evolution in ascomycete yeasts [58, 59]. However, things are more complicated concerning basidiomycetous yeasts such as *Cryptococcus* spp. and *Malassezia* spp. Cryptococcal species such as *C. wingfieldii*, *C. amyloletus* and *C. floricola* possessed some of the highest PQS frequencies in the study. Alternatively, all the *Malassezia* species investigated had much lower PQS frequencies than expected, which was surprising given their genomes were GC rich compared to many other fungi (>55% GC content). We also identified many PQS within genes, and mRNA of *A. fumigatus*, but a higher frequency of PQS within the repeat regions and mRNA. It has previously been observed that *S. cerevisiae* promoters and open reading frames are enriched with sequences capable of potentially forming intramolecular G4s [60]. A recent study by Čutová *et al.*, also demonstrated that inverted repeats were enriched in the centromeres and rDNA/rRNA regions, and G4s were enriched in the telomeres and tRNAs of *S. cerevisiae* [61].

Notably, the genomes of pathogenic aspergilli and cryptococci contained significantly lower PQS frequencies relative to their non-pathogenic/infrequently pathogenic counterparts. Conversely, there was no correlation between PQS frequency and pathogenicity between *Candida* species or within the Saccharomycotina. Interestingly, loss of PQS has




Fig. 10. Differences between PQS frequency in genes involved in the pathogenicity of *A. fumigatus* and their orthologues in the close non-pathogenic relative, *A. fischeri* (a) The differences between the distribution of PQS in *A. fumigatus* (black) and *A. fischeri* (pink) genomes. (b) The PQS frequency in genes involved in virulence, drug resistance and gliotoxin biosynthesis in *A. fumigatus* (black), compared with the PQS frequency in the respective *A. fischeri* orthologues (pink). PQS frequencies were quantified using the default G4Hunter settings. The mean genome frequencies for *A. fumigatus* and *A. fischeri* were 1.55 PQS/kbp and 1.53 PQS/kbp, respectively.

recently also been observed in pathogenic *Coronaviridae* [62]. The exact reasons underlying the loss of PQS are not currently known. Fungi may have evolved to retain the most essential quadruplex-forming sequences or those that provided them with a pathogenic advantage. It has been reported that host nucleolin (an RNA-binding protein) can bind and stabilize quadruplexes in the long terminal repeat (LTR) promoter of HIV-1, which can silence viral transcription [63]. Therefore, in this situation, loss of quadruplexes would be beneficial for immune evasion. Contrarily, it has been suggested that iMs within HIV-1 are triggered after they promote the acidification of intracellular compartments, modulating viral processes [37]. In this instance, the formation of quadruplexes could be beneficial and promote pathogenicity. In the case of *Aspergillus* and *Cryptococcus* species, this loss of PQS might also be linked with evasion of the immune response. Furthermore, the low PQS frequency observed within *Candida* and *Malassezia* species

might be associated with their ability to form commensal relationships and avoid stimulating a host response. The importance of quadruplex stabilization may therefore be situational and could be detrimental or beneficial to the fungus depending on the environment. It is known that fungal pathogens such as *A. fumigatus* and *C. neoformans* can survive the acidic conditions within intracellular organelles to propagate infection [64, 65]. Therefore, it could be interesting to investigate the dynamics/flux of quadruplex formation under conditions that fungi would experience within the host and whether these structures can contribute to fungal survival and propagation.

Although the overall genome PQS frequency in *A. fumigatus* was one of the lowest amongst the aspergilli, the average PQS frequencies in genes involved in virulence and gliotoxin synthesis, or those upregulated in response to conditions such as hypoxia, iron limitation and oxidative

stress, appeared to be much higher than the average genome frequency. GO-term enrichment analysis implicated potential roles for PQS-containing genes in functions associated with virulence in pathogenic fungi such as zinc-ion binding [66], riboflavin and tryptophan biosynthesis [67], melanin [68] and secondary metabolite synthesis [69], metallo-peptidase activity [70], and the response to environmental stimuli [71]. Moreover, PQS were predominantly found in genes associated with fungal metabolism.

It is widely acknowledged that metabolism in fungi is central to its virulence, pathogenicity and survival [72]. The ability of a fungus to rapidly adapt to the host microenvironment is achieved through processes such as metabolic remodelling, stress resistance and the utilization of amino acids [67]. This in turn has a significant effect on the triggering of important virulence traits, like the production of hyphae, biofilm growth, capsule formation in *C. neoformans* and melanization [67]. Moreover, metabolic pathways influence fungal vulnerability to innate immune defences and can regulate susceptibility to antifungal drugs [73]. Not only primary metabolism, secondary metabolism results in the release of various secondary metabolites and toxins, such as gliotoxin, aflatoxin and candidalysin [74, 75]. Furthermore, many genes containing PQS within *S. cerevisiae*, humans and bacteria were also primarily associated with metabolism [76, 77]. These data support the observations made in *A. fumigatus* and implicates an important role for quadruplexes in metabolism throughout different organisms. Thus, the enrichment of PQS within these genes is suggestive of the potential importance of quadruplexes in the regulation of key biological functions and stress responses, and it would be interesting to explore the effects quadruplex-targeting ligands could have on these processes.

There are now an ever-increasing number of G4s identified within genes linked to microbial pathogenicity. G4-forming motifs located in the *hsdS*, *recD* and *pmrA* genes of *S. pneumoniae*, and *var* genes of *P. falciparum* can modulate host-pathogen interactions [78, 79]. Moreover, targeting G4s in *espK*, *espB* and *cyp51* from *Mycobacterium tuberculosis* with the G4/iM-binding ligand TMPyP4 could inhibit transcription of these essential virulence genes [80]. Binding of several G4-targeting ligands to G4s in viruses have been shown to limit the virulence of the Herpes simplex virus-1, HIV-1, Ebola and Hepatitis C [81]. Interestingly, dinuclear ruthenium (II) complexes (well-characterized G4 and iM DNA binding agents) are active against methicillin-resistant *S. aureus* and vancomycin-resistant *Enterococcus* spp. [35, 36], and another G4/iM-ligand, berberine, has also demonstrated activity towards fluconazole-resistant *C. albicans* and *C. neoformans* [82].

Notably, sequences highly likely to form quadruplexes (G_3+L_{1-12}) were also found in important genes implicated in azole resistance (*atrR*, *cyp51A*, *cyp51B*, *abcC* and *hmg1*) and virulence (*zafA*, *pes1*, *laeA*, *pkac1*, *hasA*, *pacC*, *lysF*, *kre2*, *zrfC*, *hscA* and *mcsA*). Many of which were not found to be shared with the close non-pathogenic relative, *A. fischeri*.

Mutations in *cyp51A* are a major contributing factor to azole resistance, but there is growing evidence that mutations in *cyp51B* and the involvement other members of the ergosterol synthesis pathway contribute to this resistance phenotype [83, 84]. Moreover, genes such as *laeA* (encoding a master regulator of secondary metabolite biosynthesis) and *pacC* (encoding a pH-responsive regulator), regulate key virulent processes, such as gliotoxin production and host-cell adaption, respectively [85–87]. Thus, if the quadruplex structures in these genes form *in vivo* and if they are implicated in regulating their transcription, they could pose an interesting target to modulate fungal pathogenic processes, as observed for other pathogenic microorganisms, such as viruses.

CONCLUSIONS

Here, we provide a representative overview of potential quadruplex-forming sequences across the fungal kingdom and discuss their potential implications in fungal virulence and drug resistance. If these structures form *in vivo* and are involved in regulating transcription and biological processes in fungi, then they could have the potential to represent novel antifungal targets. Understanding whether quadruplexes could form under pathophysiological conditions, whether fungi themselves could release quadruplex-binding ligands to modulate host responses, or identifying whether there are fungal specific quadruplex-forming sequences could also be important factors to consider. Thus, targeting quadruplexes in fungi could represent a novel antifungal target, although experimental validation is necessary.

Funding information

This research was funded by the Biotechnology and Biological Sciences Research Council (grant number BB/L02229X/1). E.F.W. was supported by the Wellcome Trust Pathfinder Grant (grant number 204515/Z/16/Z). V.B. was supported by the Czech Science Foundation (grant number 18-15548S). The work conducted by the U.S. Department of Energy Joint Genome Institute, a DOE Office of Science User Facility, is supported by the Office of Science of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231. This manuscript has previously been submitted as a preprint [88].

Acknowledgements

I would like to acknowledge Phil Spence for contributing the G4 and iM structures and for critical reviewing of the manuscript.

Author contributions

Conceptualization, E.F.W. and S.B.; Methodology, E.F.W., N.B., V.B. and S.B.; Formal analysis, E.F.W., N.B., V.B. and S.B.; Investigation, E.F.W., N.B., V.B. and S.B.; Resources, N.B. and V.B.; Writing – original draft preparation, E.F.W., Z.A.E.W. and S.B.; Writing – review and editing, E.F.W., N.B., V.B., Z.A.E.W. and S.B.; Visualization, E.F.W., N.B., V.B. and S.B.; Supervision, S.B.; Project administration, S.B., Funding, V.B. and Z.A.E.W. All authors have read and agreed to the published version of the manuscript.

Conflicts of interest

The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript or in the decision to publish the results.

References

1. Correction. Stop neglecting fungi. *Nat Microbiol* 2017;2:17123.
2. Skellam E. Strategies for engineering natural product biosynthesis in fungi. *Trends Biotechnol* 2019;37:416–427.
3. Hernández VA, Machuca Ángela, Saavedra I, Chavez D, Astuya A et al. *Talaromyces australis* and *Penicillium murcianum* pigment production in optimized liquid cultures and evaluation of their cytotoxicity in textile applications. *World J Microbiol Biotechnol* 2019;35:160.
4. Hooker CA, Lee KZ, Solomon KV. Leveraging anaerobic fungi for biotechnology. *Curr Opin Biotechnol* 2019;59:103–110.
5. Gallone B, Steensels J, Prah T, Soriaga L, Saels V et al. Domestication and divergence of *Saccharomyces cerevisiae* beer yeasts. *Cell* 2016;166:1397–1410.
6. Brown GD, Denning DW, Gow NAR, Levitz SM, Netea MG et al. Hidden killers: human fungal infections. *Sci Transl Med* 2012;4:165rv13.
7. Fisher MC, Gurr SJ, Cuomo CA, Bleher DS, Jin H et al. Threats posed by the fungal Kingdom to humans, wildlife, and agriculture. *mBio* 2020;11:e00449–20 [Epub ahead of print 05 05 2020].
8. Köhler JR, Casadevall A, Perfect J. The spectrum of fungi that infects humans. *Cold Spring Harb Perspect Med* 2014;5:a019273.
9. van Arkel ALE, Rijpstra TA, Belderbos HNA, van Wijngaarden P, Verweij PE et al. COVID-19-associated pulmonary *Aspergillo*sis. *Am J Respir Crit Care Med* 2020;202:132–135.
10. Ramirez-Garcia A, Pellon A, Rementeria A, Buldain I, Barreto-Bergter E et al. *Scedosporium* and *Lomentospora*: an updated overview of underrated opportunists. *Med Mycol* 2018;56:102–125.
11. Berman J, Krysan DJ. Drug resistance and tolerance in fungi. *Nat Rev Microbiol* 2020;18:319–331.
12. World Health O. *First Meeting of the WHO Antifungal Expert Group on Identifying Priority Fungal Pathogens: Meeting Report*. Geneva: World Health Organization; 2020.
13. Varshney D, Spiegel J, Zyner K, Tannahill D, Balasubramanian S. The regulation and functions of DNA and RNA G-quadruplexes. *Nat Rev Mol Cell Biol* 2020;21:459–474.
14. Abou Assi H, Garavís M, González C, Damha MJ. i-Motif DNA: structural features and significance to cell biology. *Nucleic Acids Res* 2018;46:8038–8056.
15. Marsico G, Chambers VS, Sahakyan AB, McCauley P, Boutell JM et al. Whole genome experimental maps of DNA G-quadruplexes in multiple species. *Nucleic Acids Res* 2019;47:3862–3874.
16. Burge S, Parkinson GN, Hazel P, Todd AK, Neidle S. Quadruplex DNA: sequence, topology and structure. *Nucleic Acids Res* 2006;34:5402–5415.
17. Dai J, Dexheimer TS, Chen D, Carver M, Ambrus A et al. An intramolecular G-quadruplex structure with mixed parallel/antiparallel G-strands formed in the human BCL-2 promoter region in solution. *J Am Chem Soc* 2006;128:1096–1098.
18. Víglaský V, Bauer L, Tluczková K. Structural features of intra- and intermolecular G-quadruplexes derived from telomeric repeats. *Biochemistry* 2010;49:2110–2120.
19. Lane AN, Chaires JB, Gray RD, Trent JO. Stability and kinetics of G-quadruplex structures. *Nucleic Acids Res* 2008;36:5482–5515.
20. Brazier JA, Shah A, Brown GD. I-motif formation in gene promoters: unusually stable formation in sequences complementary to known G-quadruplexes. *Chem Commun* 2012;48:10739–10741.
21. Gehring K, Leroy JL, Guéron M. A tetrameric DNA structure with protonated cytosine-cytosine base pairs. *Nature* 1993;363:561–565.
22. Wright EP, Huppert JL, Waller ZAE. Identification of multiple genomic DNA sequences which form i-motif structures at neutral pH. *Nucleic Acids Res* 2017;45:2951–2959.
23. Rajendran A, Nakano S-ichi, Sugimoto N. Molecular crowding of the cosolutes induces an intramolecular i-motif structure of triplet repeat DNA oligomers at neutral pH. *Chem Commun* 2010;46:1299–1301.
24. Zeraati M, Langley DB, Schofield P, Moye AL, Rouet R et al. I-motif DNA structures are formed in the nuclei of human cells. *Nat Chem* 2018;10:631–637.
25. Saranathan N, Vivekanandan P. G-quadruplexes: more than just a kink in microbial genomes. *Trends Microbiol* 2019;27:148–163.
26. Balasubramanian S, Hurley LH, Neidle S. Targeting G-quadruplexes in gene promoters: a novel anticancer strategy? *Nat Rev Drug Discov* 2011;10:261–275.
27. Zhang Y, Yang M, Duncan S, Yang X, Abdelhamid MAS et al. G-quadruplex structures trigger RNA phase separation. *Nucleic Acids Res* 2019;47:11746–11754.
28. Bartas M, Čtůvá M, Brázda V, Kaura P, Štátný J et al. The presence and localization of G-quadruplex forming sequences in the domain of bacteria. *Molecules* 2019;24:E1711 [Epub ahead of print 02 May 2019].
29. Shankar U, Jain N, Mishra SK, Sharma TK, Kumar A. Conserved G-quadruplex motifs in gene promoter region reveals a novel therapeutic approach to target multi-drug resistance *Klebsiella pneumoniae*. *Front Microbiol* 2020;11:1269.
30. Shankar U, Jain N, Majee P, Kodgire P, Sharma TK et al. Exploring computational and biophysical tools to study the presence of G-quadruplex structures: a promising therapeutic solution for drug-resistant *Vibrio cholerae*. *Front Genet* 2020;11:935.
31. Jain N, Mishra SK, Shankar U, Jaiswal A, Sharma TK et al. G-quadruplex stabilization in the ions and maltose transporters gene inhibit *Salmonella enterica* growth and virulence. *Genomics* 2020;112:4863–4874.
32. Majee P, Shankar U, Pasadi S, Muniyappa K, Nayak D et al. Genome-wide analysis reveals a regulatory role for G-quadruplexes during *Adenovirus* multiplication. *Virus Res* 2020;283:197960.
33. Majee P, Kumar Mishra S, Pandya N, Shankar U, Pasadi S et al. Identification and characterization of two conserved G-quadruplex forming motifs in the Nipah virus genome and their interaction with G-quadruplex specific ligands. *Sci Rep* 2020;10:1477.
34. Majee P, Pattnaik A, Sahoo BR, Shankar U, Pattnaik AK et al. Inhibition of Zika virus replication by G-quadruplex-binding ligands. *Mol Ther Nucleic Acids* 2021;23:691–701.
35. Li F, Mulyana Y, Feterl M, Warner JM, Collins JG et al. The antimicrobial activity of inert oligonuclear polypyridylruthenium(II) complexes against pathogenic bacteria, including MRSA. *Dalton Trans* 2011;40:5032–5038.
36. Li F, Collins JG, Keene FR. Ruthenium complexes as antimicrobial agents. *Chem Soc Rev* 2015;44:2529–2542.
37. Ruggiero E, Lago S, Šket P, Nadai M, Frasson I et al. A dynamic i-motif with a duplex stem-loop in the long terminal repeat promoter of the HIV-1 proviral genome modulates viral transcription. *Nucleic Acids Res* 2019;47:11057–11068.
38. Asamitsu S, Obata S, Yu Z, Bando T, Sugiyama H. Recent progress of targeted G-Quadruplex-Preferred ligands toward cancer therapy. *Molecules* 2019;24:E429 [Epub ahead of print 24 Jan 2019].
39. Ruggiero E, Richter SN. G-Quadruplexes and G-quadruplex ligands: targets and tools in antiviral therapy. *Nucleic Acids Res* 2018;46:3270–3283.
40. Tassinari M, Zuffo M, Nadai M, Pirota V, Sevilla Montalvo AC et al. Selective targeting of mutually exclusive DNA G-quadruplexes: HIV-1 LTR as paradigmatic model. *Nucleic Acids Res* 2020;48:4627–4642.
41. Shen X-X, Steenwyk JL, LaBella AL, Ofulente DA, Zhou X et al. Genome-scale phylogeny and contrasting modes of genome evolution in the fungal phylum *Ascomycota*. *Sci Adv* 2020;6 [Epub ahead of print 04 11 2020].
42. Brázda V, Kolomazník J, Lýsek J, Bartas M, Fojta M et al. G4Hunter web application: a web server for G-quadruplex prediction. *Bioinformatics* 2019;35:3493–3495.
43. Mi H, Muruganujan A, Ebert D, Huang X, Thomas PD. Panther version 14: more genomes, a new Panther GO-slim and improvements in enrichment analysis tools. *Nucleic Acids Res* 2019;47:D419–d426.

44. Priebe S, Kreisel C, Horn F, Guthke R, Linde J. FungiFun2: a comprehensive online resource for systematic analysis of gene lists from fungal species. *Bioinformatics* 2015;31:445–446.
45. Kikin O, D'Antonio L, Bagga PS. QGRS Mapper: a web-based server for predicting G-quadruplexes in nucleotide sequences. *Nucleic Acids Res* 2006;34:W676–682.
46. Tomasello G, Armenia I, Molla G. The Protein Imager: a full-featured online molecular viewer interface with server-side HQ-rendering capabilities. *Bioinformatics* 2020;36:2909–2911.
47. Hagiwara D, Takahashi H, Kusuya Y, Kawamoto S, Kamei K et al. Comparative transcriptome analysis revealing dormant conidia and germination associated genes in *Aspergillus* species: an essential role for AtfA in conidial dormancy. *BMC Genomics* 2016;17:358.
48. Hillmann F, Linde J, Beckmann N, Cyrulies M, Strassburger M et al. The novel globin protein fungoglobulin is involved in low oxygen adaptation of *Aspergillus fumigatus*. *Mol Microbiol* 2014;93:539–553.
49. Kurucz V, Krüger T, Antal K, Dietl A-M, Haas H et al. Additional oxidative stress reroutes the global response of *Aspergillus fumigatus* to iron depletion. *BMC Genomics* 2018;19:357.
50. Stajich JE, Harris T, Brunk BP, Brestelli J, Fischer S et al. FungiDB: an integrated functional genomics database for fungi. *Nucleic Acids Res* 2012;40:D675–681.
51. Gibbons JG, Beauvais A, Beau R, McGary KL, Latgé J-P et al. Global transcriptome changes underlying colony growth in the opportunistic human pathogen *Aspergillus fumigatus*. *Eukaryot Cell* 2012;11:68–78.
52. Steenwyk JL, Shen X-X, Lind AL, Goldman GH, Rokas A. A robust phylogenomic time tree for biotechnologically and medically important fungi in the genera *Aspergillus* and *Penicillium*. *mBio* 2019;10:e00925-19 [Epub ahead of print 09 Oct 2019].
53. Bedrat A, Lacroix L, Mergny J-L. Re-evaluation of G-quadruplex propensity with G4Hunter. *Nucleic Acids Res* 2016;44:1746–1759.
54. Bedrat A, Lacroix L, Mergny J-L. Re-evaluation of G-quadruplex propensity with G4Hunter. *Nucleic Acids Res* 2016;44:1746–1759.
55. Levdansky E, Romano J, Shadkchan Y, Sharon H, Verstrepen KJ et al. Coding tandem repeats generate diversity in *Aspergillus fumigatus* genes. *Eukaryot Cell* 2007;6:1380–1391.
56. Jara-Espejo M, Fleming AM, Burrows CJ. Potential G-quadruplex forming sequences and N-Methyladenosine colocalize at human pre-mRNA intron splice sites. *ACS Chem Biol* 2020;15:1292–1300.
57. Mead ME, Knowles SL, Raja HA, Beattie SR, Kowalski CH et al. Characterizing the pathogenic, genomic, and chemical traits of *Aspergillus fischeri*, a close relative of the major human fungal pathogen *Aspergillus fumigatus*. *mSphere* 2019;4:e00018-19 [Epub ahead of print 20 Oct 2019].
58. Lopes J, Piazza A, Bermejo R, Kriegsman B, Colosio A et al. G-quadruplex-induced instability during leading-strand replication. *Embo J* 2011;30:4033–4046.
59. Guo JU, Bartel DP. RNA G-quadruplexes are globally unfolded in eukaryotic cells and depleted in bacteria. *Science*, 2016;353.
60. Hershman SG, Chen Q, Lee JY, Kozak ML, Yue P et al. Genomic distribution and functional analyses of potential G-quadruplex-forming sequences in *Saccharomyces cerevisiae*. *Nucleic Acids Res* 2008;36:144–156.
61. Čutová M, Manta J, Porubiaková O, Kaura P, Šťastný J et al. Divergent distributions of inverted repeats and G-quadruplex forming sequences in *Saccharomyces cerevisiae*. *Genomics* 2020;112:1897–1901.
62. Bartas M, Brázda V, Bohálová N, Cantara A, Volná A et al. In-depth bioinformatic analyses of nidovirales including human SARS-CoV-2, SARS-CoV, MERS-CoV viruses suggest important roles of non-canonical nucleic acid structures in their lifecycles. *Front Microbiol* 2020;11:1583.
63. Tosoni E, Frasson I, Scalabrin M, Perrone R, Butovskaya E et al. Nucleolin stabilizes G-quadruplex structures folded by the LTR promoter and silences HIV-1 viral transcription. *Nucleic Acids Res* 2015;43:8884–8897.
64. Wasylnka JA, Moore MM. *Aspergillus fumigatus* conidia survive and germinate in acidic organelles of A549 epithelial cells. *J Cell Sci* 2003;116:1579–1587.
65. Coelho C, Bocca AL, Casadevall A. The intracellular life of *Cryptococcus neoformans*. *Annu Rev Pathol* 2014;9:219–238.
66. Gerwien F, Skrahina V, Kasper L, Hube B, Brunke S. Metals in fungal virulence. *FEMS Microbiol Rev* 2018;42 [Epub ahead of print 01 Oct 2018].
67. Ene IV, Brunke S, Brown AJP, Hube B. Metabolism in fungal pathogenesis. *Cold Spring Harb Perspect Med* 2014;4:a019695.
68. Stappers MHT, Clark AE, Amanianda V, Bidula S, Reid DM et al. Recognition of DHN-melanin by a C-type lectin receptor is required for immunity to *Aspergillus*. *Nature* 2018;555:382–386.
69. Raffa N, Keller NP. A call to arms: Mustering secondary metabolites for success and survival of an opportunistic pathogen. *PLoS Pathog* 2019;15:e1007606.
70. Rementeria A, López-Molina N, Ludwig A, Vivanco AB, Bikandi J et al. Genes and molecules involved in *Aspergillus fumigatus* virulence. *Rev Iberoam Micol* 2005;22:1–23.
71. Pradhan A, Ma Q, de Assis LJ, Leaves I, Larcombe DE et al. Anticipatory stress responses and immune evasion in fungal pathogens. *Trends Microbiol* 2020 [Epub ahead of print 12 Oct 2020].
72. Brown AJP, Brown GD, Netea MG, Gow NAR. Metabolism impacts upon *Candida* immunogenicity and pathogenicity at multiple levels. *Trends Microbiol* 2014;22:614–622.
73. Parente-Rocha JA, Bailão AM, Amaral AC, Taborda CP, Paccez JD et al. Antifungal resistance, metabolic routes as drug targets, and new antifungal agents: an overview about endemic dimorphic fungi. *Mediators Inflamm* 2017;2017:9870679.
74. Bignell E, Cairns TC, Throckmorton K, Nierman WC, Keller NP. Secondary metabolite arsenal of an opportunistic pathogenic fungus. *Philos Trans R Soc Lond B Biol Sci* 2016;371 [Epub ahead of print 05 Dec 2016].
75. Moyes DL, Wilson D, Richardson JP, Mogavero S, Tang SX et al. Candidalysin is a fungal peptide toxin critical for mucosal infection. *Nature* 2016;532:64–68.
76. Eddy J, Maizels N. Gene function correlates with potential for G4 DNA formation in the human genome. *Nucleic Acids Res* 2006;34:3887–3896.
77. Rawal P, Kummarasetti VBR, Ravindran J, Kumar N, Halder K et al. Genome-wide prediction of G4 DNA as regulatory motifs: role in *Escherichia coli* global regulation. *Genome Res* 2006;16:644–655.
78. Mishra SK, Jain N, Shankar U, Tawani A, Sharma TK et al. Characterization of highly conserved G-quadruplex motifs as potential drug targets in *Streptococcus pneumoniae*. *Sci Rep* 2019;9:1791.
79. Harris LM, Monsell KR, Noulin F, Famodimu MT, Smargiasso N et al. G-Quadruplex DNA motifs in the malaria parasite *Plasmodium falciparum* and their potential as novel antimalarial drug targets. *Antimicrob Agents Chemother* 2018;62 [Epub ahead of print 23 Oct 2018].
80. Mishra SK, Shankar U, Jain N, Sikri K, Tyagi JS et al. Characterization of G-Quadruplex Motifs in espB, espK, and cyp51 genes of *Mycobacterium tuberculosis* as potential drug targets. *Mol Ther Nucleic Acids* 2019;16:698–706.
81. Ruggiero E, Richter SN. Viral G-quadruplexes: New frontiers in virus pathogenesis and antiviral therapy. *Annu Rep Med Chem* 2020;54:101–131.
82. da Silva AR, de Andrade Neto JB, da Silva CR, Campos RdeS, Costa Silva RA et al. Berberine antifungal activity in fluconazole-resistant pathogenic yeasts: action mechanism evaluated by flow cytometry and biofilm growth inhibition in *Candida* spp. *Antimicrob Agents Chemother* 2016;60:3551–3557.

83. Lestrade PPA, Meis JF, Melchers WJG, Verweij PE. Triazole resistance in *Aspergillus fumigatus*: recent insights and challenges for patient management. *Clin Microbiol Infect* 2019;25:799–806.
84. Gonzalez-Jimenez I, Lucio J, Amich J, Cuesta I, Sanchez Arroyo R et al. A Cyp51B mutation contributes to azole resistance in *Aspergillus fumigatus*. *J Fungi* 2020;6:e315 [Epub ahead of print 26 Nov 2020].
85. Bok JW, Balajee SA, Marr KA, Andes D, Nielsen KF et al. LaeA, a regulator of morphogenetic fungal virulence factors. *Eukaryot Cell* 2005;4:1574–1582.
86. Bok JW, Keller NP. LaeA, a regulator of secondary metabolism in *Aspergillus* spp. *Eukaryot Cell* 2004;3:527–535.
87. Bultman KM, Kowalski CH, Cramer RA. *Aspergillus fumigatus* virulence through the lens of transcription factors. *Med Mycol* 2017;55:24–38.
88. Warner EF. Cross Kingdom analysis of putative quadruplex-forming sequences in fungal genomes: novel antifungal targets to ameliorate fungal pathogenicity? *bioRxiv* 2020;p. 2020.09.23.310581.

Five reasons to publish your next article with a Microbiology Society journal

1. The Microbiology Society is a not-for-profit organization.
2. We offer fast and rigorous peer review – average time to first decision is 4–6 weeks.
3. Our journals have a global readership with subscriptions held in research institutions around the world.
4. 80% of our authors rate our submission process as 'excellent' or 'very good'.
5. Your article will be published on an interactive journal platform with advanced metrics.

Find out more and submit your article at microbiologyresearch.org.