

Lipid mediators from pollen act as chemoattractants and activators of polymorphonuclear granulocytes

Claudia Traidl-Hoffmann, MD,^{a*} Anna Kasche, MD,^{a*} Thilo Jakob, MD,^a Michael Huger,^a Sabine Plötz, MD,^a Ivo Feussner, PhD,^c Johannes Ring, MD, PhD,^b and Heidrun Behrendt, MD^a *Neuherberg, Munich, and Gatersleben, Germany*

Background: Under natural exposure conditions, pollen grains function as allergen carriers that release allergens from internal binding sites on contact with the aqueous phase of mucosa membranes. In addition, we recently demonstrated that pollen are a rich source of eicosanoid-like mediators, which are rapidly released on contact with the aqueous phase.

Objective: The current study was designed to characterize the biochemical nature of pollen-derived lipid mediators in more detail and to delineate their biologic activity on polymorphonuclear granulocytes (PMNs).

Methods: Aqueous and lipid extracts from *Phleum pratense* L and *Betula alba* L pollen were analyzed by means of HPLC. PMNs were exposed to aqueous extracts or lipid fractions from pollen or to HPLC-purified lipid mediators identified in pollen extracts. Effects on PMNs were tested with transwell migration, calcium mobilization, and surface expression of CD11b.

Results: Aqueous pollen extracts (APEs) contained predominantly monohydroxylated products derived of linoleic acid and linolenic acid. In chemotaxis assays PMNs displayed significant migration to APEs. Lipid extracts from pollen and the HPLC fraction containing 13-hydroxy-octadecadienoic acid/hydroxy-linoleic acid and 13-hydroxy-octadecatrienoic acid/hydroxy-linolenic acid induced migratory responses, although to a lesser degree than the APEs. In addition, APE, as well as lipid, extracts induced PMN activation, as documented by means of calcium mobilization and upregulation of CD11b. **Conclusion:** Pollen grains release mediators that recruit and activate PMNs in vitro. Similar mechanisms may be effective in vivo, suggesting that pollen-derived lipid mediators may act as adjuvants in the elicitation phase of allergic reactions.

Key words: Chemotaxis, polymorphonuclear granulocyte, pollen, allergy, oxylipins

Abbreviations used

APE: Aqueous pollen extract
HIP: Hexan-isopropanol
HODE: Hydroxy-octadecadienoic acid/hydroxy-linoleic acid
HOTE: Hydroxy-octadecatrienoic acid/hydroxy-linolenic acid
LT: Leukotriene
PMN: Polymorphonuclear granulocyte
RP: Reverse phase

The importance of grass pollen as a potent allergen source was established almost 130 years ago, when Blackley¹ performed provocation tests and documented the relationship between pollen exposure and allergic symptoms. Apart from pollen as an allergen carrier,² we recently demonstrated that pollen grains, on contact with the aqueous phase, also liberate eicosanoid-like substances, as determined by means of cross-reactivity in an ELISA for leukotriene (LT) B₄ and prostaglandin E₂.³ Prostaglandins and LTs are metabolites of arachidonic acid, which is not present in birch and grass pollen.⁴ Other unsaturated fatty acids, such as linoleic acid and linolenic acid, are major components of membrane fatty acids in plants.^{4,5} Recently, oxidized derivatives of linoleic acid have been identified in a variety of cells, and these metabolites have been shown to exhibit a number of biologic functions, including modulation of cell proliferation, apoptosis, and inflammation.⁶⁻⁸

Even though the precise role of polymorphonuclear granulocytes (PMNs) in the pathophysiology of allergic diseases is still a matter of debate, there is a general consensus that PMNs contribute to the manifestation of allergic inflammation. As cells of the innate immune system, PMNs play an important role in a number of allergic diseases.⁹⁻¹¹ In asthma PMNs are among the first cell type to enter the lung after allergen challenge,¹² and in allergic rhinitis local inflammation is associated with an accumulation of PMNs in the nasal tissue.^{13,14}

The recruitment of PMNs into tissues is a multistep process that involves selectin-mediated tethering, integrin-mediated adhesion, transendothelial migration, and directed migration along chemotactic gradients through the extracellular matrix to the site of inflammation.¹⁵ Directed migration of PMNs is regulated by a number of chemoattractants, of which there are 2 main types: chemokines and lipid mediators. LTB₄, a metabolite of arachidonic acid, is

From ^athe Division of Environmental Dermatology and Allergy TUM/GSF, Neuherberg; ^bthe Department of Dermatology and Allergy, Technical University, Munich; and ^cthe Department of Molecular Cell Biology, Institute of Plant Genetics and Crop Plant Research, Gatersleben.

*Drs Traidl-Hoffmann and Kasche contributed equally to this work.

This study was supported by a grant from the Bavarian Ministry of Environment (BSt MLU) and a grant from the Federal Ministry of Research and Education (BMBF) to C.T.-H. and T.J. The work in the laboratory of I.F. was funded by the Deutsche Forschungsgemeinschaft (SFB 363/B23).

Received for publication November 2, 2001; revised March 1, 2002; accepted for publication March 6, 2002.

Reprint requests: Claudia Traidl-Hoffmann, MD, Division of Environmental Dermatology and Allergy TUM/GSF, Biedersteinerstr. 29, 80802 Munich, Germany.

the most potent chemoattractant lipid mediator for PMNs described thus far,^{16,17} binding to the low-affinity (BLT2) and high-affinity (BLT1) receptor.^{18,19}

The present study was designed to characterize the biochemical nature of mediators released from pollen grains and to delineate their biologic activity on PMNs. We demonstrate for the first time that aqueous pollen extracts (APEs) induce a significant migration and activation of PMNs. The observed effects were independent of the donor sensitization status (ie, could be induced in PMNs from allergic, as well as nonallergic, individuals) and could not be triggered by means of stimulation with allergen alone. In addition, we provide evidence that pollen-derived lipid mediators, particularly monohydroxylated linoleic acid derivatives, contribute to the chemotactic and stimulating activity of APEs on PMNs. We hypothesize that the rapid release of bioactive mediators from pollen during contact with the aqueous phase of mucosal or respiratory membranes may act as allergen-independent proinflammatory factors contributing to the manifestation or aggravation of allergic inflammation.

METHODS

Donors

Patients included in the study had a history of allergic rhinitis, with a positive skin prick test response for grass (n = 3) or birch pollen (n = 9) and with RAST results from the respective allergen of greater than class 3. Healthy control subjects (n = 11) had no history of allergic diseases. In addition, RAST and skin prick test results for 15 common allergens, including birch and grass pollen, were negative, and total IgE was less than 20 IU/mL. All volunteers were without medication for at least 15 days before blood sampling. The ethical committee of the Technical University of Munich approved the study, and volunteers were enrolled in the study after providing written informed consent.

Antibodies and reagents

The mAb anti-CD11b/Mac-1 (clone ICRF44) and the corresponding FITC-labeled isotype control were purchased from Becton Dickinson (San Jose, Calif). LTB₄ was provided by Cayman Chemicals (Ann Arbor, Mich). The selective LTB₄ receptor antagonist LY293111 was kindly provided by Lilly Pharma (Greenfield, Calif).

Pollen grains and generation of APEs

The pollen grains studied (birch, *Betula alba* L; grass, *Phleum pratense* L) were purchased from commercial producers (Allergon, Ängelholm, Sweden) and maintained under dry conditions at 4°C until use. APEs were prepared from incubations of 30 mg/mL pollen for 30 minutes at 37°C in RPMI-1640 medium by means of centrifugation (500g) and sterile filtration (0.2 µm). This concentration was chosen because APEs of 34 mg/mL resulted, in the LTB₄ ELISA, in an average concentration of 3.9×10^{-10} mol/L LTB₄, which is known to be the concentration to induce migration in PMNs.

Hexan-isopropanol extraction of pollen and HPLC analysis

Oxidized fatty acids (oxylipins) were extracted according to the method of Weichert et al.²⁰ In case of the structural analysis of oxylipins, 0.5 g of pollen and, for preparative isolation of oxylipins,

2 g of pollen were added to 30 mL of extraction medium (isohexane-isopropanol 3:2 by volume with 0.0025% wt/vol butyl hydroxy toluol), and 150 µL of 1 mol/L HCl was added and immediately homogenized with an Ultra Turrax for 30 seconds. The extract was centrifuged for 10 minutes at 4500g at 4°C. The supernatant was collected, and the pellet was extracted an additional 3 times with 3 mL of extraction medium. To the combined organic phases, a 6.7% (wt/vol) solution of potassium sulfate was added to a total volume of 47 mL, and the upper hexane-rich layer was removed. The upper organic phase containing the oxidized fatty acid derivatives was dried, redissolved in 1 mL of methanol/water/acetic acid (85:15:0.1 by volume), and stored in argon at -80°C until analysis.

HPLC analysis was carried out on an Agilent 1100 HPLC system coupled to a diode array detector. At first, oxylipins were purified on reversed-phase (RP) HPLC. This was carried out either on an ET250/2 Nucleosil 120-5 C18 column (Macherey-Nagel, 2.1 × 250 mm, 5-µm particle size) for analytic purposes or on an ET250/2 Nucleosil 250-5 C18 column (Macherey-Nagel, 4.6 × 250 mm, 5-µm particle size) for preparative purposes with a solvent system of methanol/water/acetic acid (85:15:0.1) and a flow rate of 0.18 mL · min⁻¹. Straight-phase HPLC of the hydro(pero)xy fatty acids was carried out on a Zorbax Rx-SIL column (Agilent, 150 × 2.1 mm, 5-µm particle size), with a solvent system of *n*-hexane/isopropanol/acetic acid (100:1:0.1) and a flow rate of 0.1 mL · min⁻¹. Chiral-phase HPLC of the hydro(pero)xy fatty acids was carried out on a Chiralcel OD-H column (Daicel, 150 × 2.1 mm, 5-µm particle size) with a solvent system of *n*-hexane/isopropanol/acetic acid (100:5:0.1) and a flow rate of 0.1 mL · min⁻¹. Absorbance of conjugated diene systems at 234 nm was monitored. As an internal standard, (13S,6Z,9Z,11E)-13-hydroxy-6,9,11-octadecatrienoic acid was used.

Preparation of polymorphonuclear granulocytes

Human polymorphonuclear granulocytes were obtained from atopic and nonatopic volunteers by means of density gradient centrifugation of total blood (Histopaque 1077/1119; Sigma, Deisenhofen, Germany). Polymorphonuclear leukocytes were resuspended at 1×10^6 cells/mL in RPMI-1640 supplemented with 2 mmol/L glutamine, 1 mmol/L sodium pyruvate, and 1% nonessential amino acids (complete RPMI; Life Technologies, Chagrin Falls, Ohio) and kept at room temperature. As analyzed by means of flow cytometry, these cells were 96% pure neutrophils with a vitality of greater than 95% (data not shown). Eosinophil contamination was 2% or less (data not shown).

Migration assay

The chemotactic activity of the substances was evaluated, measuring the neutrophil migration through 5-µm pore polycarbonate filter in 24-well transwell chambers (Corning Costar, Cambridge, Mass), as previously described.²¹ Briefly, PMNs were added to the top chamber and suspended in complete RPMI plus 0.5% BSA at 1×10^6 cells/mL. Various concentrations of the different chemokines were added to the bottom chamber of the transwell (0.6 mL). After 1 hour of incubation at 37°C with 5% CO₂, cells that had transmigrated into the lower chamber were recovered and counted with a FACScan (Becton Dickinson, Heidelberg, Germany) for 60 seconds at a flow rate of 60 µL/min. Results are shown as the migration index, which represents the ratio between cells migrated to the lower chamber in the presence of the chemotactic substance and cells migrated in response to the medium alone. When indicated, cells were preincubated with a selective LTB₄ receptor antagonist (LY293111; 40, 400, and 4000 nmol/L), which was kindly provided by Lilly Pharma (Greenfield, Ind) and immediately used for the migration assay.

Checkerboard analysis

Standard checkerboard analysis was performed to analyze chemotactic versus chemokinetic activity of APEs. Briefly, various concentrations of APEs were placed in upper wells, lower wells, or both upper and lower wells of the chemotaxis chamber to determine whether the number of migrated cells was greater with a positive gradient, no gradient, or a negative gradient of APEs.

Measurement of Ca²⁺

PMNs isolated as described above were adjusted to 5×10^6 /mL and loaded with $2 \mu\text{mol/L}$ fura2/AM (Molecular Probes, Eugene, Ore), according to the instructions of the manufacturer. The calcium flux monitoring was performed on PMNs that were stimulated with APEs (30 mg of pollen/mL), LTB₄ (3.9×10^{-10} mol/L), and lipid mediators from pollen (hexan-isopropanol [HIP] extracts and RP-phase extracts). Furthermore, desensitization experiments were performed by using LTB₄, followed by pollen extracts or vice versa.

Measurement of surface expression of CD11b

PMNs were isolated as indicated above and stimulated with the different stimuli as indicated. Staining with the mAbs for CD11b (Clone ICRF44) was performed, as indicated by the manufacturer and analyzed by means of flow cytometry.

Statistical analysis

The Student paired *t* test was used to compare groups of differently stimulated cells (checkerboard analysis). The unpaired *t* test was used to compare migratory behavior of PMNs of allergic and nonallergic patients. A *P* value of less than .05 was considered significant.

RESULTS

HPLC analysis of pollen extracts

Because we could show that pollens harbor eicosanoid-like substances, we first analyzed total lipid extracts from pollen for their amount of hydro(peroxy) polyenoic fatty acids by means of HPLC analysis (Fig 1). Birch pollen contained substantial amounts of nonesterified linoleic acid (288 nmol/g fresh weight) and linolenic acid (295 nmol/g fresh weight), as quantified with RP-HPLC analysis. Moreover, we observed a strong preponderance of linoleic acid-derived hydroxy fatty acid derivatives (13-hydroxy-octadecadienoic acid [13-HODE], 9.1 nmol/g fresh weight; 9-HODE, 6.3 nmol/g fresh weight; Fig 1, A). In contrast, all hydroxy derivatives derived from linolenic acid were less prevalent (1 and 1.6 nmol/g). For grass pollen, we found a different situation (Fig 1, B). Again, we detected substantial amounts of nonesterified linoleic acid (821 nmol/g fresh weight) and linolenic acid (3.8 $\mu\text{mol/g}$ fresh weight). In addition, the amount of hydroxy fatty acids was equally distributed between linoleic acid and linolenic acid derivatives, ranging between 20 and 33 nmol/g fresh weight (Fig 1, B). Also depicted in Fig 1 (inserts) is the separation of all stereochemical isomers of hydroxy derivatives, either of linoleic acid or of linolenic acid. As described recently, the analysis, either by means of the quantitative distribution of positional isomers (Fig 1, A and B and inserts) of the hydroxy derivatives of linolenic acid or the stereochemistry of the

hydroxy fatty acids formed from all polyenoic fatty acids, are suitable measures to determine whether they are formed by autoxidation or whether they are formed enzymatically (eg, by lipoxygenases).²² In all cases we found only products derived from autoxidation.

Chemotaxis of PMNs to APEs

Previous results of our group revealed that pollen releases substances that cross-react with antibodies against LTB₄. We were interested whether those LTB₄-like substances would also have LTB₄-like effects, such as induction of chemotaxis. The concentrations of pollen extracts were chosen at a range of 0.034 to 34 mg pollen/mL because 34 mg/mL pollen elicited, in the LTB₄-ELISA, an equivalent concentration of 3.9×10^{-10} mol/L LTB₄, which is known to induce significant migration in PMNs. Interestingly, PMNs revealed an extraordinary capacity to migrate in response to APEs from birch and grass pollen independent from the allergic status of the patient (*P* = .87), pointing to chemotactic mediators released from pollen (Fig 2, A). Birch and grass pollen extracts at 34 mg/mL exhibited up to 80% of the chemotactic activity of LTB₄ (3.9×10^{-10} mol/L), suggesting a comparable potency of these LTB₄-like substances in equivalent concentrations. The LTB₄ receptor was involved in chemotaxis because LY293111, an LTB₄ receptor antagonist,^{23,24} efficiently blocked migration of PMNs (Fig 2, B). At 400 nmol/L LY293111, the migration toward LTB₄ was blocked up to 90%, whereas migration to IL-8 was not affected (Fig 2, B). Migration of APEs was blocked up to 70%, indicating that more than one signal pathway is involved in the chemotaxis induced by APEs.

Checkerboard analysis

The results of checkerboard analysis to grass pollen performed in quadruplicate are shown in Table I. Similar results were observed to birch pollen extracts (data not shown). The values along the diagonal, reflecting migratory responses to uniform concentrations of APEs on both sides of the chamber (chemokinesis), are always significantly lower than the values below the diagonal, reflecting responses to a positive gradient (chemotaxis; paired *t* test, *P* < .05). The values above the diagonal represent responses to a negative gradient (chemokinesis) and are also always significantly lower than the values under the diagonal.

Chemotaxis of PMNs in response to lipid mediators from pollen

To further characterize the substances responsible for the migratory behavior of PMNs, pollens were submitted to the HIP procedure for extraction of all lipophilic substances. Taking those HIP extracts that were quantified by means of HPLC for their amount in 13-HODE, we found a significant migration in a concentration range of 1×10^{-10} to 1×10^{-15} mol/L 13-HODE, with a maximum at the concentration of 1×10^{-15} mol/L (Fig 2, C), suggesting that lipid mediators from pollen were contribut-




FIG 1. Analysis of polyenoic fatty acid-derived oxylipin formation in pollen lipid extracts. Lipids were extracted, and free oxylipins were isolated by means of RP-HPLC. The regioisomers of hydroxy fatty acids were separated by means of straight-phase HPLC, and the enantiomers were analyzed by means of chiral-phase HPLC. The estimated ratio of the stereoisomers (S/R ratio) of HODEs and HOTEs are shown in the insets: **A**, birch; **B**, grass.

ing to the chemotactic activity of APEs. Taking the fraction from the RP-phase containing primarily 13-HODE and 13-hydroxy-octadecatrienoic acid/hydroxy-linolenic acid (13-HOTE; quantified for the concentration of 13-HODE), we still found chemotaxis of PMNs, even though migratory potency of those extracts was much lower compared with that of APEs (Fig 2, *D*). Furthermore, much higher concentrations (1×10^{-11} mol/L 13-HODE equivalent) were necessary to induce a significant migration. This indicates that the reduction of the chemotactic activity was due to lacking summative effects of multiple activating and chemotactic substances released from pollen.

Pollen extract signaling in PMNs

To address the question of whether pollen-derived mediators participate in receptor-mediated signaling, we measured calcium mobilization in PMNs. Interestingly, they induced a significant calcium release in PMNs, indicating that the induction of chemotaxis was receptor mediated. APEs from grass and birch pollen at 30 mg/mL were approximately equipotent to LTB_4 (3.9×10^{-10}) in mobilizing cytosolic calcium in PMNs (Fig 3, *A-C*). Moreover, HIP extracts, as well as RP-phase (1×10^{-7} mol/L 13-HODE equivalent) substances, were also able to induce calcium flux in PMNs (Fig 3, *F-K*).




FIG 2. Migratory response of PMN to APEs, lipid mediators from pollen, and LTB₄. **A**, Dose-dependent migration of PMNs to APEs from birch and grass pollen in comparison with migration induced by LTB₄. Data are presented as pooled migration indices from 3 donors with grass allergy, 9 donors with birch allergy, and 11 nonallergic donors (mean \pm SEM). **B**, Influence of the LTB₄ receptor antagonist LY293111 on PMN migration. For APEs, one representative experiment of 6 (3 allergic and 3 nonallergic donors) by using PMNs from one allergic donor is shown (mean and range of duplicates). For LTB₄ and IL-8, one representative experiment of 3 (3 allergic donors) is shown. **C** and **D**, PMN migration to lipid extracts of pollen. The mean \pm SEM of pooled data from 9 donors (3 nonallergic donors, 3 donors with birch allergy, and 3 donors with grass allergy) is shown.

TABLE I. Checkerboard analysis for APEs from grass pollen

	Upper chamber (mg/mL)				
	0	0.034	0.34	3.4	34
Lower chamber (mg/mL)					
0	1 \pm 0.01	1.4 \pm 0.2	0.8 \pm 0.01	0.9 \pm 0.1	0.7 \pm 0.05
0.034	1.1 \pm 0.02	1.5 \pm 0.1	1.3 \pm 0.2	1.2 \pm 0.1	0.8 \pm 0.01
0.34	1.2 \pm 0.02	1.1 \pm 0.1	1.3 \pm 0.2	1.2 \pm 0.2	0.6 \pm 0.1
3.4	2.1 \pm 0.1	2.0 \pm 0.2	1.8 \pm 0.2	1.8 \pm 0.01	1.2 \pm 0.01
34	5.6 \pm 0.3	4.6 \pm 0.2	5.6 \pm 0.1	5.0 \pm 0.1	2.7 \pm 0.1

Values represent means \pm SD of calculated chemotactic indices from one experiment performed in quadruplicate by using the transwell migration assay, as indicated in the "Methods" section.

Because an LTB₄ receptor antagonist could inhibit the migration to pollen extracts, we performed cross-desensitization studies stimulating first with LTB₄ followed by pollen extracts and vice versa. Interestingly, the responsiveness of PMNs to LTB₄ was lost on prior incubation with APEs from grass and birch pollen (Fig 3, *B* and *C*). Similarly, APEs induced only weak calcium fluxes in PMNs that had previously been exposed to LTB₄ (Fig 3,

D and *E*). These results strongly suggest that LTB₄ receptor agonists released by pollen contribute to trigger functional responses in PMNs.

APEs upregulate CD11b on PMNs

As a well-established parameter of PMN activation, we analyzed CD11b upregulation after stimulation with aqueous and lipid pollen extracts in comparison with




FIG 3. Intracellular calcium flux induced in PMNs by pollen extracts and lipid mediators. **A**, LTB_4 (3.9×10^{-8} mol/L). **B** and **C**, Aqueous extracts (AE) from grass pollen and birch pollen, respectively. **F** through **I**, Calcium mobilization in PMNs by lipid extracts from pollen: HIP extracts from birch pollen (**F**); RP-phase from birch pollen (**G**); HIP extracts from grass pollen (**H**); and RP-phase from grass pollen (**I**). For cross-sensitization experiments, PMNs were first stimulated with APEs of pollen, followed by LTB_4 (**B** and **C**) or vice versa (**D** and **E**). Shown is 1 representative experiment (donor with grass allergy) of 6 performed with PMNs from different donors: 2 from donors with birch allergy, 2 from donors with grass allergy, and 2 from nonallergic donors.

LTB_4 . Freshly isolated PMNs were resuspended in RPMI complete plus 10% FBS, and stimulators were added in the concentrations indicated. Flow cytometry was used to assess CD11b expression from 5 to 60 minutes and 2 and 12 hours after stimulation. In Fig 4 the expression after 1 hour of stimulation is shown.

APEs of birch pollen were slightly but not significantly more potent compared with those in grass pollen in inducing CD11b (Fig 4, **B** and **C**). APEs and LTB_4 (3.9×10^{-10} mol/L) were comparably potent in inducing CD11b expression. Even after 12 hours, a high expression of CD11b, usually only transiently expressed, as described by Davey et al,²⁵ was observed after stimulation with APEs (data not

shown). HIP extracts and the RP-phase from both grass (Fig 4, **E** and **F**) and birch pollen (data not shown) showed upregulation of CD11b, pointing to the importance of the lipid ingredients in pollen for the observed effects. The allergen rPhl p 5, itself in concentrations comparable with the concentration in aqueous extracts,³ did not induce any upregulation of CD11b (Fig 4, **D**).

DISCUSSION

Our recent data point to pollen as a rich source of bioactive mediators, which are rapidly released on contact with an aqueous phase.³ Here we show that mediators released by pollen (birch and grass) activate and recruit PMNs *in vitro*. Pollen-derived neutrophil chemotactic activity may be attributable to one or more lipids that are functionally and structurally closely related to LTs. Pollen extracts contained substantial amounts of linolenic acid-derived hydroxy fatty acid derivatives, namely 13-HODE and 13-HOTE, which lead to activation and migration of PMNs.

Neutrophil extravasation into the interstitium at the site of inflammation is a prerequisite for PMN defense function. This is initiated in response to lipid mediators and chemokines by means of selectin-dependent PMN rolling along the endothelial lining, followed by firm PMN adhesion and transendothelial diapedesis.²⁶ Our finding of LTB_4 -like substances in APEs prompted us to further investigate the effect of those substances on PMNs.

LTB_4 activates and recruits granulocytes into inflamed lesions.^{17,19} Furthermore, data have accumulated on the contribution of LTB_4 in inflammatory disorders,^{27,28} including bronchial asthma and allergic diseases.^{29,30} Mice lacking LT production are deficient in their response to some acute and chronic inflammatory stimuli.^{31,32} Therefore LTB_4 receptor antagonists are under development as potent anti-inflammatory and antiallergic drugs.^{23,33}

Leukotactic activity of pollen-derived (birch and grass) neutrophil chemotaxins was found to be of similar efficacy as that of well-characterized chemotaxins, such as LTB_4 and IL-8 (Fig 2). All migrating cells can be activated by certain inflammatory mediators without the induction of directional migration but increase the random movement of cells as a result of cell activation. Cells exposed to such primers typically show signs of enhanced oxidative metabolism and upregulate the expression of receptors, so that the response to bona fide chemotactic factors and the interaction with adhesion molecules on endothelial cells are enhanced.³⁴ In addition to chemotactic responses, APEs also provoke chemokinetic PMN responses (Table I). Although the chemokinetic index is significantly lower than that of chemotaxis and occurs only at high concentrations (30 mg/mL), optimal stimulation for both chemotaxis and chemokinesis occurs at the same concentration of APEs known also for other chemokines.³⁵

APEs contain considerable amounts of minor and major allergens.³ Thus it seemed possible that the




FIG 4. Upregulation of CD11b on PMNs by APEs and lipid mediators from pollen. **A-F**, Histograms of CD11b expression 1 hour after stimulation with 3.9×10^{-9} mol/L LTB₄ (**A**), APE 34 mg of grass pollen/mL (**B**), APE 34 mg of birch pollen/mL (**C**), 50 μ g/mL rPhl p 5 (**D**), HIP birch pollen extracts (**E**), and RP-phase from birch pollen (**F**). Numbers indicate net mean fluorescence intensity: left, unstimulated control; right, stimulated probe. Dotted lines indicate unstimulated control, and solid lines indicate stimulated probes. The experiment stands for 6 similar experiments performed from PMNs of 2 donors with grass allergy, 2 donors with birch allergy, and 2 nonallergic donors.

observed effects were induced by allergen-specific mechanisms. Aside from low- and high-affinity receptors for IgG, it was reported recently that PMNs express different receptors for IgE: the heterotrimeric high-affinity receptor for IgE (Fc ϵ RI)³⁶⁻³⁸; the low-affinity receptor for IgE (Fc ϵ RII/CD23)³⁹; and the Mac-2/IgE-binding protein.⁴⁰ However, the occurrence of chemotaxis in both allergic and nonallergic individuals makes a specific IgE-mediated mechanism unlikely. Additionally, we performed migration assays with recombinant Phl p 5, the major grass pollen allergen from *Phleum pratense* L, in concentrations comparable with those found in APEs.³ Here neither PMN chemotaxis nor chemokinesis could be observed, suggesting that the APE-induced migration was indeed mediated through allergen-independent mechanisms. We could also demonstrate that the migration to APEs could be inhibited by an LTB₄-receptor antagonist (LY293111). This suggests that the LTB₄ receptor is involved in the induction of migration. Recent data on hydroxyeicosanoids binding to the low-affinity LTB₄ receptor (BLT2)⁴¹ demonstrated that this receptor is rather promiscuous, binding to a broader spectrum of ligands similar to LTB₄.

The concept of LTB₄ receptor involvement in the induction of migration was supported by cross-desensitization experiments. The responsiveness of PMNs to LTB₄ by means of calcium mobilization was lost after prior desensitization with APEs. This indicates that the LTB₄ receptor was already engaged after stimulation with pollen extracts.

Full activation of PMNs is dependent on the expression of adhesion receptors of the β 2-integrin family, which is known to be critically involved in the interaction with the endothelium during cell recruitment into inflamed tissue. The upregulation of a β 2-integrin by

APEs and lipid components of pollen indicates that these substances not only act as chemoattractants but also as agonists of PMN activation.

In conclusion, we provide evidence that pollen grains release mediators that recruit and activate PMNs in vitro. It is tempting to speculate that similar mechanisms may be effective in vivo. Our observations suggest that rapid release of bioactive lipid mediators from pollen during contact with the aqueous phase of mucosal or respiratory membranes may act as allergen-independent proinflammatory factors, contributing to the manifestation or aggravation of allergic inflammation.

We thank Volker Kasche for scientific support and Gabriele Pleyl-Wisgickl, Ellis Fraust, and Johanna Grosch for excellent technical assistance.

REFERENCES

1. Blackley CH. Experimental researches on the causes and nature of Catarrhus aestivus (hayfever or hay-asthma). London: Dawson's of Pall Mall; 1959.
2. Behrendt H, Becker WM. Localization, release and bioavailability of pollen allergens: the influence of environmental factors. *Curr Opin Immunol* 2001;13:709-15.
3. Behrendt H, Kasche A, Ebner von Eschenbach C, Risse U, Huss-Marp J, Ring J. Secretion of proinflammatory eicosanoid-like substances precedes allergen release from pollen grains in the initiation of allergic sensitization. *Int Arch Allergy Immunol* 2001;124:121-5.
4. Müller MJ. Radically novel prostaglandins in animals and plants: the isoprostanes. *Chem Biol* 1998;12:R323-33.
5. Sobermann RJ, Harper TW, Betteridge D, Lewis RA, Austen KF. Characterization and separation of the arachidonic acid 5-lipoxygenase and linoleic acid omega-6 lipoxygenase (arachidonic acid 15-lipoxygenase) of human polymorphonuclear leukocytes. *J Biol Chem* 1985;260:4508-15.
6. Henricks PA, Engels J, Van der Linde HJ, Nifkamp FP. 13-hydroxylinoleic acid induces airway hyperreactivity to histamine in guinea-pigs. *Eur J Pharmacol* 1991;197:233-4.
7. Liu XH, Connolly JM, Rose DP. The 12-lipoxygenase gene-transfected MCF-7 human breast cancer cell line exhibits estrogen-independent, but

- estrogen and omega-6 fatty acid-stimulated proliferation in vitro, and enhanced growth in athymic nude mice. *Cancer Lett* 1996;109:233-40.
8. Toborek M, Blanc EM, Kaise S, Mattson MP, Hennig B. Linoleic acid potentiates TNF-mediated oxidative stress, disruption of calcium homeostasis, and apoptosis of cultured vascular endothelial cells. *J Lipid Res* 1997;38:2155-67.
 9. Henson PM, Borish LC. Neutrophil mediators in asthma. In: Busse WW, Holgate ST, editors. *Asthma and rhinitis*. Boston: Blackwell Scientific Publications; 1995. p. 367-82.
 10. Lim MC, Taylor RM, Naclerio RM. The histology of allergic rhinitis and its comparison to cellular changes in nasal lavage. *Am J Respir Crit Care Med* 1995;151:136-44.
 11. Varga EM, Jacobson MR, Masuyama K, Rak S, Till SJ, Darby Y, et al. Inflammatory cell populations and cytokine mRNA expression in the nasal mucosa in aspirin-sensitive rhinitis. *Eur Respir J* 1999;14:610-5.
 12. Teran LM, Carroll M, Frew AJ, Montefort S, Lau LC, Davies DE, et al. Neutrophil influx and interleukin-8 release after segmental allergen or saline challenge in asthmatics. *Int Arch Allergy Immunol* 1995;107:374-5.
 13. Casale TB, Costa JJ, Galli SJ. TNF alpha is important in human lung allergic reactions. *Am J Respir Cell Biol* 1997;15:35-44.
 14. Smith HR, Larsen GL, Chermiack RM, Wenzel SE, Voelkel NF, Westcott JY, et al. Inflammatory cells and eicosanoid mediators in subjects with late asthmatic responses and increase in airway responsiveness. *J Allergy Clin Immunol* 1992;89:1076-84.
 15. Wagner JG, Roth RA. Neutrophil migration mechanisms, with an emphasis on the pulmonary vasculature. *Pharmacol Rev* 2000;52:349-74.
 16. Samuelsson B, Dahlén SE, Lindgren JÅ, Rouzer CA, Serhan CN. Leukotrienes and lipoxins: structures, biosynthesis, and biological effects. *Science* 1987;237:1171-6.
 17. Ford-Hutchinson AW, Bray MA, Doig MV, Shipley ME, Smith MJ. Leukotriene B₄, a potent chemokinetic and aggregating substance released from polymorphonuclear leukocytes. *Nature* 1980;286:264-5.
 18. Yokomizo T, Kato K, Terawaki K, Izumi T, Shimizu T. A second leukotriene B₄ receptor, BLT₂. A new therapeutic target in inflammation and immunological disorders. *J Exp Med* 2000;192:421-32.
 19. Yokomizo T, Izumi T, Chang K, Takuwa Y, Shimizu T. A G-protein-coupled receptor for leukotriene B₄ that mediates chemotaxis. *Nature* 1997;387:620-4.
 20. Weichert H, Stenzel I, Berndt E, Wasternack C, Feussner I. Metabolic profiling of oxylipins upon salicylate treatment in barley leaves—preferential induction of the reductase pathway by salicylate. *FEBS Lett* 1999;464:133-7.
 21. Sebastiani S, Allavena P, Albanesi C, Nasorri F, Gincarlo B, Traidl C, et al. Chemokine receptor expression and function in CD4⁺ T lymphocytes with regulatory activity. *J Immunol* 2000;166:996-1002.
 22. Berger S, Weichert H, Porzel A, Wasternack C, Kühn H, Feussner I. Enzymatic and non-enzymatic lipid peroxidation in leaf development. *Biochim Biophys Acta* 2001;1533:266-76.
 23. Jackson WT, Froelich LL, Boyd RJ, Schrementi JP, Saussy DL, Schultz RM, et al. Pharmacologic actions of the second-generation leukotriene B₄ receptor antagonist LY293111: in vitro studies. *J Pharmacol Exp Ther* 1999;288:286-94.
 24. Sofia MJ, Floreancig P, Bach N, Baker SR, Nelson K, Sawyer JS, et al. The discovery of LY293111, a novel, potent and orally active leukotriene B₄ receptor antagonist of the biphenylphenol class. *Adv Exp Med Biol* 1997;400A:381-6.
 25. Davey PC, Zuzel M, Kamiguti AS, Hunt JA, Aziz KA. Activation-dependent proteolytic degradation of polymorphonuclear CD11b. *Br J Haematol* 2000;111:934-42.
 26. Carlos TM, Harlan JM. Leukocyte-endothelial adhesion molecules. *Blood* 1994;84:2068-101.
 27. Iversen L, Kraghalla K, Ziboh VA. Significance of leukotriene A₄ hydrolyase in the pathogenesis of psoriasis. *Skin Pharmacol* 1997;10:169-77.
 28. Sharon P, Stenson WF. Enhanced synthesis of leukotriene B₄ by colonic mucosa in inflammatory bowel disease. *Gastroenterology* 1984;86:453-60.
 29. Turner CR, Breslow R, Conklyn MJ, Andresen CJ, Patterson DK, Lopez AA, et al. In vitro and in vivo effects of leukotriene B₄ antagonism in a primate model of asthma. *J Clin Invest* 1996;97:381-7.
 30. O'Byrne PM. Why does airway inflammation persist? Is it leukotrienes? *Am J Respir Crit Care Med* 2000;161:S186-7.
 31. Byrum RS, Goulet JL, Griffiths RJ, Koller BH. Role of the 5-lipoxygenase-activating protein (FLAP) in murine acute inflammatory responses. *J Exp Med* 1997;185:1065-75.
 32. Goulet JL, Snouwaert JN, Latour AM, Coffman TM, Koller BH. Altered inflammatory responses in leukotriene-deficient mice. *Proc Natl Acad Sci U S A* 1994;91:12852-6.
 33. Showell HJ, Conklyn MJ, Alperre R, Hingorani GP, Wright KF, Smith MA, et al. The preclinical pharmacological profile of the potent and selective leukotriene B₄ antagonist CP-195543. *J Pharmacol Exp Ther* 1998;285:946-54.
 34. Kaplan AP. Chemokines, chemokine receptors and allergy. *Int Arch Allergy Immunol* 2001;124:423-31.
 35. Schröder JM, Sticherling M, Henneicke HH, Preissner WC, Christophers E. IL1 α or tumor necrosis factor- α stimulate release of three NAP-1/IL-8-related neutrophil chemotactic proteins in human dermal fibroblasts. *J Immunol* 1990;144:2223-32.
 36. Gounni AS, Lamkhioued B, Koussih L, Ra C, Renzi PM, Hamid Q. Human neutrophils express the high-affinity receptor for immunoglobulin E (Fc ϵ RI): role in asthma. *FASEB J* 2001;15:940-9.
 37. Scholl PR, Geha RS. Physical association between the high-affinity IgG receptor (Fc γ RI) and the subunit of the high-affinity IgE receptor (Fc ϵ RI). *Proc Natl Acad Sci U S A* 1993;90:8847-50.
 38. Soussi-Gounni A, Lamkhioued B, Christodoulopoulos P, Nutku E, Hamid Q. Human neutrophils express the high affinity receptor for IgE (Fc ϵ RI). *J Allergy Clin Immunol* 1998;101:S173.
 39. Yamaoka KA, Arock M, Issaly F, Dugas N, Le Goff L, Kolb JP. Granulocyte macrophage colony stimulating factor induces Fc epsilon RII/CD23 expression on normal human polymorphonuclear neutrophils. *Int Immunol* 1996;8:479-90.
 40. Yamaoka A, Kuwabara I, Frigeri LG, Liu FT. A human lectin, galectin-3 (eBP/Mac2), stimulates superoxide production by neutrophils. *J Immunol* 1995;154:3479-87.
 41. Yokomizo T, Kato K, Hagiya H, Izumi T, Shimizu T. Hydroxyeicosanoids bind to and activate the low affinity leukotriene B₄ receptor, BLT₂. *J Biol Chem* 2001;276:12454-9.