

RESEARCH ARTICLE

Psychological trauma occurring during adolescence is associated with an increased risk of greater waist circumference in Early Psychosis patients treated with psychotropic medication


Luis Alameda^{1,2,3,4^{aa}*}, Axel Levier^{5^{ab}}, Mehdi Gholam-Rezaee⁶, Philippe Golay^{2,7,8}, Frederik Vandenberghe⁵, Aurélie Delacretaz⁵, Philipp Baumann⁹, Anaïs Glatard⁵, Céline Dubath⁵, Andres Herane-Vives^{10,11,12}, Victoria Rodriguez¹, Alessandra Solida², Kim Q. Do⁹, Chin B. Eap^{5,13}, Philippe Conus²

1 Department of Psychosis Studies, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, United Kingdom, **2** Service of General Psychiatry, Treatment and Early Intervention in Psychosis, Program [TIPP-Lausanne], Lausanne University Hospital [CHUV], Lausanne, Switzerland, **3** Department of Psychiatry, Hospital Universitario Virgen del Rocío, Universidad de Sevilla, Sevilla, Spain, **4** Instituto de Investigación Sanitaria de Sevilla, IBiS, Sevilla, Spain, **5** Unit of Pharmacogenetics and Clinical Psychopharmacology, Centre for Psychiatric Neuroscience, Department of Psychiatry, Hospital of Cery, Lausanne University Hospital, Prilly, Switzerland, **6** Department of Psychiatry, Center for Psychiatric Epidemiology and Psychopathology, Lausanne University Hospital [CHUV], Lausanne, Switzerland, **7** Community Psychiatry Service, Department of Psychiatry, Consultations de Chauderon, Lausanne University Hospital and University of Lausanne, Lausanne, Switzerland, **8** Institute of Psychology, faculty of Social and Political Science, University of Lausanne, Lausanne, Switzerland, **9** Unit for Research in Schizophrenia, Center for Psychiatric Neuroscience, Department of Psychiatry, Lausanne University Hospital [CHUV], Lausanne, Switzerland, **10** Department of Psychological Medicine, Affective Disorders Research Group, Centre for Affective Disorders, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, United Kingdom, **11** Departamento de Clínicas, Facultad de Medicina, Universidad Católica del Norte, Coquimbo, Chile, **12** Affective Disorders Research Group, Centre for Affective Disorders, Department of Psychological Medicine, Institute of Psychiatry, Psychology and Neuroscience King's College London, London, United Kingdom, **13** Department of Pharmaceutical Sciences, School of Pharmacy, University of Geneva, Geneva, Switzerland

^{aa} Current address: Institute of Psychiatry, Psychology & neuroscience, Camberwell, London, United Kingdom

^{ab} Current address: Centre de Neurosciences psychiatriques, Prilly, Lausanne, Switzerland

* lualisalameda@gmail.com



OPEN ACCESS

Citation: Alameda L, Levier A, Gholam-Rezaee M, Golay P, Vandenberghe F, Delacretaz A, et al. (2020) Psychological trauma occurring during adolescence is associated with an increased risk of greater waist circumference in Early Psychosis patients treated with psychotropic medication. PLoS ONE 15(12): e0242569. <https://doi.org/10.1371/journal.pone.0242569>

Editor: Alejandra Mondragon Maya, National Autonomous University of Mexico, MEXICO

Received: April 15, 2020

Accepted: November 4, 2020

Published: December 3, 2020

Copyright: © 2020 Alameda et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: The informed consent that was used for PsyMetab did not include that the data would be made public, so unfortunately we cannot upload the original data set. Similarly, the consents given by the Ethics Committee of the Canton de Vaud to access TIPP and PsyClin data for research purposes did not include that the data could be made public. However, we can share the data set confidentially individually upon request. The president of the

Abstract

Background

It has been suggested that exposure to Childhood Trauma [CT] may play a role in the risk of obesity in Early Psychosis [EP] patients; however, whether this is independently of age at exposure to CT and the medication profile has yet to be investigated.

Methods

113 EP-patients aged 18–35 were recruited from the Treatment and Early Intervention in Psychosis Program [TIPP-Lausanne]. Body Mass Index [BMI], Weight Gain [WG] and

ethics Committee of the Canton of Vaud, Prof. Dominique Sprumont can be contacted here: dominique.sprumont@vd.ch.

Funding: This work represents an independent research part supported by the The Swiss National Science Foundation (no. 320030_135736/1 to P.C. and K.Q.D., no 320030-120686, 324730-144064 and 320030-173211 to C.B.E and PC and no 171804 to LA), the National Center of Competence in Research (NCCR) "SYNAPSY - The Synaptic Bases of Mental Diseases" from the Swiss National Science Foundation (n° 51NF40-158776) The Damm-Etienne and Alamaya Foundation were supporting the recruitment as well as the phenotyping (including psychopathology, neurocognition and neurobiological biomarkers) of our early psychosis patients cohort). The views expressed are those of the author(s) and not necessarily those of the The Swiss National Science Foundation, the Damm-Etienne or Alamaya Foundations.

Competing interests: The authors have declared that no competing interests exist.

Waist Circumference [WC] were measured prospectively at baseline and after 1, 2, 3, 6 and 12 months of weight gain inducing psychotropic treatment. Patients were classified as Early-Trauma and Late-Trauma if the exposure had occurred before age 12 or between ages 12 and 16 respectively. Generalized Linear Mixed-Models were adjusted for age, sex, socioeconomic status, baseline BMI, medication and for diagnosis of depression.

Results

Late-Trauma patients, when compared to Non-Trauma patients showed greater WCs during the follow-up [$p = 0.013$]. No differences were found in any of the other follow-up measures.

Conclusions

Exposition to CT during adolescence in EP-patients treated with psychotropic medication is associated with greater WC during the early phase of the disease. Further investigation exploring mechanisms underlying the interactions between peripubertal stress, corticoids responsiveness and a subsequent increase of abdominal adiposity is warranted.

Introduction

Patients suffering from psychosis have higher mortality rates [1] and about 18 years shorter life-expectancy when compared to the general population [2]. Risk factors such as obesity, alongside other physical health problems [smoking tobacco, hypertension, type 2 diabetes mellitus, dyslipidemia and cardiovascular disease] play an important role in the excess of mortality in patients suffering from psychosis and other severe mental illness [3–5]. Because psychotropic medications can induce strong weight gain [WG] and other metabolic disorders [6], adapted interventions targeting the well-known iatrogenic consequences of medication and life style have been proposed in the last 15 years, with positive results [7]. However, psychotropic medication is not the only weight gain predictor and some metabolic disturbances are already present in drug naïve patients in the early phase of the psychotic disorder [Early psychosis: EP] [8–10]. Identification of other risk factors could lead to a better understanding of the biological mechanisms underlying obesity in psychotic patients and allow the implementation of new preventive interventions.

Childhood trauma [CT] is a risk factor for psychosis [11], and a meta-analysis of 41 epidemiological studies in the general population revealed that exposure to trauma was associated with an increased risk of developing obesity [odds ratio: 1.36; 95% confidence interval 1.26–1.47] [12]. One study showed that traumatized EP patients had higher BMI when compared to controls and to non-traumatized EP patients [13]. Another study showed a positive association between a number of different forms of abuse and BMI in EP patients [14] and suggests a possible interplay between increased inflammation [CRP], BMI and CT. Furthermore, both inflammation [15, 16] and CT [17] have been associated with higher levels of depressive symptoms in EP patients, and depressive patients [especially those with atypical features] are at higher risk of WG independently of medication [18]. For this reason, it is thus important to take into account depression as a potential confounder when considering the link between trauma and obesity related outcomes. Moreover, different developmental periods are important for processes involved in weight gain such as energy intake, storage and expenditure [12]. Thus, if trauma exposure influences the risk of obesity, taking into account the age at which

the patient is exposed may lead to a better understanding of which developmental trajectories are operating on the link between trauma and obesity related outcomes. Considering that the studies that examined the association between CT and obesity related measures in patients with psychosis were cross sectional, a study addressing this question of the weight gain with a prospective setting is needed. This may allow the investigation into whether metabolic disturbances are already present at the beginning of the psychotropic treatment, or if they develop at a later stage.

Given the above, the present study aimed to examine in a sample of EP patient, the differential impact of trauma occurring during childhood or adolescence on the risk of obesity related outcomes during the first year of psychotropic treatment. We took into account a number of potential covariates and confounders such as a diagnosis of depression, age, sex, socio-economic status [SES], baseline BMI and medication, before and during the follow up.

Methods

Procedure and subjects

TIPP [Treatment and Early Intervention in Psychosis Program], a specialized early psychosis program, was launched in 2004 at the Department of Psychiatry, Lausanne University Hospital Switzerland [19]. Entry criteria to the program are: [I] aged between 18 and 35; [II] residing in the catchment area [Lausanne and surroundings; population about 300 000]; [III] meeting threshold criteria for psychosis, as defined by the “Psychosis threshold” subscale of the Comprehensive Assessment of At Risk Mental States [CAARMS [20]] Scale [19]. In the TIPP program, each patient is followed up by a psychiatrist and a Case manager. The program offers an integrated biopsychosocial treatment based on psychotherapy, psychoeducation, family support, cognitive assessment and remediation [when needed], social support, assistance in finding employment, psychological interventions for cannabis use, and pharmacological treatment. A specially designed questionnaire [21] is completed for all patients enrolled in the programme by case managers who have up to 100 contacts with patients during the 3 years of treatment. It allows assessment of demographic characteristics, past medical history, exposure to life events as well as symptoms and functioning. It is completed on the basis of information gathered with patients and their family over the first few weeks of treatment and can be updated during follow-up if new information emerges. Follow-up assessments exploring various aspects of treatment and co-morbidities as well as evolution of psychopathology and functional level are conducted by a research psychologist and by case managers after 2, 6, 12, 18, 24, 30 and 36 months in treatment. In particular, substance abuse diagnosis was assessed at each time-point and rated on the basis of DSM-IV criteria [22].

Patients starting a pharmacological treatment that is known to have a potential risk to induce metabolic disturbances [i.e. all antipsychotics, some antidepressants and mood stabilizers] are prospectively followed up for metabolic parameters evolution [see Biological and physical measures section below] [21, 23]. Since 2007, a study has been ongoing in the Department of Psychiatry of the Lausanne University Hospital using the routinely collected data to better characterize and understand metabolic adverse effect onset [PsyMetab-PsyClin].

Capacity to consent was determined by the case manager involved in the clinical care of the patient. This ensures that each patient has capacity when they are asked to give consent to participate in the study. If the patient is temporarily unable to give an informed consent, research questions are not presented, until his capacity to consent is recovered. In case patients are evaluated as having a permanent lack of capacity for understanding the research questions, the consent of a legal representative is asked. The Ethics Committee of the Canton of Vaud granted access to TIPP clinical data [demographic, assessment of trauma history and levels of

symptoms and other clinical variables] an also granted access to physical data [obesity-related outcomes] as part of PsyMetab cohort.

Patients included in the PsyMetab cohort were requested to give their written informed consent to participate in the study and to have their data to be used for research studies. In addition, up to end of 2015, clinical data from patients from PsyClin could be used for the current study as part of non-interventional post hoc analysis. Both PsyMetab-PsyClin protocols, as well as this particular study, were approved by the Ethics Committee of the Canton of Vaud [CER-VD].

The present analysis involves participants included in both the TIPP program and who were also part of the PsyMetab-PsyClin study between 2007 and end of 2016.

Assessment of history of past trauma

Clinicians at TIPP are trained to conduct an extensive assessment of patients, including evaluation of exposure to traumatic life events. Case managers meet patients frequently over the treatment period, which provides the framework to establish a trusting relationship, where extensive knowledge of patients' history can be gathered. If patients agree, information can also be completed with family. In the case of inconsistency between the patient's and the family's report or in case of doubt regarding the exposure to trauma or the age at the time of exposure, patients were not included in the study [24]. Case managers complete a table during the patients' 3 years of treatment, where exposure to traumatic life events can be recorded as follows: [1] Type of traumatic life event, rated as present or absent [sexual abuse, physical abuse, emotional and physical neglect, emotional abuse, among others. . .]; [2] time of occurrence in relation to psychosis stage [during the premorbid phase, during the prodrome or after onset of psychosis]; [3] age at the time of first exposure to each one of the traumas that occurred; and [4] single or repeated exposure to each one of the traumas that occurred. Considering that the clinicians who assessed exposure to life events did not rate the subjective perception of severity of the different forms of stressful events, patients were considered traumatized if they had been exposed to at least one experiences of abuse [physical, sexual, or emotional] or neglect [physical or emotional]. The consideration was that such events would undoubtedly be considered as highly traumatizing by anyone, and have been shown to be associated with risk for psychosis and functional deficits in psychotic samples [25]. Sexual abuse refers to sexual molestation and/or rape. Physical abuse refers to physical attack or assault, or being repetitively beaten by parents, relatives, or caregivers. Emotional abuse was defined as verbal assaults on a child's sense of worth or well-being or any humiliating or demeaning behaviour directed toward a child by an adult or older person. Physical neglect was defined as the failure of caretakers to provide for a child's basic physical needs, including food, shelter, clothing, safety, and health care. Emotional neglect was defined as the failure of caretakers to meet children's basic emotional and psychological needs, including love, belonging, nurturance, and support. Age at the time of first exposure was categorized as follows [1] early trauma refers to exposure between birth and age 12, according to conventions applied elsewhere [26–28] late trauma refers to exposure between ages 12 and 16. Patients who were exposed to trauma after age 16 were excluded from this study, according to other studies [17, 24] suggesting that they may already have been in the prodromal phase of their first psychotic episode.

Biological and physical measures

Data of patients in the PsyMetab-PsyClin studies were collected during hospitalization and/or in outpatient centers. Following the introduction of psychotropic treatments that have been linked to weight gain, [weight gain inducing psychotropic medications], a prospective

monitoring [at times 0, 1, 2, 3, 6, 9 and 12 months] of body mass index [BMI], waist circumference [WC] and weight gain [WG] was performed. In the case of an interruption of the above-mentioned weight gain inducing psychotropic medication for more than two weeks or in the case of replacement by another drug, the follow-up is restarted from baseline. In the case of an introduction of a second drug from the list [see list below], the follow-up is restarted. In this study, we considered only the earliest follow-up which has less missing weight information.

Other clinical variables

Main diagnosis and comorbid diagnosis of depression were based on DSM-IV [22] criteria according to others [29]. Medications were categorized into low, medium and high-risk of weight gain as follows [30] [High: Valproate, Olanzapine and Clozapine; Moderate: Quetiapine, Risperidone, Paliperidone, Lithium, Mirtazapine, Zuclopenthixol, Levomepromazine; Low: Amisulpride, Aripiprazole, Haloperidol, Lurasidone, Flupentixol]. In order to adjust by medication intake, we created two variables: [i] Medication during follow-up, which refers to the medication taken during the 1 year of follow-up, and which was categorized into High, Moderate and Low risk; [ii] Medication before the first assessment, which refers to the medication that patients had taken before being included in this study. This variable was categorized as 0 [no psychotropic treatment before the first assessment] and 1 [prescription of at least one of the medications from the list [see above] before the first assessment, with partial or total adherence]. Estimation of the adherence was based on the Treatment Adherence Scale [TAS] [31], which is assessed by the case manager based on results from the plasmatic concentration measured routinely, interviews with patients and an extensive knowledge of the patient treatment and adherence with medication and service. This is evaluated at every TIPP follow-up time points and which is scored on the basis of a 3 point scale [0 = non adherence; 1 = partial adherence, from 25% to 75% of the time during the evaluation period; and 2 = total adherence, from 75% to 100% of the time during the evaluation period]. Patients were considered as adherent to treatment if they scored 2 on the TAS. Socio-economic status [SES] were subdivided into high, intermediate and low based on the socio-economic status of the parents as previously described [32].

Statistical analysis

In order to test the distribution of continuous variables between groups we used the Kruskal Wallis test which is a non-parametric and robust test. This test uses the observed ranks of observations instead of their observed value and is unaffected by outliers if there is any. Chi-squares test of independence was employed to detect associations between categorical variables and the trauma group [trauma], considering a level below $\alpha = 0.05$ as enough evidence to reject the independence between categorical variables and trauma group. [If the test is rejected in the level of $\alpha = 0.05$ then we have enough evidence to reject the independence between the categorical variable and the group.] This test is valid unless the expected frequency is smaller than 5 in at least 20% of cells, when the Fisher Exact Test should be used instead. Generalized Linear Mixed Models [GLMMs] were used to measure the effect of trauma on the response variables, adjusting the models for potential covariates and confounders [diagnosis of depression, age, sex, socio-economic status [SES], baseline BMI and medication before and during the follow up [as detailed in previous section]]. The main advantage of GLMMs is the presence of random effects that allows adjustment of the model for interdependence among observations. For all models, we have introduced a common random intercept for observations corresponding to each individual. A Bayesian estimation approach with flat priors was used to adjust these models, due to its convenience in dealing with models of this complexity.

To do so, we used a large number of Markov Chain Monte-Carlo iterations [10 million], observing satisfactory convergence criteria for all models. These models were adjusted using the MCMCglmm package [33] developed for R [34] environment for statistical computing. Statistical significance was determined by a P value <0.05 .

Results

Patient sample

Among the 143 EP patients who accepted to participate in the study, 30 were excluded for the following reasons: [I] age at exposure to trauma was not available or there was a doubt about the date at the time of exposure [$n = 4$]; [II] first exposure to trauma occurred during the prodromal phase or after psychosis onset [$n = 3$]; [III] the physical measures were done more than 5 years after the entry into TIPP program [$n = 23$]. Therefore, analysis was carried out with the data of the remaining 113 patients.

Rates of trauma, demographics and baseline biological and physical measurements

The diagnostic breakdown, baseline demographic and clinical information including diagnosis alongside with biological measures of the sample are described in [Table 1](#).

Among 113 patients, 39 [34.5% of the total] had a history of at least one experience of trauma. Within the group with trauma, 28 [71.7% of exposed patients] had been exposed before age 12 [Early-trauma] and 11 [28.3% of exposed patients] between 12 and 16 [Late-trauma]. There were no differences in the rates of exposure to each individual traumatic experience [sexual, physical and emotional abuse; physical and emotional neglect] between patients exposed in childhood or in adolescence.

At baseline, no differences between Early-trauma, Late-trauma and Non-Trauma patients in terms of diagnostic distribution, levels of overweight, BMI, age, sex, SES, smoking status and the presence of treatment prior first assessment and during follow-up were observed.

Association of Early- and Late-Trauma with the BMI, weight gain and waist circumference during follow-up

Detailed results of GLMM models comparing the BMI, WC and WG at follow-up between Early, Late-Trauma and Non-Trauma are given in [Table 2](#) and represented in [Fig 1](#).

Compared to Non-Trauma patients, Early-Trauma patients did not differ in terms of BMI, WG and WC during follow-up. Compared to Non-Trauma patients, Late-Trauma patients did not differ in terms of BMI and WG during follow-up. However, they showed higher levels of WC [$\beta = 5.545$; $p = 0.013$] during follow-up. Results related to potential confounders and covariates are shown in [S1 Table](#), being time in treatment and baseline BMI the only variables significantly associated with BMI, WG and WC.

Discussion

We found that EP patients with trauma occurring between 12 and 16 years of age display higher WC during the first year of psychotropic medication compared to patients without trauma. In contrast, patients exposed to trauma before age 12 displayed similar levels of obesity related outcomes than non-traumatized patients. Our results support evidence suggesting that severe stress during adolescence could contribute to increased waist circumference in patients with psychosis [13, 14]. This suggests that adolescence is a critical period at which stress can

Table 1. Rates of trauma, demographics and baseline physical measurements.

	Total N = 113	Non Trauma N = 74	Early Trauma N = 28	Late Trauma N = 11	P-value
Age, median [range], years	25 [18–39]	25 [18–37]	27.5 [19–37]	24 [19–39]	0.558
Female, N [%]	36 [31.9]	21 [28.4]	13 [46.4]	2 [18.8]	0.12
Smoking, N [%]	46 [40.7]	31 [41.9]	11 [39.3]	4 [36.4]	0.955
SES, N [%]					
Low	23 [20.4]	18 [24.3]	4 [14.3]	1 [9.1]	0.092
Medium	47 [41.6]	33 [44.6]	12 [42.9]	2 [18.2]	
High	43 [38.0]	23 [31.1]	12 [42.9]	8 [72.7]	
Diagnostic, No [%]					
Schizophrenia	70 [61.9]	45 [60.8]	19 [67.9]	6 [54.5]	0.462
Schizophreniform/BPE	10 [8.9]	6 [8.1]	4 [14.3]	0 [0.0]	
Schizoaffective disorder	12 [10.6]	9 [12.2]	0 [0.0]	3 [27.3]	
Major depression with PF	4 [3.5]	3 [4.0]	1 [3.6]	0 [0.0]	
Bipolar Disorder	8 [7.1]	6 [8.1]	1 [3.6]	1 [9.1]	
Others	9 [8.0]	5 [6.8]	3 [10.7]	1 [9.1]	
Trauma subtypes					
Sexual abuse	13 [11.5]	-	9 [32.1]	4 [36.4]	0.801
Physical abuse	18 [16]	-	13 [46.4]	5 [45.5]	0.956
Emotional abuse	5 [4.4]	-	3 [10.7]	2 [18.2]	0.530
Physical neglect	5 [4.4]	-	4 [14.3]	1 [9.1]	0.662
Emotional neglect	10 [8.8]	-	8 [28.6]	2 [18.2]	0.503
Treatment prior first assessment N [%]	54 [47.8]	33 [44.6]	13 [46.4]	8 [72.7]	0.216
Treatment during follow up* N [%]					
Low	28 [24.8]	17 [23.0]	7 [25.0]	4 [36.4]	0.236
Moderate	56 [49.6]	40 [54.0]	10 [35.7]	6 [54.5]	
High	29 [25.7]	17 [23.0]	11 [39.3]	1 [9.1]	
Baseline physical measures #					
BMI, median [range] kg/ m ²	21.9 [15.6–35.4]	22.5 [15.6–33.6]	20.9 [16.8–35.4]	21.4 [17.5–30.1]	0.453
Overweight, N [%]	21 [19.6]	15 [21.7]	3 [11.1]	3 [27.3]	0.723
WC, median [range], cm	84 [45–114]	83 [58–114]	79 [45–110]	93 [82–110]	0.051
High WC, N [%] †	21 [24.1]	13 [22.4]	4 [19.1]	4 [50.0]	0.191

SES: Socio-economic-status; N = number of patients; PF: psychotic features; BPE: brief psychotic episode; BMI: Body Mass Index; WC: waist circumference

* treatment was categorized as function of risk of gaining weight into High: valproate, olanzapine and clozapine; Moderate: Quetiapine, Risperidone, Paliperidone, Lithium, Mirtazapine, Zuclopenthixol, Levomepromazine; Low: Amisulpride, Aripiprazole, Haloperidol, Lurasidone, Flupentixol;

#25 kg/m² ≥ Initial BMI < 30 kg/m², %;

†: High WC ≥ 94 cm [male], ≥ 88 cm [female] Early Trauma: refers to exposure from birth through 11 years of age; Late Trauma: refers to exposure between from age 12 years through 16 years.

<https://doi.org/10.1371/journal.pone.0242569.t001>

increase the vulnerability, in combination with other factors, to a later increase in WC, and subsequent obesity, which opens new venues for potential mechanisms explaining this link.

These results accounted for age, sex, diagnosis of depression, presence of psychotropic prescription before entry into the program and class of risk of medication intake during follow up. We did not find any association between exposure to trauma, neither during childhood nor adolescence, and increased BMI or weight change during follow-up, which suggests a stronger association of peripubertal psychological stress with abdominal adiposity.

The positive association between trauma and risk of obesity related outcomes has previously been reported in non-clinical samples [12]. Two studies have already shown an

Table 2. BMI, weight gain and waist circumference levels of patients exposed to Early and Late Trauma compared with nonexposed patients across 1 year follow-up.

	Early Trauma			Late Trauma		
	Estimate	CI [95%]	P value	Estimate	CI [95%]	P value
BMI, Kg/ m ²	0.139	[-0.62–0.89]	0.718	0.178	[-0.90–1.27]	0.742
Weight Gain in Kg	0.193	[-3.04–3.38]	0.906	0.562	[-3.99–5.17]	0.806
Waist Circumference, in cm	-1.626	[-4.67–1.41]	0.295	5.545	[1.14–9.85]	0.013*

BMI: Body Mass Index; CI: confidence interval. Early Trauma: refers to exposure from birth through 11 years of age; Late Trauma: refers to exposure between from age 12 years through 16 years.

* Significant at the 0.05 level.

<https://doi.org/10.1371/journal.pone.0242569.t002>

association between CT and greater BMI in patients with EP [13, 14]. Our study does not replicate the results regarding BMI, which could be due to a question of sample size. It may be possible that patients develop firstly a greater WC, and this contributes to later obesity and other cardiovascular risk factors. However, our stratification of traumatized patients according to the age at first exposure highlights a new finding: only patients exposed to trauma during adolescence display higher WC during the follow-up when compared to non-traumatized patients, being this association independent of a broad range of potential confounders. These results not only highlight the importance of stratifying patients according to the age of exposure but also open new venues for potential different neuro-developmental stages operating on the link between trauma and obesity related outcomes.

The Hypothalamus Pituitary Adrenal [HPA] axis is the most important regulator of the stress response [35] and stands as one of the potential mediating pathways linking trauma to psychopathology [36], including psychosis [11]. Interestingly, evidence from clinical studies in adolescent subjects to cellular and molecular studies show that elevated cortisol, particularly when combined with inhibition of sex steroids and growth hormone secretions, is causing accumulation of fat in visceral adipose tissues as well as metabolic abnormalities [37–39].

Moreover, teenagers, as a result of the pubertal surge in sex steroids, have shown increased responsiveness to stress hormones, leading to greater stress induced HPA axis responses as compared with adults [40, 41]. Taken altogether, we could speculate that trauma occurring during adolescence may induce elevated levels of glucocorticoids which, combined with inhibition of sex steroids, would lead to increased fat accumulation in visceral adipose tissue, contributing to a greater WC.

Contrary to what has been found by others [42, 43], we did not find that a diagnosis of depression accounted for changes in any of the obesity related outcomes. However, whether the depression [or sub-diagnostic depressive symptoms and different depression subtypes [44]] mediates the link between trauma and obesity related outcomes in psychosis needs to be explored in larger prospective samples. We did not find either an association between medication before the first assessment or during follow-up with our outcomes. This could be explained in part by a beneficial impact of the preventive strategies proposed by clinicians in charge of patients receiving psychotropic drugs at risk of weight gain that are part of the prospective monitoring implemented in the Lausanne University hospital [45]. Indeed, this follow up aims to reduce the negative impact of medication on weight and on metabolic parameters. These preventive measures include: [i] linking with general practitioner in the follow-up in order to activate a rapid response in case of weight gain or when an alteration in any of the metabolic disturbances appear; [ii] a continuous risk-benefit balance evaluation by the clinician in charge, considering switching to another medication appropriate to the clinical context




Fig 1. Graphical illustration of the differences in the waist circumference during follow-up [0, 2, 4, 6, 8, 10 and 12 months] between non trauma, Early Trauma and Late Trauma patients. The plot intends to capture a smooth trend in waist circumference for each subgroup of individuals using a spline fitted separately to each group. For each plot 95% Confidence bounds are also added to the plot. The plot is not adjusted for potential covariates but is rather a graphical mean to explore the variation of waist-circumference in function of time.

<https://doi.org/10.1371/journal.pone.0242569.g001>

every time a patient gains weight rapidly after initiation of a new psychotropic medication [iii] encouraging of patients to undertake healthy lifestyle habits, such as doing sport, yoga or being monitored by a nutritionist in case of rapid weight gain, especially during the first weeks of treatment [46].

Our study has various limitations. Firstly, although the obesity related outcomes were collected prospectively, exposure to trauma was assessed retrospectively, which may be particularly problematic for patients suffering from psychosis due to recall bias [47]. Moreover, although we looked at the time of exposure, we did not take into account other possible aspects such as the duration of the exposure, which may have an important modulating effect on out

measures outcomes. Despite these limitations, exposure to trauma was assessed by the case manager who followed each patient on the basis of information obtained from patients [48] and their families in the context of a 3-year therapeutic relationship. This allowed the clinician to choose the most appropriate moment when the patient did not present a mental state that could affect the recall of those experiences and may help some patients to acknowledge some experiences such as sexual abuses. Second, As we only followed patients after their first episode of psychosis and the lower age to be included in TIPP is 18, we cannot exclude the possibility that actually the increased peri abdominal adiposity in people exposed to trauma in adolescence was present prior the exposure to trauma, thus our study design does not infer causality and prospective studies following individuals from childhood to adulthood can provide a confirmation on our results. Secondly, many patients of the TIPP cohort refused to take part in the prospective metabolic follow-up, and we cannot exclude a selection bias in our sample [with patients not included in this study displaying different obesity related outcomes according to trauma exposure]. Thirdly, the sample size was relatively small and thus we cannot exclude that the lack of association between CT and BMI was due to a lack of statistical power. Furthermore, other potential confounders may have played a role in our results such as physical exercise or other aspects of psychopathology, such as anxiety. Other larger studies replicating our findings, including patients without psychotropic medication and a healthy control group exposed to CT are needed.

In conclusion, this is the first study showing that patients exposed to trauma during adolescence and being treated with AP medication have an increased WC during the early phase of the disease, independently of factors such as the previous medication or depression. These results open new venues toward possible mechanisms underlying the link between peripubertal stress and peri-abdominal adiposity, such as the possible impact of glucocorticoids related to a dysregulation of the HPA axis during the adolescence after exposure to CT. Given the high rates of cardiovascular-related mortality of patients with psychosis, there is an urgent need to further understand the association between trauma and obesity in this vulnerable population. Special attention should be paid to EPP who have been traumatized during adolescence when initiating a treatment to try to prevent an increase of WC, which is a major risk factor for metabolic syndrome [49].

Supporting information

S1 Table. Association between cofactors and confounders with BMI, weight gain and waist circumference across 1 year follow-up.
(DOCX)

Acknowledgments

We thank the Case Managers from TIPP Program for collecting this data. We thank our financial support bodies for supporting this work. Most of all, we express our gratitude to all the patients for their enduring participation.

Author Contributions

Conceptualization: Luis Alameda, Frederik Vandenberghe, Philipp Baumann, Victoria Rodriguez, Alessandra Solida, Kim Q. Do, Chin B. Eap, Philippe Conus.

Data curation: Luis Alameda, Axel Levier, Frederik Vandenberghe, Aurélie Delacretaz, Anaïs Glatard, Céline Dubath, Kim Q. Do, Chin B. Eap, Philippe Conus.

Formal analysis: Axel Levier, Mehdi Gholam-Rezaee, Philippe Golay.

Funding acquisition: Kim Q. Do, Chin B. Eap, Philippe Conus.

Investigation: Luis Alameda, Chin B. Eap.

Methodology: Luis Alameda, Mehdi Gholam-Rezaee, Kim Q. Do, Philippe Conus.

Resources: Luis Alameda, Chin B. Eap.

Supervision: Andres Herane-Vives, Alessandra Solida, Kim Q. Do, Chin B. Eap, Philippe Conus.

Validation: Victoria Rodriguez.

Visualization: Andres Herane-Vives.

Writing – original draft: Luis Alameda.

Writing – review & editing: Luis Alameda, Axel Levier, Mehdi Gholam-Rezaee, Frederik Vandenberghe, Aurélie Delacretaz, Philipp Baumann, Anaïs Glatard, Céline Dubath, Andres Herane-Vives, Victoria Rodriguez, Alessandra Solida, Kim Q. Do, Chin B. Eap, Philippe Conus.

References

1. Saha S, Chant D, McGrath J. A systematic review of mortality in schizophrenia: is the differential mortality gap worsening over time? *Arch Gen Psychiatry*. 2007; 64[10]:1123–31. Epub 2007/10/03. <https://doi.org/10.1001/archpsyc.64.10.1123> PMID: 17909124.
2. Laursen TM. Life expectancy among persons with schizophrenia or bipolar affective disorder. *Schizophr Res*. 2011; 131[1–3]:101–4. Epub 2011/07/12. <https://doi.org/10.1016/j.schres.2011.06.008> PMID: 21741216.
3. Ashworth M, Schofield P, Das-Munshi J. Physical health in severe mental illness. *The British journal of general practice: the journal of the Royal College of General Practitioners*. 2017; 67[663]:436–7. Epub 2017/10/01. <https://doi.org/10.3399/bjgp17X692621> PMID: 28963399.
4. Bioque M, Garcia-Portilla MAP, Garcia-Rizo C, Cabrera B, Lobo A, Gonzalez-Pinto A, et al. Evolution of metabolic risk factors over a two-year period in a cohort of first episodes of psychosis. *Schizophr Res*. 2017. Epub 2017/07/01. <https://doi.org/10.1016/j.schres.2017.06.032> PMID: 28663026.
5. Anjum S, Bathla M, Panchal S, Singh GP, Singh M. Metabolic syndrome in drug naive schizophrenic patients. *Diabetes & metabolic syndrome*. 2017. Epub 2017/12/02. <https://doi.org/10.1016/j.dsx.2017.11.006> PMID: 29191416.
6. Delacretaz A, Vandenberghe F, Gholam-Rezaee M, Saigi Morgui N, Glatard A, Thonney J, et al. Early changes of blood lipid levels during psychotropic drug treatment as predictors of long-term lipid changes and of new onset dyslipidemia. *Journal of clinical lipidology*. 2018; 12[1]:219–29. Epub 2017/11/13. <https://doi.org/10.1016/j.jacl.2017.10.002> PMID: 29128242.
7. Attux C, Martini LC, Elkis H, Tamai S, Freirias A, Camargo M, et al. A 6-month randomized controlled trial to test the efficacy of a lifestyle intervention for weight gain management in schizophrenia. *BMC Psychiatry*. 2013; 13:60. Epub 2013/02/20. <https://doi.org/10.1186/1471-244X-13-60> PMID: 23418863.
8. Chen DC, Du XD, Yin GZ, Yang KB, Nie Y, Wang N, et al. Impaired glucose tolerance in first-episode drug-naive patients with schizophrenia: relationships with clinical phenotypes and cognitive deficits. *Psychol Med*. 2016; 46[15]:3219–30. Epub 2016/11/03. <https://doi.org/10.1017/S0033291716001902> PMID: 27604840.
9. Gonzalez-Blanco L, Greenhalgh AMD, Garcia-Rizo C, Fernandez-Egea E, Miller BJ, Kirkpatrick B. Prolactin concentrations in antipsychotic-naive patients with schizophrenia and related disorders: A meta-analysis. *Schizophr Res*. 2016; 174[1–3]:156–60. Epub 2016/04/14. <https://doi.org/10.1016/j.schres.2016.03.018> PMID: 27068570.
10. Greenhalgh AM, Gonzalez-Blanco L, Garcia-Rizo C, Fernandez-Egea E, Miller B, Arroyo MB, et al. Meta-analysis of glucose tolerance, insulin, and insulin resistance in antipsychotic-naive patients with nonaffective psychosis. *Schizophr Res*. 2017; 179:57–63. Epub 2016/10/17. <https://doi.org/10.1016/j.schres.2016.09.026> PMID: 27743650.

11. Morgan C, Gayer-Anderson C. Childhood adversities and psychosis: evidence, challenges, implications. *World Psychiatry*. 2016; 15[2]:93–102. Epub 2016/06/07. <https://doi.org/10.1002/wps.20330> PMID: 27265690.
12. Danese A, Tan M. Childhood maltreatment and obesity: systematic review and meta-analysis. *Mol Psychiatry*. 2014; 19[5]:544–54. Epub 2013/05/22. <https://doi.org/10.1038/mp.2013.54> PMID: 23689533.
13. Heggul N, Pariante CM, Dipasquale S, DiForti M, Taylor H, Marques TR, et al. Childhood maltreatment is associated with increased body mass index and increased C-reactive protein levels in first-episode psychosis patients. *Psychol Med*. 2012; 42[9]:1893–901. Epub 2012/01/21. <https://doi.org/10.1017/S0033291711002947> PMID: 22260948.
14. Aas M, Dieset I, Hope S, Hoseth E, Mørch R, Reponen E, et al. Childhood maltreatment severity is associated with elevated C-reactive protein and body mass index in adults with schizophrenia and bipolar diagnoses. *Brain, behavior, and immunity*. 2017; 65:342–9. <https://doi.org/10.1016/j.bbi.2017.06.005> PMID: 28619247.
15. Slavich GM, Irwin MR. From stress to inflammation and major depressive disorder: a social signal transduction theory of depression. *Psychological bulletin*. 2014; 140[3]:774–815. Epub 2014/01/15. <https://doi.org/10.1037/a0035302> PMID: 24417575.
16. Pariante CM. Why are depressed patients inflamed? A reflection on 20 years of research on depression, glucocorticoid resistance and inflammation. *European neuropsychopharmacology: the journal of the European College of Neuropsychopharmacology*. 2017; 27[6]:554–9. Epub 2017/05/10. <https://doi.org/10.1016/j.euroneuro.2017.04.001> PMID: 28479211.
17. Alameda L, Golay P, Baumann PS, Progin P, Mebdouhi N, Elowe J, et al. Mild Depressive Symptoms Mediate the Impact of Childhood Trauma on Long-Term Functional Outcome in Early Psychosis Patients. *Schizophr Bull*. 2017; 43[5]:1027–35. Epub 2016/11/26. <https://doi.org/10.1093/schbul/sbw163> PMID: 27884931.
18. Lasserre AM, Glaus J, Vandeleur CL, Marques-Vidal P, Vaucher J, Bastardot F, et al. Depression with atypical features and increase in obesity, body mass index, waist circumference, and fat mass: a prospective, population-based study. *JAMA psychiatry*. 2014; 71[8]:880–8. Epub 2014/06/06. <https://doi.org/10.1001/jamapsychiatry.2014.411> PMID: 24898270.
19. Baumann PS, Crespi S, Marion-Veyron R, Solida A, Thonney J, Favrod J, et al. Treatment and early intervention in psychosis program [TIPP-Lausanne]: Implementation of an early intervention programme for psychosis in Switzerland. *Early Interv Psychiatry*. 2013; 7[3]:322–8. Epub 2013/03/01. <https://doi.org/10.1111/eip.12037> PMID: 23445318.
20. Yung AR, Yuen HP, McGorry PD, Phillips LJ, Kelly D, Dell'Olio M, et al. Mapping the onset of psychosis: the Comprehensive Assessment of At-Risk Mental States. *The Australian and New Zealand journal of psychiatry*. 2005; 39[11–12]:964–71. Epub 2005/12/14. <https://doi.org/10.1080/j.1440-1614.2005.01714.x> PMID: 16343296.
21. Choong E, Quteineh L, Cardinaux J-R, Gholam-Rezaee M, Vandenberghe F, Dobrinas M, et al. Influence of CRT1 polymorphisms on body mass index and fat mass in psychiatric patients and the general adult population. *JAMA psychiatry*. 2013; 70[10]:1011–9. <https://doi.org/10.1001/jamapsychiatry.2013.187> PMID: 23925723.
22. Association AAP. Diagnostic and statistical manual of mental disorders [DSM-IV]. Washington, DC: American Psychiatric Association; 1994.
23. Choong E, Solida A, Lechaire C, Conus P, Eap CB. Follow-up of the metabolic syndrome induced by atypical antipsychotics: recommendations and pharmacogenetics perspectives. *Revue medicale suisse*. 2008; 4[171]:1994–9. Epub 2008/10/14. PMID: 18847133.
24. Alameda L, Ferrari C, Baumann PS, Gholam-Rezaee M, Do KQ, Conus P. Childhood sexual and physical abuse: age at exposure modulates impact on functional outcome in early psychosis patients. *Psychol Med*. 2015; 45[13]:2727–36. Epub 2015/09/10. <https://doi.org/10.1017/S0033291715000690> PMID: 26350397.
25. Varese F, Smeets F, Drukker M, Lieverse R, Lataster T, Viechtbauer W, et al. Childhood adversities increase the risk of psychosis: a meta-analysis of patient-control, prospective- and cross-sectional cohort studies. *Schizophr Bull*. 2012; 38[4]:661–71. Epub 2012/03/31. <https://doi.org/10.1093/schbul/sbs050> PMID: 22461484.
26. Fisher HL, Jones PB, Fearon P, Craig TK, Dazzan P, Morgan K, et al. The varying impact of type, timing and frequency of exposure to childhood adversity on its association with adult psychotic disorder. *Psychol Med*. 2010; 40[12]:1967–78. Epub 2010/02/25. <https://doi.org/10.1017/S0033291710000231> PMID: 20178679.
27. Thornberry TP, Ireland TO, Smith CA. The importance of timing: The varying impact of childhood and adolescent maltreatment on multiple problem outcomes. *Development and psychopathology*. 2001; 13[4]:957–79. PMID: 11771916

28. Morgan C, Gayer-Anderson C, Beards S, Hubbard K, Mondelli V, Di Forti M, et al. Threat, hostility and violence in childhood and later psychotic disorder: population-based case-control study. *The British Journal of Psychiatry*. 2020;1–8. <https://doi.org/10.1192/bjp.2020.133> PMID: 32778182
29. Alameda L, Golay P, Baumann PS, Ferrari C, Do KQ, Conus P. Age at the time of exposure to trauma modulates the psychopathological profile in patients with early psychosis. *J Clin Psychiatry*. 2016; 77 [5]:e612–8. Epub 2016/06/02. <https://doi.org/10.4088/JCP.15m09947> PMID: 27249091.
30. Vandenberghe F, Saigi-Morgui N, Delacretaz A, Quteineh L, Crettol S, Ansermot N, et al. Prediction of early weight gain during psychotropic treatment using a combinatorial model with clinical and genetic markers. *Pharmacogenetics and genomics*. 2016; 26[12]:547–57. Epub 2016/10/28. <https://doi.org/10.1097/FPC.000000000000249> PMID: 27741037.
31. Alameda L, Golay P, Baumann P, Morandi S, Ferrari C, Conus P, et al. Assertive outreach for "difficult to engage" patients: A useful tool for a subgroup of patients in specialized early psychosis intervention programs. *Psychiatry Res*. 2016; 239:212–9. Epub 2016/03/28. <https://doi.org/10.1016/j.psychres.2016.03.010> PMID: 27016880.
32. Chandola T, Jenkinson C. The new UK National Statistics Socio-Economic Classification [NS-SEC]; investigating social class differences in self-reported health status. *Journal of public health medicine*. 2000; 22[2]:182–90. Epub 2000/07/27. <https://doi.org/10.1093/pubmed/22.2.182> PMID: 10912557.
33. Hadfield JD. MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R package. *Journal of Statistical Software*. 2010; 33[2]:1–22.
34. Team RC. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. 2014. 2016.
35. Lupien SJ, McEwen BS, Gunnar MR, Heim C. Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nat Rev Neurosci*. 2009; 10[6]:434–45. Epub 2009/04/30. <https://doi.org/10.1038/nrn2639> PMID: 19401723.
36. Ruby E, Polito S, McMahon K, Gorovitz M, Corcoran C, Malaspina D. Pathways Associating Childhood Trauma to the Neurobiology of Schizophrenia. *Frontiers in psychological and behavioral science*. 2014; 3[1]:1–17. Epub 2014/11/25. PMID: 25419548.
37. Bjorntorp P. Do stress reactions cause abdominal obesity and comorbidities? *Obesity reviews: an official journal of the International Association for the Study of Obesity*. 2001; 2[2]:73–86. Epub 2002/07/18. <https://doi.org/10.1046/j.1467-789x.2001.00027.x> PMID: 12119665.
38. Vicennati V, Pasquali R. Abnormalities of the hypothalamic-pituitary-adrenal axis in nondepressed women with abdominal obesity and relations with insulin resistance: evidence for a central and a peripheral alteration. *The Journal of clinical endocrinology and metabolism*. 2000; 85[11]:4093–8. Epub 2000/11/30. <https://doi.org/10.1210/jcem.85.11.6946> PMID: 11095438.
39. Steptoe A, Kunz-Ebrecht SR, Brydon L, Wardle J. Central adiposity and cortisol responses to waking in middle-aged men and women. *International journal of obesity and related metabolic disorders: journal of the International Association for the Study of Obesity*. 2004; 28[9]:1168–73. Epub 2004/06/24. <https://doi.org/10.1038/sj.ijo.0802715> PMID: 15211363.
40. Sinclair D, Purves-Tyson TD, Allen KM, Weickert CS. Impacts of stress and sex hormones on dopamine neurotransmission in the adolescent brain. *Psychopharmacology*. 2014; 231[8]:1581–99. <https://doi.org/10.1007/s00213-013-3415-z> PMID: 24481565
41. Kuhlman KR, Geiss EG, Vargas I, Lopez-Duran NL. Differential associations between childhood trauma subtypes and adolescent HPA-axis functioning. *Psychoneuroendocrinology*. 2015; 54:103–14. <https://doi.org/10.1016/j.psyneuen.2015.01.020> PMID: 25704913
42. Lasserre AM, Strippoli MF, Glaus J, Gholam-Rezaee M, Vandeleur CL, Castelao E, et al. Prospective associations of depression subtypes with cardio-metabolic risk factors in the general population. *Mol Psychiatry*. 2017; 22[7]:1026–34. Epub 2016/10/12. <https://doi.org/10.1038/mp.2016.178> PMID: 27725658.
43. Opel N, Redlich R, Grotegerd D, Dohm K, Heindel W, Kugel H, et al. Obesity and major depression: Body-mass index [BMI] is associated with a severe course of disease and specific neurostructural alterations. *Psychoneuroendocrinology*. 2015; 51:219–26. Epub 2014/12/03. <https://doi.org/10.1016/j.psyneuen.2014.10.001> PMID: 25462895.
44. Herane-Vives A, de Angel V, Papadopoulos A, Wise T, Chua KC, Strawbridge R, et al. Short-term and long-term measures of cortisol in saliva and hair in atypical and non-atypical depression. *Acta Psychiatrica Scandinavica*. 2018; 137[3]:216–30. <https://doi.org/10.1111/acps.12852> PMID: 29397570
45. Choong E, Solida A, Lechaire C, Conus P, Eap CB. [Follow-up of the metabolic syndrome induced by atypical antipsychotics: recommendations and pharmacogenetics perspectives]. *Revue medicale suisse*. 2008; 4[171]:1994–6, 8–9. Epub 2008/10/14. PMID: 18847133.
46. Association AD. Consensus development conference on antipsychotic drugs and obesity and diabetes. *Diabetes care*. 2004; 27[2]:596–601. <https://doi.org/10.2337/diacare.27.2.596> PMID: 14747245

47. Howard LM. Allegations of abuse in psychotic patients. *Am J Psychiatry*. 1993; 150[5]:839–40. Epub 1993/05/01. <https://doi.org/10.1176/ajp.150.5.839b> PMID: 8480841.
48. Bahr SM, Weidemann BJ, Castro AN, Walsh JW, deLeon O, Burnett CM, et al. Risperidone-induced weight gain is mediated through shifts in the gut microbiome and suppression of energy expenditure. *EBioMedicine*. 2015; 2[11]:1725–34. Epub 2016/02/13. <https://doi.org/10.1016/j.ebiom.2015.10.018> PMID: 26870798.
49. Després J-P, Lemieux I. Abdominal obesity and metabolic syndrome. *Nature*. 2006; 444[7121]:881. <https://doi.org/10.1038/nature05488> PMID: 17167477



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Alameda, L; Levier, A; Gholam-Rezaee, M; Golay, P; Vandenberghe, F; Delacretaz, A; Baumann, P; Glatard, A; Dubath, C; Herane-Vives, A; Rodriguez, V; Solida, A; Do, KQ; Eap, CB; Conus, P

Title:

Psychological trauma occurring during adolescence is associated with an increased risk of greater waist circumference in Early Psychosis patients treated with psychotropic medication

Date:

2020-12-03

Citation:

Alameda, L., Levier, A., Gholam-Rezaee, M., Golay, P., Vandenberghe, F., Delacretaz, A., Baumann, P., Glatard, A., Dubath, C., Herane-Vives, A., Rodriguez, V., Solida, A., Do, K. Q., Eap, C. B. & Conus, P. (2020). Psychological trauma occurring during adolescence is associated with an increased risk of greater waist circumference in Early Psychosis patients treated with psychotropic medication. PLOS ONE, 15 (12), <https://doi.org/10.1371/journal.pone.0242569>.

Persistent Link:

<http://hdl.handle.net/11343/272359>

File Description:

Published version

License:

CC BY