


Building the case for the calcitonin receptor as a viable target for the treatment of glioblastoma

Pragya Gupta, Sebastian G. B. Furness, Lucas Bittencourt , David L. Hare and Peter J. Wookey

Abstract: Researchers are actively seeking novel targeted therapies for the brain tumour glioblastoma (GBM) as the mean survival is less than 15 months. Here we discuss the proposal that the calcitonin receptor (CT Receptor), expressed in 76–86% of patient biopsies, is expressed by both malignant glioma cells and putative glioma stem cells (GSCs), and therefore represents a potential therapeutic target. Forty-two per cent (42%) of high-grade glioma (HGG; representative of GSCs) cell lines express CT Receptor protein. CT Receptors are widely expressed throughout the life cycle of organisms and in some instances promote apoptosis. Which of the common isoforms of the CT Receptor are predominantly expressed is currently unknown, but a functional response to cell stress of the insert-positive isoform is hypothesised. A model for resistant malignancies is one in which chemotherapy plays a direct role in activating quiescent stem cells for replacement of the tumour tissue hierarchy. The putative role that the CT Receptor plays in maintenance of quiescent cancer stem cells is discussed in view of the activation of the Notch–CT Receptor–collagen V axis in quiescent muscle (satellite) stem cells. The pharmacological CT response profiles of four of the HGG cell lines were reported. Both CT responders and non-responders were sensitive to an immunotoxin based on an anti-CT Receptor antibody. The *CALCR* mRNA exhibits alternative splicing commonly associated with cancer cells, which could result in the atypical pharmacology exhibited by CT non-responders and an explanation of tumour suppression. Due to the inherent instability of *CALCR* mRNA, analysis of CT Receptor protein in patient samples will lead to improved data for the expression of CT Receptor in GBM and other cancers, and an understanding of the role and activity of the splice variants. This knowledge will aid the effective targeting of this receptor for treatment of GBM.

Keywords: antibody, brain tumour, calcitonin receptor, CTR, CT Receptor, glioblastoma, glioma stem cells, G protein-coupled receptor

Received: 5 June 2020; revised manuscript accepted: 11 November 2020.

Introduction

Glioblastoma (multiforme, GBM) is the most common malignant primary brain tumour in adults. Aggressive surgical resection decreases the tumour-cell burden by 99%, leaving about 100 million cells and with cytotoxic adjuvant therapy with an agent such as temozolomide the burden is reduced to about 10 million cells, which would include cancer stem cells.¹ The mean overall survival achieved with wide-local surgical resection,

followed by adjuvant therapy of radiation and temozolomide (TMZ) plus maintenance therapy with TMZ, remains only 14.6 months.²

Glioblastoma (GBM) are highly heterogeneous tumours thought to be derived from oligodendrocyte-type 2-astrocyte (O-2A) progenitors of the astroglial lineage⁴ as malignant glioma cells express glial fibrillary acidic protein (GFAP) and respond to similar mitogens and differentiation factors.⁵

Ther Adv Med Oncol

2020, Vol. 12: 1–15

DOI: 10.1177/
1758835920978110

© The Author(s), 2020.
Article reuse guidelines:
[sagepub.com/journals-](https://sagepub.com/journals-permissions)
[permissions](https://sagepub.com/journals-permissions)

Correspondence to:
Peter J. Wookey
Department of Medicine,
University of Melbourne,
Level 10, Lance Townsend
Building, Austin Health,
Studley Road, Heidelberg,
Victoria 3084, Australia
pwookey@unimelb.edu.au

Pragya Gupta
Lucas Bittencourt
David L. Hare
Department of Medicine
(Austin Health,
Heidelberg), University of
Melbourne, Melbourne,
Victoria, Australia

Sebastian G. B. Furness
Drug Discovery Biology
Laboratory, Monash
Institute of Pharmaceutical
Sciences & Department
of Pharmacology, Monash
University (Parkville),
Victoria, Australia




Figure 1. (A) Illustration of the major anatomical features of glioblastomas (GBM); (B) Haematoxylin and eosin stain of a tissue section of GBM with an example of vascular mimicry (central vessel) and oedema; and (C) A serial section showing staining of malignant glioma cells³ with the anti-human CT Receptor antibody mAb31/01-1H10. Nuclei are stained blue with haematoxylin.

The tumour architecture (illustrated in Figure 1) includes regions of necrosis and oedema (where the blood–brain barrier is compromised), pro-inflammatory micro-environments with cellular infiltrates and proliferative domains. Vascular-like structures referred to as vascular mimicry have been described.¹ Within these latter hyperplastic structures pericyte precursors^{1,6} and endothelial cells^{7–9} share the same genetic modifications as glioma stem-like cells (GSCs), supporting plasticity with regard to differentiation lineages and the existence of dominant GSC clones.^{3,10–13} It is noteworthy that pericyte precursors are multipotent,¹⁴ are evident in the GBM vasculature in the close vicinity to the zones of proliferation, and a percentage express GFAP as noted in a stroke model.¹⁵ This raises the possibility that re-emerged GBM post-treatment might be derived from the pericyte lineage.

Comprehensive studies have detailed the gene expression profiles of GBM tissue samples^{10–13} and this is supplemented by an anatomical transcriptional atlas.¹⁶

The current inability to improve or predict patient outcomes based on genetic profiling or histopathological features points to a deficit in our understanding of the driving forces for tumorigenesis of GBM and hence targets for tumour reduction or potential therapy.

GSCs play a role as precursor cells for GBM.¹⁷ They display inherent functional diversity,^{1,3} convey relative resistance to conventional treatments such as chemo and radiotherapy,¹⁸ and provide invasive potential.^{19–21} GSCs may also contribute to tumour survival and expansion in a number of hostile (hypoxic, inflammatory) micro-environments. As described for normal stem cell populations, GSCs are likely to be associated with dense vascular beds,²² which are generally located towards the periphery of GBM, and GSCs are believed to be present within the surrounding neuropil perhaps in a quiescent state. GSCs, also known as brain tumour-propagating cells¹⁹ in contrast to brain tumour-initiating cells,^{20,23} can be perpetuated and expanded serum free²⁴ *in vitro* as high grade glioma (HGG) cell lines.^{25–27} When formed as xenografts in the brains of immune-compromised mice, HGG cell lines form orthotopic, intracranial (*ic*) tumours that retain their invasive potential and recapitulate much of the pathology of the original tumour.^{17,19,20,23,24,27}

Some evidence has been presented to support the idea that some GSC populations are normally quiescent (G_0 phase of the cell cycle) and re-enter G_1 phase for self-renewal, but during quiescence are resistant to radiotherapy and conventional chemotherapy, designed to target proliferating cells.

Furthermore, evidence suggests that dying cells targeted by chemotherapy release mitogens that stimulate quiescent stem cells to repopulate the tumour in between cycles of chemotherapy.^{3,18} Thus, the biology of quiescent stem cells represents further complexity when considering the evaluation of new drugs. GSCs are currently recognised as potential targets for therapy, in which case specific molecular targets on quiescent GSCs should be validated and corresponding therapeutics developed. In support of the idea that quiescent stem cells are responsible for GBM relapse, in childhood lymphoblastic leukaemia, quiescent leukaemic (stem) cells appear to account for relapse-causing minimal residual disease (MRD).²⁸

In GBM, genetic profiles generated from patient biopsies with considerable cellular hierarchy might not provide enough definition to characterise targets on small subpopulations of quiescent GSCs which have the potential to become dominant clones following expansion after treatment cycles.

In a study that investigated CT Receptor protein expression in a small number of GBM patient biopsies, 86% showed expression that is restricted to glioma cells (example in Figure 2) and smaller cells bearing the GSC associated marker, CD-133.²⁹ The fact that the surrounding neuropil is negative for CT Receptor²⁹ in those regions of the brain in which GBM is common suggests CT Receptor is a GBM restricted biomarker. Another report described expression of *CALCR* mRNA found in 115/152 (76%) of primary tumours³⁰ as calculated from previous reported data.¹²

The widespread expression of the calcitonin receptor (CT Receptor)

CT Receptor isoforms are expressed in a wide range of tissues throughout the life cycle of mammals, under conditions such as cell stress, inflammation and wound healing, and in a range of diseases (Table 1).³¹ In spite of this wide expression, it is not normally expressed in the cerebrum/cortex where GBMs typically arise but is restricted to specific neuronal networks in the limbic system, and in the mid and hindbrain.

There are two common isoforms, CT Receptor_a, insert-negative and CT Receptor_b, insert-positive.⁸⁰ The insert positive form was originally isolated from a breast cancer cell line and has an additional 16-amino acid sequence within the first




Figure 2. An example of a Glioblastoma (GBM) tumour stained with an anti-human CT Receptor monoclonal antibody mAb30/07-9B4. Malignant glioma cells are stained brown and nuclei are stained blue with haematoxylin.²⁹

intracellular loop. The insert negative form has the more extensively characterised pharmacology and appears to be the relevant isoform for the well-defined physiology of calcitonin signaling and calcium metabolism.

Data from transfected COS (monkey kidney) cell lines that express either of the two human isoforms show CT Receptor_a is located predominantly in the plasma membrane (Figure 3) whereas CT Receptor_b is largely intracellular, located in the perinuclear domain presumably in small membranous elements and has a lower molecular weight suggesting an unglycosylated form that has not been processed for normal cell-surface expression.⁸¹ In transfected cell lines flow cytometry experiments confirm that lesser amounts of CT Receptor_b are found on the plasma membrane, but the relative distribution is cell line dependent.⁸²

The CT Receptor_b isoform is expressed more widely than previously thought across the mammalian order. In the majority of mammalian species examined so far (exception *Muroidea*), the position of insertion is identical at the beginning of the second transmembrane span; however, the insertion varies in amino acid sequence and length (16–18), and would be predicted to interfere with the interaction of the receptor with its primary transducers. The recent description of the widespread expression of insert-positive isoforms

Table 1. Updated and abbreviated list of tissues that express CTR from several mammalian species. A more comprehensive list with references is listed in Wookey *et al.*³¹

Life cycle entity, condition or disease	Tissue	References
Blastula		Wang <i>et al.</i> ³²
Gastrula		Burgess ³³
Foetus		Jagger <i>et al.</i> ^{34,35}
Embryonic	Brain	Wookey <i>et al.</i> , ³¹ Tolcos <i>et al.</i> ³⁶
Perinatal	Gut (transient expression)	Wookey <i>et al.</i> ³⁷
	Kidney	Tikellis <i>et al.</i> ³⁸
Adult tissues	Brain	Fischer <i>et al.</i> , ³⁹ Sexton <i>et al.</i> , ⁴⁰ Becskei <i>et al.</i> , ⁴¹ Paxinos <i>et al.</i> , ⁴² Goda <i>et al.</i> , ⁴³ Walker <i>et al.</i> ⁴⁴
	Bone (osteoclasts, osteocytes)	Marx <i>et al.</i> , ⁴⁵ Nicholson <i>et al.</i> , ⁴⁶ Hattersley and Chalmers ⁴⁷ , Zaidi <i>et al.</i> , ⁴⁸ Gooi <i>et al.</i> ⁴⁹
	Kidney	Marx <i>et al.</i> , ⁴⁵ Sexton <i>et al.</i> ⁵⁰
	Thyroid	Hanna <i>et al.</i> ⁵¹
	Satellite stem cells of muscle	Fukada <i>et al.</i> , ⁵² Gnocchi <i>et al.</i> , ⁵³ Yamaguchi <i>et al.</i> , ⁵⁴ Baghdadi and Tajbakhsh ⁵⁵
	Spermatozoa	Silvestroni <i>et al.</i> , ⁵⁶ Adeoya-Osiguwa <i>et al.</i> ⁵⁷
	Placenta	Nicholson <i>et al.</i> , ⁵⁸ Kovacs <i>et al.</i> ⁵⁹
	Prostate	Wu <i>et al.</i> ⁶⁰
Conditions	Cell stress	Furness <i>et al.</i> ⁶¹
	Inflammatory cytokines	Meeuwssen <i>et al.</i> ⁶²
	Wound healing	Wookey <i>et al.</i> ³¹
Cardiovascular disease	Blood vessels	Wookey <i>et al.</i> ^{63,64}
Cancers	Glioblastoma	Wookey <i>et al.</i> , ²⁹ Gilabert-Oriol <i>et al.</i> , ⁶⁵ Pal <i>et al.</i> ⁶⁶
	Prostate	Ritchie <i>et al.</i> , ⁶⁷ Thomas <i>et al.</i> ⁶⁸
	Medullary thyroid	Frendo <i>et al.</i> , ^{69,70} Cappagli <i>et al.</i> ⁷¹
	Thyroid carcinomas	Boot <i>et al.</i> ⁷²
	Bone (osteoclastoma, giant cell)	Nicholson <i>et al.</i> , ⁷³ Gorn <i>et al.</i> ⁷⁴
	Multiple myeloma	Silvestris <i>et al.</i> ⁷⁵
	Leukaemia (ALL, AML), K562	Wookey <i>et al.</i> , ³¹ Mould and Pondel ⁷⁶
	Ovarian cell line BIN-67	Gorn <i>et al.</i> ⁷⁷
Breast	Findlay <i>et al.</i> , ⁷⁸ Gillespie <i>et al.</i> ⁷⁹	

ALL, Acute Lymphoblastic Leukaemia; AML, Acute Myeloid Leukaemia.




Figure 3. COS-7 stable transfectants expressing CT Receptor_a and CT Receptor_b⁸¹ stained with primary anti-human CT Receptor antibody mAb31/01-1H10, secondary goat anti-mouse IgG2a:AF568 and imaged using confocal microscopy (Zeiss LSM 800). Nuclei are stained blue with DAPI (4',6 diamidino-2-phenylindole).

throughout the mammalian species does, however, suggest a specific function of CT Receptor_b in cell physiology.⁸¹

There are no conclusive data yet that demonstrates in which normal and diseased tissues CT Receptor_b is more highly expressed and what its function might be. However, there is some evidence that this isoform is more highly expressed in ovary and placenta⁸⁰ and CT Receptor_b mRNA predominates in a group of samples of mononuclear blood cells.⁸³ The recent validation of a mouse monoclonal anti-human CT Receptor_b antibody⁸¹ will aid in the resolution of questions about expression in normal tissues, during inflammation and diseases. Unpublished data from our group on the expression of CT Receptor_b upregulated in cell stress (see also Adeoya-Osiguwa and Fraser)⁵⁷ are consistent with a cytosolic function.

Pharmacology of CT Receptor_a and CT Receptor_b

The pharmacology of these isoforms is quite different in terms of the second messenger outputs which, for CT Receptor_a, includes adenylyl cyclase activation, phosphorylation of ERK1/2 and p38 MAP kinase, as well as mobilisation of intracellular calcium (summarised in Figure 4 below). This might be expected given the location of the insert in relation to the binding of CT Receptor to its primary




Figure 4. Typical signaling pathways from CT Receptor_a and CT Receptor_b.

transducer.⁸⁴ In heterologous systems (COS-7 and HEK 293), second messenger coupling of CT Receptor_b is substantially reduced: the potency for adenylyl cyclase activation is more than 100-fold weaker, ligand stimulated phosphorylation of ERK1/2 is reduced in its maximum with some reduction in potency and CT Receptor-dependent mobilisation of intracellular calcium is undetectable.⁸² Such outputs are also cell type dependent^{82,85} and mediated by different G protein complexes. However, it was noted that stimulation with salmon CT of HEK-293 transfectants (both CT Receptor isoforms compared to vector control) resulted in acidification of the media⁸⁶ suggesting possible metabolic outputs or involvement in efflux mechanisms from acidified intracellular compartments.

The CT Receptor ligand promoted conformation of Receptor bound G protein complexes influences their residency on the plasma membrane and contributes to ligand-mediated biased agonism of CT Receptor_a.⁸⁷ However, little is known about these mechanisms in the context of insert-positive CT Receptor_b and how the peptide insert might perturb residency, and thus coupling of G protein complexes.

CT Receptor protein was detected in the 5/12 (42%) of HGG cell lines (isolated from human biopsies) tested,²⁷ including JK2 (mesenchymal), PB1 (proneural/classic), SB2b (mesenchymal/classic) and WK1 (classic), using immunoblotting with the monoclonal antibody mAb31/01-1H10³⁰ which binds an intracellular epitope. In contrast to the major, broad CT Receptor band of 70–80 kD from COS-7/CT Receptor_a transfectants,⁸¹ the upper band is tight, with an apparent molecular weight (MW) of approximately 57 kD, consistent with an unglycosylated form. Unglycosylated CT Receptor is largely confined to the cytosolic domain and is characteristic of the CT Receptor_b isoform⁸¹ (Figure 3 above). However, as discussed above a small proportion of CT Receptor_b is located on the cell surface, which is cell line dependent,⁸² and in the case of HGG cell lines JK2, SB2b and WK1, CT Receptor is found in the membrane fraction determined from immunoblot.³⁰ In PB1 there is much less CT Receptor protein found in the membrane fraction. Rapid turnover of unglycosylated human CT Receptor_a protein located in intracellular compartments has been described.⁸⁸ Further studies including nanopore sequencing to identify splicing events will be important to improve our understanding of the biology and functional consequences of these observations.

The pharmacology of CT Receptor was studied in these HGG lines³⁰ and only SB2b had functional CT Receptor as determined by responses to classic ligands which stimulated adenylyl cyclase, calcium mobilisation and ERK^{1/2} phosphorylation. Pharmacological studies are discussed in more detail below.

Hypothesis on the mechanism of the CT Receptor_b isoform

This hypothesis is built on several separate ideas. Firstly, intracellular CT Receptors are folded in the membrane of intracellular structures attached to the cytoskeleton.⁸⁹ Secondly, unglycosylated CT Receptor_b is confined to the cytosol (Figure 3 above) and is concentrated in a structure thought to be the microtubule organising centre (MTOC). In experiments similar to those described by Safaei *et al.*⁹⁰ we found cytotoxin concentrated in the exosomes harvested from MG63 cells treated with staurosporine and these exosomes were also positive for CT Receptor (unpublished results). As these exosomes originate from lysosomes (pH 3–4) their release is likely to contribute to acidification of the media as described previously⁸⁶ for CT Receptor_b.

The upregulation of *CALCR* mRNA in response to cytokines TNF α and IL1 β by primary cultures of human astrocytes has been described,⁶² demonstrating a response to cell stress.⁶¹ Furthermore, in U87 MG cells treated with staurosporine, nanopore sequencing of long cDNA products from CT Receptor mRNA has established alternative splicing events that include exon 10 such that the CT Receptor_b isoform (insert-positive) is significantly upregulated,⁸¹ although total *CALCR* mRNA remains unchanged.

Taken together there is evidence for the upregulation of CT Receptor_b in the cytosol of stressed cells and the production of CT Receptor-positive exosomes laden with cytotoxins, which as hypothesised, amounts to new important mechanism together with a number of other cellular stress responses.

In the context of the expression of CT Receptor by malignant glioma cells and GSCs, targeting CT Receptor might provide an opportunity to overcome resistance to chemotherapeutics and treat the pool of GSCs thought to be responsible for relapse.

Calcitonin/CT Receptor, cell survival/apoptosis and the cell cycle

There are several reports that describe data showing CT or CT Receptor promote proliferation/survival or apoptosis resulting in decreased survival in model cancer cell lines. The effect of CT Receptor expression and calcitonin-dependent CT Receptor activation differs according to the cell line under investigation.

CT stimulates proliferation early in treatment protocol of T47D cells (derived from human breast cancer) and then later inhibits proliferation in the log phase.⁹¹ In serum deprived LLC-PK cells, derived from porcine kidney, CT reduced cell survival⁹² perhaps by induction of apoptosis.

In a study of serum-starved transfected HEK-293 cells induced to express either human hCT Receptor_a compared to hCT Receptor_b or vector alone, treatment of these lines with salmon CT resulted in decreased proliferation, in the accumulation of cells in G₂ phase of the cell cycle in the hCT Receptor_a transfectant but not the hCT Receptor_b transfectant.⁸⁶ In the hCT Receptor_a transfectant, ERK^{1/2} activation mediates modulation of the cell cycle *via* p21^{Cip1}. However,

G₂-arrest was unexpected as p21^{Cip} has been shown to inhibit CDK2 rather than CDK1 (cdc 2)⁹³ and results in a cycle block at G₁.^{94,95} This anomaly has never been resolved as far as we are aware and raises questions about the validity of CT Receptor transfectants to probe the real biological functions of CT Receptor.

In a subsequent report CT induced apoptosis in an hCT Receptor_a transfectant under conditions of low serum which were not observed with replete serum.⁹⁶ While one explanation might be a factor present in serum that overcomes the CT effect, another possibility is metabolic reprogramming driven by starvation and cell stress associated with autophagy which is independent of caspase 3.⁹⁶ In this regard, in a *p53*^{-/-} mouse model with thymic lymphoma, CT Receptor is essential for the transmission of the effects of amylin for metabolic reprogramming, induction of apoptosis and tumour regression.⁹⁷

SiRNA mediated knockdown of CT Receptor in TT2609-C02 cells derived from follicular thyroid carcinoma, resulted in G₁ arrest, a decrease in proliferation and an increase in caspase 3 activity.⁷² The authors conclude that CT Receptor is one of several genes important for survival of thyroid carcinomas and therefore can be classified as a putative oncogene.

In human prostate cancer cell lines and mouse models, CT and CT Receptor activate survival of cells following cytotoxic insult⁶⁸ and enhanced tumour growth⁹⁸ consistent with the induction of apoptosis following knockdown of CT Receptor.⁹⁹

Consistent with these results are further reports that CT prevents apoptosis in osteocytes and osteoblasts¹⁰⁰ and promotes survival of osteoclasts.¹⁰¹

Overall in the context of different cell lines maintained in normal serum (unstarved), CT/CT Receptor promotes cell survival and proliferation, while promotion of apoptosis is associated with P53 status, metabolic reprogramming and nutrient deprivation.

Stem cell quiescence and a role for a surrogate ligand of CT Receptor

In skeletal muscle stem (satellite) cells maintenance of quiescence is dependent on activation of CT Receptor⁵⁴ and it is now proposed that an active cell autonomous Notch–Collagen V (COL

V)–CT Receptor axis maintains the quiescent muscle stem cells in their niche.⁵⁵ Notch also contributes to the stem-like character of glioma cells.¹⁰²

The mechanism of how COL V acts as a surrogate ligand of CT Receptor is yet to be described. We have modelled [unpublished] the carboxy cleaved peptide of collagen V and found similarities when arranged as a 3₁₀ helix with salmon CT. In this case the carboxyl tail of nascent COL V might act as a surrogate ligand for CT Receptor after it is cleaved. However, the researchers⁵⁵ used a commercial collagen V isolated from human placenta and it is unclear whether this is pure mature collagen V or contains pre-pro-collagen V and processed peptides.⁵⁵ If this work can be corroborated this discovery will have a profound effect on our understanding of CT Receptor as expressed in the context of many tissues. As presented in Table 1, there is a wide expression of CT Receptor throughout the life cycle in many tissues, sometimes expression is transient and sometimes persistent, and expression is found in cells that play a prominent role in a range of diseases.

For instance, this axis might well be a driving factor in atherosclerosis in which we have identified expression of CT Receptor^{63,64,103,104} and expression of collagen V is also known.^{103,104}

Formation of granulation tissue is a histological hallmark of wound healing following tissue injury.¹⁰⁵ COLV immunostaining predominates in the blood vessel walls of the granulation tissue and COLV transcript is expressed in the fibroblast-like cells of the granulation tissue.^{105,106} Similar cells and blood vessels express CT Receptor in a mouse model of wound healing.³¹

Genotyped profiles of GBM

Extensive genetic profiles of GBM patient biopsies have been published,^{10–12} including regions of chromosomal aberrations both broad and focal¹⁰ together with the somatic genomic landscape.¹²

Gliomas with broad amplification of chromosome 7 have properties different from those with overlapping focal *EGFR* amplification (chr7p11) and the broad events around chr7q32 (*CALCR* gene location chr 7q21.3) which appear to act in part through effects on *MET* (CXCR4) and its ligand *HGF* and correlate with *MET* dependence *in vitro*.¹⁰ It should be noted that *CALCR* maps close to *MET*.

Alternative splicing has also been recognised as a driving force of tumorigenesis.¹⁰⁷ Aberrant profiles observed in tumours mostly reflect the selection of endogenous alternative splicing variants with different functional properties that allow the malignant progression of initiated tumour cells. Selected functions relate, for example, to sustained proliferation, evasion of apoptosis, metabolic adaptation, or angiogenesis.¹⁰⁸ Results from our group investigating the U87 MG cell line have demonstrated increased frequency of exon 10 (insert-positive sequence) splicing-in when these cells are stressed with cytotoxin treatment.⁸¹ The possibility of other splicing events that might result in receptor inactivation are currently under investigation.

In a study on oncocytic thyroid carcinomas⁷² hemi-methylation in the region of the 5'UTR of *CALCR* was characterised, upregulation of *CALCR* mRNA was found in the majority of samples and in the cell line TT2609-C02 reduction in the expression of CTR leads to a pause in the cell cycle, upregulation of caspase 3 mRNA and enhanced cell death. Consistently, there was no loss of heterozygosity of chromosome 7 which was interpreted as a role for maternal and paternal genes imprinted on this chromosome including *CALCR*. These data support a role for CT Receptor as an oncogene.

Regulation of *CALCR* transcription and stability of mRNA

The structure of human *CALCR* gene has been described in BIN-67 cells (human ovarian carcinoma) cell line,⁷⁷ human osteoclasts and MCF-7 cells (human breast adenocarcinoma).¹⁰⁹ Promoters P1 and P2 were demonstrated in transfected T47D cells (human breast cancer)^{110,111} and a further promoter (POc) is specifically active in human osteoclasts.¹¹² This paper also described the tissue-specific splicing of the *CALCR* 5'UTR and the transcripts that are generated. Analysis of the 5'UTR of human *CALCR* reveals multiple Sp1 binding sites¹¹³ and CpG islands. The regulation by Sp1 is of particular interest because many genes upregulated during stress are also regulated by this transcription factor which has been proposed to drive the adaptive response of cancer cells to hypoxia.¹¹⁴

The structure of the murine *CALCR* gene from brain has been determined¹¹⁵ and later *CALCR* mRNA shown to be transcribed from three promoters (P1, P2 and P3) of which P3 is osteoclast

specific and the promoters become functional through alternative splicing of exons in the 5'UTR.¹¹⁶ A comparison of the exon structures in human and mouse *CALCR* has been published.¹¹² The structure of porcine *CALCR* gene has been reported from LLC-PK1 cells.¹¹⁷

In the 3'UTR of human *CALCR* mRNA there are eight AUUUA sequences and five in porcine *CALCR*.¹¹⁸ These sequences have been demonstrated to increase the instability of transcripts.^{119,120} *CALCR* mRNA levels are generally low in tissues and HGG cell lines³⁰ perhaps reflecting the inherent instability of this mRNA due to multiple (AUUUA) sequences. This instability could explain the low representation of *CALCR* mRNA in some genetic studies of GBM patients.¹¹

Targeting inactivated CT Receptor in GBM

The pharmacology of responses to CT ligands was reported in these four HGG cell lines.³⁰ Only SB2b responded to CT-like ligands (human CT, salmon CT, rat amylin) indicating that CT Receptor is non-functional in the cell lines JK2, PB1 and WK1. It is likely that this inactivation results from alternative splicing common in cancers including the possibility that expression of CT Receptor_b predominates. A conclusion drawn from this study is that pharmacological intervention of CT Receptor is unlikely to provide an avenue for treatment of GBM.

Our group⁶⁵ has published a study with HGG cell lines which characterises the potency of an anti-hCT Receptor antibody conjugated to cytotoxins monomethyl-aurostatin E (MMAE) or the plant toxins diathin-30¹²¹ and gelonin (ribosome-inactivating proteins). The anti-hCT Receptor antibody deployed in the ADC or immunotoxin binds an extracellular epitope and is internalised. The potency of the immunotoxin (EC₅₀ 10–20 pM) is greatly increased with saponins (triterpene glycoside SO1861) which enhance the release of the toxin from the acidic lysosomes.⁶⁵

Three of the HGG cell lines (JK2, SB2b, WK1, all classic/mesenchymal)²⁷ were equally sensitive to the immunotoxin and expressed high levels of the CT Receptor protein in the membrane fraction as determined by immunoblot³⁰ using an anti-CT Receptor antibody that recognises an intracellular epitope. Although all four HGG cell lines displayed similar levels of CT Receptor

using whole cell lysates,⁶⁵ one HGG cell line PB1 (proneural) expresses low levels of CT Receptor in the membrane fraction³⁰ and is relatively resistant to the immunotoxin.⁶⁵

These data suggest that a potential treatment of GBM stem-like cells is possible using immunotoxin directed against CT Receptor regardless of the pharmacological status of the receptor.

The proposed mechanism⁶⁵ of this therapy (refer to Figure 5) involves binding to the extracellular domain of CT Receptor on the plasma membrane, uptake of the immunotoxin *via* the endosomes into the lysosomes (with low pH) and cleavage of the immunotoxin (either protease or pH mediated depending on the linker). The escape of the toxin from the lysosomal compartment⁶⁵ is enhanced by the triterpene glycoside SO1861.¹²² The toxin targets the ribosomes.¹²¹

Whether the HGG cell lines express CT Receptor_a, CT Receptor_b or an alternatively spliced isoform is yet to be established. From our current analysis elevated CT Receptor expression in the membrane fraction is necessary for sensitivity to the immunotoxin as PB1 is deficient in this respect and is resistant to the immunotoxin.

Animal studies are planned to test the immunotoxins *in vivo* and further refinements of the prototype immunotoxin, to improve tumour penetration, are also in progress.

Evidence for CTR as a tumour suppressor and/or an oncogene

Tumour suppressor genes regulate a diverse range of cellular activities including cell cycle checkpoint responses and mitogenic signaling.¹²³ Classic tumour suppressors have three principal characteristics, firstly they are recessive and undergo biallelic inactivation in tumours, secondly inheritance of a mutant allele potentiates tumour initiation, and thirdly, the same gene is frequently inactivated in sporadic tumours.¹²³

CT Receptor has been shown to influence the cell cycle with induction of quiescence (G₀) for satellite stem cells and checkpoint G₁ for cell lines as discussed above. Two reports have identified inactivated CT Receptor expressed in GBM.^{30,66} As CT Receptor is upregulated with cell stress⁶¹ and ligand activation results in apoptosis (discussed above) then inactivation of CT Receptor by mutation or




Figure 5. Proposed mechanism of CT Receptor_b in the excretion of cell debris and cytotoxins *via* exosomes.

alternative splicing (U87 MG, discussed above) could result in survival of tumour cells. It should be noted at this point that U87 MG might not have originated from GBM.¹²⁴ CT Receptor protein is expressed by HGG cell lines in which CT Receptor activation is either non-canonical or inhibited by mutation or deletion, or the CT Receptor_b isoform is preferentially expressed.

Pal *et al.*⁶⁶ describe the downregulated levels of *CALCR* transcripts in GBMs from data sets TCGA, GSE7696 and the Indian cohorts. However, as described above, low levels might be expected to result as the transcripts include eight repeat AUUUA sequences in the 3'UTR that are responsible for instability. The steady state levels are not an indication of the rates of transcription.

They also found altered activity of mutated CT Receptor derived from a small cohort⁶⁶ of GBM patients and drew conclusions about survival compared to normal CT Receptor. While much of the data are not conclusive, the idea that CT Receptor might act as a tumour suppressor and/or oncogene warrants further exploration.

Evidence for a role as an oncogene includes knockdown of CT Receptor in tumour cell lines as outlined above, which results in apoptosis and suggests, in the case of mutant CT Receptor, a further function of ligand-insensitive CT Receptor for tumour survival as proposed in Figure 5. It remains to be demonstrated that knockdown of inactivated (mutant) CT Receptor leads to cell death in the case of GBM.

Conclusion

CT Receptor is a G protein-coupled receptor (GPCR) that is highly expressed in patient biopsies between 76% (*CALCR* mRNA) and 86% (CT Receptor protein). Furthermore, it is expressed by malignant glioma cells, in cells that express markers of brain tumour initiating cells and 42% of HGG cell lines that represent tumour stem cells.

Quiescent cancer stem cells represent minimal resistant disease and enter the cell cycle in response to treatment to re-establish tumour malignancy. We postulate that the maintenance of quiescence results from activation of the Notch–COL V–CT Receptor axis as has been proposed for skeletal muscle satellite (stem) cells.

There are likely to be several potential mechanisms for the upregulation of CT Receptor in GBM. Firstly, the region around chromosome 7q21.3 (*CALCR* gene) is frequently amplified. Secondly, the transcription factor Sp1 regulates genes important for stress responses in many tissues including GBM and Sp1 has been shown to stimulate *CALCR* mRNA transcription in cell lines.

The potential role of the CT Receptor_b isoform is discussed in a mechanism that provides resistance to cytotoxins and drug resistance in glioblastoma and possibly other cancers.

In HGG cell lines CT Receptor is frequently pharmacologically inactive although the protein is detected by immunoblotting. Inactivation might result either by inactivating mutations/deletions/

insertions or alternative splicing resulting in inactivation or a switch to the CT Receptor_b isoform. The inactivation of CT Receptor would be consistent with its role as a tumour suppressor. Furthermore, knockdown of CT Receptor promotes apoptosis consistent with a role as an oncogene.

The antibody (mAb2C4) that binds the extracellular epitope of human CT Receptor has been developed as an immunotoxin to study its potency with HGG cell lines. The mAb2C4:dianthin immunotoxin has an effective EC₅₀ of 10–20pM as compared to an equivalent ADC (mAb2C4:MMAE) which is 250 times less potent.

Targeting CT Receptor expressed by quiescent cancer stem cells with the immunotoxin is expected to provide an additional weapon, in combination with traditional therapy that targets dividing malignant glioma cells, for the treatment of glioblastoma. This novel therapy aims to eradicate minimal residual disease associated with this cancer.

Acknowledgements

The authors would like to thank Apop Biosciences Pty Ltd for access to the mAb2C4 antibody (CalRexin).

Conflict of interest statement

PJW and DLH own shares in Apop Biosciences (formerly Apop Imaging) which owns intellectual property around the anti-CTR antibodies mAb2C4 and mAb9B4. The other authors declare no conflict of interest.

Funding

The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: SGBF is an ARC Future Fellow (FT180100543).

ORCID iD

Lucas Bittencourt  <https://orcid.org/0000-0002-4815-0548>

References

1. Das S and Marsden PA. Angiogenesis in glioblastoma. *N Engl J Med* 2013; 369: 1561–1563.
2. Stupp R, Mason WP, van den Bent MJ, *et al.* Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *N Engl J Med* 2005; 352: 987–996.

3. Kreso A, O'Brien CA, van Galen P, *et al.* Variable clonal repopulation dynamics influence chemotherapy response in colorectal cancer. *Science* 2013; 339: 543–548.
4. Raff MC, Miller RH and Noble M. A glial progenitor cell that develops in vitro into an astrocyte or an oligodendrocyte depending on culture medium. *Nature* 1983; 303: 390–396.
5. Noble M, Gutowski N, Bevan K, *et al.* From rodent glial precursor cell to human glial neoplasia in the oligodendrocyte-type-2 astrocyte lineage. *Glia* 1995; 15: 222–230.
6. Cheng L, Huang Z, Zhou W, *et al.* Glioblastoma stem cells generate vascular pericytes to support vessel function and tumor growth. *Cell* 2013; 153: 139–152.
7. Wang R, Chadalavada K, Wilshire J, *et al.* Glioblastoma stem-like cells give rise to tumour endothelium. *Nature* 2010; 468: 829–833.
8. Ricci-Vitiani L, Pallini R, Biffoni M, *et al.* Tumour vascularization via endothelial differentiation of glioblastoma stem-like cells. *Nature* 2010; 468: 824–828.
9. Soda Y, Marumoto T, Friedmann-Morvinski D, *et al.* Transdifferentiation of glioblastoma cells into vascular endothelial cells. *Proc Natl Acad Sci* 2011; 108: 4274–4280.
10. Beroukhi R, Getz G, Nghiemphu L, *et al.* Assessing the significance of chromosomal aberrations in cancer: methodology and application to glioma. *Proc Natl Acad Sci* 2007; 104: 20007–20012.
11. Cancer Genome Atlas Research Network. Comprehensive genomic characterization defines human glioblastoma genes and core pathways. *Nature* 2008; 455: 1061–1068.
12. Brennan CW, Verhaak RG, McKenna A, *et al.* The somatic genomic landscape of glioblastoma. *Cell* 2013; 155: 462–477.
13. Lee JK, Wang J, Sa JK, *et al.* Spatiotemporal genomic architecture informs precision oncology in glioblastoma. *Nat Genet* 2017; 49: 594–599.
14. Dore-Duffy P, Mehedi A, Wang X, *et al.* Immortalized CNS pericytes are quiescent smooth muscle actin-negative and pluripotent. *Microvasc Res* 2011; 82: 18–27.
15. Sharma V, Ling TW, Rewell SS, *et al.* A novel population of alpha-smooth muscle actin-positive cells activated in a rat model of stroke: an analysis of the spatio-temporal distribution in response to ischemia. *J Cereb Blood Flow Metab* 2012; 32: 2055–2065.
16. Puchalski RB, Shah N, Miller J, *et al.* An anatomic transcriptional atlas of human glioblastoma. *Science* 2018; 360: 660–663.
17. Singh SK, Clarke ID, Terasaki M, *et al.* Identification of a cancer stem cell in human brain tumors. *Cancer Res* 2003; 63: 5821–5828.
18. Chan KS. Molecular pathways: targeting cancer stem cells awakened by chemotherapy to abrogate tumor repopulation. *Clin Cancer Res* 2016; 22: 802–806.
19. Lathia JD, Heddleston JM, Venere M, *et al.* Deadly teamwork: neural cancer stem cells and the tumor microenvironment. *Cell Stem Cell* 2011; 8: 482–485.
20. Singh SK, Hawkins C, Clarke ID, *et al.* Identification of human brain tumour initiating cells. *Nature* 2004; 432: 396–401.
21. Rath BH, Fair JM, Jamal M, *et al.* Astrocytes enhance the invasion potential of glioblastoma stem-like cells. *PLoS One* 2013; 8: e54752.
22. Calabrese C, Poppleton H, Kocak M, *et al.* A perivascular niche for brain tumor stem cells. *Cancer Cell* 2007; 11: 69–82.
23. Galli R, Binda E, Orfanelli U, *et al.* Isolation and characterization of tumorigenic, stem-like neural precursors from human glioblastoma. *Cancer Res* 2004; 64: 7011–7021.
24. Lee J, Kotliarova S, Kotliarov Y, *et al.* Tumor stem cells derived from glioblastomas cultured in bFGF and EGF more closely mirror the phenotype and genotype of primary tumors than do serum-cultured cell lines. *Cancer Cell* 2006; 9: 391–403.
25. Verhaak RG, Hoadley KA, Purdom E, *et al.* Integrated genomic analysis identifies clinically relevant subtypes of glioblastoma characterized by abnormalities in PDGFRA, IDH1, EGFR, and NF1. *Cancer Cell* 2010; 17: 98–110.
26. Day BW, Stringer BW, Al-Ejeh F, *et al.* EphA3 maintains tumorigenicity and is a therapeutic target in glioblastoma multiforme. *Cancer Cell* 2013; 23: 238–248.
27. Stringer BW, Day BW, D'Souza RCJ, *et al.* A reference collection of patient-derived cell line and xenograft models of proneural, classical and mesenchymal glioblastoma. *Scientific Rep* 2019; 9: 4902.
28. Lutz C, Woll PS, Hall G, *et al.* Quiescent leukaemic cells account for minimal residual disease in childhood lymphoblastic leukaemia. *Leukemia* 2013; 27: 1204–1207.
29. Wookey PJ, McLean CA, Hwang P, *et al.* The expression of calcitonin receptor detected in malignant cells of the brain tumour glioblastoma

- multiforme and functional properties in the cell line A172. *Histopathology* 2012; 60: 895–910.
30. Ostrovskaya A, Hick C, Hutchinson DS, *et al.* Expression and activity of the calcitonin receptor family in a sample of primary human high-grade gliomas. *BMC Cancer* 2019; 19: 157–168.
 31. Wookey P, Zulli A, Lo C, *et al.* Calcitonin receptor (CTR) expression in embryonic, foetal and adult tissues: developmental and pathophysiological implications. In: Hay D and Dickerson I (eds) *The calcitonin gene-related peptide family; form, function and future perspectives*. Netherlands: Springer, 2010, pp.199–233.
 32. Wang J, Rout UK, Bagchi IC, *et al.* Expression of calcitonin receptors in mouse preimplantation embryos and their function in the regulation of blastocyst differentiation by calcitonins. *Development* 1998; 125: 4293–4302.
 33. Burgess AM. The effect of calcitonin on the prechordal mesoderm, neural plate and neural crest of *Xenopus* embryos. *J Anat* 1985; 140: 49–55.
 34. Jagger C, Chambers T and Pondel M. Transgenic mice reveal novel sites of calcitonin receptor gene expression during development. *Biochem Biophys Res Commun* 2000; 274: 124–129.
 35. Jagger C, Gallagher A, Chambers T, *et al.* The porcine calcitonin receptor promoter directs expression of a linked reporter gene in a tissue and developmental specific manner in transgenic mice. *Endocrinology* 1999; 140: 492–499.
 36. Tolcos M, Tikellis C, Rees S, *et al.* Ontogeny of calcitonin receptor mRNA and protein in the developing central nervous system of the rat. *J Comp Neurol* 2003; 456: 29–38.
 37. Wookey PJ, Turner K and Furness JB. Transient expression of the calcitonin receptor by enteric neurons of the embryonic and early post-natal mouse. *Cell Tis Res* 2012; 347: 311–317.
 38. Tikellis C, Xuereb L, Casley D, *et al.* Calcitonin receptor isoforms expressed in the developing rat kidney. *Kidney Int* 2003; 63: 416–426.
 39. Fischer JA, Tobler PH, Kaufmann M, *et al.* Calcitonin: regional distribution of the hormone and its binding sites in the human brain and pituitary. *Proc Natl Acad Sci* 1981; 78: 7801–7805.
 40. Sexton PM, McKenzie JS and Mendelsohn FAO. Evidence for a new subclass of calcitonin/calcitonin gene-related peptide binding sites in rat brain. *Neurochem Int* 1988; 12: 323–335.
 41. Becskei C, Riediger T, Zund D, *et al.* Immunohistochemical mapping of calcitonin receptors in the adult rat brain. *Brain Res* 2004; 1030: 221–233.
 42. Paxinos G, Chai SY, Christopoulos G, *et al.* In vitro autoradiographic localization of calcitonin and amylin binding sites in monkey brain. *J Chem Neuroanat* 2004; 27: 217–236.
 43. Goda T, Doi M, Umezaki Y, *et al.* Calcitonin receptors are ancient modulators for rhythms of preferential temperature in insects and body temperature in mammals. *Genes Dev* 2018; 32: 140–155.
 44. Walker CS, Eftekhari S, Bower RL, *et al.* A second trigeminal CGRP receptor: function and expression of the AMY1 receptor. *Ann Clin Transl Neurol* 2015; 2: 595–608.
 45. Marx SJ, Woodward CJ and Aurbach GD. Calcitonin receptors of kidney and bone. *Science* 1972; 178: 999–1001.
 46. Nicholson GC, Moseley JM, Sexton PM, *et al.* Abundant calcitonin receptors in isolated rat osteoclasts. Biochemical and autoradiographic characterization. *J Clin Invest* 1986; 78: 355–360.
 47. Hattersley G and Chalmers TJ. Calcitonin receptors as markers for osteoclast differentiation: correlation between generation of bone-resorptive cells and cells that express calcitonin receptors in mouse bone marrow cultures. *Endocrinology* 1989; 125: 1606–1612.
 48. Zaidi M, Pazianas M, Shankar VS, *et al.* Osteoclast function and its control. *Exp Physiol* 1993; 78: 721–739.
 49. Gooi JH, Pompolo S, Karsdal MA, *et al.* Calcitonin impairs the anabolic effect of PTH in young rats and stimulates expression of sclerostin by osteocytes. *Bone* 2010; 46: 1486–1497.
 50. Sexton P, Adam W, Moseley J, *et al.* Localization and characterization of renal calcitonin receptors by *in vitro* autoradiography. *Kidney Int* 1987; 32: 862–868.
 51. Hanna FW, Smith DM, Johnston CF, *et al.* Expression of a novel receptor for the calcitonin peptide family and a salmon calcitonin-like peptide in the alpha-thyrotropin thyrotroph cell line. *Endocrinology* 1995; 136: 2377–2382.
 52. Fukada S, Uezumi A, Ikemoto M, *et al.* Molecular signature of quiescent satellite cells in adult skeletal muscle. *Stem Cells* 2007; 25: 2448–2459.
 53. Gnocchi VF, White RB, Ono Y, *et al.* Further characterisation of the molecular signature of quiescent and activated mouse muscle satellite cells. *PLoS One* 2009; 4: e5205.

54. Yamaguchi M, Watanabe Y, Ohtani T, *et al.* Calcitonin receptor signaling inhibits muscle stem cells from escaping the Quiescent State and the Niche. *Cell Rep* 2015; 13: 302–314.
55. Baghdadi MB, Castel D, Machado L, *et al.* Reciprocal signalling by Notch-Collagen V- CALCR retains muscle stem cells in their niche. *Nature* 2018; 557: 714–718.
56. Silvestroni L, Menditto A, Frajese G, *et al.* Identification of calcitonin receptors in human spermatozoa. *J Clin Endocrinol Metab* 1987; 65: 742–746.
57. Adeoya-Osiguwa SA and Fraser LR. Calcitonin acts as a first messenger to regulate adenylyl cyclase/cAMP and mammalian sperm function. *Mol Reprod Dev* 2003; 65: 228–236.
58. Nicholson GC, D’Santos CS, Evans T, *et al.* Human placental calcitonin receptors. *Biochem J* 1988; 250: 877–882.
59. Kovacs C, Chafe L, Woodland M, *et al.* Calcitropic gene expression in the murine placenta suggests a role for the intraplacental yolk sac in maternal-fetal calcium exchange. *Am J Physiol* 2002; 282: E721–E732.
60. Wu G, Burzon DT, di Sant’Agnese PA, *et al.* Calcitonin receptor mRNA expression in human prostate. *Urology* 1996; 47: 376–381.
61. Furness SGB, Hare DL, Kourakis A, *et al.* A novel ligand of calcitonin receptor reveals a potential new sensor that modulates programmed cell death. *Cell Death Disc* 2016; 2: 16062.
62. Meeuwsen S, Persoon-Deen C, Bsibsi M, *et al.* Cytokine, chemokine and growth factor gene profiling of cultured human astrocytes after exposure to proinflammatory stimuli. *Glia* 2003; 43: 243–253.
63. Wookey PJ, Zulli A, Buxton BF, *et al.* Calcitonin receptor immunoreactivity associated with specific cell types in diseased radial and internal mammary arteries. *Histopathology* 2008; 52: 605–612.
64. Wookey PJ, Zulli A and Hare DL. The elevated expression of calcitonin receptor by cells recruited into the endothelial layer and neointima of atherosclerotic plaque. *Histochem Cell Biol* 2009; 132: 181–189.
65. Gilabert-Oriol R, Furness SGB, Stringer BW, *et al.* Dianthin-30 or gelonin versus monomethyl auristatin E, each configured with an anti-calcitonin receptor antibody, are differentially potent in vitro in high grade glioma cell lines derived from glioblastoma. *Cancer Immunol Immunother* 2017; 66: 1217–1228.
66. Pal J, Patil V, Kumar A, *et al.* Loss-of-function mutations in Calcitonin Receptor (CALCR) identify highly aggressive glioblastoma with poor outcome. *Clin Cancer Res* 2018; 24: 1448–1458.
67. Ritchie CK, Thomas KG, Andrews LR, *et al.* Effects of the calcitrophic peptides calcitonin and parathyroid hormone on prostate cancer growth and chemotaxis. *Prostate* 1997; 30: 183–187.
68. Thomas S, Chigurupati S, Anbalagan M, *et al.* Calcitonin increases tumorigenicity of prostate cancer cells: evidence for the role of protein kinase A and urokinase-type plasminogen receptor. *Mol Endocrinol* 2006; 20: 1894–1911.
69. Frenzo JL, Delage-Mourroux R, Cohen R, *et al.* Calcitonin receptor mRNA is expressed in human medullary thyroid carcinoma. *Thyroid* 1998; 8: 141–147.
70. Frenzo JL, Pichaud F, Mourroux RD, *et al.* An isoform of the human calcitonin receptor is expressed in TT cells and in medullary carcinoma of the thyroid. *FEBS Lett* 1994; 342: 214–216.
71. Cappagli V, Potes CS, Ferreira LB, *et al.* Calcitonin receptor expression in medullary thyroid carcinoma. *PeerJ* 2017; 5: e3778.
72. Boot A, Oosting J, de Miranda NF, *et al.* Imprinted survival genes preclude loss of heterozygosity of chromosome 7 in cancer cells. *J Pathol* 2016; 240: 72–83.
73. Nicholson GC, Horton MA, Sexton PM, *et al.* Calcitonin receptors of human osteoclastoma. *Horm Metab Res* 1987; 19: 585–589.
74. Gorn AH, Rudolph SM, Flannery MR, *et al.* Expression of two human skeletal calcitonin receptor isoforms cloned from a giant cell tumor of bone. The first intracellular domain modulates ligand binding and signal transduction. *J Clin Invest* 1995; 95: 2680–2691.
75. Silvestris F, Cafforio P, De Matteo M, *et al.* Expression and function of the calcitonin receptor by myeloma cells in their osteoclast-like activity in vitro. *Leuk Res* 2008; 32: 611–623.
76. Mould R and Pondel MD. Calcitonin receptor gene expression in K562 chronic myelogenous leukemic cells. *Cancer Cell Int* 2003; 3: 6–12.
77. Gorn AH, Lin HY, Yamin M, *et al.* Cloning, characterization, and expression of a human calcitonin receptor from an ovarian carcinoma cell line. *J Clin Invest* 1992; 90: 1726–1735.
78. Findlay DM, Michelangeli VP, Moseley JM, *et al.* Calcitonin binding and degradation by two cultured human breast cancer cell lines (MCF 7 and T 47D). *Biochem J* 1981; 196: 513–520.

79. Gillespie M, Thomas R, Pu Z, *et al.* Calcitonin receptors, bone sialoprotein and osteopontin are expressed in primary breast cancers. *Intern J Cancer* 1997; 73: 812–815.
80. Kuestner RE, Elrod RD, Grant FJ, *et al.* Cloning and characterization of an abundant subtype of the human calcitonin receptor. *Mol Pharmacol* 1994; 46: 246–255.
81. Furness SGB, Gupta P, Zhong M, *et al.* Strategy to develop and validate antibodies against linear epitopes of human calcitonin receptor: identification of insert-positive isoforms and ubiquity across mammalian species suggests new physiological functions. Unpublished data.
82. Dal Maso E, Just R, Hick C, *et al.* Characterization of signalling and regulation of common calcitonin receptor splice variants and polymorphisms. *Biochem Pharmacol* 2018; 148: 111–129.
83. Beaudreuil J, Balasubramanian S, Chenais J, *et al.* Molecular characterization of two novel isoforms of the human calcitonin receptor. *Gene* 2004; 343: 143–151.
84. Liang YL, Khoshouei M, Radjainia M, *et al.* Phase-plate cryo-EM structure of a class B GPCR-G-protein complex. *Nature* 2017; 546: 118–123.
85. Cohen DP, Thaw CN, Varma A, *et al.* Human calcitonin receptors exhibit agonist-independent (constitutive) signaling activity. *Endocrinology* 1997; 138: 1400–1405.
86. Raggatt LJ, Evdokiou A and Findlay DM. Sustained activation of Erk1/2 MAPK and cell growth suppression by the insert-negative, but not the insert-positive isoform of the human calcitonin receptor. *J Endocrinol* 2000; 167: 93–105.
87. Furness SGB, Liang YL, Nowell CJ, *et al.* Ligand-dependent modulation of G protein conformation alters drug efficacy. *Cell* 2016; 167: 739–749.e711.
88. Dermer SJ, Cohen DP, Thaw CN, *et al.* Intracellular retention and rapid degradation of human calcitonin receptors overexpressed in COS cells. *Endocrinology* 1996; 137: 5502–5508.
89. Seck T, Baron R and Horne WC. Binding of filamin to the C-terminal tail of the calcitonin receptor controls recycling. *J Biol Chem* 2003; 278: 10408–10416.
90. Safaei R, Larson BJ, Cheng TC, *et al.* Abnormal lysosomal trafficking and enhanced exosomal export of cisplatin in drug-resistant human ovarian carcinoma cells. *Mol Cancer Ther* 2005; 4: 1595–1604.
91. Ng KW, Livesey SA, Larkins RG, *et al.* Calcitonin effects on growth and on selective activation of type II isoenzyme of cyclic adenosine 3':5'-monophosphate-dependent protein kinase in T 47D human breast cancer cells. *Cancer Res* 1983; 43: 794–800.
92. Dayer JM, Vassalli JD, Bobbitt JL, *et al.* Calcitonin stimulates plasminogen activator in porcine renal tubular cells: LLC-PK1. *J Cell Biol* 1981; 91: 195–200.
93. Evdokiou A, Raggatt LJ, Atkins GJ, *et al.* Calcitonin receptor-mediated growth suppression of HEK-293 cells is accompanied by induction of p21WAF1/CIP1 and G2/M arrest. *Mol Endocrinol* 1999; 13: 1738–1750.
94. LaBaer J, Garrett MD, Stevenson LF, *et al.* New functional activities for the p21 family of CDK inhibitors. *Genes Dev* 1997; 11: 847–862.
95. Santra M, Mann DM, Mercer EW, *et al.* Ectopic expression of decorin protein core causes a generalized growth suppression in neoplastic cells of various histogenetic origin and requires endogenous p21, an inhibitor of cyclin-dependent kinases. *J Clin Invest* 1997; 100: 149–157.
96. Findlay DM, Raggatt LJ, Bouralexis S, *et al.* Calcitonin decreases the adherence and survival of HEK-293 cells by a caspase-independent mechanism. *J Endocrinol* 2002; 175: 715–725.
97. Venkatanarayan A, Raulji P, Norton W, *et al.* IAPP-driven metabolic reprogramming induces regression of p53-deficient tumours in vivo. *Nature* 2015; 517: 626–630.
98. Shah GV, Thomas S, Muralidharan A, *et al.* Calcitonin promotes in vivo metastasis of prostate cancer cells by altering cell signaling, adhesion, and inflammatory pathways. *Endocr Relat Cancer* 2008; 15: 953–964.
99. Thomas S, Muralidharan A and Shah GV. Knock-down of calcitonin receptor expression induces apoptosis and growth arrest of prostate cancer cells. *Int J Oncol* 2007; 31: 1425–1437.
100. Plotkin LI, Weinstein RS, Parfitt AM, *et al.* Prevention of osteocyte and osteoblast apoptosis by bisphosphonates and calcitonin. *J Clin Invest* 1999; 104: 1363–1374.
101. Selander KS, Harkonen PL, Valve E, *et al.* Calcitonin promotes osteoclast survival in vitro. *Mol Cell Endocrinol* 1996; 122: 119–129.
102. Shih AH and Holland EC. Notch signaling enhances nestin expression in gliomas. *Neoplasia* 2006; 8: 1072–1082.
103. Morton LF and Barnes MJ. Collagen polymorphism in the normal and diseased blood

- vessel wall. Investigation of collagens types I, III and V. *Atherosclerosis* 1982; 42: 41–51.
104. Ooshima A. Collagen alpha B chain: increased proportion in human atherosclerosis. *Science* 1981; 213: 666–668.
 105. Mak KM, Png CY and Lee DJ. Type V collagen in health, disease, and fibrosis. *Anat Rec (Hoboken)* 2016; 299: 613–629.
 106. Inkinen K, Wolff H, Von Boguslawski K, *et al.* Type V collagen in experimental granulation tissue. *Connect Tiss Res* 1998; 39: 281–294.
 107. Golan-Gerstl R, Cohen M, Shilo A, *et al.* Splicing factor hnRNP A2/B1 regulates tumor suppressor gene splicing and is an oncogenic driver in glioblastoma. *Cancer Res* 2011; 71: 4464–4472.
 108. Goncalves V, Pereira JFS and Jordan P. Signaling pathways driving aberrant splicing in cancer cells. *Genes (Basel)* 2017; 9: 9.
 109. Nishikawa T, Ishikawa H, Yamamoto S, *et al.* A novel calcitonin receptor gene in human osteoclasts from normal bone marrow. *FEBS Lett* 1999; 458: 409–414.
 110. Hebden C, Smalt R, Chambers T, *et al.* Multiple promoters regulate human calcitonin receptor gene expression. *Biochem Biophys Res Commun* 2000; 272: 738–743.
 111. Pondel MD, Jagger C, Hebden C, *et al.* Transcriptional regulation of the calcitonin receptor gene. *Biochem Soc Trans* 2002; 30: 423–427.
 112. Shen Z, Crotti TN, Flannery MR, *et al.* A novel promoter regulates calcitonin receptor gene expression in human osteoclasts. *Biochim Biophys Acta* 2007; 1769: 659–667.
 113. Pondel MD, Partington GA and Mould R. Tissue-specific activity of the proximal human calcitonin receptor promoter is mediated by Sp1 and an epigenetic phenomenon. *FEBS Lett* 2003; 554: 433–438.
 114. Koizume S and Miyagi Y. Diverse mechanisms of Sp1-dependent transcriptional regulation potentially involved in the adaptive response of cancer cells to oxygen-deficient conditions. *Cancers* 2015; 8: 2.
 115. Yamin M, Gorn AH, Flannery MR, *et al.* Cloning and characterization of a mouse brain calcitonin receptor complementary deoxyribonucleic acid and mapping of the calcitonin receptor gene. *Endocrinology* 1994; 135: 2635–2643.
 116. Anusaksathien O, Laplace C, Li X, *et al.* Tissue-specific and ubiquitous promoters direct the expression of alternatively spliced transcripts from the calcitonin receptor gene. *J Biol Chem* 2001; 276: 22663–22674.
 117. Lin HY, Harris TL, Flannery MS, *et al.* Expression cloning of an adenylate cyclase-coupled calcitonin receptor. *Science* 1991; 254: 1022–1024.
 118. Zolnierowicz S, Cron P, Solinas-Toldo S, *et al.* Isolation, characterization, and chromosomal localization of the porcine calcitonin receptor gene. Identification of two variants of the receptor generated by alternative splicing. *J Biol Chem* 1994; 269: 19530–19538.
 119. Malter JS. Identification of an AUUUA-specific messenger RNA binding protein. *Science* 1989; 246: 664–666.
 120. Shaw G and Kamen R. A conserved AU sequence from the 3' untranslated region of GM-CSF mRNA mediates selective mRNA degradation. *Cell* 1986; 46: 659–667.
 121. Fuchs H. Dianthin and its potential in targeted tumor therapies. *Toxins (Basel)* 2019; 11: 592.
 122. von Mallinckrodt B, Thakur M, Weng A, *et al.* Dianthin-EGF is an effective tumor targeted toxin in combination with saponins in a xenograft model for colon carcinoma. *Future Oncol* 2014; 10: 2161–2175.
 123. Sherr CJ. Principles of tumor suppression. *Cell* 2004; 116: 235–246.
 124. Allen M, Bjerke M, Edlund H, *et al.* Origin of the U87MG glioma cell line: good news and bad news. *Sci Transl Med* 2016; 8: 354re353.



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Gupta, P; Furness, SGB; Bittencourt, L; Hare, DL; Wookey, PJ

Title:

Building the case for the calcitonin receptor as a viable target for the treatment of glioblastoma

Date:

2020-12-01

Citation:

Gupta, P., Furness, S. G. B., Bittencourt, L., Hare, D. L. & Wookey, P. J. (2020). Building the case for the calcitonin receptor as a viable target for the treatment of glioblastoma. THERAPEUTIC ADVANCES IN MEDICAL ONCOLOGY, 12, <https://doi.org/10.1177/1758835920978110>.

Persistent Link:

<http://hdl.handle.net/11343/272465>

File Description:

Published version

License:

CC BY-NC