

Transglycosylating -d-galactosidase and -l-fucosidase from *Paenibacillus* sp. 3179 from a hot spring in East Greenland

Thøgersen, Mariane S.; Christensen, Stefan J.; Jepsen, Morten; Pedersen, Lars H.; Stougaard, Peter

Published in:
MicrobiologyOpen

DOI:
[10.1002/mbo3.980](https://doi.org/10.1002/mbo3.980)

Publication date:
2020

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (APA):

Thøgersen, M. S., Christensen, S. J., Jepsen, M., Pedersen, L. H., & Stougaard, P. (2020). Transglycosylating -d-galactosidase and -l-fucosidase from *Paenibacillus* sp. 3179 from a hot spring in East Greenland. *MicrobiologyOpen*, 9(3), [e980]. <https://doi.org/10.1002/mbo3.980>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact rucforsk@ruc.dk providing details, and we will remove access to the work immediately and investigate your claim.

Transglycosylating β -D-galactosidase and α -L-fucosidase from *Paenibacillus* sp. 3179 from a hot spring in East Greenland

Mariane S. Thøgersen¹  | Stefan J. Christensen¹  | Morten Jepsen¹  |
Lars H. Pedersen²  | Peter Stougaard¹ 

¹University of Copenhagen, Frederiksberg C, Denmark

²Aalborg University, Aalborg, Denmark

Correspondence

Peter Stougaard, University of Copenhagen, Thorvaldsensvej 40, DK-1871 Frederiksberg C, Denmark.
Email: pst@envs.au.dk

Present address

Mariane S. Thøgersen and Peter Stougaard, Aarhus University, Roskilde, Denmark

Stefan J. Christensen, Roskilde University, Roskilde, Denmark

Morten Jepsen, Novo Nordisk A/S, Bagsværd, Denmark

Funding information

Seoul National University; Danish Council for Strategic Research, Grant/Award Number: 1308-00014B and 0652-20140008

Abstract

Thermal springs are excellent locations for discovery of thermostable microorganisms and enzymes. In this study, we identify a novel thermotolerant bacterial strain related to *Paenibacillus dendritiformis*, denoted *Paenibacillus* sp. 3179, which was isolated from a thermal spring in East Greenland. A functional expression library of the strain was constructed, and the library screened for β -D-galactosidase and α -L-fucosidase activities on chromogenic substrates. This identified two genes encoding a β -D-galactosidase and an α -L-fucosidase, respectively. The enzymes were recombinantly expressed, purified, and characterized using oNPG (2-nitrophenyl- β -D-galactopyranoside) and pNP-fucose (4-nitrophenyl- α -L-fucopyranoside), respectively. The enzymes were shown to have optimal activity at 50°C and pH 7–8, and they were able to hydrolyze as well as transglycosylate natural carbohydrates. The transglycosylation activities were investigated using TLC and HPLC, and the β -D-galactosidase was shown to produce the galactooligosaccharides (GOS) 6'-O-galactosyllactose and 3'-O-galactosyllactose using lactose as substrate, whereas the α -L-fucosidase was able to transfer the fucose moiety from pNP-fuc to lactose, thereby forming 2'-O-fucosyllactose. Since enzymes that are able to transglycosylate carbohydrates at elevated temperature are desirable in many industrial processes, including food and dairy production, we foresee the potential use of enzymes from *Paenibacillus* sp. 3179 in the production of, for example, instant formula.

KEYWORDS

galactooligosaccharides, *Paenibacillus*, transglycosylation, α -L-fucosidase, β -D-galactosidase

1 | INTRODUCTION

Thermal springs are hot spots of microbial biodiversity that may be utilized within agriculture, food and feed industry, textile industry, and biotechnology (Gupta, Gupta, Capalash, & Sharma, 2017; Mahajan & Balachandran, 2017; Sahay et al., 2017). Thermostable enzymes

from such environments have attracted increased interest from industry and academia due to the opportunities for high-temperature processes and interesting molecular structures (DeCastro, Rodríguez-Belmonte, & González-Siso, 2016; Elleuche, Schäfers, Blank, Schröder, & Antranikian, 2015; Mehta, Singhal, Singh, Damle, & Sharma, 2016; Ribeiro, Sánchez, Hidalgo, & Berenguer, 2017; Urbietta et al., 2015).

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2019 The Authors. *MicrobiologyOpen* published by John Wiley & Sons Ltd.

Especially within dairy production, processes at high or low temperatures are required due to low growth capabilities of food spoilers (Hervert, Martin, Boor, & Wiedmann, 2017; Ruusunen, Surakka, Korkeala, & Lindström, 2012). Furthermore, in some processes like oligosaccharide synthesis high temperatures are desired to obtain higher solubility of the reactants (Hassan et al., 2015; Yu & O'Sullivan, 2018). Galactooligosaccharides (GOS) in particular have been studied thoroughly (Singla & Chakkaravarthi, 2017). GOS are indigestible oligosaccharides, which have a documented positive effect on development and growth of beneficial gut bacteria such as bifidobacteria and lactobacilli in the human gastrointestinal tract (Gibson, Probert, Jan, Rastall, & Roberfroid, 2004; Knol, Scholtens, et al., 2005; Ladirat et al., 2014). Additionally, they have been shown to prevent adherence of pathogenic bacteria such as enteropathogenic *Escherichia coli* and *Salmonella enterica* to the cell walls of the colon (Knol, Boehm, et al., 2005; Searle et al., 2009; Shoaf, Mulvey, Armstrong, & Hutkins, 2006). Synthesis of oligosaccharides can be carried out by either chemical synthesis using acid hydrolysis, which is a relatively nonspecific, energy-demanding, and time-consuming process, or by enzymatic synthesis carried out by glycosyltransferases (EC 2.4) or glycoside hydrolases (EC 3.2.1). Glycoside hydrolases (GH) are the preferred catalysts due to high availability, natural hydrolytic function, ability to glycosylate a wide variety of hydroxyl group-containing molecules, and the use of cheap and simple donor substrates (Crout & Vic, 1998; de Roode, Franssen, Padt, & Boom, 2003; Watt, Lowden, & Flitsch, 1997). One of the cheapest donor substrates for GOS synthesis is lactose (β -D-galactopyranosyl-(1 \rightarrow 4)-D-glucopyranose) which is a by-product from cheese production. Great efforts have been spent on converting this low-price disaccharide into carbohydrates of higher value, especially enzymatic conversion of lactose to lactobionic acid (Goderska, Szwengiel, & Czarnecki, 2014; Gutiérrez, Hamoudi, & Belkacemi, 2012), to the sweetener D-tagatose (Jørgensen, Hansen, & Stougaard, 2004; Wanarska & Kur, 2012; Zhan et al., 2014) and to GOS (Arreola et al., 2014; Jørgensen, Hansen, & Stougaard, 2001; Splechtna et al., 2006). Synthesis of GOS through enzymatic transglycosylation results in linear oligomers of galactose ($n = 2$ – 6) joined by β 1-2, β 1-3, β 1-4, or β 1-6 linkages with a glucose at the reducing end (Ishikawa et al., 2015; Li et al., 2009; Torres, Gonçalves, Teixeira, & Rodrigues, 2010; Wu et al., 2013). Due to the risk of contamination by food-spoiling bacteria during GOS production, focus has been on enzymes that are active at either high or low temperatures. Cold-active enzymes that are able to catalyze the formation of GOS have been described from, for example, *Pseudoalteromonas* sp. 22b (Cieśliński et al., 2005) and *Alkalilactibacillus ikkensis* (Schmidt & Stougaard, 2010), whereas thermostable GOS-producing enzymes have been described from, for example, *Sulfolobus* spp. (Park, Kim, Lee, & Oh, 2010; Park, Kim, Lee, Kim, & Oh, 2008), *Halothermothrix orenii* (Hassan et al., 2015), and hot environment metagenomes (Gupta, Govil, Capalash, & Sharma, 2012). In addition to production of GOS, where lactose is used as donor as well as acceptor, some β -D-galactosidases may use other acceptors than lactose (Díez-Municio, Herrero, Olano, & Moreno, 2014). Mu, Chen, Wang, Zhang, and Jiang (2013) and Li et al. (2009) have described that a β -D-galactosidase from *Bacillus circulans* forms

the trisaccharide lactosucrose (O- β -D-galactopyranosyl-(1 \rightarrow 4)-O- α -D-glucopyranosyl-(1 \rightarrow 2)- β -D-fructofuranoside) from lactose and sucrose, and Guerrero, Vera, Acevedo, and Illanes (2015), Wang, Yang, Hua, Zhao, and Zhang (2013), and Tang et al. (2011) have reported the enzymatic synthesis of the disaccharide lactulose (4-O- β -D-galactopyranosyl- β -D-fructofuranose). GOS, lactosucrose, and lactulose all show prebiotic effects and are presumed to play an important role in stimulating growth of beneficial bifidobacteria in the human intestine (Gibson et al., 2004; Mu et al., 2013; Wang et al., 2013). Furthermore, previous studies have shown that several bifidobacterial species can grow on GOS as sole carbon source (Garrido et al., 2013; Gavlighi, Michalak, Meyer, & Mikkelsen, 2013; Rada et al., 2008).

Another group of prebiotics belong to the fucosylated oligosaccharides. L-fucose (6-deoxy-L-galactose) is a monosaccharide which is a component of many glycoproteins and glycolipids, and is often located at the nonreducing end of oligosaccharide chains via an α -(1 \rightarrow 2) linkage to a D-galactosyl residue or via α -(1 \rightarrow 3), α -(1 \rightarrow 4), or α -(1 \rightarrow 6) linkage to N-acetyl-D-glucosamine (reviewed by Becker & Lowe, 2003). In addition, fucosylated oligosaccharides are major components in human breast milk, while almost absent in bovine milk (Kunz, Kuntz, & Rudloff, 2014; Kunz & Rudloff, 1993; Kunz, Rudloff, Baier, Klein, & Strobel, 2000; Thurl, Henker, Taut, Tovar, & Sawatzki, 1993), which is the basis for infant formula. By supplementing bovine instant formula with fucosylated oligosaccharides as well as GOS, it has been shown that growth of bifidobacteria and lactobacilli in the fecal microflora of infants was improved (Boehm et al., 2002; Moro et al., 2002), and α -linked fucosylated oligosaccharides have furthermore been shown to prevent infection with the pathogenic bacteria *E. coli* and *Campylobacter jejuni* (Morrow, Ruiz-Palacios, Jiang, & Newburg, 2005). Fucosylated oligosaccharides such as 2'-O-fucosyllactose (2FL) and 3'-O-fucosyllactose (3FL) can be synthesized enzymatically through transglycosylation by the action of transfucosylating fucosidases (Lezyk et al., 2016; Zeuner et al., 2018) or by using fucosyltransferases, where an L-fucose unit is transferred from guanosine 5'-phosphate (GDP)-L-fucose to an acceptor substrate (Chin, Kim, Kim, Jung, & Seo, 2017; Chin, Kim, Lee, & Seo, 2015; Sprenger, Baumgärtner, & Albermann, 2017).

In this study, we screened hitherto unexplored hot springs in East Greenland for bacteria expressing thermostable glycoside hydrolases. One of the isolates, affiliated to *Paenibacillus* sp., was shown to produce a transglycosylating β -D-galactosidase and a transglycosylating α -L-fucosidase. The two enzymes were recombinantly expressed in *E. coli*, purified, and functionally characterized, and here, we report on the characterization of the β -D-galactosidase and the α -L-fucosidase from *Paenibacillus* sp. 3179.

2 | MATERIALS AND METHODS

2.1 | Bacterial strains and chemicals

All chemicals were purchased from Sigma-Aldrich (St. Louis) unless otherwise stated. All enzymes were purchased from New England Biolabs (Ipswich) unless otherwise stated.

For construction of a genomic library and initial protein expression and characterization, the plasmid pUC18 Δ lacZ (Schmidt & Stougaard, 2010) was used in *E. coli* TOP10 (F- *mcrA* Δ (*mrr-hsdRMS-mcrBC*) Φ 80lacZ Δ M15 Δ lacX74 *recA1 araD139* Δ (*ara leu*) 7,697 *galU galK rpsL* (StrR) *endA1 nupG*) (Thermo Fisher Scientific, Waltham, MA, USA). For enzyme expression, the cloning vector pET9a-USER-1 (Schultz-Johansen et al., 2018) was used in the lactose-deficient strain *E. coli* BL21(DE3) Δ lacZ, which was kindly provided by Professor Jin-Ho Seo from Seoul National University, Seoul, the Republic of Korea.

2.2 | Samples, isolation, and growth conditions

Samples were collected from a hot spring in Knighton Fjord in East Greenland (69°21'60.4"N; 24°39'67.1"W) in 2001, where more than 30 discharges were observed with temperatures around 54°C. Gravel samples from the bottom and the sides of the hot springs were plated directly on R2 agar, which contained per liter: 0.5 g yeast extract, 0.5 g Bacto peptone, 0.5 g casamino acids, 0.05 g MgSO₄•7H₂O, 0.1 g NaCl, 0.5 g soluble starch, 0.3 g sodium pyruvate, 0.3 g KH₂PO₄, and 20 g agar. The substrate was buffered to pH 7 with 0.01 M NaH₂PO₄/Na₂HPO₄ buffer. Plates were incubated at 50°C for 1–2 days, and bacterial colonies were isolated by re-inoculating on R2 agar.

2.3 | Screening for enzymatic activities

For screening of native bacterial isolates, the R2 agar was supplemented with 0.2% lactose and either 20 μ g/ml 5-bromo-4-chloro-3-indolyl β -D-galactopyranoside (X-gal) for β -D-galactosidase activity or 20 μ g/ml 5-bromo-4-chloro-3-indolyl β -D-fucopyranoside (X-fuc) for α -L-fucosidase activity. Cultures were incubated at 50°C overnight, and positive colonies were re-inoculated until pure colonies with enzymatic activity were obtained.

Crude protein extract was made by bead beating of cells harvested from an overnight culture (50°C, 200 rpm) in a MP FastPrep Homogenizer (MP Biomedicals) at speed 5.5 for 3 \times 25 s in 2-ml Micro tubes PP (Sarstedt, Nümbrecht) supplemented with approx. 250 μ l acid-washed glass beads (212–300 μ m, 425–600 μ m, ratio 1:1). Extracts were tested for hydrolytic activity with 1 mM oNPG and 1 mM pNP-fucose at 50°C until visible yellow color was developed (10–35 min). Reactions were stopped with 0.5 M Na₂CO₃, and OD₄₀₅ was measured in an EL808 Ultra Microplate Reader (BioTek Instruments, Inc.).

2.4 | Characterization of bacterial growth

Bacterial growth experiments were carried out in triplicates in R2 broth supplemented with 0.2% lactose. Growth and enzyme activity was tested at 20, 28, 37, 50, and 60°C at 200 rpm. Optimal pH for

growth was tested at pH 5.0, 5.6, 6.0 (50 mM citrate buffer), 7.0, 8.0 (50 mM Na-phosphate buffer), 9.0, and 10.0 (50 mM glycine-NaOH buffer) at 37°C, 200 rpm. Growth was monitored as OD₆₀₀ using an UVmini 1,240 spectrophotometer (Shimadzu). Phenotypes were furthermore investigated on a nutrient-poor substrate containing 0.8% (w/v) agar and 0.2% (w/v) peptone. After 48 hr incubation at 50°C, the agar plates were stained with 0.1% (w/v) Coomassie Brilliant Blue R-250 dissolved in a solution of 20% EtOH (v/v) and 5% (v/v) acetic acid.

2.5 | Construction and screening of genomic library

A genomic library was generated in *E. coli* TOP10 using the pUC18 Δ lacZ plasmid. Genomic DNA from *P. dendritiformis* sp. 3179 was extracted using the Puregene Yeast/Bact. Kit B (Qiagen, Hilden) according to the manufacturer's protocol. The genomic DNA was partially digested with Sau3AI for 25 min at 37°C. The digest was applied to a 1% SeaPlaque® GTG agarose gel (Lonza, Visp), and fragments corresponding to 3–6 kb were cut out and purified using 1X GELase™ Reaction Buffer and GELase™ Enzyme (Epicentre) followed by precipitation with 70% EtOH.

The pUC18 Δ lacZ vector was linearized by digestion with BamHI for 1 hr at 37°C, applied to a 1% SeaPlaque® GTG agarose gel, and purified using the GeneJET™ Gel Extraction Kit (Fermentas, Waltham, MA, USA) according to the manufacturer's protocol. The vector DNA was phosphorylated using the Shrimp Alkaline Phosphatase (New England Biolabs) for 45 min at 37°C followed by inactivation for 10 min at 65°C. Ligation of the partially digested genomic DNA and the digested vector was carried out using the T4 DNA Ligase at 16°C overnight. The ligation reaction was transformed into chemically competent *E. coli* TOP10. The transformation reaction was subsequently plated onto LB agar supplemented with 100 μ g/ml ampicillin and either 20 μ g/ml X-gal + 0.3 mM isopropyl β -D-1-thiogalactopyranoside (IPTG) or 20 μ g/ml X-fuc. Plates were incubated for 37°C overnight, and blue colonies, indicating hydrolytic activity on the X-linked substrates, were selected for further investigation.

2.6 | Identification and cloning of enzyme-coding genes

Plasmids from blue colonies were purified using the QIAprep® Spin Miniprep Kit (Qiagen), and inserts were sequenced at GATC Biotech (Constance). Sequences were analyzed for open reading frames (ORFs) using CLC Main Workbench v7.9.1 (Qiagen), translated into amino acid sequences, and blasted using the NCBI BLASTp Tool (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>). ORFs with the highest similarity to β -galactosidases and α -fucosidases were selected for further analysis. Two sequences of 1,749 bp and 1,281 bp corresponding to a β -D-galactosidase and α -L-fucosidase, respectively, were identified as best candidates and amplified with specific primers

(PdenGal_USER_F: 5'-GGCTTAAUATGACAACGCTTTCTTATAAC-3', PdenGal_USER_R: 5'-GGTTTAAUTTTCGATAAAGAAGGGGC-3', PdenFuc_USER_F: 5'-GGCTTAAUATGCGTCGGTCACG-3', PdenFuc_USER_R: 5'-GGTTTAAUTTGCCATCTCCA-3'); uracil-specific excision reagent (USER) overhangs are underlined). Reverse primers were designed without stop codons to utilize the 6xHis tag included in the cloning vector for protein purification.

The β -D-galactosidase gene was amplified using the following PCR protocol: 1 \times 98°C for 30 s; 30 \times (98°C for 10 s, 55°C for 20 s; and 72°C for 1 min); and 72°C for 10 min, while the gene encoding the α -L-fucosidase was amplified using the PCR protocol: 1 \times 98°C for 30 s; 30 \times (98°C for 10 s, 56°C for 20 s; and 72°C for 1 min 30 s); and 72°C for 10 min.

The cloning vector, pET9a-USER-1 vector, was prepared by digestion with PaeI and nt.BbvCI at 37°C overnight followed by purification from a 1% SeaPlaque® GTG agarose gel using the GeneJET™ Gel Extraction Kit. The amplified ORFs were inserted into pET9a-USER-1 by including specific USER overhangs during the PCR (Table I) and ligating the PCR product into the vector by means of USER enzyme (New England Biolabs). The reaction was carried out at 37°C for 20 min followed by 25°C for 20 min and subsequently used for transformation into chemically competent *E. coli* BL21(DE3) Δ lacZ. Colonies were grown on LB agar supplemented with 50 μ g/ml kanamycin (Km) at 37°C overnight. The plasmids from one randomly selected colony from each transformation were purified and verified by sequencing (as above).

2.7 | Phylogenetic analyses

The 16S rRNA gene was analyzed against BLASTn (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>), and the closest relatives were aligned using Clustal W (Thompson, Higgins, & Gibson, 1994). A phylogenetic tree was made in MEGA7 (Kumar, Stecher, & Tamura, 2016) using the maximum likelihood method based on the Tamura-Nei model (Tamura & Nei, 1993).

The two ORFs identified as a putative β -D-galactosidase and α -L-fucosidase, respectively, were analyzed using BLASTp against PDB to identify the phylogenetically closest relatives. When this only gave very few results, the search was expanded to the nonredundant (nr) protein database in combinations with the CAZy database. Sequences were aligned using Clustal W, and a phylogenetic tree was made using the maximum likelihood method based on the JTT matrix-based model (Jones, Taylor, & Thornton, 1992).

2.8 | Protein expression and purification

For initial transglycosylation experiments, the construct encoding the β -D-galactosidase gene was expressed and purified as described by Schmidt and Stougaard (2010) with minor modifications. In short, the gene containing the native SD sequence was amplified using specific primers β -galFpUC

(5'-TCTCGGTACCAGAAGGAGAGAAGCCAATG-3') and β -galRpUC (5'-CTCTAAGCTTATTTTCGATAAAGAGGGGC-3'), and cloned into the KpnI and HindIII site of pUC18 Δ LacZ. Overnight expression at 37°C in *E. coli* TOP10 cells was followed by protein extraction by bead beating (as described below), heating to 60°C for 15 min, and subsequent purification by anion exchange chromatography at pH 6.2 and eluted with a NaCl gradient, resulting in apparent homogeneity on a SDS gel.

For the remaining characterization and transglycosylation experiments, recombinant strains in *E. coli* BL21(DE3) Δ lacZ were used for protein expression. Cells were inoculated in 10 ml LB with 50 μ g/ml Km and incubated at 37°C with 200 rpm overnight. The overnight culture was used to inoculate the expression culture in 100 ml LB with 50 μ g/ml Km, which was incubated at 37°C with 200 rpm until OD₆₀₀ ~ 0.3–0.4. The culture was then supplemented with 0.2 mM IPTG and 0.1% D-L-rhamnose to induce expression, and incubation was continued at room temperature with 200 rpm overnight to increase correct protein folding.

After overnight expression, cells were cooled on ice for 10 min and harvested by centrifugation for 10 min at 4,700 rpm, 4°C, and the supernatant was discarded. The cell pellet was resuspended in 10 ml binding buffer [20 mM sodium phosphate, 500 μ M NaCl, and 20 mM imidazole, pH 7.4] and transferred to 50-ml centrifuge tubes with screw caps and a gasket (VWR, Radnor) supplemented with approx. 2 ml acid-washed glass beads (212–300 μ m, 425–600 μ m, ratio 1:1) for bead beating. Bead beating was carried out in an MP FastPrep Homogenizer at speed 4.0 for 3 \times 25 s. The samples were incubated on ice for 1 min in between rounds. Samples were centrifuged for 5 min at 4,700 rpm to remove glass beads and cell debris, and the supernatant was transferred to a clean tube and kept on ice until purification.

Protein purification was carried out using the His GraviTrap Kit (GE Healthcare) following the manufacturer's protocol for purifications under native conditions. Elution was carried out using 500 mM imidazole. Expression was checked by SDS-PAGE on RunBlue SDS 4%–12% Protein Gels (Expedeon).

2.9 | Characterization of recombinant proteins

Concentration of purified enzyme was determined in a Bradford assay using BradfordUltra reagent (Expedeon Protein Solutions) with a dilution series of BSA (bovine serum albumin) as model protein. Absorbance was measured at 595 nm using an Epoch Microplate Spectrophotometer running Gen5 software ver. 3.03 (BioTek Instruments Inc.), and the slope was used to estimate the concentrations of purified enzymes.

To determine pH and temperature optimum, the purified β -D-galactosidase was characterized using 0.5 mM oNPG as a substrate, while the purified α -L-fucosidase was characterized using 0.5 mM pNP-fucose. pH optimum was determined (with six replicates) at 50°C at pH 5.0, 5.6, and 6.0 (50 mM citrate buffer); pH 6.8, 7.4, and 8.0 (50 mM Na-phosphate buffer); and pH 9.0, 10.0, and 10.7 (50 mM

glycine-NaOH buffer). Temperature optimum was determined (in triplicates) from 25 to 65°C for 5 min in 50 mM Na-phosphate buffer at pH 7.4. OD_{420} was determined using an Epoch Microplate Spectrophotometer running Gen5 software ver. 3.03.

Enzyme kinetics was investigated at 50°C. Reactions of 200 μ l [in 50 mM Na-phosphate buffer pH 7.4] with a fixed amount of enzyme (18 μ g for the α -L-fucosidase and 1,200 μ g for the β -D-galactosidase) were incubated with increasing concentrations of the substrate (0.25–1 mM oNPG/pNP-fucose). The product formation was monitored by measuring the absorbance at 420 nm every 30 s in real time by using a FLOUstar Omega microplate reader (BMG Labtech GmbH). Blanks contained buffer and substrate alone to represent the autohydrolysis of the substrate. All samples were performed in triplicates. Data were analyzed using the Omega Mars data analysis software ver. 3.20 (BMG Labtech). Molar extinction coefficients for oNP and pNP of 3.47 $\text{mM}^{-1} \text{cm}^{-1}$ and 18.4 $\text{mM}^{-1} \text{cm}^{-1}$, respectively (Bowers, McComb, Christensen, & Schaffer, 1980; Zeyer & Kocher, 1988), were used to calculate concentrations of the product. The steady-state rates were determined from the slope of the initial (linear) part of the resulting progress curves. The experimental data points were fitted to the Michaelis–Menten equation to obtain the kinetic parameters K_m and V_{max} . Nonlinear regression and estimations of kinetic parameters were performed in the software OriginPro (OriginLab).

2.10 | Transglycosylation

Transglycosylation by the β -D-galactosidase was tested using 10% (w/v) lactose as galactosyl donor and acceptor, and sucrose or fructose was used as additional galactosyl acceptors. The α -L-fucosidase was tested using 5–50 mM pNP-fucose (dissolved in 10% dimethylformamide) as donor and 5% (w/v) lactose as acceptor. Reactions were incubated at 50°C for 0, 15, 30 min, 1, and 3 hr. Reactions were stopped at 99°C for 15 min. For transglycosylation reactions by the β -D-galactosidase using lactose, sucrose, and fructose as acceptors, a control reaction with inactivated enzyme (99°C for 15 min) was included.

Reactions were analyzed by thin-layer chromatography (TLC) on TLC Silica gel 60 F₂₅₄ (Merck, Darmstadt, Germany) developed in ethyl acetate:acetic acid:H₂O (6:6:1, v/v/v). Saccharides were detected by 0.1M 2-methylresorcinol dissolved in a 5% (v/v) solution of sulfuric acid in ethanol followed by heating of the silica gel.

Formation of oligosaccharides was analyzed by HPLC using an ICS-5000 system (Dionex) with an AS autosampler and a pulsed amperometric detector (carbohydrate four-potential waveform, sampling rate 2 Hz), with a gold electrode (Au) and an Ag/AgCl reference electrode. The column used was a Dionex CarboPac PA1 (250 \times 4 mm, with 50 \times 4 mm Guard, Thermo Scientific) maintained at 22°C. Eluents consisted of water and aqueous solutions of sodium hydroxide (NaOH) and sodium acetate (NaOAc), in the combinations: (a) water; (b) 1 M NaOH; (c) 500 mM NaOH + 750 mM NaOAc; and (d) 500 mM NaOAc. The flow: 1.0 ml/min, injection volume: 10 μ l, and samples were maintained at 5°C. For identification

of peaks, qualitative standards of 3'-O-galactosyllactose, 4'-O-galactosyllactose, and 6'-O-galactosyllactose were used.

3 | RESULTS

3.1 | Isolation and characterization of glycoside hydrolase-producing isolate 3179

Water and sediment samples from hot springs in Knighton Fjord, East Greenland, were screened for isolates producing β -D-galactosidase and α -L-fucosidase enzymes on growth media supplemented with chromogenic substrates. One isolate, #3179, that showed good β -D-galactosidase and α -L-fucosidase activity was selected for further analysis.

Phylogenetical analysis of the 16S rRNA gene sequence of isolate 3179 (GenBank accession number MK616148) showed a high degree of similarity to the genus *Paenibacillus* with 98.4% identity to *Paenibacillus thiaminolyticus* LMG 17412^T, 97.8% identity to *Paenibacillus dendritiformis* DSM 18844^T, and 94.7% identity to *P. alvei* DSM29^T (Figure 1a). Culturing isolate 3179 on thin agar plates with peptone and subsequent staining with Coomassie Brilliant Blue showed a unique complex growth pattern (Figure 1b), which was similar to that reported for *P. dendritiformis* (Tcherpakov, Ben-Jacob, & Gutnick, 1999). Thus, we assume that isolate 3179 was affiliated to the genus *Paenibacillus*, and in the following, the strain is denoted *Paenibacillus* sp. 3179.

When *Paenibacillus* sp. 3179 was cultivated in lysogenic broth (LB) medium, growth optimum was determined to be at 50°C. No growth was observed at 28°C nor at 60°C. Optimal growth was observed at pH 8, and no growth was observed below pH 5.6; good growth was still maintained at pH 9–10.

3.2 | *Paenibacillus* sp. 3179 crude extract contains α -L-fucosidase and β -D-galactosidase activity

A crude extract of the native *Paenibacillus* sp. 3179 was subjected to analysis of α -L-fucosidase and β -D-galactosidase hydrolytic and transglycosylation activity. No activity was observed in the membrane fraction, but the cytosolic fraction showed α -L-fucosidase and β -D-galactosidase hydrolytic activity as determined from hydrolysis of pNP-fucose (4-nitrophenyl- α -L-fucopyranoside) and oNPG (2-nitrophenyl- β -D-galactopyranoside) substrates, respectively. In addition, the crude cytosolic extracts showed transgalactosylation activity using lactose as substrate as well as transfucosylation activity using pNP-fucose as substrate (Figure 2).

3.3 | Identification of enzyme-encoding genes

A functional expression library was constructed by inserting partially digested genomic DNA (gDNA) fragments from *Paenibacillus* sp. 3179




FIGURE 1 (a) Phylogeny of 16S rRNA genes generated using the maximum likelihood method based on the Tamura–Nei algorithm, showing the phylogenetic relationships between *Paenibacillus* sp. 3179 and related taxa. Numbers at nodes are bootstrap values shown as percentages of 100 replicates. (b). *Paenibacillus* sp. 3,170 cultured at 50°C for 48 hr on agar (0.8% w/v) with peptone (0.2% w/v) and subsequently stained with Coomassie Brilliant Blue and destained in ethanol (20% v/v) and acetic acid (5% v/v)





FIGURE 2 TLC analysis of hydrolysis and transglycosylation reactions of crude, intracellular extract from *Paenibacillus* sp. 3179. (a) Reaction with lactose; standard sugars in lanes 1–3; lane 4, extract from *Paenibacillus* sp. 3179 incubated with lactose for 16 hr at 37°C. (b) Reaction with pNP-fucose; lane 5, extract from *Paenibacillus* sp. 3179 incubated with pNP-fucose and fucose for 16 hr at 37°C; lane 6, pNP-fucose and fucose. Transglycosylation products are shown with a red line (reaction with lactose) and a star (reaction with pNP-fucose)

into plasmid pUC18ΔlacZ and transformed into *E. coli* TOP10. The library was screened for β-galactosidase-encoding genes by plating the expression library onto agar plates supplemented with the chromogenic substrate X-gal. One inserted DNA fragment from a blue colony contained a 1,749 bp open reading frame (ORF) next to a seven bp putative Shine–Dalgarno sequence (AGGAGAG) located seven bp upstream

of the ORF. BLASTp analysis of the translated ORF showed that the amino acid sequence displayed 92.5% identity to an uncharacterized putative β-D-galactosidase from *P. dendritiformis* (WP_133381740.1), 91.6% identity to a characterized β-D-galactosidase from *P. thiaminolyticus* (CAZ44333) (Benešová, Lipovová, Dvořáková, & Králová, 2010), and 79.7% identity to an uncharacterized putative β-D-galactosidase

P. alvei (EPY10792.1) as its closest relatives (Figure 3a). A conserved domain search (CDD) (Marchler-Bauer et al., 2015) showed the same domains in the *Paenibacillus* sp. 3179 β -D-galactosidase as in enzymes from *P. thiaminolyticus* and *P. dendritiformis*: a conserved GH35-domain (Pfam01301) at position 10–322 and a β -D-galactosidase jelly roll domain (Pfam13364) at position 527–562. A BLAST search at UniProt (www.uniprot.org) confirmed the affiliation to the glycoside hydrolase family 35. A HMM search and in silico folding analysis over known structures showed that the *Paenibacillus* sp. 3179 β -D-galactosidase displayed high similarity to the characterized structure of the bgaC β -D-galactosidase from *Bacillus circulans* (4MAD) (5.6e-95) (Henze et al., 2014): The *Paenibacillus* sp. 3179 amino acid sequence including the catalytic residues (nucleophile Glu233 and acid/base Glu157, according to the *B. circulans* numbering) could be superimposed onto the corresponding *B. circulans* structure, 4MAD (Figure A1).

The expression library was subsequently screened for α -L-fucosidase activity on agar plates containing the chromogenic substrate X-fuc. DNA sequence analysis of an inserted DNA fragment from a blue colony showed a 1,281 bp ORF. BLASTp analysis showed that the amino acid sequence encoded by the ORF displayed 97.4% amino acid sequence identity to a characterized α -L-fucosidases from *P. thiaminolyticus* (aLfuk1, FN869117) (Benešová, Lipovová, Dvořáková, & Králová, 2013) and 97.2% sequence identity to a putative uncharacterized α -L-fucosidase from *P. dendritiformis* (WP_006676648.1) (Figure 3b). The conserved domain search revealed a conserved glycoside hydrolase superfamily D (GH-D) domain (pfam01120), which comprises the families GH27, GH31, and GH36 containing enzymes with structurally and mechanistically similar activities, but none of them are known to show α -L-fucosidase activity. However, a BLAST analysis at UniProt showed that the most closely related α -L-fucosidases all belong to GH29. Thus, we conclude that the α -L-fucosidase from *Paenibacillus* sp. 3179 belongs to GH29. An HMM search and modeling over the structure with highest similarity (2.2e-86), which was an α -L-fucosidase from *P. thiaminolyticus* (6GN6) (Koval'ová, 2019), showed that the *Paenibacillus* sp. 3179 sequence including the catalytic residues (nucleophile Asp186 and acid/base Glu239, according to the *P. thiaminolyticus* numbering) could be superimposed onto the corresponding 6GN6 structure (Figure A1). The DNA sequences encoding the α -L-fucosidase and the β -D-galactosidase from *Paenibacillus* sp. 3179 have been uploaded to GenBank with the accession numbers MK625194 and MK625195, respectively.

3.4 | Characterization of recombinant enzymes

The *Paenibacillus* sp. 3179 β -D-galactosidase and α -L-fucosidase were recombinantly produced in *E. coli* as 6xHis-tagged enzymes. SDS-PAGE showed that purified β -D-galactosidase and α -L-fucosidase migrated as 70 kDa and 50 kDa protein, respectively (calculated theoretical molecular weights 65.7 kDa and 49.5 kDa) (Figure A2).

Using oNPG as a substrate, the purified β -D-galactosidase showed optimal activity at 50°C and at pH 8.0 (Figure 4a). Stability

was investigated at 37, 50, and 70°C, and the β -D-galactosidase was stable for up to 5 hr at 37°C or 50°C, while it was completely inactivated after 30 min at 70°C (data not shown). K_m for the β -D-galactosidase with oNPG as substrate was estimated to be 0.57 ± 0.12 mM and $V_{max} = 0.11 \pm 0.01$ mmoles/min/mg enzyme (Table 1, Figure A3). The β -D-galactosidase was also tested with *p*-nitrophenyl- β -D-galactopyranoside (pNPG) as substrate, but no activity was detected (data not shown).

Activity of the purified α -L-fucosidase was similarly determined with pNP-fucose as substrate. Optimal activity was observed at 50°C and at pH 7.4 (Figure 4b). Stability was investigated at 37°C, 50°C, and 70°C, and the α -L-fucosidase was stable for up to 5 hr at 37°C and 50°C, while it was completely inactivated after 30 min at 70°C (data not shown). K_m with pNP-fucose as substrate was estimated to be 1.11 ± 0.75 mM and $V_{max} = 3.88 \pm 1.58$ mmoles/min/mg enzyme (Table 1, Figure A3).

3.5 | Transglycosylation of recombinant enzymes

Both of the investigated enzymes exhibited transglycosylation activity. Transglycosylation by the β -D-galactosidase was analyzed with lactose as galactosyl donor and lactose, sucrose, or fructose as accepting carbohydrates. When lactose was used as donor and acceptor, hydrolysis products (galactose and glucose) appeared in the TLC analysis together with unhydrolyzed lactose and galactooligosaccharides (GOS) (Figure 5a). Two of these GOS were identified as 6'-O-galactosyllactose and 3'-O-galactosyllactose by HPLC analysis (Figure 6a), and the enzyme was able to transform lactose into 6'-galactosyllactose, 0.14 mol/mol, and 3'-galactosyllactose, 0.027 mol/mol (Figure 6b). Hydrolysis and transgalactosylation were seen in the range from pH 5 to 8 (data not shown). When sucrose or fructose was used as acceptors, reddish spots appeared in the TLC analysis (white arrows in Figure 5b). In these transgalactosylation experiments, both dark blue/purple spots corresponding to GOS, lactose, galactose, and glucose and reddish spots were observed, indicating that transgalactosylation occurred with sucrose and fructose as galactosyl acceptors (Figure 5b, red and yellow arrows, respectively). Similarly, transglycosylation with the α -L-fucosidase was analyzed using pNP-fucose as fucosyl donor and lactose as fucosyl acceptor. With TLC, an oligosaccharide corresponding in size to 2FL was detected (Figure 5c, black arrow). This finding was confirmed by HPLC, which showed that the formed oligosaccharide eluted as 2FL (Figure 6c). Further products were formed at about the same rate as 2FL, but these were not characterized further (Figure 6d).

4 | DISCUSSION

A β -D-galactosidase and α -L-fucosidase producing strain, #3179, was isolated from a thermal spring in Greenland. 16S rRNA gene analysis showed that strain 3179 was affiliated to the genus *Paenibacillus* with




FIGURE 3 Phylogenetic analysis of (a) *Paenibacillus* sp. 3179 β -D-galactosidase and (b) *Paenibacillus* sp. 3179 α -L-fucosidase. Sequences that showed similarities to the *Paenibacillus* sp. 3179 enzymes were identified by BLASTp analysis against Protein Data Bank (PDB) and supplemented with closely related enzymes from the nonredundant (nr) protein sequence database. The latter were only selected if also identified among GH35 and GH29 family enzymes, respectively, in the CAZy database (www.cazy.org). Phylogenetic relationships were inferred by the maximum likelihood method based on the JTT matrix-based model. The trees with the highest log likelihood are shown. Bootstrap values ($n = 100$) above 50 are shown. All positions containing gaps and missing data were eliminated. Evolutionary analyses were conducted in MEGA7

P. dendritiformis and *P. thiaminolyticus* as closest relatives. Strain 3179 displayed dendritic growth similar to *P. dendritiformis* why we conclude that strain 3179 most likely could be classified as *P. dendritiformis*. However, as no other phylogenetical or chemotaxonomical analyses were carried out, we denote the strain *Paenibacillus* sp. 3179.

The two genes encoding a β -D-galactosidase and an α -L-fucosidase, respectively, were isolated and analyzed phylogenetically and functionally. Sequence analysis of the β -D-galactosidase showed close relationship to β -galactosidases from *P. dendritiformis* (92.5% sequence identity) and from *P. thiaminolyticus* (91.6% identity) (Benešová et al., 2010). Further distantly related enzymes showed below 80% identity for hypothetical enzymes and below 55% identity for classified β -galactosidase like *Bacillus circulans* BgaC (BAA21669) (Ito & Sasaki, 1997). All related β -D-galactosidase sequences were classified as GH35 according to the glycoside hydrolase classification system (Henrissat, 1991), and thus, the β -D-galactosidase from *Paenibacillus* sp. 3179 was hypothesized to belong to GH35, which primarily consists of β -galactosidases (EC 3.2.1.23), but also contain exo- β -glucosaminidases (EC 3.2.1.165), exo- β -1,4-galactanases (EC

3.2.1.-), and β -1,3-galactosidases (EC 3.2.1.-). GH35 enzymes can be found in more than 2,700 different organisms spanning archaea, bacteria, and eukaryotes, including plants and animals. GH35 β -galactosidases have an $(\alpha/\beta)_8$ barrel as catalytic domain (www.cazypedia.org), and the mechanism is catalysis of the hydrolysis of terminal nonreducing β -galactosyl residues in, for example, lactose, oligosaccharides, glycolipids, and glycoproteins. In silico analysis showed that the overall structure, including the catalytic residues, could be modeled over a known structure of a β -D-galactosidase from *B. circulans* (Figure A1), making it possible that the β -D-galactosidase from *Paenibacillus* sp. 3179 has the same functional activity as that from *B. circulans*.

The α -L-fucosidase sequence was analyzed in a similar manner. The closest related sequence was an α -L-fucosidase from *P. thiaminolyticus* (FN869117; 6GN6) (Benešová et al., 2013; Koval'ová et al., 2019). The phylogenetically closest related α -L-fucosidases were all shown to belong to GH29, which consists of exo-acting α -fucosidases (EC 3.2.1.51) and α -1 \rightarrow 3/1 \rightarrow 4-L-fucosidases (EC 3.2.1.111). Enzymes in GH29 also consist of an $(\alpha/\beta)_8$ -like barrel domain plus

FIGURE 4 Temperature optimum determined at pH 9.0 (a) and pH optimum determined at 50°C (b) for *Paenibacillus* sp. 3179 β -D-galactosidase (solid line) and α -L-fucosidase (dashed line) based on reaction rates. Error bars indicate standard error ($n = 3$)




TABLE 1 Characteristics of the β -D-galactosidase and α -L-fucosidase from *Paenibacillus* sp. 3179 compared to the closest related enzymes from *P. thiaminolyticus* (CCM 3,599)

Enzyme	Km (mM)	Substrate	Temperature optimum (°C)	pH optimum	Ref.
<i>Paenibacillus</i> sp. 3179 β -D-galactosidase	0.57 \pm 0.12	oNPG	50	8.0	This work
<i>P. thiaminolyticus</i> β -D-galactosidase	250	pNPG	65	5.5	^a
<i>Paenibacillus</i> sp. 3179 α -L-fucosidase	1.11 \pm 0.75	pNP-L-fucose	50	7.4	This work
<i>P. thiaminolyticus</i> α -L-fucosidase iso1	0.44	pNP-L-fucose	48	8.2	^b
<i>P. thiaminolyticus</i> α -L-fucosidase iso2	0.52	pNP-L-fucose	50	6.5	^c

^aBenešová et al. (2010).

^bBenešová et al. (2013).

^cBenešová et al. (2015).




FIGURE 5 Transglycosylation activities by *Paenibacillus* sp. 3179 enzymes. (a) Transgalactosylation by *Paenibacillus* sp. 3179 β -D-galactosidase preliminarily purified over time (0–21 hr) with lactose as donor and acceptor. (b) Transgalactosylation by preliminarily purified *Paenibacillus* sp. 3179 β -D-galactosidase with lactose (lac) as donor, and sucrose (suc), fructose (fruc), and lactose (lac) as acceptors. White arrows show the acceptor molecules sucrose and fructose. Red and yellow arrows show the transglycosylation products using sucrose and fructose as galactosyl acceptors, respectively. * indicates where the enzyme was inactivated before addition of substrates. (c) Transfucosylation by *Paenibacillus* sp. 3179 α -L-fucosidase with increasing concentration of *p*-nitrophenyl- α -L-fucopyranoside as donor and lactose as acceptor. Black arrow shows the transglycosylation product corresponding in size to 2FL.

a C-terminal β -sandwich domain (www.cazypedia.org) and function by classical retaining mechanisms involving double displacement via a covalent intermediate. In silico structural analysis showed that the α -L-fucosidase sequence from *Paenibacillus* sp. 3179 could be superimposed onto the corresponding 6GN6 structure, including catalytic residues, indicating a functional relationship (Figure A1).

Sequence analyses revealed that there are a lot of potentially unclassified β -galactosidases and α -fucosidases from strains within the *Paenibacillus* genus ranging in temperature profile from the psychrotolerant *P. antarcticus* which shows optimal growth at 10–15°C (Montes, Mercadé, Bozal, & Guinea, 2004) to the described thermotolerant *Paenibacillus* sp. 3179 with optimal growth at 50°C. However, since most of the proteins are only annotated as potential enzymes with no documented functions, the genus *Paenibacillus* appears to have a lot of potential for discovery of cold-active as well as thermostable enzymes for a range of industrial applications.

Functional characterization of recombinant purified enzymes from *Paenibacillus* sp. 3179 produced by *E. coli* showed that both enzymes possessed hydrolytic as well as transglycosylating activities. Chromatography analyses showed that the β -D-galactosidase was able to form 6'-O-galactosyllactose and 3'-O-galactosyllactose using lactose as substrate, and the α -L-fucosidase was able to transfer the fucose moiety from *p*NP-fuc to lactose, thereby forming 2FL and also

indication of 3FL was observed. The chromatogram in Figure 6c identified the transglycosylation product 2FL and a peak (no 3), which relative to elution of 2FL and lactose corresponded to the retention of 3FL (Tan, Chai, & Zhang, 2015). This compound was observed in increasing concentration with reaction progress, indicating that 3FL was a product of the transglycosylation reaction. Enzyme catalyzing production of 2FL and 3FL is known, and analyses of these oligosaccharides are well established (Tan et al., 2015; Zeuner, Teze, Muschiol, & Meyer, 2019), but 6FL as an enzymatic transglycosylation product is not described and the present investigation cannot establish its presence. The compound corresponding to peak 6 was not identified, but increased concentration with reaction time indicated it was a product of the reaction.

In studies carried out on the β -D-galactosidase and two α -L-fucosidases from *P. thiaminolyticus* (Benešová et al., 2010, 2013, 2015), it is described that all three enzymes were able to transfucosylate a wide range of *p*-nitrophenyl substrates and alcohols; however, little attention has been focused on transgalactosylation. In this study, we showed that the β -D-galactosidase from *Paenibacillus* sp. 3179 is able to transgalactosylate a range of natural substrates such as lactose, sucrose, and fructose, which can be of use for industrial purposes when, for example, upgrading disaccharides to oligosaccharides in food items. We furthermore showed that the α -L-fucosidase from *Paenibacillus* sp. 3179 is able to transfucosylate using lactose as acceptor, making it




FIGURE 6 HPLC analyses of transglycosylation reactions. (a) Transgalactosylation profile after 4 hr with *Paenibacillus* sp. 3179 β -D-galactosidase using lactose as sole substrate. Peaks are identified as follows: 1: glucose and galactose monomers. 2: unidentified (not a part of the reaction, present in all samples). 3: lactose. 4: 6'-O-galactosyl-lactose (6-gal-lac). 5: 3'-O-galactosyl-lactose (3-gal-lac). After 120 min, 24% of the lactose has been converted and results in 0.14 mol/mol 6-gal-lac and 0.027 mol/mol 3-gal-lac. (b) Diagram showing the utilization of lactose (blue squares, secondary axis) and production of 6-gal-lac (black circles, primary axis) and 3-gal-lac (black triangles, primary axis) over time (120 min.). (c) Transfucosylation profile after 3 hr with *Paenibacillus* sp. 3179 α -L-fucosidase using pNP-fucose as donor and lactose as acceptor. Peaks are identified as follows: 1: fucose. 2: unidentified (not a part of the reaction, present in all samples). 3: product of the reaction with increasing concentration over reaction time and a relative elution to 2FL and lactose corresponding retention of 3FL. 4: lactose. 5: 2'-O-fucosyllactose (2FL). 6 and 7: unidentified. (d) Diagram showing utilization of lactose (blue squares, secondary axis) and production of monomeric fucose (open circles, primary axis), 2FL (open squares, primary axis), and the compound with retention corresponding to 3FL (peak 3 in panel C) (open triangles, primary axis) over time (180 min.)

a good candidate for production of fucosylated oligosaccharides like 2FL for infant formula (Aldredge et al., 2013; Chen, 2015) due to its transfucosylating activity at elevated temperatures.

As the interest for thermostable transglycosylating enzymes has gained interest, similar enzymatic reactions have been reported. For example, Petrova and Kujumdzieva (2010) isolated several strains of thermotolerant yeasts from a range of milk products with transgalactosylation activities, and Fai, Silva, Andrade, Bution, and Pastore (2014) showed transgalactosylation activity by *Pseudozyma tsukubaensis* and *Pichia kluyveri* using lactose as a substrate. Lezyk et al. (2016) created a metagenome library from soil and discovered several putative α -fucosidases, which were recombinantly expressed and proven to have hydrolytic as well as transfucosylating activities, potentially suitable for formation of fucosylated oligosaccharides for supplementing infant formula to mimic the natural composition of human breast milk.

5 | CONCLUSIONS

In the current study, it was shown that the bacterium *Paenibacillus* sp. 3179 isolated from a thermal spring in East Greenland produced two thermostable enzymes, a β -D-galactosidase and an α -L-fucosidase, with hydrolytic as well as transglycosylating activities. It was confirmed that the β -D-galactosidase was able to produce 6'-O-galactosyllactose and

3'-O-galactosyllactose at its optimal temperature at 50°C and that the α -L-fucosidase was able to produce 2FL along with some uncharacterized oligosaccharides, speculated to include 3FL. Future studies will be aimed at characterizing the remaining oligosaccharides and to expand the range of investigated substrates for transglycosylation.

ETHICS STATEMENT

None required.

ACKNOWLEDGMENTS

We thank Professor Jin-Ho Seo from Seoul National University, Seoul, The Republic of Korea, for kindly providing us with the lactose-deficient *E. coli* BL21(DE3) Δ lacZ and Susanne Iversen for her excellent technical skills. This work was supported by The Danish Council for Strategic Research under grant no. 1308-00014B / 0652-20140008.

CONFLICTS OF INTEREST

None declared.

AUTHOR CONTRIBUTION

Mariane Thøgersen: data curation—lead, formal analysis—lead, investigation—equal, methodology—equal, writing—original draft—equal, and writing—review and editing—equal; Stefan Christensen: data curation—supporting, formal analysis—supporting, and

writing—review and editing—supporting; Morten Jepsen: data curation—supporting; Lars Pedersen: data curation—supporting, formal analysis—supporting, and writing—review and editing—supporting; and Peter Stougaard: conceptualization—lead, funding acquisition—lead, methodology—lead, project administration—lead, writing—original draft—equal, and writing—review and editing—equal.

DATA AVAILABILITY STATEMENT

All data generated in this study are available from the corresponding author upon reasonable request.

The 16S rRNA gene sequence is available at <https://www.ncbi.nlm.nih.gov/genbank/> under accession number MK616148, and the sequences encoding the β -D-galactosidase and an α -L-fucosidase under accession numbers MK625195 and MK625194, respectively.

ORCID

Mariane S. Thøgersen  <https://orcid.org/0000-0003-0799-6894>

Stefan J. Christensen  <https://orcid.org/0000-0001-9164-2225>

Morten Jepsen  <https://orcid.org/0000-0001-5307-7448>

Lars H. Pedersen  <https://orcid.org/0000-0003-3400-714X>

Peter Stougaard  <https://orcid.org/0000-0002-2796-7137>

REFERENCES

- Aldredge, D. L., Geronimo, M. R., Hua, S., Nwosu, C. C., Lebrilla, C. B., & Barile, D. (2013). Annotation and structural elucidation of bovine milk oligosaccharides and determination of novel fucosylated structures. *Glycobiology*, 23, 664–676. <https://doi.org/10.1093/glycob/cwt007>
- Arreola, S. L., Intanon, M., Suljic, J., Kittl, R., Pham, N. H., Kosma, P., ... Nguyen, T. H. (2014). Two β -galactosidases from the human isolate *Bifidobacterium breve* DSM 20213: Molecular cloning and expression, biochemical characterization and synthesis of galacto-oligosaccharides. *PLoS ONE*, 9, 1–13. <https://doi.org/10.1371/journal.pone.0104056>
- Becker, D. J., & Lowe, J. B. (2003). Fucose: Biosynthesis and biological function in mammals. *Glycobiology*, 13, 41R–53R. <https://doi.org/10.1093/glycob/cwg054>
- Benešová, E., Lipovová, P., Dvořáková, H., & Králová, B. (2010). β -D-galactosidase from *Paenibacillus thiaminolyticus* catalyzing transfucosylation reactions. *Glycobiology*, 20, 442–451. <https://doi.org/10.1093/glycob/cwp196>
- Benešová, E., Lipovová, P., Dvořáková, H., & Králová, B. (2013). α -L-fucosidase from *Paenibacillus thiaminolyticus*: Its hydrolytic and transglycosylation abilities. *Glycobiology*, 23, 1052–1065. <https://doi.org/10.1093/glycob/cwt041>
- Benešová, E., Lipovová, P., Krejzová, J., Kovalová, T., Buchtová, P., Spiwok, V., & Králová, B. (2015). Alpha-L-fucosidase isoenzyme iso2 from *Paenibacillus thiaminolyticus*. *BMC Biotechnology*, 15, 1–7. <https://doi.org/10.1186/s12896-015-0160-x>
- Boehm, G., Lidestri, M., Casetta, P., Jelinek, J., Negretti, F., Stahl, B., & Marini, A. (2002). Supplementation of a bovine milk formula with an oligosaccharide mixture increases counts of faecal bifidobacteria in preterm infants. *Archives of Disease in Childhood. Fetal and Neonatal Edition*, 86, F178–F181. <https://doi.org/10.1136/fn.86.3.F178>
- Bowers, G. N., McComb, R. B., Christensen, R. G., & Schaffer, R. (1980). High-purity 4-nitrophenol: Purification, characterization, and specifications for use as a spectrophotometric reference material. *Clinical Chemistry*, 26, 724–729.
- Chen, X. (2015). Human milk oligosaccharides (hmos): Structure, function, and enzyme-catalyzed synthesis. *Advances in Carbohydrate Chemistry and Biochemistry*, 72, 113–190. <https://doi.org/10.1016/bs.accb.2015.08.002>
- Chin, Y. W., Kim, J. Y., Kim, J. H., Jung, S. M., & Seo, J. H. (2017). Improved production of 2'-fucosyllactose in engineered *Escherichia coli* by expressing putative α -1,2-fucosyltransferase, WcFB from *Bacteroides fragilis*. *Journal of Biotechnology*, 257, 192–198. <https://doi.org/10.1016/j.jbiotec.2016.11.033>
- Chin, Y. W., Kim, J. Y., Lee, W. H., & Seo, J. H. (2015). Enhanced production of 2'-fucosyllactose in engineered *Escherichia coli* BL21star(DE3) by modulation of lactose metabolism and fucosyltransferase. *Journal of Biotechnology*, 210, 107–115. <https://doi.org/10.1016/j.jbiotec.2015.06.431>
- Cieśliński, H., Kur, J., Białkowska, A., Baran, I., Makowski, K., & Turkiewicz, M. (2005). Cloning, expression, and purification of a recombinant cold-adapted β -D-galactosidase from antarctic bacterium *Pseudoalteromonas* sp. 22b. *Protein Expression and Purification*, 39, 27–34. <https://doi.org/10.1016/j.pep.2004.09.002>
- Crout, D. H., & Vic, G. (1998). Glycosidases and glycosyl transferases in glycoside and oligosaccharide synthesis. *Current Opinion in Chemical Biology*, 2, 98–111. [https://doi.org/10.1016/S1367-5931\(98\)80041-0](https://doi.org/10.1016/S1367-5931(98)80041-0)
- de Roode, B. M., Franssen, M. C. R., van der Padt, A., & Boom, R. M. (2003). Perspectives for the industrial enzymatic production of glycosides. *Biotechnology Progress*, 19, 1391–1402. <https://doi.org/10.1021/bp030038q>
- DeCastro, M. E., Rodríguez-Belmonte, E., & González-Siso, M. I. (2016). Metagenomics of thermophiles with a focus on discovery of novel thermozymes. *Frontiers in Microbiology*, 7, 1–21. <https://doi.org/10.3389/fmicb.2016.01521>
- Díez-Municio, M., Herrero, M., Olano, A., & Moreno, F. J. (2014). Synthesis of novel bioactive lactose-derived oligosaccharides by microbial glycoside hydrolases. *Microbial Biotechnology*, 7, 315–331. <https://doi.org/10.1111/1751-7915.12124>
- Elleuche, S., Schäfers, C., Blank, S., Schröder, C., & Antranikian, G. (2015). Exploration of extremophiles for high temperature biotechnological processes. *Current Opinion in Microbiology*, 25, 113–119. <https://doi.org/10.1016/j.mib.2015.05.011>
- Fai, A. E. C., da Silva, J. B., de Andrade, C. J., Bution, M. L., & Pastore, G. M. (2014). Production of prebiotic galactooligosaccharides from lactose by *Pseudozyma tsukubaensis* and *Pichia kluyveri*. *Biocatalysis and Agricultural Biotechnology*, 3, 343–350. <https://doi.org/10.1016/j.bcab.2014.04.005>
- Garrido, D., Ruiz-Moyano, S., Jimenez-Espinoza, R., Eom, H.-J., Block, D. E., & Mills, D. A. (2013). Utilization of galactooligosaccharides by *Bifidobacterium longum* subsp. *infantis* isolates. *Food Microbiology*, 33, 262–270. <https://doi.org/10.1016/j.fm.2012.10.003>
- Gavighi, H. A., Michalak, M., Meyer, A. S., & Mikkelsen, J. D. (2013). Enzymatic depolymerization of gum tragacanth: Bifidogenic potential of low molecular weight oligosaccharides. *Journal of Agriculture and Food Chemistry*, 61, 1272–1278. <https://doi.org/10.1021/jf304795f>
- Gibson, G. R., Probert, H. M., Jan, V. L., Rastall, R. A., & Roberfroid, M. B. (2004). Dietary modulation of the human colonic microbiota: Introducing the concept of prebiotics. *Nutrition Research Reviews*, 17, 259–275. <https://doi.org/10.1093/jn/125.6.1401>
- Goderska, K., Szwengiel, A., & Czarnecki, Z. (2014). The utilization of *Pseudomonas taetrolens* to produce lactobionic acid. *Applied Biochemistry and Biotechnology*, 173, 2189–2197. <https://doi.org/10.1007/s12010-014-1024-x>
- Guerrero, C., Vera, C., Acevedo, F., & Illanes, A. (2015). Simultaneous synthesis of mixtures of lactulose and galacto-oligosaccharides and their selective fermentation. *Journal of Biotechnology*, 209, 31–40. <https://doi.org/10.1016/j.jbiotec.2015.06.394>
- Gupta, R., Govil, T., Capalash, N., & Sharma, P. (2012). Characterization of a glycoside hydrolase family 1 β -D-galactosidase from hot spring metagenome with transglycosylation activity. *Appl Biochemistry Biotechnol*, 168, 1681–1693. <https://doi.org/10.1007/s12010-012-9889-z>

- Gupta, V., Gupta, N., Capalash, N., & Sharma, P. (2017). Bio-prospecting bacterial diversity of hot springs in Northern Himalayan region of India for laccases. *Indian J Microbiol*, 57, 285–291. <https://doi.org/10.1007/s12088-017-0656-2>
- Gutiérrez, L. F., Hamoudi, S., & Belkacemi, K. (2012). Lactobionic acid: A high value-added lactose derivative for food and pharmaceutical applications. *International Dairy Journal*, 26, 103–111. <https://doi.org/10.1016/j.idairyj.2012.05.003>
- Hassan, N., Nguyen, T. H., Intanon, M., Kori, L. D., Patel, B. K. C., Haltrich, D., ... Tan, T. C. (2015). Biochemical and structural characterization of a thermostable β -glucosidase from *Halothermothrix orenii* for galacto-oligosaccharide synthesis. *Applied Microbiology and Biotechnology*, 99, 1731–1744. <https://doi.org/10.1007/s00253-014-6015-x>
- Henrissat, B. (1991). A classification of glycosyl hydrolases based on amino acid sequence similarities. *The Biochemical Journal*, 280, 309–316. <https://doi.org/10.1042/bj2800309>
- Henze, M., You, D. J., Kamerke, C., Hoffmann, N., Angkawidjaja, C., Ernst, S., ... Elling, L. (2014). Rational design of a glycosynthase by the crystal structure of β -D-galactosidase from *Bacillus circulans* (BgaC) and its use for the synthesis of N-acetylglucosamine type 1 glycan structures. *Journal of Biotechnology*, 191, 78–85. <https://doi.org/10.1016/j.jbiotec.2014.07.003>
- Hervet, C. J., Martin, N. H., Boor, K. J., & Wiedmann, M. (2017). Survival and detection of coliforms, *Enterobacteriaceae*, and gram-negative bacteria in Greek yogurt. *Journal of Dairy Science*, 100, 950–960. <https://doi.org/10.3168/jds.2016-11553>
- Ishikawa, K., Kataoka, M., Yanamoto, T., Nakabayashi, M., Watanabe, M., Ishihara, S., & Yamaguchi, S. (2015). Crystal structure of β -D-galactosidase from *Bacillus circulans* ATCC 31382 (BgaD) and the construction of the thermophilic mutants. *FEBS Journal*, 282, 2540–2552. <https://doi.org/10.1111/febs.13298>
- Ito, Y., & Sasaki, T. (1997). Cloning and characterization of the gene encoding a novel β -galactosidase from *Bacillus circulans*. *Bioscience, Biotechnology, and Biochemistry*, 61, 1270–1276. [https://doi.org/10.1016/0378-1119\(95\)00758-X](https://doi.org/10.1016/0378-1119(95)00758-X)
- Jones, D. T., Taylor, W. R., & Thornton, J. M. (1992). The rapid generation of mutation data matrices from protein sequences. *Computer Applications in the Biosciences*, 8, 275–282. <https://doi.org/10.1093/bioinformatics/8.3.275>
- Jørgensen, F., Hansen, O. C., & Stougaard, P. (2001). High-efficiency synthesis of oligosaccharides with a truncated β -D-galactosidase from *Bifidobacterium bifidum*. *Applied Microbiology and Biotechnology*, 57, 647–652. <https://doi.org/10.1007/s00253-001-0845-z>
- Jørgensen, F., Hansen, O. C., & Stougaard, P. (2004). Enzymatic conversion of D-galactose to D-tagatose: Heterologous expression and characterisation of a thermostable L-arabinose isomerase from *Thermoanaerobacter mathranii*. *Applied Microbiology and Biotechnology*, 64, 816–822. <https://doi.org/10.1007/s00253-004-1578-6>
- Knol, J., Boehm, G., Lidestri, M., Negretti, F., Jelinek, J., Agosti, M., ... Mosca, F. (2005). Increase of faecal bifidobacteria due to dietary oligosaccharides induces a reduction of clinically relevant pathogen germs in the faeces of formula-fed preterm infants. *Acta Paediatrica Supplement*, 94, 31–33. <https://doi.org/10.1111/j.1651-2227.2005.tb02152.x>
- Knol, J., Scholtens, P., Kafka, C., Steenbakkens, J., Gro, S., Helm, K., ... Wells, J. (2005). Colon microflora in infants fed formula with galacto- and fructo-oligosaccharides: More like breast-fed infants. *Journal of Pediatric Gastroenterology and Nutrition*, 40, 36–42. <https://doi.org/10.1097/00005176-200501000-00007>
- Koval'ová, T., Koval, T., Benešová, E., Vodičková, P., Spiwok, V., Lipovová, P., & Dohnálek, J. (2019). Active site complementation and hexameric arrangement in the GH family 29; a structure–function study of α -L-fucosidase isoenzyme 1 from *Paenibacillus thiaminolyticus*. *Glycobiology*, 29, 59–73.
- Kumar, S., Stecher, G., & Tamura, K. (2016). MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution*, 33, 1870–1874. <https://doi.org/10.1093/molbev/msw054>
- Kunz, C., Kuntz, S., & Rudloff, S. (2014). Bioactivity of human milk oligosaccharides. In F. J. Moreno, & M. L. Sanz (Eds.), *Food Oligosaccharides: Production, Analysis and Bioactivity* (pp. 5–20). Hoboken, NJ: John Wiley & Sons Ltd.
- Kunz, C., & Rudloff, S. (1993). Biological functions of oligosaccharides in human milk. *Acta Paediatrica*, 82, 903–912. <https://doi.org/10.1111/j.1651-2227.1993.tb12597.x>
- Kunz, C., Rudloff, S., Baier, W., Klein, N., & Strobel, S. (2000). Oligosaccharides in human milk: Structural, functional, and metabolic aspects. *Annual Review of Nutrition*, 20, 699–722. [10.1146/annurev.nutr.20.1.699](https://doi.org/10.1146/annurev.nutr.20.1.699)
- Ladirat, S. E., Schuren, F. H. J., Schoterman, M. H. C., Nauta, A., Gruppen, H., & Schols, H. A. (2014). Impact of galacto-oligosaccharides on the gut microbiota composition and metabolic activity upon antibiotic treatment during *in vitro* fermentation. *FEMS Microbiology Ecology*, 87, 41–51.
- Lezyk, M., Jers, C., Kjaerulf, L., Gotfredsen, C. H., Mikkelsen, M. D., & Mikkelsen, J. D. (2016). Novel α -L-fucosidases from a soil metagenome for production of fucosylated human milk oligosaccharides. *PLoS ONE*, 11. <https://doi.org/10.1371/journal.pone.0147438>
- Li, Y., Wang, H., Lu, L., Li, Z., Xu, X., & Xiao, M. (2009). Purification and characterization of a novel β -D-galactosidase with transglycosylation activity from *Bacillus megaterium* 2–37-4-1. *Applied Biochemistry and Biotechnology*, 158, 192–199. <https://doi.org/10.1007/s12010-008-8310-4>
- Mahajan, G. B., & Balachandran, L. (2017). Sources of antibiotics: Hot springs. *Biochemical Pharmacology*, 134, 35–41. <https://doi.org/10.1016/j.bcp.2016.11.021>
- Marchler-Bauer, A., Derbyshire, M. K., Gonzales, N. R., Lu, S., Chitsaz, F., Geer, L. Y., ... Bryant, S. H. (2015). CDD: NCBI's conserved domain database. *Nucleic Acids Research*, 43, D222–D226. <https://doi.org/10.1093/nar/gku1221>
- Mehta, R., Singhal, P., Singh, H., Damle, D., & Sharma, A. K. (2016). Insight into thermophiles and their wide-spectrum applications. *3 Biotech*, 6, 1–9. <https://doi.org/10.1007/s13205-016-0368-z>
- Montes, M. J., Mercadé, E., Bozal, N., & Guinea, J. (2004). *Paenibacillus antarcticus* sp. nov., a novel psychrotolerant organism from the Antarctic environment. *International Journal of Systematic and Evolutionary Microbiology*, 54, 1521–1526. <https://doi.org/10.1099/ijs.0.63078-0>
- Moro, G., Minoli, I., Mosca, M., Fanaro, S., Jelinek, J., Stahl, B., & Boehm, G. (2002). Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *Journal of Pediatric Gastroenterology and Nutrition*, 34, 291–295. <https://doi.org/10.1097/00005176-200203000-00014>
- Morrow, A. L., Ruiz-Palacios, G. M., Jiang, X., & Newburg, D. S. (2005). Human-milk glycans that inhibit pathogen binding protect breast-feeding infants against infectious diarrhea. *Journal of Nutrition*, 135, 1304–1307. <https://doi.org/10.1093/jn/135.5.1304>
- Mu, W., Chen, Q., Wang, X., Zhang, T., & Jiang, B. (2013). Current studies on physiological functions and biological production of lactosucrose. *Applied Microbiology and Biotechnology*, 97, 7073–7080. <https://doi.org/10.1007/s00253-013-5079-3>
- Park, A. R., Kim, H. J., Lee, J. K., & Oh, D. K. (2010). Hydrolysis and transglycosylation activity of a thermostable recombinant β -glucosidase from *Sulfolobus acidocaldarius*. *Applied Biochemistry and Biotechnology*, 160, 2236–2247. <https://doi.org/10.1007/s12010-009-8705-x>
- Park, H. Y., Kim, H. J., Lee, J. K., Kim, D., & Oh, D. K. (2008). Galactooligosaccharide production by a thermostable β -D-galactosidase from *Sulfolobus solfataricus*. *World Journal of Microbiology & Biotechnology*, 24, 1553–1558. <https://doi.org/10.1007/s11274-007-9642-x>

- Petrova, V. Y., & Kujumdzieva, A. V. (2010). Thermotolerant yeast strains producers of galacto-oligosaccharides. *Biotechnology and Biotechnological Equipment*, 24, 1612–1619. <https://doi.org/10.2478/V10133-010-0014-6>
- Rada, V., Nevorál, J., Trojanová, I., Tomá Nková, E., Mehilová, M. Š., & Killer, J. (2008). Growth of infant faecal bifidobacteria and clostridia on prebiotic oligosaccharides in in vitro conditions. *Anaerobe*, 14(4), 205–208. <https://doi.org/10.1016/j.anaerobe.2008.05.003>
- Ribeiro, A. L., Sánchez, M., Hidalgo, A., & Berenguer, J. (2017). Stabilization of enzymes by using thermophiles. In J.-L. Barredo, & I. Herráiz (Eds.), *Microbial Steroids: Methods and Protocols - Methods in Molecular Biology* (pp. 297–312). New York, NY: Springer New York. https://doi.org/10.1007/978-1-4939-7183-1_21
- Ruusunen, M., Surakka, A., Korkeala, H., & Lindström, M. (2012). *Clostridium tyrobutyricum* strains show wide variation in growth at different NaCl, pH, and temperature conditions. *Journal of Food Protection*, 75, 1791–1795. <https://doi.org/10.4315/0362-028X.JFP-12-109>
- Sahay, H., Yadav, A. N., Singh, A. K., Singh, S., Kaushik, R., & Saxena, A. K. (2017). Hot springs of Indian Himalayas: Potential sources of microbial diversity and thermostable hydrolytic enzymes. *3 Biotech*, 7, 1–11. <https://doi.org/10.1007/s13205-017-0762-1>
- Schmidt, M., & Stougaard, P. (2010). Identification, cloning and expression of a cold-active β -D-galactosidase from a novel Arctic bacterium, *Alkalilactibacillus ikkense*. *Environmental Technology*, 31, 1107–1114. <https://doi.org/10.1080/09593331003677872>
- Schultz-Johansen, M., Bech, P. K., Hennessy, R. C., Glaring, M. A., Barbeyron, T., Czjzek, M., & Stougaard, P. (2018). A novel enzyme portfolio for red algal polysaccharide degradation in the marine bacterium *Paraglaciecola hydrolytica* S66T encoded in a sizeable polysaccharide utilization locus. *Frontiers in Microbiology*, 9, 1–15. <https://doi.org/10.3389/fmicb.2018.00839>
- Searle, L. E. J., Best, A., Nunez, A., Salguero, F. J., Johnson, L., Weyer, U., ... La Ragione, R. M. (2009). A mixture containing galactooligosaccharide, produced by the enzymic activity of *Bifidobacterium bifidum*, reduces *Salmonella enterica* serovar Typhimurium infection in mice. *Journal of Medical Microbiology*, 58, 37–48. <https://doi.org/10.1099/jmm.0.004390-0>
- Shoaf, K., Mulvey, G. L., Armstrong, G. D., & Hutkins, R. W. (2006). Prebiotic galactooligosaccharides reduce adherence of enteropathogenic *Escherichia coli* to tissue culture cells. *Infection and Immunity*, 74, 6920–6928. <https://doi.org/10.1128/IAI.01030-06>
- Singla, V., & Chakkaravarthi, S. (2017). Applications of prebiotics in food industry: A review. *Food Science and Technology International*, 23, 649–667. <https://doi.org/10.1177/1082013217721769>
- Splechna, B., Nguyen, T. H., Steinböck, M., Kulbe, K. D., Lorenz, W., & Haltrich, D. (2006). Production of prebiotic galacto-oligosaccharides from lactose using β -galactosidases from *Lactobacillus reuteri*. *Journal of Agriculture and Food Chemistry*, 54, 4999–5006. <https://doi.org/10.1021/jf053127m>
- Sprenger, G. A., Baumgärtner, F., & Albermann, C. (2017). Production of human milk oligosaccharides by enzymatic and whole-cell microbial biotransformations. *Journal of Biotechnology*, 258, 79–91. <https://doi.org/10.1016/j.jbiotec.2017.07.030>
- Tamura, K., & Nei, M. (1993). Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution*, 10, 512–526. <https://doi.org/10.1093/oxfordjournals.molbev.a040023>
- Tan, J., Chai, F., & Zhang, C. (2015). Determination of human milk oligosaccharides in human breast milk by HPAE-PAD with on-line sample cleanup. *Costumer Application Note 119*, *Abbott Nutrition / Thermo Fisher Scientific*, 1–6.
- Tang, L., Li, Z. A., Dong, X. X., Yang, R. J., Zhang, J. H., & Mao, Z. G. (2011). Lactulose biosynthesis by β -D-galactosidase from a newly isolated *Arthrobacter* sp. *Journal of Industrial Microbiology and Biotechnology*, 38, 471–476. <https://doi.org/10.1007/s10295-010-0897-0>
- Tcherpakov, M., Ben-Jacob, E., & Gutnick, D. L. (1999). *Paenibacillus dendritiformis* sp. nov., proposal for a new pattern-forming species and its localization within a phylogenetic cluster. *International Journal of Systematic Bacteriology*, 49, 239–246. <https://doi.org/10.1099/00207713-49-1-239>
- Thompson, J. D., Higgins, D. G., & Gibson, T. J. (1994). CLUSTAL W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acid Research*, 22, 4673–4680. <https://doi.org/10.1093/nar/22.22.4673>
- Thurl, S., Henker, J., Taut, H., Tovar, K., & Sawatzki, G. (1993). Variations of neutral oligosaccharides and lactose in human milk during the feeding. *Zeitschrift Fur Ernährungswissenschaft*, 32, 262–269. <https://doi.org/10.1007/BF01611164>
- Torres, D. P. M., Gonçalves, M. D. P. F., Teixeira, J. A., & Rodrigues, L. R. (2010). Galacto-oligosaccharides: Production, properties, applications, and significance as prebiotics. *Compr Rev Food Sci Food Saf*, 9, 438–454. <https://doi.org/10.1111/j.1541-4337.2010.00119.x>
- Urbiet, M. S., Donati, E. R., Chan, K. G., Shahar, S., Sin, L. L., & Goh, K. M. (2015). Thermophiles in the genomic era: Biodiversity, science, and applications. *Biotechnology Advances*, 33, 633–647. <https://doi.org/10.1016/j.biotechadv.2015.04.007>
- Wanarska, M., & Kur, J. (2012). A method for the production of D-tagatose using a recombinant *Pichia pastoris* strain secreting β -D-galactosidase from *Arthrobacter chlorophenolicus* and a recombinant L-arabinose isomerase from *Arthrobacter* sp. 22c. *Microbial Cell Factories*, 11, 1–15. <https://doi.org/10.1186/1475-2859-11-113>
- Wang, H., Yang, R., Hua, X., Zhao, W., & Zhang, W. (2013). Enzymatic production of lactulose and 1-lactulose: Current state and perspectives. *Applied Microbiology and Biotechnology*, 97, 6167–6180. <https://doi.org/10.1007/s00253-013-4998-3>
- Watt, G. M., Lowden, P. A., & Flitsch, S. L. (1997). Enzyme-catalyzed formation of glycosidic linkages. *Current Opinion in Structural Biology*, 7, 652–660. [https://doi.org/10.1016/S0959-440X\(97\)80074-7](https://doi.org/10.1016/S0959-440X(97)80074-7)
- Wu, Y., Yuan, S., Chen, S., Wu, D., Chen, J., & Wu, J. (2013). Enhancing the production of galacto-oligosaccharides by mutagenesis of *Sulfolobus solfataricus* β -galactosidase. *Food Chemistry*, 138, 1588–1595. <https://doi.org/10.1016/j.foodchem.2012.11.052>
- Yu, L., & O'Sullivan, D. J. (2018). Immobilization of whole cells of *Lactococcus lactis* containing high levels of a hyperthermostable β -D-galactosidase enzyme in chitosan beads for efficient galacto-oligosaccharide production. *Journal of Dairy Science*, 101, 2974–2983. <https://doi.org/10.3168/jds.2017-13770>
- Zeuner, B., Muschiol, J., Holck, J., Lezyk, M., Gedde, M. R., Jers, C., ... Meyer, A. S. (2018). Substrate specificity and transglucosylation activity of GH29 α -L-fucosidases for enzymatic production of human milk oligosaccharides. *N Biotechnol*, 41, 34–45. <https://doi.org/10.1016/j.nbt.2017.12.002>
- Zeuner, B., Teze, D., Muschiol, J., & Meyer, A. S. (2019). Synthesis of human milk oligosaccharides: Protein engineering strategies for improved enzymatic transglycosylation. *Molecules*, 24(2033), 1–22. <https://doi.org/10.3390/molecules24112033>
- Zeyer, J., & Kocher, H. P. (1988). Purification and characterization of a bacterial nitrophenol oxygenase which converts ortho-nitrophenol to catechol and nitrite. *Journal of Bacteriology*, 170, 1789–1794. <https://doi.org/10.1128/jb.170.4.1789-1794.1988>
- Zhan, Y., Xu, Z., Li, S., Liu, X., Xu, L., Feng, X., & Xu, H. (2014). Coexpression of β -D-galactosidase and L-arabinose isomerase in the production of D-tagatose: A functional sweetener. *Journal of Agriculture and Food Chemistry*, 62, 2412–2417. <https://doi.org/10.1021/jf4042485>

How to cite this article: Thøgersen MS, Christensen SJ, Jepsen M, Pedersen LH, Stougaard P. Transglycosylating β -d-galactosidase and α -l-fucosidase from *Paenibacillus* sp. 3179 from a hot spring in East Greenland. *MicrobiologyOpen*. 2020;9:e980. <https://doi.org/10.1002/mbo3.980>

APPENDIX




FIGURE A1 Three dimension structure of the catalytic residues of (a) the *Paenibacillus* sp. 3179 β -D-galactosidase (amino acid backbone in green, nucleophile Glu and acid/base Glu in yellow) superimposed onto the corresponding *B. circulans* structure, 4MAD, (amino acid backbone in blue, nucleophile Glu233 and acid/base Glu157 (*B. circulans* numbering) in red), and (b) the *Paenibacillus* sp. 3179 α -L-fucosidase sequence (amino acid backbone in green) including the catalytic residues (nucleophile Asp and acid/base Glu in yellow) superimposed onto the corresponding *P. thiaminolyticus* structure 6GN6 (amino acid backbone in blue, nucleophile Asp186 and acid/base Glu239, *P. thiaminolyticus* numbering, in red)




FIGURE A2 SDS-PAGE of Ni-NTA-purified enzymes. Lanes 1–4: *Paenibacillus* sp. 3179 β -D-galactosidase at approx. 70 kDa [1:1,200 μ g; 2:240 μ g; 3:120 μ g; 4:24 μ g]. Lanes 5–8: *Paenibacillus* sp. 3179 α -L-fucosidase at approx. 50 kDa [5:450 μ g; 6:90 μ g; 7:45 μ g; 8:9 μ g]. Lane 9: 5 μ l PageRuler Unstained Protein Ladder (Thermo Fisher Scientific #26614). Enzyme concentrations were determined by Bradford assay with BSA as standard protein




FIGURE A3 Nonlinear regression and estimation of kinetic parameters for (a) the *Paenibacillus* sp. 3179 β -D-galactosidase and (b) the *Paenibacillus* sp. 3179 α -L-fucosidase. The experimental data points (black squares) were fitted to the Michaelis–Menten equation (red lines) to obtain the kinetic parameters K_m and V_{max} . The two curves did not reach full saturation, which is reflected in the relatively large errors on the parameters. Therefore, the values given here are rough estimates of the kinetic parameters