



UNIVERSIDADE ESTADUAL DE CAMPINAS
SISTEMA DE BIBLIOTECAS DA UNICAMP
REPOSITÓRIO DA PRODUÇÃO CIENTÍFICA E INTELLECTUAL DA UNICAMP

Versão do arquivo anexado / Version of attached file:

Versão do Editor / Published Version

Mais informações no site da editora / Further information on publisher's website:

https://www.scielo.br/scielo.php?script=sci_arttext&pid=S1517-83822016000100102

DOI: 10.1016/j.bjm.2015.11.006

Direitos autorais / Publisher's copyright statement:

© by Springer. All rights reserved.

DIRETORIA DE TRATAMENTO DA INFORMAÇÃO

Cidade Universitária Zeferino Vaz Barão Geraldo

CEP 13083-970 – Campinas SP

Fone: (19) 3521-6493

<http://www.repositorio.unicamp.br>



Industrial Microbiology

Structure of xanthan gum and cell ultrastructure at different times of alkali stress



Márcia de Mello Luvielmo^{a,*}, Caroline Dellinghausen Borges^b,
Daniela de Oliveira Toyama^c, Claire Tondo Vendruscolo^d,
Adilma Regina Pippa Scamparini^c

^a School of Chemistry and Food, Federal University of Rio Grande, RS, Brazil

^b Center of Chemical, Pharmaceutical and Food Sciences, Universidade Federal de Pelotas, RS, Brazil

^c Department of Food Sciences, Faculdade de Engenharia de Alimentos, Universidade Estadual de Campinas, SP, Brazil

^d Center of Technological Development, Biopolymers, Universidade Federal de Pelotas, Pelotas, RS, Brazil

ARTICLE INFO

Article history:

Received 7 April 2014

Accepted 29 July 2015

Associate Editor: Susana Marta Isay Saad

Keywords:

Xanthomonas campestris

Xanthan production

Alkali stress

Transmission microscopy

Scanning electron microscopy

ABSTRACT

The effect of alkali stress on the yield, viscosity, gum structure, and cell ultrastructure of xanthan gum was evaluated at the end of fermentation process of xanthan production by *Xanthomonas campestris* pv. *manihotis* 280-95. Although greater xanthan production was observed after a 24 h-alkali stress process, a lower viscosity was observed when compared to the alkali stress-free gum, regardless of the alkali stress time. However, this outcome is not conclusive as further studies on gum purification are required to remove excess sodium, verify the efficiency loss and the consequent increase in the polymer viscosity. Alkali stress altered the structure of xanthan gum from a polygon-like shape to a star-like form. At the end of the fermentation, early structural changes in the bacterium were observed. After alkali stress, marked structural differences were observed in the cells. A more vacuolated cytoplasm and discontinuities in the membrane cells evidenced the cell lysis. Xanthan was observed in the form of concentric circles instead of agglomerates as observed prior to the alkali stress.

© 2015 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Xanthan gum is a polysaccharide produced by bacterium species *Xanthomonas campestris*.¹ *Xanthomonas* cells are single straight rods, Gram-negative, 0.4–0.7 μm wide and 0.7–1.8 μm long.² Several biological functions, attributed to

the exopolysaccharide produced by these pathogenic bacteria, include protection against adverse environmental conditions such as drying, temperature oscillations, radiation, certain chemical products, and adhesion.³

The production and commercialization of xanthan gum as thickener and stabilizer has progressively increased at an annual rate of 5–10%,⁴ due to its better physicochemical

* Corresponding author.

E-mail: mmluvielmo@gmail.com (M. de Mello Luvielmo).

<http://dx.doi.org/10.1016/j.bjm.2015.11.006>

1517-8382/© 2015 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

properties as compared to other available polysaccharides. The most significant properties of xanthan gum include its high viscosity at low concentrations and stability for a wide range of temperatures and pH (even in the presence of salts).^{5,6}

Both microorganism growth and xanthan production are influenced by several factors that include the type of bioreactor, operation mode (batch or continuous), medium composition, cultivation conditions (temperature, pH, stirrer speed, air flow rate), fermentation time, strain of the microorganisms and post-fermentation conditions (heat treatment, recovery, purification). Due to the wide application of xanthan gum and its worldwide market, several studies have been performed to optimize xanthan gum production.^{4,7–13} However, the effect of this polysaccharide on the ultrastructure of cells and behavior of gum at different production stages have not yet been addressed properly, and thus the present study is relevant and may provide greater understanding on this process.

Peter et al.¹⁴ observed accumulation of xanthan around the cells cultured on agar using transmission electron microscopy, while Contreras et al.¹⁵ have reported plant-bacterial cell interactions during pathogenesis.

The present study aimed to evaluate the effect of alkali stress on yield, viscosity, gum structure, and cell ultrastructure, at the end of fermentation process of xanthan production by *X. campestris* pv. *manihotis* 280-95.

Material and methods

Microorganism

The lyophilized culture *X. campestris* pv. *manihotis* 280-95 was kindly gifted from the Culture Collection of the Phytopathological Bacteriology Section of the Biological Institute of Campinas (IBSBF), Campinas SP Brazil, and maintained in solid Yeast-Malt (YM) medium after activation.

Culture media

Following media were used for the culture of the bacterium: standard solid and liquid YM medium,¹⁶ medium I⁷ and nutrient medium.¹⁷

The standard YM medium consisted of (w/v) yeast extract 0.3%, malt extract 0.3%, peptone 0.5%, glucose 1.0%, and agar (for the solid medium) 2.5%.

The fermentation medium (Medium MP-I) comprised of (w/v) $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.02%, KH_2PO_4 0.5%, H_3BO_3 0.0006%, $(\text{NH}_4)_2\text{SO}_4$ 0.2%, FeCl_3 0.00024%, $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ 0.0002%, ZnSO_4 0.0002%, citric acid 0.20% and 5% sucrose. The pH of Medium I was adjusted to 7.0 with 0.1 M HCl or 0.1 M NaOH.

Culture media were sterilized in a vertical autoclave (Phoenix VA-75) at 121 °C for 20 min. Sucrose was sterilized separately and added to the media prior to use under aseptic conditions.

Inoculum production

An aliquot of 50 mL of standard YM medium was placed in three 250 mL-Erlenmeyer flasks with vented caps. Ten standard scoops (platinum 3.0 mm diameter-scoop) of culture

grown in standard solid YM for 48 h at 28 °C were added to each flask. The flasks were then incubated on an orbital shaker (New Brunswick Scientific Co. G27) at 28 °C and stirred at 250 rpm for 24 h. Cell suspensions with at least 2.6×10^9 CFU mL⁻¹¹⁸ were used as fermentation inoculum.

Biopolymer production

After 24 h, the inoculum was transferred aseptically to a 2 L-reactor vessel containing 1.35 L of fermentation medium (Medium MP-I) and 3 drops of pure Tween 80. New Brunswick's MULTIGEN fermenter monitored to provide an airflow at 1.5 vvm, at 28.5 ± 0.5 °C and 500 rpm for 72 h was used. The pH was initially adjusted to 7.0, and a change in pH was not followed throughout 72 h.

Alkali stress

After 72 h of fermentation, 15 mL of 10 M sodium hydroxide was added to the fermentation medium until an alkali stress at pH 12 was reached. The fermented medium was sampled after 1 h, 24 h, and 48 h. The inoculum production, fermentation process, and alkali stress were performed in triplicate.

Biopolymer separation

The culture medium was centrifuged at $22,300 \times g$ and 20 °C for 15 min to separate the biomass. Ethanol (92.8° GL) was added to the supernatant at a ratio of 4:1 to precipitate the biopolymer. The culture medium was stirred for 5 min, and the gum was collected on a wire mesh, placed on a glass Petri dish, frozen, and lyophilized. Then, the material was weighed and ground.

Xanthan yield

The lyophilized gum was placed in a desiccator and weighed on an analytical balance, and the gum yield was expressed by gum weight per volume of precipitated broth (g L⁻¹).

Viscosity

The rheological behavior of the samples was evaluated by a HAAK rheometer RS 150, using coaxial cylinder type sensor DG 41. The viscosity of the samples was determined at 25 °C. For that, 1.0% (w/v) xanthan solutions were prepared using distilled water, the mixture was stirred for 2 h, and then heated at 60 °C for 20 min.¹⁹ The rheological analysis was performed in triplicate, using shear rate from 0.01 s⁻¹ to 60 s⁻¹, with 5100 mm slot for 300 s.

Scanning electron microscopy

The structure of xanthan gum powder produced without alkali stress (W/AS) and with alkali stress (AS) at pH 12 for 1 h was assessed by scanning electron microscopy.

Gum samples were first fixed overnight in a solution containing 0.1 M sodium cacodylate buffer pH 7.4, 25% glutaraldehyde and tannic acid, and then post-fixed in osmium tetroxide. Pellets were washed thrice with sodium cacodylate

buffer pH 7.4, between the first and second fixation procedure, and at the end of the second fixation. The samples were then subjected to a dehydration series in 30%, 50%, 70%, and 90% (v/v) ethanol solutions for about 20 min each. Critical point drying of the samples was accomplished via Balzers CPD 030. The samples were fixed to metal stubs and gold coated with a sputter coater (Balzers CD 050 Sputter 5), followed by measurements at 15 kV in a scanning electron microscope JSM 5800 LV JEOL.^{20,21}

Transmission electron microscopy

The samples subjected to transmission electron microscopy were: pure inoculum (PI), initial fermentation (IF), final fermentation (W/AS) (after 72 h fermentation), and alkali stress for 1 h (AS1h), 24 h (AS24h) and 48 h (AS48h).

The samples were collected at their appropriate time point according to fermentation pathway, and stored at 4 °C. Then they were centrifuged at 12,200 × g for 5 min at room temperature and re-suspended in ultra-pure water, stirred, and centrifuged again under the same conditions. The procedure was repeated thrice. The pellets were fixed overnight in a solution containing 0.1 M sodium cacodylate buffer pH 7.4, 25% glutaraldehyde, and tannic acid, and then post-fixed in osmium tetroxide for 2 h. The pellets were washed thrice with sodium cacodylate buffer between the first and second fixation procedures and at the end of the second fixation. The samples were then subjected to a dehydration series in ethanol solutions, absolute ethanol, a mixture of propylene oxide and absolute ethanol (1:1), and propylene oxide alone, for approximately 20 min each, with the exception of propylene oxide. The resin and propylene oxide (1:1) remained on the rotator apparatus for 2 h. The oxide was removed and blended with the pure resin. The Eppendorf tubes remained inside the rotator overnight. After inclusion, polymerization was carried out at 60 °C for 48 h. The material was sectioned using ultramicrotome Leica Ultra Cut, and observed under a Zeiss LEO 906 Transmission Electron Microscope, at a magnification between 6000 and 27800×.^{20,22,23}

Results and discussion

Fermentation and alkali stress process

As shown in Fig. 1, 9.4 gL⁻¹ of xanthan gum was obtained after 72 h of fermentation while 11.8 gL⁻¹ of xanthan gum was obtained after 1 h of alkali stress, which was 25.7% higher than the amount obtained at the end of the fermentation. After 24 h of alkali stress, the amount of biopolymer per liter of fermentation medium increased to 16.5 gL⁻¹, translating to a 74.8% increase in xanthan production as compared to the conventional process without alkali stress, which after 48 h decreased to 14.2 gL⁻¹.

The present results indicate that the alkali stress might have induced the cells to produce a greater amount of exopolysaccharides as a protective mechanism against adverse conditions of the medium, while the lower yield after 48 h of alkali stress could be due to the alkaline hydrolysis of the polymer.




Fig. 1 – Effect of alkali stress time on xanthan production (g L⁻¹) by *X. campestris* pv. *manihotis* 280-95. W/AS, final fermentation without alkali stress; AS1h, alkali stress for 1 h; AS24h, alkali stress for 24 h; AS48h, alkali stress for 48 h.

Low production rates were observed when pH was not controlled during processing. Papagianni et al.¹¹ evaluated xanthan production by *X. campestris* ATCC 1395 under non-controlled pH conditions at different stirring speeds. The highest yield obtained by them was approximately 6.5 gL⁻¹ at 600 rpm, which is lower than that obtained in the current study. In contrast, Borges et al.²⁴ studied the xanthan production by *Xanthomonas arboricola* pv. *pruni* 106 under non-controlled pH conditions, and found slightly higher production rate of 10.2 gL⁻¹ at 400 rpm.

Different behaviors have been observed under non-controlled pH conditions, as the medium pH might either decrease from neutral to near pH 5 or even increase, depending on several factors including medium composition, strain used and processing conditions.^{11,24,25}

However, the present study demonstrates that the gum production tends to increase with alkali stress or pH control. Gupte and Kamat²⁶ evaluated xanthan production by *X. campestris* ICA-125 at different pH values (4–9), and observed the highest yield at pH 6 with a production rate of 18.6 gL⁻¹, which decreased and reached 3.1 gL⁻¹ at pH 9. In contrast, Borges et al.²⁴ found a slight reduction in xanthan production at pH 9 when compared to pH 7, with values of 16.05 and 16.9 gL⁻¹, respectively. Liakopoulou-Kyriakides et al.²⁷ investigated xanthan production by *X. campestris* XLM1521 and found an increase from pH 6 to 8, reaching 17.3 gL⁻¹ at pH 8. These values were close to those obtained in the current study using alkali stress.

As shown in Fig. 2, the 1% solutions prepared from xanthan gums obtained from the conventional fermentation alkali stress free (W/AS) and with alkali stress process for 1 h (AS1h), 24 h (AS24h), and 48 h (AS48h) exhibited a pseudoplastic fluid behavior, as also observed by other authors.^{6,28-31} The samples not subjected to alkali stress (W/AS) exhibited apparent viscosity of 317 mPa s at a shear rate of 10 s⁻¹, while the samples subjected to AS for 1 h, 24 h, and 48 h yielded viscosity values of approximately 117 mPa s, under the same shear rate. It is worth noting that the quality of the gum obtained after the alkali stress was 64% lower as compared to the gum obtained




Fig. 2 – Apparent viscosity (mPa s) of xanthan solutions (1.0%, w/v) as a function of shear rate (s^{-1}) at 25°C. W/AS, final fermentation without alkali stress; AS1h, alkali stress for 1 h; AS24h, alkali stress for 24 h; AS48h, alkali stress for 48 h.

without the alkali stress process while the time of alkali stress did not affect the quality of the gum obtained in the present study.

Similar results were observed by Gupte and Kamat²⁶ and Borges et al.,²⁴ who also evaluated the viscosity of the gum produced at different pH values.

As reported by Gupte and Kamat,²⁶ the viscosity of the xanthan solution increased by 1% when produced at pH 4–7 (153 mPa s to 559 mPa s, respectively), with a decrease at pH 8 and pH 9 (280 mPa s to 124 mPa s, respectively).

Borges et al.²⁴ have shown that the gum produced under non-controlled pH conditions and at pH 5 (76.3 mPa s and 77.7 mPa s, respectively) had higher viscosity rates when compared to that produced at pH 7 and pH 9 (52.8 mPa s and 50.4 mPa s, respectively), at a shear rate of $19 s^{-1}$, using 0.5% solution. According to these authors, this phenomenon could be due to the lower molecular weight of the xanthan produced at higher pH values.

Besides molecular weight, the concentration of xanthan might affect the viscosity when associated with salts. Zhong

et al.³² demonstrated that at low xanthan concentration, addition of salt (Na^+ and Ca^{2+}) lowered the solution viscosity, while at higher xanthan concentration (5000 mg/L), the salt addition increased the viscosity.

Thus, a possible explanation for the lower viscosity of xanthan produced under alkali stress conditions might be the hydrolysis of gum under these conditions and electrostatic repulsion in the polymer, as shown by the microscopy results.

Gum structure with and without alkali stress

As depicted in Fig. 3A and B, the gum without alkali stress showed a polygonal structure, whereas the gum subjected to alkali stress had star-shaped structure. While comparing these results with those found by Borges et al.,²⁴ it was assumed that no reduction in molecular weight occurred. It has also been suggested that the high sodium concentration due to the alkali might lead to the structure compaction due to the repulsion caused by the positively charged ions. The present results corroborate with those obtained by Carrington et al.³³ on the effect of salts on the extension flow properties. In fact, high salt concentration may change xanthan molecule from an expanded to a compact conformation. The changes in the molecule hydrodynamic size can lead to a decrease in viscosity, as observed in the current study.

However, opposite behavior was observed by Savi-Junior et al.,³⁴ who found that the viscosity decreased at low salt concentration as the polymer coil became contracted, providing a smaller hydrodynamic volume.

Ultrastructure analysis

Pure inoculum (PI)

The cells showed presence of the features typical for *Xanthomonas* (Fig. 4A). Although the bacterium appeared as rods, the histological ultracuts by ultramicrotome revealed that the bacterium might appear in longitudinal, sagittal, and transversal cuts. In fact, Fig. 4A revealed that the outer membrane and cytoplasmic structures of the bacterial cells were intact. No bacterial lysis was observed, thus indicating that cells from the pure inoculum were viable. Cells division can also be observed through a bifurcation followed by stretching in the middle of the cell.




Fig. 3 – Scanning electron microscopy of xanthan gum obtained by fermentation with *X. campestris* pv. *manihotis* 280-95, at 15 kV and 10,000 \times magnification: (A) final fermentation without alkali stress and (B) with 1 h alkali stress.




Fig. 4 – Transmission electron microscopy of: (A) pure inoculum, at 60 kV and 12,930× magnification; (B) initial fermentation, at 60 kV and 6000× magnification; (C) final fermentation, at 60 kV and 16,700× magnification; (D) alkali stress for 1 h, at 60 kV and 27,800× magnification.

The production of xanthan gum around some cells in very low quantity was also observed. During the preparation of inoculum, an increase in cell concentration was observed, and only a small amount of gum was produced. The gum surrounding the cell prevented the transport of nutrients and extended the growth phase.^{35,36}

Initial fermentation (IF)

The bacterial cells were intact at the beginning of the fermentation (Fig. 4B), which suggests that the initial stages of the fermentation process occurred as expected. As shown in Fig. 4B, a large amount of gum was found around the bacterial cells.

Since the fermentation medium (medium I) contained various salts, it was believed that xanthan gum surrounding the bacteria moved away during fermentation (Fig. 4B), probably due to the changes in osmolality.

The extracellular polysaccharides may be classified into two types: (I) polymers physically attached to the microbial cell in a capsule form, and (II) polymers excreted by the microbial cells in the medium in the form of loose slime.³⁷ The present results suggest a revision of such classification, since xanthan

gum can be classified either as type (I) or type (II) depending on the medium conditions.

Final fermentation without alkali stress (W/AS)

Early structural changes in bacteria, such as cytoplasmic vacuolization and membrane discontinuity indicated an early stage of bacterial lysis (Fig. 4C).

The prolonged fermentation time of 72 h at this stage led to different gum behavior, since xanthan gum was present as agglomerates rather than around the bacteria, possibly due to the different osmolality than that is ideal for bacterial growth.

Alkali stress after 1 h (AS1h)

The alkali stress conditions of this study consisted in raising pH to 12.0. After 1 h (AS1h), marked structural differences in the bacterial cells were observed (Figs. 4D, 5A and B). A more vacuolated cytoplasm and the discontinuity in the cell membranes as evidence of cell lysis were observed (Fig. 4D).

Despite the considerable decrease in cells number, the image at lowest magnification (Fig. 5B) revealed the presence of certain bacterial cells.




Fig. 5 – Transmission electron microscopy of alkali stress: (A) for 1 h, at 60 kV and 12,930 \times magnification; (B) at 60 kV and 6000 \times magnification; (C) alkali stress for 24 h, at 60 kV and 10,000 \times magnification; (D) alkali stress for 48 h, at 60 kV and 7750 \times magnification.

Xanthan gum was found in an organized shape (Fig. 5B) rather than agglomerates as reported earlier. This was quite similar to water–oil emulsion system with a hydrophobic interior and a polysaccharide xanthan gum on the outer layer. Such an organized conformation, in the form of concentric circles, had never been reported in earlier studies.

Alkali stress after 24 h (AS24h)

After the alkali stress of 24 h, no bacterial cells were observed, which indicates a possibility of complete disintegration of the bacterial membranes (Fig. 5C). A more organized shape with concentric circles was observed in this case (Fig. 5C) in contrast to that obtained after 1 h of alkali stress.

Co-occurrence of acetic and pyruvic acid molecules with xanthan molecules results in anion-type of polysaccharides.^{5,37,38} It is believed that when this biopolymer is subjected to strong alkaline conditions, the charges are neutralized prior to the electrostatic repulsion. In the absence of repulsion, there might be some changes in the conformation of the molecule (Fig. 3B) and in the way the gum is deposited (Fig. 5C).

Alkali stress after 48 h (AS48h)

We did not see much difference between the results obtained after 48 h and 24 h of alkali stress (Fig. 5D).

Conclusion

Xanthan production was enhanced by a 24 h-alkali stress process in *X. campestris* pv. *manihotis* 280-95 strain while grown in a 2 L bioreactor. However, regardless the alkali stress time, xanthan gum was observed to be of lower viscosity as compared to the alkali stress-free gum. Nevertheless, this result is not conclusive as further studies are required such as gum purification to remove the excess sodium, and verification of the efficiency loss and consequent increase in the polymer viscosity.

Alkali stress altered the structure of xanthan from polygon-like shape to a star-like form. At the end of the fermentation, early structural changes in the cells were observed. Whereas, marked structural differences were observed in the cells after the alkali stress process. A more vacuolated cytoplasm and discontinuities in the cell membranes indicated occurrence of the cell lysis. The gum was seen to be in the form of concentric

circles after the alkali stress instead of the agglomerates as seen prior to the alkali stress.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgments

The authors would like to thank CNPq for the financial support and the Biological Institute of Campinas for providing the bacterial strain.

REFERENCES

- Sutherland IW, Xanthan. In: Swings JG, Civerolo EL, eds. *Xanthomonas*. London: Chapman & Hall; 1993:363–388; Rudolph K. Infection of the plant by *Xanthomonas*. In: Swings JG, Civerolo EL, eds. *Xanthomonas*. London: Chapman & Hall; 1993:121–146.
- Bradbury JF. Genus II: *Xanthomonas*. In: Krieg NR, Holt CG, eds. *Manual of Systematic Bacteriology*. Baltimore: Williams & Wilkins; 1984:199–210.
- Romeiro RS. *Bactérias fitopatogênicas*. Viçosa: UFV Impr. Univer.; 1995.
- Salah RB, Chaari K, Besbes S, et al. Optimisation of xanthan gum production by palm date (*Phoenix dactylifera* L.) juice by-products using response surface methodology. *Food Chem*. 2010;121:627–633.
- García-Ochoa F, Santos VE, Casas JA, Gómez E. Xanthan gum: production, recovery, and properties. *Biotechnol Adv*. 2000;18:549–579.
- Faria S, Petkowicz CLO, Sérgio Morais AL, et al. Characterization of xanthan gum produced from sugar cane broth. *Carb Polym*. 2011;86:469–476.
- Souw P, Demain AL. Nutritional studies on xanthan production by *X. campestris* NRRL B-1459. *Appl Environ Microbiol*. 1979;37:1186–1192.
- Kennedy JF, Jones P, Baker A. Factors affecting microbial growth and polysaccharides production during the fermentation of *Xanthomonas campestris* cultures. *Enzyme Microb Technol*. 1982;4:39–43.
- Esgalhado ME, Roseiro JC, Collaço MTA. Interactive effects of pH and temperature on cell growth and polymer production by *Xanthomonas campestris*. *Process Biochem*. 1995;30:667–671.
- Moreira AS, Vendruscolo JLS, Gil-Turnes C, Vendruscolo CT. Screening among 18 novel strains of *Xanthomonas campestris* pv. *pruni*. *Food Hydrocoll*. 2001;15:469–474.
- Papagianni M, Psomas SK, Batsilas L, Paras SV, Kyriakidis DA, Liakopoulou-Kyriakides M. Xanthan production by *Xanthomonas campestris* in batch cultures. *Process Biochem*. 2001;37:73–80.
- Borges CD, da Moreira AS, Vendruscolo CT, Ayub MAZ. Influence of agitation and aeration in xanthan production by *Xanthomonas campestris* pv. *pruni* strain 101. *Rev Argent Microbiol*. 2008;40:81–85.
- Mabrouk MEM, Ahwany MADE, Beliah MMB, Sabry SA. Xanthan production by a novel mutant strain of *Xanthomonas campestris*: application of statistical design for optimization of process parameters. *Life Sci J*. 2013;10:1660–1667.
- Peters HU, Herbst H, Hesselink PG, Lünsdorf H, Schumpe A, Deckwer WD. The influence of agitation rate on xanthan production by *Xanthomonas campestris*. *Biotechnol Bioeng*. 1989;34:1393–1397.
- Contreras N, Trujillo G, Borges O, Centeno F. Análisis ultraestructural de la interacción de *Xanthomonas axonopodis* pv. *phaseoli* con genotipos resistentes, moderadamente resistentes y susceptibles de *Phaseolus vulgaris* L. *Interciencia*. 2001;26:554–557.
- Haynes WC, Wickerham LJ, Hesseltine CW. Maintenance of cultures of industrially important microorganisms. *Appl Microbiol*. 1955;3:361–368.
- Pelczar MJ, Reid R, Chan ECS. *Microbiologia*. São Paulo: Mc Graw-Hill do Brasil; 1981.
- Pan A, Moreira AS, Vendruscolo CT. Efeito da concentração inicial do inóculo no crescimento celular e qualidade de biopolímeros de *X. campestris* pv. *pruni* cepa 06. *Rev Bras Agrociência*. 2000;6:273–277.
- Xuewu Z, Xin L, Dexiang G, Wei Z, Tong X, Yonghong M. Rheological models for xanthan gum. *J Food Eng*. 1996;27:203–209.
- Mercer EH, Birbeck MSC. *Manual de microscopia eletrônica para biólogos*. Madrid, España: Ed. Blume; 1972.
- Santi-Gadelha T, de Almeida Gadelha CA, Aragão KS, et al. Purification and biological effects of *Araucaria angustifolia* (Araucariaceae) seed lectin. *Biochem Biophys Res Commun*. 2006;350:1050–1055.
- Johannessen JV. *Electron Microscopy in Human Medicine, Instrumentation and Techniques*. Great Britain: McGraw Hill International Book Company; 1978.
- Rádis-Baptista G, Moreno FBMB, Nogueira LL, et al. Crotacetin a novel snake venom of c-type lectin, is homolog convulxin. *Cell Biochem Biophys*. 2006;44:412–423.
- Borges CD, Paula RCM, Feitosa JPA, Vendruscolo CT. The influence of thermal treatment and operational conditions on xanthan produced by *X. arboricola* pv. *pruni* strain 106. *Carb Polym*. 2009;75:262–268.
- Psomas SK, Liakopoulou-Kyriakides M, Kyriakidis DA. Optimization study of xanthan gum production using response surface methodology. *Biochem Eng J*. 2007;35:273–280.
- Gupte MD, Kamat MY. Isolation of wild *Xanthomonas* strains from agricultural produce, their characterization and potential related to polysaccharide production. *Folia Microbiol (Praha)*. 1997;42:621–628.
- Liakopoulou-Kyriakides M, Tzanakakis ES, Kiparissidis C, Ekaterianiadou LV, Kyriakidis DA. Kinetics of xanthan gum production from whey by constructed strain of *X. campestris* in batch fermentations. *Chem Eng Technol*. 1997;20:354–360.
- Nitschke M, Thomas RWSP, Knauss C. Xanthan production by *X. campestris* in a whey-based medium. *Rev Microbiol*. 1997;28:148–151.
- Katzbauer B. Properties and applications of xanthan gum. *Polym Degrad Stabil*. 1998;59:81–84.
- Cacik F, Dondo RG, Marqués D. Optimal control of a batch bioreactor for the production of xanthan gum. *Comput Chem Eng*. 2001;25:409–418.
- Bueno SM, Garcia-Cruz CH. The influence of fermentation time and the presence of salts in the rheology of the fermentation broth of a polysaccharide-producing bacteria free of soil. *J Food Eng*. 2001;50:41–46.
- Zhong L, Oostrom M, Truex MJ, Vermeul VR, Szecsody JE. Rheological behavior of xanthan gum solution related to shear thinning fluid delivery for subsurface remediation. *J Hazard Mater*. 2013;244–245:160–170.
- Carrington S, Odell J, Fisher L. Polyelectrolyte behaviour of dilute xanthan solutions: salt effects on extensional rheology. *Polymer*. 1996;37:2871–2875.
- Savi-Junior R, Freitas RA, Sassak GL, Koop HS, Silveira JLM. Real-time monitoring of the change in stiffness of single-strand xanthan gum induced by NaCl. *Food Hydrocoll*. 2015;44:191–197.

-
35. Pons A, Dussap CG, Gros JB. Modelling *Xanthomonas campestris* batch fermentations in a bubble column. *Biotechnol Bioeng.* 1989;33:394–405.
 36. Pons A, Dussap CG, Gros JB. Xanthan batch fermentations: compared performance of a bubble column and a stirred tank fermentor. *Biop Eng.* 1990;5:107–114.
 37. Kennedy JF, Bradshaw IJ. Production, properties and applications of xanthan. *Prog Ind Microbiol.* 1984;19: 319–371.
 38. Abbaszadeh A, Lad M, Janin M, et al. A novel approach to the determination of the pyruvate and acetate distribution in xanthan. *Food Hydrocoll.* 2015;44:162–171.