



UNIVERSIDADE ESTADUAL DE CAMPINAS
SISTEMA DE BIBLIOTECAS DA UNICAMP
REPOSITÓRIO DA PRODUÇÃO CIENTÍFICA E INTELLECTUAL DA UNICAMP

Versão do arquivo anexado / Version of attached file:

Versão do Editor / Published Version

Mais informações no site da editora / Further information on publisher's website:

<https://isident.org/DOIx.php?id=10.5624/isd.2020.50.2.181>

DOI: 10.5624/isd.2020.50.2.181

Direitos autorais / Publisher's copyright statement:

©2020 by Korean Academy of Oral and Maxillofacial Radiology. All rights reserved.

DIRETORIA DE TRATAMENTO DA INFORMAÇÃO

Cidade Universitária Zeferino Vaz Barão Geraldo

CEP 13083-970 – Campinas SP

Fone: (19) 3521-6493

<http://www.repositorio.unicamp.br>

COVID-19 and oral radiology

Wilson Gustavo Cral^{1,*}, Carlos Augusto Souza Lima², Dagmar de Paula Queluz¹

¹Department of Oral Diagnosis, Division of Oral Radiology, Piracicaba Dental School, University of Campinas (UNICAMP), Piracicaba, São Paulo, Brazil

²Faculdade São Leopoldo Mandic, Instituto de Pesquisas São Leopoldo Mandic, Division of Oral Radiology and Department of Stomatology, Public Health and Forensic Dentistry, School of Dentistry of Ribeirão Preto - University of São Paulo, São Paulo, Brazil

COVID-19 and oral radiology

Dear Editors,

The global coronavirus disease 2019 (COVID-19) pandemic and its impact on people's quality of life and the practice of health professionals have been the subject of many studies aiming to establish protective procedures in hospital and clinical practice environments. In dentistry, more specifically in the oral radiology field, certain procedures must be strictly followed when performing radiographic techniques to reduce the risk of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) transmission.

Initially, it is important to follow the basic requirements for radiological protection in radiodiagnosis, and all health care workers and patients should be thoroughly encouraged to meticulously wash their hands for at least 20 seconds, along with disinfection with 70% gel alcohol whenever possible.¹

Radiological clinic staff must use a full array of individual protective equipment, that is, a head cap, shoe protectors, goggles, a surgical gown, and gloves. Patients and their companions should always wear masks, protective aprons, and shoe protectors in the clinical setting, and should be carefully asked about previous risk exposure (i.e., travel and contact with possible infected persons) and whether they have experienced any symptoms in the last 14 days. Additionally, their body temperature should be measured upon entrance. It is also recommended that pa-

tients should be selected by a screening service, via internet programs and applications, that prioritizes clinical conditions for which a radiographic examination is considered crucial.¹ These conditions include dental urgency, such as extractions and coronary openings of irreversible pulpitis. Non-emergency dental procedures should be postponed, with the intention of reducing the number of patients treated per day to avoid cross-contact and proximity as much as possible.

COVID-19 has identified in the saliva of infected patients,² which is considered the main route of human-to-human transmission. In light of the risk of intraoral radiographic techniques causing cough stimuli, saliva secretion, and vomiting reflex in some patients,³ extrabuccal radiographic techniques, such as panoramic radiography and cone-beam computed tomography, are preferable when possible during the COVID-19 pandemic.³ When the use of intraoral techniques is strictly necessary, the patient should perform mouth-washing (e.g., 9 mL of 1.0%-1.5% hydrogen peroxide for 30 seconds) to reduce the number of microorganisms in the oral cavity.^{1,4}

SARS-CoV-2 may persist on surfaces for a few hours or up to several days depending on the temperature of the environment.⁵ For this reason, it is necessary to completely disinfect all surfaces of the X-ray room and adjacent areas with 0.5% sodium hypochlorite before and after all services, including the floor, door handles, light switch, table, support bench, and chairs. It is recommended to disinfect components of the radiographic device, such as the X-ray control panel and tube head, according to the manufacturers' guidance.¹

Although aerosol-generating procedures are not performed in oral radiology, contact with the patient's oral cavity may result in the transmission of SARS-CoV-2. In

Received May 18, 2020; Revised May 22, 2020; Accepted May 27, 2020

*Correspondence to : Prof. Wilson Gustavo Cral

Department of Oral Diagnosis, Division of Oral Radiology, Piracicaba Dental School, University of Campinas (UNICAMP), Av. Limeira, 901 - Areião, Piracicaba - SP, 13414-903, Brazil
Tel) 55-19-2106-5209, E-mail) wgcral@gmail.com

Copyright © 2020 by Korean Academy of Oral and Maxillofacial Radiology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Imaging Science in Dentistry · pISSN 2233-7822 eISSN 2233-7830

order to minimize this risk, it is necessary to comply with essential biosafety procedures in the clinical environment and to pay careful attention to the criteria for prescribing radiographic examinations, as well as reducing the number of employees working and people circulating in the clinical environment.

References

1. Wan YL, Schoepf UJ, Wu CC, Giovagnoli DP, Wu MT, Hsu HH, et al. Preparedness and best practice in Radiology Department for COVID-19 and other future pandemics of severe acute respiratory infection. *J Thorac Imaging* (in press).
2. Sabino-Silva R, Jardim AC, Siqueira WL. Coronavirus COVID-19 impacts to dentistry and potential salivary diagnosis. *Clin Oral Investig* 2020; 24: 1619-21.
3. Vandenberghe B, Jacobs R, Bosmans H. Modern dental imaging: a review of the current technology and clinical applications in dental practice. *Eur Radiol* 2010; 20: 2637-55.
4. Meng L, Hua F, Bian Z. Coronavirus disease 2019 (COVID-19): emerging and future challenges for dental and oral medicine. *J Dent Res* 2020; 99: 481-7.
5. World Health Organization. Report of the WHO-China joint mission on Coronavirus disease 2019 (COVID-19) [Internet]. Geneva: World Health Organization; 2020 [cited 2020 May 17]. Available from [https://www.who.int/publications-detail/report-of-the-who-china-joint-mission-on-coronavirus-disease-2019-\(covid-19\)](https://www.who.int/publications-detail/report-of-the-who-china-joint-mission-on-coronavirus-disease-2019-(covid-19)).