

CHEMICAL WEED AND BRUSH CONTROL



SUGGESTIONS FOR RANGELAND

2009 UPDATE

*This publication provides an update for some of the herbicide recommendations contained in B-1466.
The herbicide control recommendations contained in this update supercede any information contained in the original publication.*

Common, Chemical and Product Names of Herbicides*

Herbicide common name	Chemical name	Product name	Active ingredient or acid equivalent
Aminopyralid	2-pyridine carboxylic acid, 4-amino-3, 6-dichloro-2-pyridine carboxylic acid, triisopropanolammonium salt	Milestone	2 lbs./gal.
Aminopyralid:2,4-D (1:8)	See Aminopyralid and 2,4-D	GrazonNext	3 lbs./gal.
Clopyralid	3,6-dichloro-2-pyridinecarboxylic acid	Reclaim, Pyramid R&P, Clopyralid 3	3 lbs./gal.
2,4-D	(2,4-dichlorophenoxy) acetic acid	Weedar 64, Formula 40, Hi-Dep, Weedone LV4, Esteron 99 and others	amine salts, free acids and esters of variable concentration
Dicamba	3,6-dichloro-2-methoxybenzoic acid	Banvel, Clarity, Vision	4 lbs./gal.
Dicamba:2,4-D(1:3)	See Dicamba and 2,4-D	Weedmaster, Banvel + 2,4-D, RangeStar, Outlaw	4 lbs./gal.
Diesel fuel oil or kerosene	refined petroleum fractions	Several manufacturers	
Fluroxypyr	1-methylheptyl ester: ((4-amino-3,5-dichloro-6-fluoropyridin-2-yl)oxy)acetic acid	Vista	1.5 lbs./gal.
Glyphosate	N-(phosphonomethyl) glycine	Several including Rodeo**, Roundup, Roundup Ultradry, Glyphosate 417	isopropylamine salt, concentration varies depending on the product
Hexazinone	3-cyclohexyl-6-(dimethylamino)-1-methyl-1,3,5-triazine-2,4(IH, 3H)-dione	Velpar L, Pronone Power Pellet	2 lbs./gal. (Velpar L) 75% (Pronone Power Pellet)
Imazapyr	2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid	Arsenal, Habitat**	2 lbs./gal.
Metsulfuron methyl	methyl 2[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]amino] sulfonyl] benzoate	Escort, Clean Pasture, MSM 60DF	60%
Metsulfuron:Chlorosulfuron (3:1)	See Metsulfuron methyl + 2-Chloro-N-[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)aminocarbonyl] benzenesulfonamide	Cimarron Plus	48% metsulfuron 15% Chlorosulfuron
Metsulfuron:Chlorosulfuron (1:1)	See Metsulfuron: Chlorosulfuron	Cimarron X-Tra	30% Metsulfuron 37.5% Chlorosulfuron
Metsulfuron methyl Dicamba:2,4-D(1:3)	See Metsulfuron methyl, dicamba and 2,4-D	Cimarron Max	60% (Part A) 3.87 lbs./gal. (Part B)
Picloram	4-amino-3,5,6-trichloro-2-pyridinecarboxylic acid	Tordon 22K, Triumph 22K, Picloram 22K	2 lbs./gal.
Picloram:Fluroxypyr (1:1)	See Picloram and Fluroxypyr	Surmount	1.34 lbs./gal.
Picloram:2,4-D(1:4)	See Picloram and 2,4-D	Grazon P+D, Gunslinger, Picloram + D	2.5 lbs./gal.
Tebuthiuron	N-[5-(1,1-dimethylethyl)-1,3,4-thiadiazol-2-yl]-N-N'-dimethylurea	Spike 20P, Spike 80 DF	20% (Spike 20P) 80% (Spike 80 DF)
Triclopyr	{(3,5,6-trichloro-2-pyridinyl)oxy}acetic acid	Clear Pasture, Pathfinder II, Triclopyr R&P Remedy Ultra, Triclopyr 4 EC	0.75 lbs./gal. (Pathfinder II) 4 lbs./gal. (all others)
Triclopyr:Fluroxypyr (3:1)	See Triclopyr and 1-methylheptyl ester:((4-amino-3,5-dichloro-6-fluoropyridin-2-yl)oxy)acetic acid	PastureGard	2.0 lbs./gal.
Triclopyr:2,4-D(1:2)	See Triclopyr and 2,4-D	Crossbow	3 lbs./gal.

*Herbicides have been identified by the accepted Weed Science Society of America common name, and when practical, one or more product names.

**Aquatic label

Items in gray boxes are recent changes approved by the Rangeland Herbicide Use Committee.

Page	Weed Controlled	Herbicide (common and chemical names – page 4)	Herbicide quantity (active ingredient rate in parenthesis)		Spray volume (per acre for broadcast, as described for individual plant)	Time to apply	Remarks
			Broadcast rate per acre	Individual plant/spot treatment			
6	African rue	Velpar L		VH** 2 ml/plant	Use an exact delivery handgun to apply undiluted herbicide to soil surface at the edge of the plant canopy.	Spring or summer.	Do not use on heavy clay or caliche soils.
		Pronone Power Pellets Spike 20 P		1 pellet/plant			
		Arsenal	H 1 qt. (0.5 lbs.)	VH 0.5%	10 to 25 gals./acre for ground broadcast. Thoroughly wet foliage for individual plant treatment. Add 1 to 2 qts. surfactant per 100 gals. water.	Late September through October (to first frost).	Applications should be made to fall regrowth that is in good growing condition. Recommend using individual plant treatment when growing with desirable vegetation to reduce non-target damage. Arsenal is a non-selective herbicide and will kill or damage desirable vegetation if sprayed.
6	Berlander lobelia, bitter sneezeweed, (annual or common), buffalobur, camphorweed, cocklebur, croton, horehound, marshelder (sumpweed or sulfaweed), plantain (tallowweed), prairie gerardia (see remarks), ragweed, smartweed, sunflower, thistles, Western bitterweed (see remarks), Western ragweed, wild carrot and others	2,4-D amine or low volatile ester	VH 1 pt. to 1 qt. (½ to 1 lb.) 4 lbs./gal. product	VH 1% (4 lbs./gal. product)	2 to 4 gals. water for aerial spray; 10 to 25 gals. water for ground broadcast application. Thoroughly wet foliage for individual plant treatment. Add 1 to 2 qts. surfactant per 100 gals. of water.	Spring, weed 4 to 6 inches high, good moisture conditions. Spray thistles in rosette stage.	Use 2,4-D amine in areas with 25 inches of rainfall or more. Use 2,4-D low volatile ester in drier areas where no susceptible crops are nearby. For Western bitterweed control use 2,4-D low volatile ester or amine at 1 qt./acre before plants flower and temperature (above 72°F) and soil moisture favors plant growth. When three-fourths of plants are blooming and/or temperature is less than 60°F, use Weedmaster, 2,4-D plus Banvel, Grazon P+D, GrazonNext or 2,4-D plus Tordon 22K For prairie gerardia control, when plants are 4 to 6 inches high, use 1½ qts./acre of 2,4-D or the low rates of Weedmaster, Banvel plus 2,4-D, Grazon P+D, or Tordon 22K + 2,4-D. When plants are 6 to 10 inches high, use the high rates, but spray before flowering.
		Weedmaster	VH 1 pt. to 1 qt. (½ to 1 lb.)	VH 1%			
		Tank mix Banvel with 2,4-D amine or low volatile ester	VH ¼ to ½ pt. (⅛ to ¼ lb.) Banvel + ¾ to 1 ½ pts. (⅜ to ¾ lb.) 2,4-D	VH ¼% Banvel + ¾% 2,4-D (4 lbs./gal. product)			
		Grazon P+D	VH 1 pt. to 1 ½ qts. (0.3 to 0.9 lb.)	VH 1%			
		Tank mix Tordon 22K with 2,4-D or low volatile ester	VH ¼ to ¾ qts. (⅙ to ¾ lb.) Tordon 22K + ½ to 1 ½ pts. (¼ to ¾ lb.) 2,4-D	VH ¼% Tordon 22K + ½% 2,4-D (4 lbs./gal. product)			
		Cimarron Max	VH Rate 1 to Rate 2				
		Surmount	VH 1.5 pt to 1 qt. (0.25 to 0.33 lb.)	VH 1%			
		PastureGard	H 1 to 1.5 qt. (0.5 to 0.75 lb.)	VH 1%			
		GrazonNext	VH 2 pts. (0.75 lb.)				
		Milestone	VH 4 ozs. (0.125 lb.)				

** Treatment control ratings: VH - Very High; H - High; M - Moderate; L - Low.

Page	Weed Controlled	Herbicide (common and chemical names – page 4)	Herbicide quantity (active ingredient rate in parenthesis)		Spray volume (per acre for broadcast, as described for individual plant)	Time to apply	Remarks
			Broadcast rate per acre	Individual plant/ spot treatment			
7	Broomweed (annual or common), plantain (tallow weed), wild carrot	Escort or Cimarron	VH** 0.1 oz.		2 to 4 gals. water for aerial spray; 10 to 25 gals. water for ground broadcast application. Add 1 to 2 qts. surfactant per 100 gals. of water.	Spring, weeds less than 4 inches tall.	
		Cimarron Plus	VH 0.125 oz.				
		Cimarron X-tra	VH 0.2 oz.				
8	Broom snakeweed (perennial broomweed)	Escort or Cimarron	VH 0.6 oz.		2 to 4 gals. water for aerial spray; 10 to 25 gals. water for ground broadcast application. Add 1 to 2 qts. of surfactant per 100 gals. of water.	Optimum time is in the fall, but may be applied in the spring.	
		Cimarron Plus	VH 0.75 oz.				
		Cimarron X-tra	VH 1.2 oz.				
		Cimarron Max	H to VH Rate 1 to Rate 2				
		Spike 20P	VH 3.75 lbs. of pellets (¾ lb.)	VH ¼ oz. of pellets (1/30 oz.) per 100 sq. ft.	Any time – optimum period is Oct. 1 to April 1, except in the Trans-Pecos where optimum period is May 1 to July 1.	Use only on sand, loamy sand, sandy loam, loam, silt loam, silt or sandy clay loam soils.	
NEW	Flathead Sedge	Weedmaster	VH 64 oz. (2 lbs.)		Ground broadcast 20 to 30 gals. per acre with the addition of 1 to 2 qts. of surfactant per 100 gals. of water.	Spring or fall.	Control may be enhanced if the stand is burned and/or shredded and allowed to regrow to a height of 12 to 15 inches before spraying.
		Cimarron Max	VH Rate 2				
10	Lespedeza	Remedy	VH 1 to 2 pts. (½ to 1 lb.)		Ground broadcast 20 to 20 gals. per acre with 1 to 2 qts. of surfactant per 100 gals. of water.	June through August under good growing conditions.	Plants need to be 12 to 18 inches tall before spraying. Use the higher rate if plants are large, approaching maturity, or if the infestation level is high. Begin application at flower bud initiation through full bloom.
		Escort or Cimarron	H 0.5 oz.				
		Cimarron Plus	H 0.7 oz.				
		Cimarron X-tra	H 1.0 oz.				
		Cimarron Max	H Rate 2				
		Surmount	VH 1.5 to 2 pt. (0.25 to 0.33 lb.)	VH 1%			
		PastureGard	VH 1.5 to 2 pt. (0.38 to 0.5 lb.)	VH 0.75%			

** Treatment control ratings: VH - Very High; H - High; M - Moderate; L - Low.

Page	Weed Controlled	Herbicide (common and chemical names – page 4)	Herbicide quantity (active ingredient rate in parenthesis)		Spray volume (per acre for broadcast, as described for individual plant)	Time to apply	Remarks
			Broadcast rate per acre	Individual plant/ spot treatment			
11	Rayless goldenrod (jimmyweed)	Escort or Cimarron	VH** 0.75 oz.		2 to 4 gals. water for aerial spray; 10 to 25 gals. water for ground broadcast application. Add 1 to 2 qts. surfactant per 100 gals. of water. Thoroughly wet foliage for individual plant treatment. Add 1 to 2 qts. surfactant per 100 gals. water.	Fall.	
		Cimarron Plus	VH 1.0 oz.				
		Cimarron X-tra	VH 1.5 oz.				
		Surmount	VH 6 pt. (1.0 lb.)	VH 2%			
		Tordon 22K	VH 1 qt. (½ lb.)	VH 1%			
11	Threadleaf Groundsel	Escort or Cimarron	VH 0.4 oz.		2 to 4 gals. water for aerial spray; 10 to 25 gals. water for ground broadcast application. Add 1 to 2 qts. surfactant per 100 gals. of water.		
		Cimarron Plus	VH 0.5 oz.				
		Cimarron X-tra	VH 0.8 oz.				
		Cimarron Max	VH Rate 2				
12	Upright Prairie- coneflower	Escort or Cimarron	VH 0.2 oz.		2 to 4 gals. water for aerial spray; 10 to 25 gals. water for ground broadcast application. Add 1 to 2 qts. surfactant per 100 gals. of water.	Spring, before flower stalk development.	
		Cimarron Plus	VH 0.25 oz.				
		Cimarron X-tra	VH 0.4				
13	Baccharis (dryland willow, Roosevelt willow, seep willow or willow baccharis)	2,4-D low volatile ester	H 3 pts. to 3 qts. (1.5 to 3 lbs.) 4 lbs./gal. product	H 1%	4 to 5 gals. of water for aerial spray. 15 to 20 gals. water for ground broadcast. For individual plant treatment thoroughly wet the entire plant— foliage, stems and trunks. Add 1 to 2 qts. surfactant per 100 gals. water. For individual plant treatment thoroughly wet the entire plant— foliage, stems and trunks. Add 1 to 2 qts. surfactant per 100 gals. water.	Spring, when leaves are fully expanded and dark green in color.	
		Grazon P+D		H 1%			
		Weedmaster		H 1%			
		Remedy		VH 1%			
		PastureGard		VH 1%			
		Surmount		VH 1%			
NEW	Giantreed	Habitat	VH 64 oz. (1 lb.)	VH ½%	Use a minimum 20 to 30 gals./acre total volume for ground broadcast applications and a minimum 15 gals./ acre for aerial. When using individual plant applications spray plants to runoff. Add 1% MSO to individual plant treatments and 1 qt./acre for broadcast applications.	Spray when plants are actively growing during the summer or fall with a minimum 3 feet of plant height.	Do not mow plants for 3 to 4 months after treatment.

** Treatment control ratings: VH - Very High; H - High; M - Moderate; L - Low.

Woody Plants

Page	Weed Controlled	Herbicide (common and chemical names – page 4)	Herbicide quantity (active ingredient rate in parenthesis)		Spray volume (per acre for broadcast, as described for individual plant)	Time to apply	Remarks
			Broadcast rate per acre	Individual plant/ spot treatment			
22	Mesquite (suppression and weed control)	2,4-D amine or low volatile ester	L** 2 to 4 qts. (2 to 4 lbs.)	M 2% (4 lbs./gal. product)	<p>For aerial applications suggested total spray volume is 4 gals./acre. Use oil-in-water emulsion (1 to 5 oil to water ratio considered optimum), or water plus surfactant, crop oil or methylated seed oil.</p> <p>For ground broadcast applications the suggested total spray volume is 10 to 25 gals./acre. Use oil-in-water emulsion (1 to 5 oil to water ratio considered optimum), or water plus surfactant, crop oil or methylated seed oil.</p> <p>Thoroughly wet foliage for individual plant treatments. Add 1 to 2 qts. surfactant per 100 gals. of water or an oil-in-water emulsion (5% diesel + 95% water).</p>	<p>Late spring to mid-summer with mature leaves (dark green color). Optimum period of application begins when soil temperature at a depth of 12 inches reaches 75°F and continues for 45 days.</p> <p>If treatment is applied prior to optimum soil temperatures, efficacy will be decreased and multiple applications over a period of years may be necessary to maintain less than 10% canopy.</p>	<p>Treatments will control many weeds. When using oil-in-water emulsion use emulsifier. Use of a treatment with a low control rating may result in multi-stem growth that may be more difficult to control in the future.</p>
		Grazon P+D	L 1 to 1.5 qts. (0.6 to 0.9 lbs.)				
		Weedmaster	L 1 to 1.5 qts. (1 to 1.5 lbs.)				
		Cimarron Max	L Rate 1 to Rate 2				
		Tordon 22K + 2,4-D amine or low volatile ester	L 0.5 to 0.75 pt. (1/8 to 3/16 lb.) Tordon 22K + 1 to 1.5 qts. (1 to 1.5 lbs.) 2,4-D, 4 lbs./gal. product				
		Banvel + 2,4-D amine or low volatile ester	L 0.5 to 0.75 pt. (1/4 to 3/8 lb.) Banvel + 3/4 to 1 1/8 qts. (3/4 to 1 1/8 lbs.) 2,4-D, 4 lbs./gal. product				
		Remedy	L 1 pt. to 1 qt. (0.5 lb. to 1 lb.)	M 1%			
		Banvel	L 1 pt. to 1 qt. (0.5 lb. to 1 lb.)	M 1%			
22	Mesquite, Christ thorn and other hardwoods (cut stumps)	Remedy		VH 15% in diesel fuel oil	<p>Spray the sides of the stump and the outer portion of the cut surface, including the cambium, immediately after cutting, to thoroughly wet the stem and root collar area, but not to the point of runoff.</p>	<p>Any season of the year, except when snow or water prevents spraying to the ground line.</p>	<p>This is commonly called the cut stump application method. Apply with a backpack or knapsack sprayer using low pressures and a solid cone or flat fan nozzle. This is an excellent treatment to use after cutting mesquite with hydraulic shears.</p>
		PastureGard		VH 20% in diesel fuel oil			
		Pathfinder II		VH Undiluted			

** Treatment control ratings: VH - Very High; H - High; M - Moderate; L - Low.

Page	Weed Controlled	Herbicide (common and chemical names – page 4)	Herbicide quantity (active ingredient rate in parenthesis)		Spray volume (per acre for broadcast, as described for individual plant)	Time to apply	Remarks	
			Broadcast rate per acre	Individual plant/spot treatment				
23	Mesquite	Remedy		VH** 2% in diesel fuel oil	Apply to the base of the trunk from 12 to 18 inches above the soil surface. Apply until solution puddles on soil surface.	Any time soil is dry and pulled away from the trunk.		
		Reclaim	M to H 1/3 qt. to 2/3 qt. (1/4 to 1/2 lb.)	VH 1%	For aerial applications suggested total spray volume is 4 gals./acre. Use oil-in-water emulsion (1 to 5 oil to water ratio considered optimum), or water plus surfactant, crop oil or methylated seed oil.	Late spring to mid-summer with mature leaves (dark green color). Optimum period of application begins when soil temperature at a depth of 12 inches reaches 75°F and continues for 45 days thereafter. When Reclaim is used alone or in a tank mix the period should continue for 60 days. For optimum root kill do not spray if white flowers or bean elongation are observable, if over 25% of the leaf canopy is damaged due to insects, disease or hail, if soil temperatures are less than 75°F, 1 ft. deep, or if new vegetative growth is present due to recent rains.	Use 1 pt./acre Tordon 22K plus 1/2 pt./acre Remedy, 1/2 pt./acre Banvel plus 1/2 pt./acre Reclaim, 1 pt./acre Tordon 22K plus 1/2 pt./acre Banvel, 1 pt./acre Tordon 22K plus 1/3 qt./acre Reclaim and 1/3 qt./acre Reclaim only in West Texas. Banvel and Banvel mixtures have been more effective in West Texas than in other parts of the state. Use mixtures that include 1/4 pt./acre Remedy and 1/3 pt./acre Reclaim only in Montague, Wise, Parker, Hood, Somervell, Bosque, Coryell, Lampasas, Burnet, Blanco, Kendall, Bandera, Real, Edwards and Val Verde counties and those counties north and west of the named counties. Mixtures that include 1/2 pt./acre Remedy and 2/3 pt./acre Reclaim will give better control than mixtures with 1/4 pt./acre Remedy and 1/3 pt./acre Reclaim. When using oil-in-water emulsion, use emulsifier added to oil for proper emulsion. Use of a treatment with a low-control rating may result in a multi-stem growth form that may be more difficult to control in the future.	
		Remedy + Tordon 22K	M 1/2 to 1 pt. (1/4 to 1/2 lb.) Remedy + 1 to 2 pts. (1/4 to 1/2 lb.) Tordon 22K	M to H 1/2% Remedy + 1/2% Tordon 22K	For ground broadcast applications the suggested total spray volume is 10 to 25 gals./acre. Use oil-in-water emulsion (1 to 5 oil to water ratio considered optimum), or water plus surfactant, crop oil or methylated seed oil.			
		Remedy + Banvel	L 1/2 to 1 pt. (1/4 to 1/2 lb.) Remedy + 1/2 to 1 pt. (1/4 to 1/2 lb.) Banvel	M 1/2% Remedy + 1/2% Banvel				
		Remedy + Reclaim	M to H 1/4 to 1 pt. (1/8 to 1/2 lb.) Remedy + 1/3 to 2/3 pt. (1/8 to 1/4 lb.) Reclaim	VH 1/2% Remedy + 1/2% Reclaim	Thoroughly wet foliage for individual plant treatments. Add 1 to 2 qts. surfactant per 100 gals. of water or an oil-in-water emulsion (5% diesel + 95% water).			
		Tordon 22K + Banvel	M 1 to 2 pts. (1/4 to 1/2 lb.) Tordon 22K + 1/2 to 1 pt. (1/4 to 1/2 lb.) Banvel	H 1/2% Remedy + 1/2% Banvel				
		Tordon 22K + Reclaim	M to H 1 to 2 pts. (1/4 to 1/2 lb.) Tordon 22K + 1/3 to 2/3 qt. (1/4 to 1/2 lb.) Reclaim	VH 1/2% Tordon 22K + 1/2% Reclaim				
26	Pricklypear, tasajillo	Tordon 22K	H 1 pt. to 1 qt. (1/4 lb. to 1/2 lb.)	VH 1%	For aerial applications suggested total spray volume is 2 to 4 gals./acre. Use oil-in-water emulsion (1 to 5 oil to water ratio considered optimum), or water plus surfactant, crop oil or methylated seed oil.	Any time; best results have been obtained with late summer through fall applications.	Use emulsifier when adding diesel fuel oil to water. Use 1 pt./acre Tordon 22K only on High Plains where no brush overstory is present. Late summer or fall applications, especially with Vista, will provide best results, but aerially spray in the winter or early spring if heavy overstory of woody plants is present or if damage to live oak is a concern.	
		Surmount	H 4 pt. (0.67 lb.)	VH 1%				
		Grazon P+D	H 1 gal. (2.5 lbs.)	VH 2%				
		Vista		VH 1%	For ground broadcast applications the suggested total spray volume is 10 to 25 gals./acre. Use oil-in-water emulsion (1 to 5 oil to water ratio considered optimum), or water plus surfactant, crop oil or methylated seed oil.			
		Gramoxone Extra	These two treatments are deleted from B-1466.					
		Tank mix Tordon 22K with Gramoxone Extra						

** Treatment control ratings: VH - Very High; H - High; M - Moderate; L - Low.

Page	Weed Controlled	Herbicide (common and chemical names – page 4)	Herbicide quantity (active ingredient rate in parenthesis)		Spray volume (per acre for broadcast, as described for individual plant)	Time to apply	Remarks	
			Broadcast rate per acre	Individual plant/spot treatment				
29	Saltcedar	Arsenal	VH** 2 qts. (1 lb.)	VH 1%	Minimum 10 gals./acre for aerial or ground broadcast sprays. Thoroughly wet foliage for individual plant treatment. Add 1 to 2 qts. surfactant per 100 gals. water.	July through September or until leaves begin to turn yellow.	When exposure to aquatic environments is possible use aquatic labels of Arsenal and Roundup (see table “Common, Chemical and Product Names of Herbicides”). Arsenal alone or in combination with Roundup will cause damage to desirable plants if contacted by the spray mix.	
		Arsenal + Roundup	VH 1 qt. (½ lb.) Arsenal + 1 pt. (½ lb.) Roundup	VH ½% Arsenal + ½% Roundup				
		Remedy		VH 25% in diesel fuel oil	Apply to lower 12 to 18 inches of trunk to wet the trunk; do not spray to point of runoff. Apply completely around the trunk.	Growing season when trees have mature leaves.	This is commonly called the low volume basal application method. Use a hollow cone nozzle with X1 orifice.	
		Tank mix Arsenal with Rodeo (glyphosate)	This recommendation is deleted, and moved to the remarks section.					
NEW	Saw Palmetto	Cimarron Max	H to VH Rate 2 or Rate 3 (see remarks)		For ground broadcast applications the suggested total spray volume is 15 to 30 gals./acre. Use water plus surfactant, crop oil or methylated seed oil.	Mid to late summer (August).	Rate 2 applications generally result in “high” control levels the first year after treatment, improving to “very high” by the second year after treatment. Rate 3 applications can be expected to produce “very high” control levels by the first year after treatment. Do not mow treated areas for at least 1 year.	
30	Yucca	Remedy		H 2% in diesel fuel oil	Spray the center of each individual whorl of leaves to the point of runoff.	Any time.	Complete coverage of leaves is not necessary. The crown of each plant must be thoroughly wetted with the herbicide mixture.	
				H 2% in 1:5 diesel fuel oil:water emulsion				May through September.
				VH 15% in diesel fuel oil		Use an adjustable cone nozzle (X1 orifice) and spray a 2-second burst into each whorl.	Spring and summer.	Direct spray into the center of each whorl.
				H Undiluted 2 to 4 ml./plant whorl		Use an exact delivery handgun set at 2 or 4 ml. per whorl dose.		
		Pathfinder II		VH Undiluted	Use an adjustable cone nozzle (X1 orifice) and spray a 2-second burst into each whorl.	Spring through fall prior to frost.	A second follow-up application of Cimarron Max at Rate 1 or Rate 2 plus 2,4-D low volatile ester at 0.5 to 0.75 lbs. a.i./acre within 2 years of the initial application may be required to control yucca seedlings and regrowth from rootstocks.	
		Cimarron Max + 2,4-D low volatile ester	H Rate 2 Cimarron Max + 1 pt. to 1.5 pts. 2,4-D (4 lbs. a.i./gal. product)		Crop oil concentrate, modified seed oil or modified seed oil/organosilicone are the preferred adjuvants. For aerial applications a minimum of 4 gals./acre total spray volume is recommended.			
		Cimarron Plus + 2,4-D low volatile ester	H 0.675 oz. Cimarron Plus + 1 qt. 2,4-D (4 lbs. a.i./gal. product)					
NEW	Burrobrush	Tordon 22K		VH 1%	Thoroughly wet foliage for individual plant treatment.	April through July.		
		Grazon P+D		VH 2%				

** Treatment control ratings: VH - Very High; H - High; M - Moderate; L - Low.

Funding for this publication was provided in part by Dow AgroSciences, Dupont Agricultural Products and BASF.

The information given herein is for educational purposes only. Reference to commercial products or trade names is made with the understanding that no discrimination is intended and no endorsement by the Texas AgriLife Extension Service is implied.

Produced by AgriLife Communications, The Texas A&M System
Extension publications can be found on the Web at: <http://agriflifebookstore.org>.
Visit Texas AgriLife Extension Service at <http://agriflifeextension.tamu.edu>.

Educational programs of the Texas AgriLife Extension Service are open to all people without regard to socioeconomic level, race, color, sex, disability, religion, age, or national origin.

Issued in furtherance of Cooperative Extension Work in Agriculture and Home Economics, Acts of Congress of May 8, 1914, as amended, and June 30, 1914, in cooperation with the United States Department of Agriculture. Edward G. Smith, Director, Texas AgriLife Extension Service, The Texas A&M System.
1500, Revision