

Free subgroups of the parametrized modular group

Subgrupos libres del grupo modular parametrizado

CHRISTIAN POMMERENKE¹, MARGARITA TORO^{0,2,✉}

¹Institut für Mathematik, Berlin, Germany

²Universidad Nacional de Colombia, Medellín, Colombia

ABSTRACT. We study free subgroups of index four of the parametrized modular group Π , the subgroup of $SL(2, \mathbb{Z}[\xi])$ generated by $\begin{pmatrix} 1 & \xi \\ 0 & 1 \end{pmatrix}$ and $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. There are eight free subgroups, four of which are normal and four are non-normal. Then we study the intersections of the normal subgroups. We give canonical presentations in terms of generators and relations. At the end of the paper we study connections between Π and the Bianchi groups, the two-parabolic group and a group from relativity theory.

Key words and phrases. Parametrized modular group, free subgroups, Bianchi groups, Picard group, discrete relativity theory.

2010 Mathematics Subject Classification. 11R65, 14C22.

RESUMEN. Estudiamos los subgrupos libres de índice cuatro del grupo modular parametrizado Π , que es el subgrupo de $SL(2, \mathbb{Z}[\xi])$ generado por $\begin{pmatrix} 1 & \xi \\ 0 & 1 \end{pmatrix}$ y $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. Hay ocho subgrupos libres, cuatro de los cuales son normales y los otros cuatro no lo son. Luego estudiamos las intersecciones de estos subgrupos. Damos presentaciones canónicas en término de generadores y relaciones. Al final del artículo estudiamos conexiones entre Π y los grupos de Bianchi, el grupo dos-parabólico y un grupo de la teoría de la relatividad.

Palabras y frases clave. Grupo modular parametrizado, subgrupos libres, grupos de Bianchi, grupo de Picard, teoría de la relatividad discreta.

⁰The second author was partially supported by the project “Matemáticas y computación”, Hermes code 20305, Universidad Nacional de Colombia, Sede Medellín.

1. Introduction

The parametrized modular group Π is defined in [10] as the subgroup of $SL(2, \mathbb{Z}[\xi])$ generated by

$$A = \begin{pmatrix} 1 & \xi \\ 0 & 1 \end{pmatrix}, B = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \tag{1}$$

where $\mathbb{Z}[\xi]$ is the polynomial ring over \mathbb{Z} with ξ as indeterminate. In the last section we describe some connections with the Picard group and other Bianchi groups using the results of R. G. Swan [14]. Furthermore we sketch the relation to discrete relativity theory and knot theory.

The previous paper [10] studied analytical properties of the singular set of Π and the enumeration of the elements of Π , see Lemma 2.1 below. The present paper investigates Π more in the spirit of combinatorial group theory [9] [7].

The exponent sums of a word $W \in \Pi$ with respect to the generators (1) are

$$\sigma(W) := (\text{sum of exponents of } A \text{ in } W), \tag{2}$$

which defines a homomorphism of Π into the additive group \mathbb{Z} , and

$$\tau(W) := (\text{sum modulo 4 of exponents of } B \text{ in } W), \tag{3}$$

which defines a homomorphism of Π into the additive group $\mathbb{Z}/4\mathbb{Z}$, note that $B^4 = I$.

In particular we shall study the subgroups


$$\Pi_k := \{W \in \Pi : \tau(W) \equiv (k - 1)\sigma(W) \pmod{4}\} \quad (k = 1, 2, 3, 4) \tag{4}$$

and their common subgroup

$$\Pi_0 := \{W \in \Pi : \sigma(W) \equiv \tau(W) \equiv 0 \pmod{4}\}. \tag{5}$$

We prove that each Π_k is a rank two free normal subgroup of index four in Π and that Π_0 is a rank five free normal subgroup of index 4 in Π_k ($k = 1, 2, 3, 4$).

Our main results are summarized in the following subgroup diagram where $[A, B]$ denotes the commutator.



See Theorem 2.2 for the first row, Theorem 3.3 for the second row and Theorem 3.1 for the third row. The other four intersections $\Pi_1 \cap \Pi_2$ and so on are equal to Π_0 by Proposition 3.2.

The presentations in this diagram are canonical in the sense of [9, p.140]: If X is a free group of rank n and Y is a subgroup of rank $m > n$ then there are generators x_1, \dots, x_n of X and generators y_1, \dots, y_m of Y such that

$$y_\nu = x_\nu^{d_\nu} z_\nu \quad (1 \leq \nu \leq n), \quad y_\nu = z_\nu \quad (n < \nu \leq m)$$

where z_ν is a word in Y and

$$\sigma_{x_\nu}(z_\mu) = 0 \quad \text{for } 1 \leq \nu \leq n, 1 \leq \mu \leq m,$$

where $\sigma_{x_\nu}(z_\mu)$ is the exponent sums of x_ν in the word z_μ .

We study other index four free subgroups of Π that are not normal subgroups.

2. The subgroups Π_i for $1 \leq i \leq 8$

The derivation of our presentations relies on the following result. See formulas (2.6) and (2.7) in [10], note that any negative sign in W is absorbed in $l \in \mathbb{Z}$ because $B^2 = -I$.

Lemma 2.1. *All words $W \in \Pi$ with $W \neq \pm I, \pm B$ have the form*

$$W = B^e A^{j_n} V \quad \text{with } V = BA^{j_{n-1}} \dots A^{j_1} B^l \tag{6}$$

where $e \in \{0, 1\}$, $l \in \{0, 1, 2, 3\}$, $j_\nu \in \mathbb{Z}$ and $j_n \neq 0$.

First we study the groups Π_k defined in (4). See Section 4.3 for the connection of Π_1 with the two-parabolic group.

Theorem 2.2. *Let $k = 1, 2, 3, 4$. The group Π_k is a free normal subgroup of index 4 in Π with the free presentation*

$$\Pi_k = \langle AB^{k-1}, [A, B] \rangle. \tag{7}$$

Remark 2.3. Let $\Gamma_k = \langle AB^{k-1}, [A, B] \rangle$. The generators in (7) can be replaced by other pairs of generators which we state in the following four lines.

- $\Gamma_1 = \langle A, BAB^{-1} \rangle$ because $[A, B] = A \cdot (BAB^{-1})^{-1}$,
- $\Gamma_2 = \langle AB, BA \rangle$ because $[A, B] = AB \cdot (BA)^{-1}$,
- $\Gamma_3 = \langle -A, -BAB^{-1} \rangle$ because $[A, B] = (-A) \cdot (-BAB^{-1})^{-1}$,
- $\Gamma_4 = \langle -AB, -BA \rangle$ because $[A, B] = (-AB) \cdot (-BA)^{-1}$.

Proof. (a) Since σ and τ are homomorphisms into additive groups it is clear from (4) that Π_k is a normal subgroup. Furthermore

$$\Pi_k B^j := \{W \in \Pi : \tau(W) \equiv (k - 1)\sigma(W) + j \pmod{4}\} \quad (j = 0, 1, 2, 3)$$

are distinct cosets of Π_k and their union is Π . Hence Π_k has index 4.

(b) Since $\sigma(B^j) = 0$ and $\tau(B^j) = j \not\equiv 0$ for $j = 1, 2, 3$, it follows from (4) that $B^j \notin \Pi_k$ for all k . If $W \in \Pi_k, W \neq \pm I, \pm B$, then W has the form (6). It follows from [10, Lemma 2.2] that all these words are different. Hence Π_k is a free group. Let $\Gamma_k = \langle AB^{k-1}, [A, B] \rangle$.

(c) Now we show that, for $W \in \Pi$,

$$W \in \Gamma_k \implies \tau(W) \equiv (k - 1)\sigma(W) \pmod{4}. \tag{8}$$

We have $\sigma(AB^{k-1}) = 1, \tau(AB^{k-1}) = k - 1$ and $\sigma([A, B]) = \tau([A, B]) = 0$, which proves (8).

(d) Now we prove the converse, namely that, for $W \in \Pi$,

$$\tau(W) \equiv (k - 1)\sigma(W) \pmod{4} \implies W \in \Gamma_1. \tag{9}$$

We shall however use the alternative forms of the generators listed in Remark 2.3 above. We proceed by induction on the number n of occurrences of the symbol A in the representation (6). If $n = 0$ then $\sigma(W) = 0$ and (9) is trivial. Suppose that (9) holds when the number of occurrences of A is $< n$. Now let there be n occurrences of the symbol A . Below we shall show that there exists $U \in \Gamma_k$ such that $W' = U^{-1}W$ has less than n occurrences of the symbol A . By (4) applied to U we have $\tau(U) \equiv (k - 1)\sigma(U) \pmod{4}$. Using the left-hand side of (9) we conclude that $\tau(W') \equiv (k - 1)\sigma(W') \pmod{4}$. By the induction hypothesis we have $W' \in \Gamma_k$. It follows that $W = UW' \in \Gamma_k$. Now we turn to the construction of U for the different four cases. Let $W = B^e A^{j_n} V$ with $V = BA^{j_{n-1}} \dots A^{j_1} B^l$ as described in (6). (d1) Let $k = 1$. We define

$$U := B^e A^{j_n} B^{-e} = (B^e A B^{-e})^{j_n} \in \Pi_1,$$

and we have $W' = U^{-1}W = B^e V$ which is shorter than W . (d2) Let $k = 2$. We define

$$U := B^e A^{j_n} B^{j_n - e}$$

and use that $B^2 = -I$. If $j_n = 2q$

$$U = B^e (AA)^q (B^2)^q B^{-e} = B^e (A(-I)A)^q B^{-e} = B^e (AB \cdot BA)^q B^{-e} \in \Pi_2.$$

If $j_n = 2q + 1$ then

$$U := B^e (AA)^q A (B^2)^q B B^{-e} = B^e (AB \cdot BA)^q (AB) B^{-e} \in \Pi_2.$$

Therefore, in both cases

$$W' = B^{e-j_n} A^{-j_n} B^{-e} B^e A^{j_n} V = B^{e-j_n} V$$

so W' is shorter than W . (d3) Finally let $k = 3$ or $k = 4$. We see from the remark after Theorem 2.2 that the generators to be obtained are the same as for the cases $k = 1$ and $k = 2$ except for different signs. This difference only changes the exponent l of B in (6) so that we can argue as above. \checkmark

Proposition 2.4. *Let $j = 1, 2, 3, 4$. The group*

$$\Pi_{j+4} := \langle AB^{j-1}, -[A, B] \rangle \tag{10}$$

is a free subgroup of index 4 in Π and satisfies

$$B\Pi_{j+4}B^{-1} = \Pi_{j'+4} \tag{11}$$

with $j' = j + 2 \pmod 4$ if $j \neq 2$ and $j' = 4$ if $j = 2$.

Proof. We write $C := [A, B]$. Since Π_j is a free group there is a unique homomorphism $\varphi_j : \Pi_j \rightarrow \Pi$ such that

$$\varphi_j(AB^{j-1}) = AB^{j-1}, \varphi_j(C) = -C, \tag{12}$$

see [9, p.48]. Hence we have $\Pi_{j+4} = \varphi_j(\Pi_j)$ by (10). Since the generators of Π_j and Π_{j+4} only differ by the sign of C we have $\varphi_j(W) = \pm W$ for $W \in \Pi_j$. Now suppose that $\varphi_j(W) = I$. Then $W = \pm I$ where $-I$ is not possible because Π_j is a free group. It follows that $W = I$. Hence φ_j is an isomorphism so that Π_{j+4} is also a free group. As in part (a) of the proof of Theorem 7 we can prove that Π_{j+4} has index 4 in Π . Since $BCB^{-1} = C^{-1}$ it follows from (10) that

$$B\Pi_{j+4}B^{-1} = \langle BAB^{j-2}, -C^{-1} \rangle = \langle BAB^{j-2}, -C \rangle,$$

in the last step we replaced the generator $-C^{-1}$ by its inverse. By another Tietze transformation we can replace BAB^{j-2} by

$$-C \cdot BAB^{j-2} = ABA^{-1}B^{-1} \cdot BAB^{j-2} \cdot B^2 = AB^{j+1} = AB^{j-1+2}$$

and (11) follows from (7). \checkmark

Remark 2.5. We mention, without proof, that the eight groups Π_i , for $1 \leq i \leq 8$, are the only index four free subgroups of Π .

3. The intersections of these subgroups

Now we turn to the subgroups of Π_k ($k = 1, 2, 3, 4$). First we study the group defined by (5).

Theorem 3.1. *The group Π_0 is a free normal subgroup of index 16 in Π and has the free presentation*

$$\Pi_0 = \langle A^4, [A, B], [A^{-1}, B], [A^2, B], [A^{-2}, B] \rangle. \quad (13)$$

Proof. (a) It follows from (4) and (5) that Π_0 is a normal subgroup. The 16 sets $\{W : \sigma(W) = j, \tau(W) = k\}$ ($j, k = 0, 1, 2, 3$) form a complete coset system of Π_0 in Π . Hence Π_0 has index 16 in Π . Since Π_1 has index 4 in Π it follows that Π_0 has index 4 in Π_1 . The free group Π_1 has rank 2 by (8). Hence it follows [9, Th.2.10] that Π_0 is free of rank $4(2-1) + 1 = 5$. Therefore the 5 generators in (13) are free generators.

(b) Let Γ be the group with the presentation in (13). Each of the words W in (13) satisfies $\sigma(W) \equiv \tau(W) \equiv 0 \pmod{4}$. Hence it follows from (5) that $\Gamma \subset \Pi_0$. All $W \in \Pi$ with $W \neq \pm I, \pm B$ have the form (6). We shall prove $\Pi_0 \subset \Gamma$ again by induction on the number n of occurrences of the symbol A . In view of (13) we have $A^4 \in \Gamma$ and

$$BA^4B^{-1} = BA^2 \cdot A^2B^{-1} = [A^2, B]^{-1}A^4[A^{-2}, B] \in \Gamma.$$

It follows that

$$A^{4q} \in \Gamma, \quad BA^{4q}B^{-1} \in \Gamma \quad (q \in \mathbb{Z}). \quad (14)$$

Suppose that $W \in \Pi_0 \implies W \in \Gamma$ is true if W has $< n$ occurrences of A . Let $W = B^e A^{j_n} V$, with $V = BA^{j_{n-1}} B \dots A^{j_1} B^l$ as described in (6). We write

$$j_n = 4q + r, \quad q \in \mathbb{Z}, \quad r = -1, 0, 1, 2.$$

If $e = 0$, then

$$W = A^{4q+r} V = A^{4q} \cdot A^r B A^{-r} B^{-1} \cdot B A^r A^{j_{n-1}} B \dots A^{j_1} B^l = A^{4q} [A^r, B] V'$$

where $V' = A^{r+j_{n-1}} B \dots A^{j_1} B^l$ has the form (6) with $n-1$ occurrences of A . Since the factors of V' belong to Γ by (14), it follows that $W \in \Gamma$. If $e = 1$, then

$$W = BA^{4q+r} V = BA^{4q} B^{-1} \cdot B A^r B^{-1} A^{-r} \cdot A^r B V = BA^{4q} B^{-1} \cdot [A^r, B]^{-1} V'$$

where $V' = A^r B V = A^r B^2 A^{j_{n-1}} B \dots A^{j_1} B^l = A^{r+j_{n-1}} B \dots A^{j_1} B^{l+2}$ has the form (6) with $n-1$ occurrences of A . By (14) the factors before V' are in Γ . Hence $W \in \Gamma$. \square

Proposition 3.2. *The group Π_0 satisfies*

$$\Pi_0 = \Pi_1 \cap \Pi_2 = \Pi_2 \cap \Pi_3 = \Pi_3 \cap \Pi_4 = \Pi_4 \cap \Pi_1 \tag{15}$$

and is a normal subgroup of index 4 in each Π_k ($k = 1, 2, 3, 4$).

Proof. We abbreviate

$$\{\sigma \equiv m, \tau \equiv n\} := \{W \in \Pi : \sigma(W) \equiv m, \tau(W) \equiv n \pmod{4}\}. \tag{16}$$

It follows from (8) that

$$\begin{aligned} \Pi_1 \cap \Pi_2 &= \{\tau \equiv 0\} \cap \{\tau \equiv \sigma\} = \{0 \equiv \tau \equiv \sigma\} = \Pi_0, \\ \Pi_2 \cap \Pi_3 &= \{\tau \equiv \sigma\} \cap \{\tau \equiv 2\sigma\} = \{\sigma \equiv 0 \equiv \tau\} = \Pi_0, \\ \Pi_3 \cap \Pi_4 &= \{\tau \equiv 2\sigma\} \cap \{\tau \equiv 3\sigma\} = \{\sigma \equiv 0, \tau \equiv 0\} = \Pi_0, \\ \Pi_4 \cap \Pi_1 &= \{\tau \equiv 3\sigma\} \cap \{\tau \equiv 0\} = \{\tau \equiv 0, \sigma \equiv 0\} = \Pi_0. \end{aligned}$$

We see from (15) that Π_0 is a subgroup of all Π_k which is normal because all definitions are in terms of σ and τ . ☑

We see from Proposition 3.2 that four of the six possible intersections of the groups Π_k are equal to Π_0 . The remaining two intersections however lead to new groups.

Theorem 3.3. *We have the free presentations*

$$\Pi_1 \cap \Pi_3 = \langle A^2, [A, B], [A^2, B] \rangle, \quad \Pi_2 \cap \Pi_4 = \langle A^2 B^2, [A, B], [A^2, B] \rangle \tag{17}$$

and Π_0 has index 2 in these two groups.

Proof. (a) First we show that \supset holds in (17). It follows from (4) that, with the notation (16),

$$\Pi_1 \cap \Pi_3 = \{\sigma \equiv 0, 2, \tau \equiv 0\}, \quad \Pi_2 \cap \Pi_4 = \{\sigma \equiv 0, 2, \tau \equiv \sigma\}. \tag{18}$$

The generators that occur in (17) satisfy these conditions. (b) Now we prove that \subset holds in (17). To simplify the proof we write

$$\begin{aligned} \Gamma &:= \langle A^2, [A, B], [A^2, B] \rangle, \quad s = +1 && \text{in the case } \Pi_1 \cap \Pi_3 \\ \Gamma &:= \langle A^2 B^2, [A, B], [A^2, B] \rangle, \quad s = -1 && \text{in the case } \Pi_2 \cap \Pi_4. \end{aligned} \tag{19}$$

Then the assertion (17) becomes

$$\Gamma = \{W \in \Pi : \sigma(W) \equiv 0, 2, \tau(W) \equiv \frac{1}{2}(1-s)\sigma(W) \pmod{4}\}. \tag{20}$$

First we derive an identity. Since

$$sA^{-2}[A^2, B] = sA^{-2}A^2BA^{-2}B^{-1} = B(sA^{-2})B^{-1}$$

we obtain from (19) and (20) that

$$B(sA^{-2})^q B^{-1} \in \Gamma (q \in \mathbb{Z}). \quad (21)$$

We shall use the notation of Lemma 6 and proceed by induction on n . Let $n = 1$. In the case $e = 0$ it follows from (20) with $q \in \mathbb{Z}$ that $W = A^{j_1} B^{l_1} = A^{2q} B^{(1-s)q} \in \Gamma$. In the case $e = 1$ we obtain from (19) and (21) that $W = BA^{j_1} B^{l_1} = B(sA^2)^q B^{-1} \in \Gamma$. Now we assume that our assertion holds for $n - 1$. There are four cases where always $q \in \mathbb{Z}$.

If $W = A^{2q} BA^{j_{n-1}} B \dots$ then we write

$$W = (sA^2)^q V, V = s^q BA^{j_{n-1}} \dots .$$

If $W = A^{2q+1} BA^{j_{n-1}} B \dots$ then we write

$$W = (sA^2)^q [A, B] V, V = s^q BA^{j_{n-1}+1} \dots .$$

If $W = BA^{2q} BA^{j_{n-1}} B \dots$ then we write

$$W = (sA^2)^q B^{-1} V, V = s^q BA^{j_{n-1}} \dots .$$

If $W = BA^{2q+1} BA^{j_{n-1}} B \dots$ then we write

$$W = (sA^2)^q B^{-1} [A, B] V, V = s^q BA^{j_{n-1}+1} \dots .$$

We check that V satisfies (20) in all four cases so that $V \in \Gamma$. In the first two cases, the factor before V lies in Γ because of (19). In the last two cases we also use (21) to obtain the same conclusion. Since $V \in \Gamma$ by the induction hypothesis we see that $W \in \Gamma$ holds in all cases. (c) Since the groups $\Pi_1 \cap \Pi_3$ and $\Pi_2 \cap \Pi_4$ lie properly between the groups Π_k on the one hand and their subgroup Π_0 of index 4 on the other hand, it follows that Π_0 has index 2 in our groups. \checkmark

Proposition 3.4. *The commutator subgroup Π' has infinite index in Π .*

Proof. Let $\Gamma := \{W \in \Pi : \sigma(W) = \tau(W) = 0\}$, here we do not consider congruences. Then $\Pi' \subset \Gamma$ and all cosets ΓA^{4k} are disjoint. We conclude that $|\Pi : \Pi'| \geq |\Gamma : \Pi'| = \infty$. \checkmark

The situation is often quite different in other contexts. For instance for the Picard group (see below), the first three commutator subgroups have finite index [3, Th.1].

4. Connection with other groups

The groups $\Pi(\zeta)$ are obtained by replacing the indeterminate ξ in Π by the complex number ζ . For instance $\Pi(1)$ and $\Pi(2)$ lead to classical modular groups. Many groups are obtained by combining groups $\Pi(\zeta)$ with different values of ζ as we will see below.

4.1. *The Bianchi groups.* Let $d \in \mathbb{N}$ be square-free and let O_d be the ring of integers of the quadratic field $\mathbb{Q}(\sqrt{-d})$. We now consider the Bianchi groups $SL(2, O_d)$ but only for the d that allow an euclidean algorithm, namely $d = 1, 2, 3, 7, 11$, see [14] [2, Chapter 4]. The case $d = 1$ is the Picard group, here considered in $SL(2, \mathbb{C})$, see for instance [2, Chapt. 5] [3] [5]. It will be convenient to enlarge our groups $\Pi(\zeta)$ by adding the generator $A(1)$.

Proposition 4.1. *Let $\Pi^+(\zeta)$ be the group generated by $A(1), A(\zeta)$ and B .*

- (1) *If $d = 1$ then $\Pi^+(1 + i) = SL(2, O_1)$.*
- (2) *If $d = 3$ then $\Pi^+(\omega) = SL(2, O_3)$ where $\omega = \frac{1}{2}(-1 + i\sqrt{3})$.*
- (3) *If $d = 2, 7, 11$ then $SL(2, O_d) = \Pi^+(\omega)$ where $\omega = i\sqrt{d}$ for $d = 2$ and where $\omega = \frac{1}{2}(1 + i\sqrt{d})$ for $d = 7, 11$.*

Proof. The cases $d = 2, 7, 11$ are due to Swan but the cases $d = 1, 3$ are new, Swan had an additional generator, see [14, p.64-71] or [2, Chapt.4]. In a later paper it will be proved the cases $d = 1, 3$. □

4.2. *Discrete space-time.* The Schild group Σ is the subgroup of $SL(2, \mathbb{C})$ that leaves the discrete space-time grid $\{(t, x, y, z) : t, x, y, z \in \mathbb{Z}\}$ invariant under the Lorentz transformation

$$X = \begin{pmatrix} t + z & x + iy \\ x - iy & t - z \end{pmatrix} \mapsto SXS^*, \quad S \in SL(2, \mathbb{C}). \tag{22}$$

See [13][6][4]. The Schild group Σ is generated by

$$A(1 + i), \begin{pmatrix} (1 - i)/2 & (1 - i)/2 \\ -(1 + i)/2 & (1 + i)/2 \end{pmatrix}, \begin{pmatrix} 0 & -(1 - i)/\sqrt{2} \\ (1 + i)/\sqrt{2} & 0 \end{pmatrix}.$$

The last two matrices generate a subgroup of order 24.

Proposition 4.2. *The group*

$$\Sigma_1 := \langle A(1 + i), A(1 - i), B, Q \rangle, \quad Q := \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$$

is a subgroup of index 6 in the Schild group Σ and is also a subgroup of index 3 in the Picard group $SL(2, O_1)$. Furthermore $\langle A(1 + i), A(1 - i), B \rangle$ is a subgroup of index 2 in Σ_1 .

Proof. The first two assertions were proved in [4, Sect.4]. The group Σ_1 is similar to the Picard group discussed in Section 4.1, only the generators $A(1), A(i)$ are replaced by $A(1+i), A(1-i)$. This will be proved in a later paper. \square

For the Lorentz transformation (22), the generators B and Q of Σ_1 have a simple physical interpretation, namely

$$B : x \mapsto -x, y \mapsto y, z \mapsto -z, \quad Q : x \mapsto -x, y \mapsto -y, z \mapsto z$$

whereas the time t remains unchanged. But $t \mapsto 2t + x \mp y - z$ holds for $A(1 \pm i)$. We remark that the groups $A(1)$ and $A(i)$ are not subgroups of Σ_1 .

4.3. *The two-parabolic group.* We consider the polynomial ring $\mathbb{Z}[x]$. The two-parabolic group is the subgroup of $\mathrm{SL}(2, \mathbb{Z}[x])$ generated by $X := \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix}$ and $Y := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$; see e.g. [12][11]. Up to conjugation in $\mathrm{SL}(2, \mathbb{Z}[x])$, this is the only subgroup generated by two parabolic matrices with distinct fixed points, see [11].

The subgroup Π_1 of Π was studied in Section 2. It is generated by A and $D := BA^{-1}B^{-1}$. If $V := \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Pi_1$ then b and c are odd polynomials [10, (2.15)]. Hence

$$\varphi(V) := \begin{pmatrix} a & b/\xi \\ c\xi & d \end{pmatrix} \in \mathrm{SL}(2, \mathbb{Z}[x]), \quad x = \xi^2 \quad (23)$$

is well-defined and it can be checked that φ is a homomorphism. It follows from (23) that $\varphi(D) = X, \varphi(A) = Y$. Hence $\varphi(\Pi_1)$ is the two-parabolic group.

The two-parabolic group is of interest in knot theory, see e.g. [1] [8, Chapt.4]. A knot K is a Jordan curve in \mathbb{R}^3 . In many cases a discrete group $\Gamma \subset \mathrm{SL}(2, \mathbb{C})$ is associated with the knot complement $\mathbb{R}^3 \setminus K$. For a 2-bridge knot a ‘‘Wirtinger word’’ $W \in \varphi(\Pi_1)$ partially describes the knot K . To get Γ one chooses x such that the relation $XW = WY$ is satisfied.

For example, for the knot $4_1 = (5, 3)$ we have $W = YX^{-1}Y^{-1}X$ and $x = \frac{1}{2}(1 + i\sqrt{3})$ [11, Example 8]. Going back to Π_1 we have the relation $DAD^{-1}A^{-1}D = AD^{-1}A^{-1}DA$ and obtain $\Gamma = \varphi(\Pi_1(-\frac{1}{2} + \frac{1}{2}i\sqrt{3}))$.

References

- [1] G. H. Burde and H. Zieschang, *Knots*, Walter de Gruyter, 1985.
- [2] B. Fine, *Algebraic theory of Bianchi groups*, Marcel Dekker, New York, 1989.
- [3] B. Fine and M. Newman, *The normal subgroup structure of the picard group*, Trans. Amer. Math. Soc. **302** (1987), 769–786.

- [4] G. Jensen and Ch. Pommerenke, *Discrete space-time and Lorentz transformations*, to appear 2015 in the Canadian Mathematical Bulletin.
- [5] N. Yilmaz and N. Cangül, *Conjugacy classes of elliptic elements in the Picard group*, Turk. J. Math. **24** (2000), 209–220.
- [6] M. Lorente and P. Kramer, *Representations of the discrete inhomogeneous Lorentz group and Dirac wave equation on the lattice*, J. Phys A. **32** (1999), 2481–2497.
- [7] R. C. Lyndon and P. E. Schupp, *Combinatorial group theory*, Springer, New York, 1977.
- [8] C. MacLachlan and A. W. Reid, *The arithmetic of hyperbolic 3-manifolds*, Springer, New York, 2003.
- [9] W. Magnus, A. Karrass, and D. Solitar, *Combinatorial group theory*, 2. ed, Dover Publ., New York, 1976.
- [10] D. Mejia, Ch. Pommerenke, and M. Toro, *On the parametrized modular group*, to appear in J. Analyse Math.
- [11] Ch. Pommerenke and M. Toro, *On the two-parabolic subgroups of SL_2, \mathbb{C}* , Rev. Col. Mat. **45** (2011), 37–50.
- [12] R. Riley, *Parabolic representations of knot groups I*, Proc. London Math. Soc. **3** (1972), 217–242.
- [13] A. Schild, *Discrete space-time and integral Lorentz transformations*, Canad. J. Math. **1** (1949), 29–47.
- [14] R. G. Swan, *Generators and relations for certain special linear groups*, Advances Math. **6** (1971), 1–77.

(Recibido en mayo de 2015. Aceptado en octubre de 2015)

TECHNISCHE UNIVERSITÄT BERLIN
INSTITUT FÜR MATHEMATIK
D-10623 BERLIN, GERMANY
e-mail: pommeren@math.tu-berlin.de

ESCUELA DE MATEMÁTICAS
UNIVERSIDAD NACIONAL DE COLOMBIA - SEDE MEDELLÍN
CALLE 59A #63-20, MEDELLÍN, COLOMBIA
e-mail: mmtoro@unal.edu.co