

# **Validity and reliability of the ‘Isometric Exercise Scale’ (IES) for measuring ratings of perceived exertion during continuous isometric exercise**

John W. D. Lea<sup>1</sup>, Jamie M. O’Driscoll<sup>1</sup>, Damian A. Coleman<sup>1</sup> & Jonathan D. Wiles<sup>1,\*</sup>

*1-School of Human and Life Sciences, Canterbury Christ Church University, Kent, UK*

*\*Corresponding Author – Email Address: Jim.Wiles@canterbury.ac.uk*

## **Abstract**

**Purpose** Isometric exercise (IE) interventions have been shown to be effective at reducing resting arterial blood pressure (BP). IE intensity is generally determined using force, electromyography (EMG), heart rate (HR), or knee joint angle. However, ratings of perceived exertion (RPE) may provide a more accessible means of determining exercise intensity. Therefore, the aim of this study was to assess the validity of a specific Isometric Exercise Scale (IES) during a continuous incremental isometric wall squat test.

**Methods** Twenty-nine male participants completed 4 incremental IE tests. Each test consisted of 5 isometric wall squat intensities, determined by knee joint angles from 135° to 95°, each held for up to 2 minutes. The tests were continuous until volitional fatigue or completion of the 5 work stages. Throughout the exercise protocol, RPE (IES and Borg’s CR-10), HR and blood pressure were recorded.

**Results** The IES produced a strong positive linear relationship with the CR-10 ( $r = 0.967$ ). Additionally, strong positive relationships between the IES and wall squat duration ( $r = 0.849$ ), HR ( $r = 0.819$ ) and BP ( $r = 0.841$ ) were seen. Intra-class correlation coefficients and coefficients of variations for the IES ranged from  $r = 0.81$  to  $0.91$  and  $4.5\%$  to  $54\%$  respectively, with greater reliability seen at the higher workloads.

**Conclusions** The IES is a valid and reliable measure of RPE, exercise intensity, and it accurately represents the changes in physiological measures of exertion during IE; as such, the IES can be used as a useful additional measure of exercise intensity during IET interventions.

**Key Words:** RPE, Perceived Effort, Exercise Intensity, CR-10 scale, Resistance Exercise

## **Abbreviations**

ANOVA	Analysis of Variance
BP	Blood Pressure
DBP	Diastolic Blood Pressure
EMG	Electromyography
HR	Heart Rate
IE	Isometric Exercise
IES	Isometric Exercise Scale
IET	Isometric Exercise Training
MAP	Mean Arterial Pressure
RPE	Rating of Perceived Exertion
SBP	Systolic Blood Pressure

## **Introduction**

Hypertension, characterised by a sustained elevation in arterial blood pressure ( $\geq 140$  mmHg systolic and/or  $\geq 90$  mmHg diastolic), is the leading attributable risk factor for increased CVD mortality [1]. Exercise has been recommended as a non-pharmacological lifestyle modification for the treatment of hypertension [2]. Isometric exercise training (IET) interventions have been shown to be an effective and time efficient methodology to reduce resting [3,4,5,6] and ambulatory blood pressure [6].

The control of exercise intensity is a key factor in ensuring the safety and efficacy of physical activity in any context, including athletic, recreational, and therapeutic settings [7,8]. Previous methods of administering IET and monitoring its intensity have tended to require expensive equipment such as isokinetic dynamometers [3,9,10], hand grip dynamometers [11] and electromyography (EMG) [5]. It has been suggested that the need for expensive equipment and time-consuming testing protocols, may present unnecessary barriers that could ultimately limit the effectiveness of these interventions [12]. Consequently, more accessible modes of IET that could be implemented in the home have been explored. One such intervention, is the use of the isometric wall squat, where intensity is controlled by manipulation of the knee joint angle [13]. A 4-week home based isometric wall squat intervention, with exercise intensity derived from a maximal isometric wall squat test [6,14], was shown to produce significant reductions in resting [6,15] and ambulatory arterial BP [6]. While these methods of training are more accessible than previous iterations, the ability to accurately monitor exercise intensity without laboratory testing and the use of additional equipment could help to further promote uptake of this type of IE intervention.

Ratings of perceived exertion (RPE) could provide an easy to use and accessible alternative means of assessing and monitoring exercise intensity [16,17]. Indeed, it has long been established that RPE provides an accurate estimation of exercise intensity and physiological exertion during cardiovascular exercise [18]. In addition, there is now a growing body of evidence that indicates that various RPE scales provide a valid measure of exercise intensity during resistance exercise, including the Borg 6-20 [19], Omni-res [20], and the Borg CR-10 [21] scales. The validity of these scales has been shown to be independent of participant age [22,23] or sex [21,24,25]. Additionally, the Borg CR-10 scale has largely been adopted within IE research up to this point [14], despite its intended application being for rating pain with no numerical ceiling effect [26]. In an exercise setting, where the average participant's understanding of RPE is likely to be limited, the open-ended nature of the CR-10 scale may make monitoring and prescribing IE intensity more difficult. There are currently no RPE scales that have been specifically designed and validated for IET; it has been suggested that it is important to design and validate scales for specific populations, exercise types and modalities [16], and that caution should be taken when using RPE scales with modalities and materials other than those they have been validated for [27]. It has also been proposed that for an RPE scale to be considered a valid measure for use in the clinical and/or health-fitness setting, it must demonstrate both concurrent and construct validity, evidenced by strong positive correlations with physiological variables (e.g. HR) and a previously validated criterion scale respectively [28].

Therefore, the aims of this research were to: (1) assess the construct validity of a new isometric exercise scale (IES) as a measure of RPE during isometric wall squat exercise, using the frequently adopted CR-10 scale as a criterion measure; (2) examine the validity of the IES to measure changes in isometric wall squat intensity during a continuous maximal incremental test; (3) explore the concurrent validity of the IES using criterion measures of physiological exertion (HR and BP); and (4) examine the reliability of the IES responses over time.

## **Methods**

### **Participants**

Twenty-nine healthy, normotensive males (age:  $23.2 \pm 4.0$  years; stature:  $180.9 \pm 7.8$  cm; body mass:  $82.7 \pm 17.3$  kg) volunteered to participate in this study. Before testing, written informed consent was obtained from all participants. Additionally, participants completed a health and medical questionnaire and self-reported that they were not suffering from any injury or disease. All participants were physically active, non-smokers and not taking any medication during the investigation. Participants agreed to maintain regular dietary and physical activity habits throughout the testing period.

### **Study Design**

All participants were required to attend the laboratory on 4 separate occasions, separated by a minimum of 48 hours. Each session followed the same procedures; starting with resting measurements before completion of a continuous maximal incremental wall squat tests. Participants were asked to abstain from food 4 hours, caffeine 12 hours, alcohol and strenuous exercise 24 hours pre-testing. All participants verbally confirmed adherence to the testing requirements prior to the start of each testing session. This study was approved by the University Ethics Committee and conducted according to the 1964 Declaration of Helsinki.

### **Procedures**

*Familiarisation.* Prior to the first testing session participants received an information pack outlining the testing protocols and measurement procedures included in the study. At the start of the first laboratory session participants had the study design, resting and exercise measurements, and exercise protocols explained to them verbally. As part of this explanation, participants were shown the equipment that would be used and were given a demonstration of the wall squat, including the correct wall squat position. Finally, participants were shown the RPE scales and received the standardised instructions and anchoring. Following this, if the participant wished to be part of the study, written informed consent was collected and resting measurements were taken.

*Resting measures.* Upon arrival to the laboratory, participants rested in a seated position for 10 minutes. After 10 minutes rest, HR, systolic BP (SBP), diastolic BP (DBP) and mean arterial pressure (MAP) were recorded using an oscillometric BP monitor on the participants left arm (Dinamap® Pro, GEMedical Systems, Slough, UK). Three measurements were taken, each separated by 1-minute [29]. Following the seated measurements, participants rested in a supine position for 15 minutes. After an initial 10-minute period, HR and BP were measured continuously for 5 minutes using a plethysmographic device (Task Force® Monitor, CNSystems, Graz, Austria). Resting HR and BP values were calculated as the mean of the 5-minute supine measurement period.

*Maximal isometric wall squat test protocol.* Following the resting measures, participants were required to complete a maximal incremental isometric wall squat test, as described in [13]. The test required participants to lower their back down a fixed wall and make small adjustments to their feet position until the required knee joint angle was reached, while maintaining a vertical lower limb and an erect trunk. Knee joint angle was measured using a clinical goniometer, secured to the participants lower and upper leg using elasticated Velcro strapping. The fulcrum was aligned with the lateral epicondyle of the femur, the moving arm was placed on the lateral midline of the femur using the greater trochanter for reference and the stationary arm on the lateral midline of the fibula using the lateral malleolus and fibular head for reference. A spirit level was attached to the stationary arm to ensure that the lower leg was kept vertical during exercise. The internal angle between the femur and fibula was measured [13]. The test consisted of five consecutive 2-min stages, beginning at a knee joint angle of  $135^\circ$  and guided to reduce the angle by  $10^\circ$  every 2 min ( $125^\circ$ ,  $115^\circ$ ,  $105^\circ$ , and  $95^\circ$ ) (Figure 1). Each participant's foot position was measured from the back of the left heel to the wall and their back position was measured as the distance from the ground to the lower back, which was defined as the lowest point of contact that the participants back had with the wall. Participants were not permitted to stand or rest between angles. The test continued until volitional exhaustion, the participant was unable to maintain the required knee angle, or completion of the 10-minute test [8]. Participants were monitored for signs of physical

distress throughout the test and were instructed to terminate the exercise at the first signs of dizziness or feeling unwell. Verbal encouragement was given throughout, with instructions to maintain normal breathing to avoid the Valsalva manoeuvre. Heart rate and BP were recorded continuously during the test; mean HR and BP (SBP, DBP and MAP) were calculated for the last 5 seconds of each minute of the test.

**INSERT FIGURE 1 HERE**

**Figure 1:** Knee joint angles used for the five consecutive 2-min stages of the incremental isometric exercise test (left to right: 135°, 125°, 115°, 105°, and 95°).

*Ratings of perceived exertion.* Participants were asked to rate the perceived exertion in their active muscles using the IES and Borg's CR-10 scales, 50 seconds in to each minute of the test. Participants were cued to give their ratings using the standardised questions: "How hard do you feel your leg muscles are working". The participants were randomly assigned into one of two groups: Group 1 were asked to rate their perceived exertion using the IES first followed by the CR-10; group 2 rated using the CR-10 first and then the IES. The scales were in full view of the participant for the entire test, presented one above the other, either IES on top or CR-10 on top (Figure 2) depending on group allocation. The scales were presented in this way, rather than side-to-side, to stop participants picking a rating on the first scale and then moving sideways to the corresponding value on the second scale, without consideration of differences in the two values.

**INSERT FIGURE 2 HERE**

**Figure 2:** RPE Scales Presentation Formats

*Scale instructions and anchoring.* Standardised scaling instructions and anchoring procedures were explained for both scales before testing. The standardised instructions and anchoring procedures for the CR-10 scale were used, as previously published in Borg's work [26]. The instructions for the IES were as follows: "This scale is used to rate how hard you think your active muscles are working. This scale has 3 different columns: Rating, Description and TTF. The 'Rating' numbers are from 0-10 and are used to rate the exertion or effort in the active muscle group(s). The 'Description' words and 'TTF' are used to help you choose a rating. 0 (Rest) is absolutely no effort, as felt during complete rest. 5 (Moderate) is right in the middle of 0 and 10. It's not especially hard and it is no problem to continue; but it no longer feels comfortable. 10 (Maximal) is maximum effort; your muscles are working as hard as they can, and you can only maintain this for seconds before you will have to stop.

*TTF (Time to Failure) indicates the amount of time remaining, during an isometric contraction, before you will be unable to continue. In other words, this describes how much you have left in your 'fuel tank'. 100% - your muscles are fresh; you haven't started the contraction yet (fuel tank is full). 50% - means you can continue to hold the contraction for the same amount of time that you have already completed (fuel tank is half full). 0% - your muscles are failing/have failed (fuel tank is empty). When you give your rating: focus only on the muscle group(s) that is working. You can use the 'Description' words, the Time to Failure (TTF), and/or you can simply rate the exertion out of ten".*

### **Data analysis**

All data were analysed using the statistical package for social sciences (SPSS 22 release version for Windows; SPSS Inc., Chicago IL). Before analysis, all data were checked for conformity with the parametric assumptions [30]. Construct validity between the IES and CR-10 ratings was determined using linear regression analysis and Spearman's Rank-Order Correlation. Concurrent validity of the IES and CR-10 results with the criterion measures of exercise intensity was assessed using Spearman's Rank-Order Correlations. To test for differences in concurrent validity, between the IES and CR-10, validity coefficients underwent Fishers Z score transformation followed by statistical analysis for differences in the Z-Scores. Reliability of the IES, CR-10, HR and BP results across the 4 testing sessions were examined separately using: Two-factor (Session x Intensity) repeated measures ANOVA's or Friedman's test (normal distribution dependant); Intraclass Correlation Coefficients (ICC); Standard Error of the Mean (SEM) and Coefficient of Variations (COV). For the difference

tests, the 'Session' factor had 4 levels (testing sessions 1-4), and the 'Intensity' factor had 5 levels (knee angle - 135°, 125°, 115°, 105° & 95°). Where main effects were found, post-hoc testing was conducted with Bonferroni adjustment for multiple comparisons. The ICC (3,1) [31] model was used to assess the agreement between the repeated measures taken during consecutive sessions. Within-participant variance was calculated as the SEM from the ICC analyses, and as COVs with 95% confidence interval, derived from log-transformed two-way ANOVA for each variable [30, 32]. ICC and COV results, for the IES and CR-10 scales, were considered to be significantly different if the mean results for each scale lay outside of the 95% confidence interval of the other. An alpha level of  $< 0.05$  was set as the threshold for statistical significance. All data are expressed as mean  $\pm$  S.D. unless otherwise indicated.

## **Results**

### **Resting Measures**

At the start of each of the 8 testing sessions, resting measures were recorded for each participant. The mean resting values for HR, SBP, DBP and MAP were:  $61 \pm 8 \text{ b} \cdot \text{min}^{-1}$ ,  $109 \pm 7 \text{ mmHg}$ ,  $63 \pm 5 \text{ mmHg}$ , and  $80 \pm 6 \text{ mmHg}$  respectively. There were no significant differences in any resting measures between trials ( $P > 0.05$ ). The Intra-class correlation coefficients for the resting measures ranged from  $r = 0.52$  to  $0.91$ . Coefficients of Variation (with 95% confidence intervals) were 4.0% (3.5-4.9%) for resting HR, 2.9% (2.5-3.5%) for SBP, 6.2% (5.3-7.6%) for DBP, and 4.2% (3.6-5.1%) for MAP.

### **Construct Validity**

The construct validity of the IES was established by correlation and linear regression analysis of the ratings from the IES and CR-10 scale. Spearman's Rank-Order Correlation showed a strong positive linear relationship ( $r = 0.97$ ,  $P < 0.001$ ) between the CR-10 and IES ratings of exertion. Likewise, linear regression analysis to assess the ability of the CR-10 results to predict the IES results, showed a significant linear regression equation ( $F(1, 977) = 13958$ ,  $p < 0.000$ ) with an  $r$  value of  $0.97$  (Figure 3).

**INSERT FIGURE 3 HERE**

**Figure 3:** The relationship between the IES and CR-10 ratings of perceived exertion

### **RPE and Exercise Intensity**

The validity of the IES and CR-10 to represent isometric exercise intensity was assessed by correlating the RPE ratings against percentage of maximum exercise intensity (workload x wall squat duration). Strong positive correlations were shown for the IES ( $r = 0.89$ ,  $P < 0.001$ ) and CR-10 ( $r = 0.88$ ,  $P < 0.001$ ) with exercise intensity (Figure 4).

**INSERT FIGURE 4 HERE**

**Figure 4:** The relationships of the IES and CR-10 ratings with isometric exercise intensity (as a percentage of maximum)

### **RPE and Measures of Physiological Exertion**

Significant positive relationships were found when correlating the physiological measures of exercise intensity: HR ( $r = 0.82$  &  $r = 0.81$ ,  $P < 0.001$ ), SBP ( $r = 0.84$  &  $r = 0.84$ ,  $P < 0.001$ ), DBP ( $r = 0.81$  &  $r = 0.80$ ,  $P < 0.001$ ), and MAP ( $r = 0.84$  &  $r = 0.83$ ,  $P < 0.001$ ) with RPE ratings from the IES and CR-10 respectively (Figure 5).

**INSERT FIGURE 5 HERE**

**Figure 5:** The relationships of the IES and CR-10 ratings with the physiological measures of exercise intensity

## Reliability of Exercise Measures

There were no significant differences in IES or CR-10 ratings between sessions at any knee angle ( $P > 0.05$ ). The ICCs and COVs between sessions 1 and 2, fell outside of the confidence intervals for the between sessions 2-4 comparisons; therefore, the reliability data for sessions 1 and 2 are presented separately to sessions 2-4. ICCs for sessions 1 and 2, across all knee joint angles, ranged from  $r = -0.49$  to  $0.76$  ( $SEM = 0.67-1.07$ ) and  $r = 0.30$  to  $0.76$  ( $SEM = 0.71-1.39$ ) for the IES and CR-10 respectively. Additionally, the COVs between sessions 1 and 2 ranged from 42.1% to 10.5% for the IES and 77.2% to 12.1% for the CR-10. The ICCs for sessions 2 to 4 ranged from  $r = 0.81$  to  $0.91$  and  $r = 0.79$  to  $0.90$  for the IES and CR-10 respectively (Table 1). The ICC for the 135° knee angle was significantly higher for the CR-10 scale when compared to the IES; no other differences in ICCs were found. The SEMs calculated for Sessions 2-4 ranged from 0.37 – 0.87 for the IES and 0.35 – 0.84 for the CR-10 Scale. As such, The COVs ranged from 54% to 4.5% and 41.9% to 7% for the IES and CR-10 respectively, with the lowest variances seen at the higher intensity levels (Table 1).

**Table 1:** IES and CR-10 results at each knee angle between sessions and the corresponding reliability statistics.

	Knee Angle	Session Number				ICC	SEM	COV
		1	2	3	4			
IES	135°	1.9 ± 1.5	1.4 ± 1.3	1.3 ± 1.1	1.7 ± 1.5	0.81 (0.70-0.89)	0.67	54.0% (44.9-70.0%)
	125°	3.9 ± 1.8	3.5 ± 1.8	3.3 ± 1.9	3.7 ± 2.2	0.86 (0.78-0.92)	0.87	33.1% (27.8-42.1%)
	115°	6.6 ± 1.7	6.2 ± 2.0	6.1 ± 2.0	6.1 ± 2.3	0.91 (0.85-0.95)	0.72	14.2% (12.0-17.6%)*
	105°	8.6 ± 1.2	8.5 ± 1.2	8.2 ± 1.4	8.1 ± 1.5	0.83 (0.72-0.91)	0.56	8.1% (6.8-10.4%)*
	95°	9.4 ± 0.5	9.3 ± 0.8	9.3 ± 0.8	9.1 ± 1.0	0.84 (0.66-0.93)	0.37	4.5% (3.5-6.7%)*
CR-10	135	1.2 ± 1.0	1.0 ± 1.0	0.9 ± 0.9	1.0 ± 0.9	0.90 (0.83-0.94)☒	0.35	41.9% (35.0-53.7%)*
	125	3.0 ± 1.7	3.0 ± 1.7	2.7 ± 1.8	3.0 ± 2.1	0.87 (0.78-0.92)	0.76	38.9% (32.6-49.7%)
	115	5.7 ± 2.3	5.4 ± 2.1	5.3 ± 2.3	5.4 ± 2.5	0.90 (0.83-0.94)	0.84	18.7% (15.9-23.5%)
	105	7.6 ± 1.8	8.0 ± 1.5	7.5 ± 1.8	7.5 ± 1.7	0.79 (0.64-0.88)	0.79	13.7% (11.4-17.7%)
	95	8.6 ± 1.1	9.0 ± 1.2	9.0 ± 1.5	8.9 ± 1.2	0.86 (0.69-0.94)	0.48	7.0% (5.4-10.3%)

ICC = Intraclass correlation coefficients (95% confidence intervals), COV = Coefficient of variation (95% confidence intervals). ICC and COV values for comparisons of sessions 2, 3 and 4 only.

☒ Significantly greater  $r$ -value than the other RPE scale at the same Knee angle

\* Significantly lower variance than the other RPE scale at the same Knee angle

## **Discussion**

This study demonstrated that the IES is a valid and reproducible measure of RPE, exercise intensity and the physiological responses to isometric wall squat exercise. The construct validity of the IES was assessed using the Borg CR-10 scale, which has previously been shown to be a valid and reliable measure of RPE and exercise intensity during resistance exercise [33] and is the scale most commonly adopted for this type of exercise intervention [14]. The IES ratings showed excellent agreement with the ratings from the Borg CR-10 ( $r = 0.97$ ), during correlation and linear regression analysis. In addition, there were no significant differences in the relationships shown with exercise intensity and physiological measures of exertion (HR and BP) between the IES and CR-10 scales. The Borg CR-10 scale has previously been used to validate the OMNI Elliptical Ergometer Scale, during aerobic exercise, yielding similar construct validity coefficients to the present study ( $r = 0.96-0.98$ ) [28]. During dynamic resistance exercise, the CR-10 scale was used to validate a novel Estimated Repetitions to Failure Scale, giving strong validity coefficients ranging from  $r = 0.86$  to  $r = 0.96$ , depending on the specific exercise used [34]. Likewise, the now widely used OMNI-RES scale was validated during resistance exercise using the Borg 6-20 scale [35]; this analysis showed validity coefficients from  $r = 0.94$  to  $0.97$ . The construct validity results of the current study, when compared to previous research, suggest that the IES is a valid measure of RPE during isometric exercise.

Ratings from the IES were shown to be an accurate representation of exercise intensity during maximal isometric wall squat exercise. This was shown through correlation of the IES results with wall squat duration, which in this case represents an increase in both contraction time and workload throughout the test. The results of this analysis showed a strong positive relationship between the two variables ( $r = 0.89$ ). Previous research has shown the OMNI-RES scale to be a valid measure of exercise intensity, with correlations ranging from  $r = 0.89$  to  $0.91$  in males and females [27]. Similarly, the CR-10 was shown to be a valid measure of exercise intensity with validity coefficients of  $r = 0.77$  at baseline and  $r = 0.91$  following a 12-week training intervention [36]. Likewise, in production mode, the Borg 6-20 scale has also been shown to be valid when used by sedentary, active and strength trained individuals alike ( $r = 0.83 - 0.92$ ) [37]. As such, the validity coefficient shown for the IES is comparable to those shown in previous resistance exercise research, suggesting it is an accurate measure of exercise intensity.

The IES also showed strong positive relationships with HR ( $r = 0.82$ ) and BP ( $r = 0.81 - 0.84$ ), indicating that the IES can accurately represent the changes in physiological exertion caused during the incremental isometric test, to the same extent as the CR-10 scale. The CR-10 scale has previously been shown to produce strong positive correlations with HR ( $r = 0.76$ ) and blood lactate ( $r = 0.730$ ) during dynamic weight training [38], and with HR ( $r = 0.71$ ) during bodyweight suspension training [39]. Likewise, the CR-10 has shown comparable relationships during aerobic training [18]. The strong relationships seen between the IES and physiological measures could allow it to be used as an important additional safety measure during IET, to ensure that participants are work at intensities that keep them within safe HR and BP limits [8,40].

Reliability of the IES measures across the 4 testing sessions was examined by correlating the results from consecutive sessions using intraclass correlation coefficients. The ICC measures of reliability between sessions 1-2 were significantly different, when compared to between sessions 2-3 and 3-4. These results show a learning effect following the first session and suggests that physical familiarisation with this maximal incremental exercise and RPE scale may be required before a stable relationship is achieved. The ICC results for the IES showed excellent agreement ( $r = 0.81$  to  $0.91$ ) across sessions 2 – 4, indicating that the IES is a reliable measure of RPE and exercise intensity over time. These reliability coefficients were closely matched by the ICCs for HR and BP, suggesting that the relationship between the IES results and physiological exertion is stable over time. These results are comparable to those shown previously for the OMNI-RES Thera-band ( $r = 0.72 - 0.76$ ) [41] and Borg CR-10 ( $r = 0.88$ ) [42] scales during different forms of resistance exercise. Additionally, in production mode, where the participant selects/modifies the exercise intensity to elicit a specific RPE response, reliability coefficients ranged from  $r = 0.69 - 0.95$  for OMNI-RES RPE ratings of 3, 6 and 9 [43].

The within participant variance was assessed using the SEM and COV. The CR-10 scale showed lower variance at the 135-workload ( $0.35$  vs  $0.67$ ) when compared to the IES, corresponding to the significantly higher ICC



result seen with the CR-10 at that level. Whereas, the IES showed lower SEMs at the 4 higher workloads, when compared to the CR-10. This translated to statistically lower COVs at the 3 highest workloads with the IES. The COV results for the IES showed higher percentage variance at the lighter intensities and lower variances at the higher intensities; this is to be expected as the lower average RPE values seen in the earlier stages of the test (IES = 1.6 and CR-10 = 1.0 in the first stage) mean that even the smallest possible change between session (0.5) would elicit 30-50% variance. Arguably, this high variance makes comparison of COVs from RPE with COVs from other measurement methods, e.g. HR, inappropriate; however, this can still be a useful measure of the variance/reliability when comparing two like measures, such as two RPE scales. The CR-10 scale produced significantly lower variance at the lowest intensity (135 degrees), possibly due to the increased number of lower value numbers and therefore smaller differences between values at the lower end of that scale (0, 0.3, 0.5 and 1). Whereas, The IES showed significantly lower variance at the highest three intensities (115°, 105° and 95°), possibly due to the simpler closed-ended nature of the IES as opposed to the open-ended CR-10; these intensities are more representative of the intensities used during IET for reducing resting blood pressure [14] and may therefore favour the IES during this type of intervention. Previous analysis of the reliability of the CR-10 showed a COV of 17%, for exercise eliciting an average RPE of approximately 6 for females and 7 males [44]. This RPE is approximately equivalent to the mean IES results seen at the 115° knee angle (6.3) which showed between session variance of 14.2%. Similarly, Day et al. [41] assessed the reliability of the CR-10 across 2-sessions at 3 difference intensities, giving a COV of 14.5%. The three intensities used in Day's study gave mean RPE scores of 3.7, 5.6 and 6.9, approximating the mean IES ratings achieved across the 125° and 115° knee angles in the current study (3.6 – 6.3); The IES COVs for these knee angles ranged from 13.2-33.1%. These results suggest that the IES is reliable across sessions following habituation, especially at the higher workloads associated with BP training interventions.

Further research is required to explore the validity of IES during discontinuous isometric wall squat exercise at submaximal intensities, as is currently used during isometric wall squat training for the reduction of resting blood pressure [15]. Additionally, research is needed to explore the potential use of RPE as a means of prescribing and monitoring IET intensity, especially in pre-hypertensive and hypertensive populations.

## **Conclusion**

The IES provides a valid and reliable measure of RPE and exercise intensity during maximal isometric wall squat exercise. Additionally, the IES results produced strong positive relationships with the criterion measures of physiological exertion (HR and BP). As such, the IES can be used as a valid measure of RPE and could be useful in the selection and monitoring of workloads during IET interventions for the reduction of resting blood pressure.

## References

1. Mancia, G. et al. ESH/ESC guidelines for the management of arterial hypertension: The Task Force for the Management of Arterial Hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *European Heart Journal*. **34**, 2159-2219 (2013).
2. Brook, R.D., Jackson, E.A., Giorgini, P. & McGowan, C.L. When and how to recommend “alternative approaches” in the management of high blood pressure. *Am J Med*. **128**, 567–570 (2015).
3. Wiles, J.D., Coleman, D.A. & Swaine, I.L. The effects of performing isometric training at two exercise intensities in healthy young males. *European journal of applied physiology*. **108**(3), 419-428 (2010).
4. Cornelissen, V.A. & Smart, N.A. Exercise Training for Blood Pressure: A Systematic Review and Meta-analysis. *Journal of the American Heart Association*. **2**, e004473 (2013).
5. Devereux, G. R., Wiles, J. D. & Howden, R. Immediate post-isometric exercise cardiovascular responses are associated with training-induced resting systolic blood pressure reductions. *European journal of applied physiology*. **115**(2), 327-333 (2015).
6. Taylor, K. A., Wiles, J. D., Coleman, D. A., Leeson, P., Sharma, R. & O’Driscoll, J. M. Neurohumoral and ambulatory haemodynamic adaptations following isometric exercise training in unmedicated hypertensive patients. *Journal of hypertension*. **37**(4), 827-836 (2018).
7. Robertson, R.J., Goss, F.L., Dubé, J., Rutkowski, J., Dupain, M. Brennan, C. & Andreacci, J. Validation of the adult OMNI scale of perceived exertion for cycle ergometer exercise. *Medicine and Science in Sports and Exercise*. **36**, 102-108 (2004).
8. Wiles, J.D., Taylor, K., Coleman, D., Sharma, R. & O’driscoll, J.M. The safety of isometric exercise: Rethinking the exercise prescription paradigm for those with stage 1 hypertension. *Medicine*. **97**,10 (2018).
9. Devereux, G. R., Wiles, J. D., & Swaine, I. L. Reductions in resting blood pressure after 4 weeks of isometric exercise training. *European journal of applied physiology*. **109**(4), 601-606 (2010).
10. Baross, AW, Wiles, JD & Swaine, IL. Double-leg isometric exercise training in older men. *Open Access Journal of Sports Medicine*. **4**, 33-40 (2013).
11. McGowan, C.L., Levy, A.S., McCartney, N. & MacDonald, M.J. Isometric handgrip training does not improve flow-mediated dilation in subjects with normal blood pressure. *Clinical Science*. **112**, 403-409 (2007).
12. Millar, P.J., Paashuis, A. & McCartney, N. Isometric handgrip effects on hypertension. *Current Hypertension Reviews*. **5**, 54-59 (2009).
13. Goldring, N, Wiles, JD & Coleman, D. The effects of isometric wall squat exercise on heart rate and blood pressure in a normotensive population. *Journal of Sports Science*. **32**, 129-36 (2014).
14. Wiles, J.D., Goldring, N., O’Driscoll, J.M., Taylor, K.A. & Coleman, D.A. An alternative approach to isometric exercise training prescription for cardiovascular health. *Translational Journal of the American College of Sports Medicine*. **3**(2), 10-18 (2018).
15. Wiles, J.D., Goldring, N. & Coleman, D. Home-based isometric exercise training induced reductions resting blood pressure. *European Journal of Applied Physiology*. **117**, 8393 (2017).
16. Colado, J.C., Garcia-Masso, X., Triplett, N.T., Calatayud, J., Flandez, J., Behm, D. & Rogers, M.E. Construct and concurrent validation of a new resistance intensity scale for exercise with Thera-Band® elastic bands. *Journal of sports science & medicine*. **13**(4), 758 (2014).
17. Morrin, N.M., Stone, M.R., Swaine, I.L. & Henderson, K.J. The use of the CR-10 scale to allow self-regulation of isometric exercise intensity in pre-hypertensive and hypertensive participants. *European journal of applied physiology*. **118**(2), 339-347 (2018).
18. Chen, M.J., Fan, X. & Moe, S.T. Criterion-related validity of the Borg ratings of perceived exertion scale in healthy individuals: a meta-analysis. *Journal of sports sciences*. **20**(11), 873-899 (2002).
19. Lagally, K.M., McCaw, S.T., Young, G.T., Medema, H.C. & Thomas, D.Q. Ratings of perceived exertion and muscle activity during the bench press exercise in recreational and novice lifters. *The Journal of Strength & Conditioning Research*. **18**(2), 359-364 (2004).
20. Aniceto, R.R., Ritti-Dias, R.M., dos Prazeres, T.M., Farah, B.Q., de Lima, F.F. & do Prado, W.L. Rating of perceived exertion during circuit weight training: a concurrent validation study. *The Journal of Strength & Conditioning Research*. **29**(12), 3336-3342 (2015).

21. Buckley, J.P. & Borg, G.A. Borg's scales in strength training; from theory to practice in young and older adults. *Applied physiology, nutrition, and metabolism*. **36**(5), 682-692 (2011).
22. Robertson, R.J., Goss, F.L., Andreacci, J.L., Dub , J.J., Rutkowski, J.J., Snee, B.M., Kowallis, R.A., Crawford, K., Aaron, D.J. & Metz, K.F. Validation of the children's OMNI RPE scale for stepping exercise. *Medicine & Science in Sports & Exercise*. **37**(2), 290-298 (2005).
23. Pincivero, D.M., Polen, R.R. & Byrd, B.N. Gender and contraction mode on perceived exertion. *International journal of sports medicine*. **31**(05), 359-363 (2010).
24. Eston, R. & Evans, H.J.L. The validity of submaximal ratings of perceived exertion to predict one repetition maximum. *Journal of sports science & medicine*. **8**(4), 567 (2009).
25. Gearhart, R.F., Riechman, S.E., Lagally, K.M., Andrews, R.D. & Robertson, R.J. Safety of using the adult OMNI Resistance Exercise Scale to determine 1-RM in older men and women. *Perceptual and motor skills*. **113**(2), 671-676 (2011).
26. Borg, G., 1998. Perceived Exertion and Pain Scales. Human Kinetics, Champaign, Illinois, USA.
27. Robertson, R.J. et al. Concurrent validation of the OMNI perceived exertion scale for resistance exercise. *Medicine and science in sports and exercise*. **35**(2), 333-341 (2003).
28. Mays, R.J., Goss, F.L., Schafer, M.A., Kim, K.H., Nagle-Stilley, E.F. & Robertson, R.J. Validation of adult OMNI perceived exertion scales for elliptical ergometry. *Perceptual and motor skills*. **111**(3), 848-862 (2010).
29. Whelton, P.K., Carey, R.M., Aronow, W.S., Casey, D.E., Collins, K.J., Dennison Himmelfarb, C., DePalma, S.M., Gidding, S., Jamerson, K.A., Jones, D.W., MacLaughlin, E.J., Muntner, P., Ovbigele, B., Smith, S.C., Spencer, C.C., Stafford, R.S., Taler, S.J., Thomas, R.J., Williams, K.A., Williamson, J.D. & Wright, J.T. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. *A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines*. (2017).
30. Atkinson, G. & Nevill, A. M. Selected issues in the design and analysis of sport performance research. *Journal of sports sciences*. **19**(10), 811-827 (2001).
31. Shrout PE, Fleiss JL (1979). Intraclass correlations: uses in assessing rater reliability. *Psychological Bulletin* **86**, 420-428
32. Field, A. *Discovering statistic using SPSS for Windows* (3rd ed.). London: (Sage 2009).
33. Tate, R. F. & Klett, G. W. Optimal confidence intervals for the variance of a normal distribution. *Journal of the American statistical Association*. **54**(287), 674-682 (1959).
34. Brown, D.M. & Bray, S.R. Isometric exercise and cognitive function: an investigation of acute dose-response effects during submaximal fatiguing contractions. *Journal of sports sciences*. **33**(5), 487-497 (2015).
35. Hackett, D.A., Johnson, N.A., Halaki, M. & Chow, C.M. A novel scale to assess resistance-exercise effort. *Journal of sports sciences*. **30**(13), 1405-1413 (2012).
36. Lagally, K.M. & Robertson, R.J. Construct validity of the OMNI resistance exercise scale. *The Journal of Strength & Conditioning Research*. **20**(2), 252-256 (2006).
37. Desgorces, F.D., Thomasson, R., Aboueb, S., Toussaint, J.F. & Noirez, P. Prediction of one-repetition maximum from submaximal ratings of perceived exertion in older adults pre-and post-training. *Aging clinical and experimental research*. **27**(5), 603-609 (2015).
38. Tiggemann, C.L., Korzenowski, A.L., Brentano, M.A., Tartaruga, M.P., Alberton, C.L. & Kruel, L.F. Perceived exertion in different strength exercise loads in sedentary, active, and trained adults. *The Journal of Strength & Conditioning Research*. **24**(8), 2032-2041 (2010).
39. Hollander, D.B., Durand, R.J., Trynicki, J.L., Larock, D. Castracane, V.D., Hebert, E.P. & Kraemer, R.R. RPE, pain, and physiological adjustment to concentric and eccentric contractions. *Medicine and science in sports and exercise*. **35**(6), 1017-1025 (2003).
40. Giancotti, C.f., Foster, C., Pezzotta, C., Lecce, D., Rodio, A., Capranica, L. & Cortis, C. Evaluation of training load during Suspension Training: Is session-RPE a valid method? *Conference: 20th Annual Congress of the European College of Sport Science (ECSS)*, Malm  (Sweden), (2015).
41. Williams, M.A. et al. American Heart Association Council on Clinical Cardiology; American Heart Association Council on Nutrition, Physical Activity, and Metabolism. Resistance exercise in

individuals with and without cardiovascular disease: 2007 update: a scientific statement from the American Heart Association Council on Clinical Cardiology and Council on Nutrition, Physical Activity, and Metabolism. *Circulation*. **31**, 572-584 (2007).

42. Colado, J. C., Garcia-Masso, X., Triplett, T. N., Flandez, J., Borreani, S. & Tella, V. Concurrent validation of the OMNI-resistance exercise scale of perceived exertion with Thera-band resistance bands. *The Journal of Strength & Conditioning Research*. **26**(11), 3018-3024 (2012).
43. Day, M. L., McGuigan, M. R., Brice, G., & Foster, C. Monitoring exercise intensity during resistance training using the session RPE scale. *The Journal of Strength & Conditioning Research*. **18**(2), 353-358 (2004).
44. Lagally KM, Amorose AJ, Rock B. Selection of Resistance Exercise Intensity Using Ratings of Perceived Exertion from the OMNI—RES. *Percept Mot Skills*. **108**(2), 573-586 (2009).
45. Egan, A.D. Session rating of perceived exertion during high intensity and low intensity bouts of resistance exercise. *UW-L J Undergrad Res*. **6**, 1-6 (2003).