

## INFLUENCE OF PLANTS DENSITY ON PRODUCTIVITY OF CORN FOR SILAGE IN THE CONDITIONS OF DEPRESSION JIJIA - BAHLUI

Marius SCARLAT<sup>1</sup>, Adrian-Ilie NAZARE<sup>1</sup>, Costel SAMUIL<sup>1</sup>, Mihai STAVARACHE<sup>1</sup>,  
Vasile VÎNTU<sup>1</sup>

email: [scarlatmmarius@gmail.com](mailto:scarlatmmarius@gmail.com)

### Abstract

Corn silage is the most efficient way of feeding the animals during the grazing period, with a very high degree of consumability. The success of corn silage cultivation depends largely on the proper choice of hybrid, but also on the pedoclimatic conditions of the crop area. The characteristics and the suitability of the hybrid have an important role in the maize culture for silage, but in order to obtain superior productions from a qualitative and quantitative point of view, the technological elements and their application must be respected. The density of corn crop for silage is considered the most important technological element. All the improvements of the maize crop, the introduction of the most productive maize hybrids, the fertilization, the irrigation, etc., have led to new levels of production, changing each time the optimum plants density. The research was conducted in 2018 in the pedoclimatic conditions corresponding to the Depression Jijia – Bahlui. Three experimental factors were studied, as followed: the influence of plants density, the distance between the rows and the hybrid on the production of dry matter (DM). The obtained results showed that the technological factors (the density of plants and the distance between rows) caused changes in the biomass production, the quantity of dry matter being influenced by the hybrid, as well as the plants density and the distance between rows. As the density of plants per hectare increases, the amount of dry matter decreases thus registering at all three densities, reduced production differences.

**Key words:** maize hybrids, corn silage, dry matter production, row spacing

Corn is one of the most important cultivated plants that has a special role in the food of humans and animals. The success of a corn crop for silage involves some requirements that refer to establishing the optimum plants of density, choosing the hybrids, as well as the correct choice of the technological elements (Roth G.W. and Heinrichs A.J., 2017).

Corn silage provides high palatability with relatively constant quality and higher energy content than other forages (Oliveira I.L. *et al*, 2017). Chaudhary D.P. *et al*, 2013, argue that corn silage has a rapid growth, producing a very high quantity of biomass and of high quality. It contains sufficient amounts of protein and minerals and has a high digestibility compared to other forages.

The density of plants as a factor of production is important and influences the production of dry matter in the silage maize crop (Vîntu V. *et al*, 2010). The susceptibility to corn silage depends on several factors, among which: the choice of cultivated hybrid, climatic conditions, plants densities, relations between cultural, quantitative and generative factors (Ceclan O.A.,

2010). Numerous studies showed that maize crop density is considered the most important technological element in maize crop for silage (Çarpici Budakli E. *et al*, 2010; Ramezani M. *et al*, 2011; Ferreira G. *et al*, 2014; Mandić V. *et al*, 2015; Sharifi R.S. and Namvar A., 2016).

Cañadas Á. *et al*, 2016, show that increasing the density of plants in corn cultivation for silage resulted in thinner and longer stems. According to the results, the highest production of dry matter was recorded at the plants density, of 125.000 plants·ha<sup>-1</sup> with a production increase of 20.11% higher than the density of 65.500 plants·ha<sup>-1</sup>. Ferreira G. and Teets C.L., 2017, show that increasing the density of plants increases the production of fodder by affecting the nutritional quality minimally. All the improvements of the corn crop, had led to new levels of production, changing each time the optimum plants density.

The study followed the production of dry matter on the surface unit and its modification depending on the density of plants, hybrid and the distance between rows.

<sup>1</sup> “Ion Ionescu de la Brad” University of Agricultural Sciences and Veterinary Medicine, Iasi

## MATERIAL AND METHOD

The studies were conducted in 2018 in Moldavian Forest Steppe. From the administrative point of view, the area under study is located in the Iasi county, on the coordinates (47°02' North latitude and 27°22' East longitude).

From the climatic point of view, the territory belongs to the temperate-continental climate zone under the influence of Atlantic and Euro-Asian anticyclones. The year of experience was characterized by normal weather conditions, no extreme weather phenomena were recorded during the whole vegetation period of the experience. The soil corresponds to the forest steppe zone types and is a cambic chernozem, formed on clay deposits. The research has followed the influence of plants density, distance between rows and three maize hybrids on the production of dry matter in the maize for silage. In order to attain the proposed objectives, a polyfactorial experience was set up. The experiment was established, with a design type 2x3x3, arranged in subdivided plots in three replications, which have a 22,4 m<sup>2</sup>. The factors were as follows: A - distance between rows, with two graduations (a<sub>1</sub> - 70 cm between rows and a<sub>2</sub> - 50 cm between rows), B - cultivated maize hybrid, with three graduations (b<sub>1</sub> - H<sub>1</sub>, b<sub>2</sub> - H<sub>2</sub> and b<sub>3</sub> - H<sub>3</sub>) and C - plants density, with three graduations (c<sub>1</sub> - 70,000 plants·ha<sup>-1</sup>, c<sub>2</sub> - 83,000 plants·ha<sup>-1</sup> and c<sub>3</sub> - 100,000 plants·ha<sup>-1</sup>).

The precursor culture was rape, the experience being in the second year of research in the same location. In the autumn, a basic fertilization of 200 kg·ha<sup>-1</sup> (N<sub>18</sub>P<sub>46</sub>) was carried out and in the pre-sowing seedbed was fertilized throughout the experiment with 200 kg·ha<sup>-1</sup> of urea (N<sub>46</sub>).

Production was expressed in dry matter (DM); the results were statistically analyzed by the analyses of variance and limit differences.

## RESULTS AND DISCUSSIONS

From the analysis of the influence of the distance between rows on the production of dry matter (*table 1*), it appears that the distance between rows had influenced insignificantly the production of dry matter. Thus, reducing the interval between plants to 50 cm resulted in the highest production of 15.71 Mg·ha<sup>-1</sup> DM, the increase being 9.9% compared to the control variant. The lowest production, of 14.29 Mg·ha<sup>-1</sup> DM was recorded in the control variant, respectively at the distance between rows of 70 cm. From the analysis of the influence of the hybrid on the production of dry matter (*table 2*), it appears that the genotypes analyzed behaved differently, the differences from the control (b<sub>1</sub>-H<sub>1</sub>) being very insignificant and distinctly significant. Within the experience, in 2018, the H<sub>3</sub> hybrid adapted best to the pedoclimatic conditions in the studied area, achieving the highest production of dry matter, of 18.49 Mg·ha<sup>-1</sup> DM, with a production increase of 33.7% compared to the control variant (b<sub>1</sub> -H<sub>1</sub>), the difference being a distinctly significant one. The lowest production, of 12.68 Mg·ha<sup>-1</sup> DM was recorded in the H<sub>2</sub> hybrid, recording a lower output than the control variant (*table 2*).

As a result of the improvement brought to the corn crop, the introduction of more productive hybrids, fertilization, irrigation, the density of the crop is considered the most valuable technological element.

Considering the separate influence of the plants density on the dry matter production in the studied maize hybrids, average dry matter productions with values between 13.94 Mg·ha<sup>-1</sup> DM at the density of 83000 plants·ha<sup>-1</sup> and 16.23 Mg·ha<sup>-1</sup> DM at the density of 70000 plants·ha<sup>-1</sup> (*table 3*).

Table 1

Experimental plot	DM production (Mg·ha <sup>-1</sup> )	Diferences		Statistical significance
		Mg·ha <sup>-1</sup>	%	
a <sub>1</sub> - 70 cm between rows (c)	14.29	Control	100	Control
a <sub>2</sub> - 50 cm between rows	15.71	1.42	109.9	ns
	LSD 5%	2.14		
	LSD 1%	3.56		
	LSD 0.1%	6.65		

Table 2

Experimental plot	DM production (Mg·ha <sup>-1</sup> )	Diferences		Statistical significance
		Mg·ha <sup>-1</sup>	%	
b <sub>1</sub> - H <sub>1</sub> (control)	13.83	Control	100	Control
b <sub>2</sub> - H <sub>2</sub>	12.68	-1.15	91.7	ns
b <sub>3</sub> - H <sub>3</sub>	18.49	4.66	133.7	**
	LSD 5%	2.51		
	LSD 1%	4.17		
	LSD 0.1%	7.79		

The increase of the density from 70000 plants·ha<sup>-1</sup> to 100000 plants·ha<sup>-1</sup> has caused a significant increase in the production of dry matter. Of the studied variants, the density of 70000 plants·ha<sup>-1</sup> determined to obtain the highest production of dry matter, respectively of 16.23 Mg·ha<sup>-1</sup> DM.

Analyzing the influence of the studied factors (the distance between rows, the density of plants and the cultivated hybrid) on the production of dry matter (table 4), it is found that the values of production of maize for silage, are different depending on the hybrid, and on the same hybrid, depending on the density of plants and the distance between the rows. Thus, at the distance of 70 cm

between rows, at all three density plants studied, the hybrids recorded different values of the production of dry matter in relation to the control variant (table 4), the highest production of dry matter, of 19.26 Mg·ha<sup>-1</sup> DM being recorded at the interaction between a<sub>1</sub> - 70 cm x b<sub>3</sub> - H<sub>3</sub> x c<sub>3</sub> - 70000 plants·ha<sup>-1</sup>, with a very significant difference compared to the control.

As can be seen (table 4), at the distance between rows of 70 cm, the values of the dry matter production registered a slight decrease with the increase of the density in all the three hybrids, with positive and negative differences compared to the control (table 4).

Table 3

**Influence of plants density on dry matter production**

Experimental plot	DM production (Mg·ha <sup>-1</sup> )	Diferences		Statistical significance
		Mg·ha <sup>-1</sup>	%	
c <sub>1</sub> - 70000 plants·ha <sup>-1</sup> (c)	16.23	Control	100	Control
c <sub>2</sub> - 83000 plants·ha <sup>-1</sup>	14.83	-1.40	91.4	ns
c <sub>3</sub> - 100000 plants·ha <sup>-1</sup>	13.94	-2.29	85.9	ns
	LSD 5%	2.48		
	LSD 1%	4.10		
	LSD 0.1%	7.67		

Table 4

**Influence of interaction between experimental factors on dry matter production**

Experimental plot		Dry matter production (Mg·ha <sup>-1</sup> )	Diferences		Statistical significance	
			Mg·ha <sup>-1</sup>	%		
a <sub>1</sub> - 70 cm between rows (c)	b <sub>1</sub> -H <sub>1</sub>	c <sub>1</sub> - 70000 plants·ha <sup>-1</sup> (c)	14.97	Control	100	Control
		c <sub>2</sub> - 83000 plants·ha <sup>-1</sup>	12.59	-2.38	84.1	o
		c <sub>3</sub> - 100000 plants·ha <sup>-1</sup>	11.98	-2.99	80.0	oo
	b <sub>2</sub> -H <sub>2</sub>	c <sub>1</sub> - 70000 plants·ha <sup>-1</sup>	12.37	-2.60	82.6	oo
		c <sub>2</sub> - 83000 plants·ha <sup>-1</sup>	12.21	-2.76	81.6	oo
		c <sub>3</sub> - 100000 plants·ha <sup>-1</sup>	11.65	-3.32	77.8	ooo
	b <sub>3</sub> -H <sub>3</sub>	c <sub>1</sub> - 70000 plants·ha <sup>-1</sup>	19.26	4.29	128.7	***
		c <sub>2</sub> - 83000 plants·ha <sup>-1</sup>	17.58	2.61	117.4	**
		c <sub>3</sub> - 100000 plants·ha <sup>-1</sup>	16.01	1.04	106.9	ns
a <sub>2</sub> - 50 cm between rows	b <sub>1</sub> -H <sub>1</sub>	c <sub>1</sub> - 70000 plants·ha <sup>-1</sup>	16.32	1.35	109.0	ns
		c <sub>2</sub> - 83000 plants·ha <sup>-1</sup>	13.85	-1.12	92.5	ns
		c <sub>3</sub> - 100000 plants·ha <sup>-1</sup>	13.30	-1.67	88.8	ns
	b <sub>2</sub> -H <sub>2</sub>	c <sub>1</sub> - 70000 plants·ha <sup>-1</sup>	13.48	-1.49	90.1	ns
		c <sub>2</sub> - 83000 plants·ha <sup>-1</sup>	13.43	-1.54	89.7	ns
		c <sub>3</sub> - 100000 plants·ha <sup>-1</sup>	12.93	-2.04	86.4	o
	b <sub>3</sub> -H <sub>3</sub>	c <sub>1</sub> - 70000 plants·ha <sup>-1</sup>	20.99	6.02	140.2	***
		c <sub>2</sub> - 83000 plants·ha <sup>-1</sup>	19.34	4.37	129.2	***
		c <sub>3</sub> - 100000 plants·ha <sup>-1</sup>	17.77	2.80	118.7	**
		LSD 5%	1.88			
		LSD 1%	2.52			
		LSD 0.1%	3.32			

The distance between rows of 50 cm, caused a slight increase in the production of dry matter in most variants compared to the distance between rows of 70 cm, the values obtained being different depending on the plants density, with insignificant differences, distinctly significant and very significant compared to the control variant

(table 4). Thus, at the distance of 50 cm between rows, the studied hybrids recorded higher values of dry matter production, the highest production, of 20.99 Mg·ha<sup>-1</sup> DM being recorded at the interaction between a<sub>2</sub> - 50 cm x b<sub>1</sub> - H<sub>3</sub> x c<sub>1</sub>-70000 plants·ha<sup>-1</sup>, with a very significant difference compared to the control. Also, high values of the

dry matter production, of 19.34 Mg·ha<sup>-1</sup> DM and 17.77 Mg·ha<sup>-1</sup> DM respectively were recorded in the same H<sub>3</sub> hybrid at the density of 83000 plants·ha<sup>-1</sup>, as well as in the interaction between a<sub>2</sub> - 50 cm x b<sub>3</sub> - H<sub>3</sub> x c<sub>3</sub> - 100000 plants·ha<sup>-1</sup>, the differences being distinct and very significant compared to the control. According to the data obtained, it can be seen that the plants of density and the distance between the rows significantly influenced the production of dry matter (table 4).

## CONCLUSIONS

The success of maize for silage crop implies some requirements that refer to establishing the optimum plants of density, choosing the hybrids, choosing the technological works and establishing the optimal harvest time.

The density of corn crop for silage is considered the most important technological element.

The obtained results showed that the technological factors (the density of plants and the distance between rows) caused changes in the biomass production, the quantity of dry matter being influenced by the hybrid, as well as the plants density and the distance between rows. Sowing corn for silage at a distance of 50 cm between rows, compared to 70 cm, increased the production of dry matter.

As the density of plants per hectare increases, the amount of dry matter decreases thus registering at all three densities, reduced production differences.

## REFERENCES

Çarpıcı Budaklı E., Çelik N., Bayram G., 2010 - Yield and quality of forage maize as influenced by plant density and nitrogen rate. Turkish Journal of Field Crops, vol. 15(2): 128-132.

Cañadas Á., Molina C., Rade D., Fernández F., 2016 - Seasons and planting densities interaction on

forage production of eight hybrids maize Ecuador. Revista MVZ Córdoba, vol. 21(1).

Ceclan O.A., 2010 - Studiul comportării în câmp și a pretabilității la însilozare a unor hibrizi de porumb în condițiile SCDA - Turda. Teză de doctorat, USAMV Cluj-Napoca.

Chaudhary D.P. Jat S.L., Kumar R., Kumar A., Kumar B., 2013 - Fodder quality of maize: its preservation. Maize: Nutrition Dynamics and Novel Uses: 153-160.

Ferreira G., Alfonso M., Depino S., Alessandri E., 2014 - Effect of planting density on nutritional quality of green-chopped corn for silage, Journal of Dairy Science, vol. 97, nr. 9: 5918-5921.

Ferreira G. and Teets C.L., 2017 - Effect of planting density on yield, nutritional quality, and ruminal in vitro digestibility of corn for silage grown under on-farm conditions. American Registry of Professional Animal Scientists, vol. 33, nr. 4: 420-425.

Mandić V., Krnaja V., Bijelic Z., Tomic Z., Simic A., Stanojkovic A., Petricevic V., Petrovic-Caro V., 2015 - The effect of crop density on yield of forage maize. Biotechnology in Animal Husbandry 31 (4): 567-575.

Oliveira I.L., Lima L.M., Casagrande D.R., Lara M.A., Bernardes T.F., 2017 - Nutritive value of corn silage from intensive dairy farms in Brasil. Revista Brasileira de Zootecnia, vol. 46(6).

Ramezani M., Abandani Soukht Rezaie R., Mobasser H.R., Amiri E., 2011 - Effects of row spacing and plant density on silage yield of corn (*Zea mays L.*) in two plant pattern in North of Iran. African Journal of Agricultural Research, vol. 6(5): 1128-1133.

Roth G.W. and Heinrichs A.J., 2017 - Corn silage production and management. Nutrition Workshop Proceedings.

Sharifi R.S. and Namvar A., 2016 - Plant density and intra-row spacing effects on phenology, dry matter accumulation and leaf area index of maize in second cropping. BIOLOGIA, vol. 62, nr. 1: 46-57.

Vintu V., Samuil C., Stavarache M., Muntianu I., Popovici C.I., 2010 - Research on the influence of some technological factors on production components of biomass in maize (*Zea Mays L.*) for bioenergy under the forest-steppe conditions of Moldova. Lucrări Științifice, seria Agronomie vol. 53, nr. 2.