## PARTIAL RESULTS REGARDING THE TYPES OF SUBSTRATES USED FOR OBTAINING MICROGREENS VEGETABLE PRODUCTS

## REZULTATE PARTIALE PRIVIND TIPURILE DE SUBSTRATURI FOLOSITE PENTRU OBȚINEREA CULTURILOR LEGUMICOLE DE TIP MICROGREENS

MAFTEI (HRISCU) Adriana<sup>1</sup>, MUNTEANU N.<sup>1\*</sup>, STOLERU V.<sup>1</sup>, TELIBAN G.C.<sup>1</sup>, GACHE (LUNGU) Mirabela<sup>1</sup> \*Corresponding author e-mail: nmunte@uaiasi.ro

Abstract. The paper addresses research on the use of four types of substrates (coconut substrate, 80% peat + 20% coconut waste, 80% peat + 20% sand and 80% peat + 20% pearlite) to obtain the products microgreens type vegetables, in order to establish the best substrate variant. The best results were obtained on the coconut waste substrate.

Key words: substrate, coconut waste, peat, microgreens

**Rezumat.** Lucrarea abordează cercetări privind folosirea a patru tipuri de substraturi (substrat pe bază de nucă de cocos, 80% turbă +20% deşeu de nucă de cocos, 80% turbă + 20% nisip şi 80% turbă+20% perlit) pentru obținerea produselor legumicole de tip microgreens, în vederea stabilirii celei mai bune variante de substrat. Cele mai bune rezultate au fost obținute pe substratul ce a avut la bază fibra de nucă de cocos.

Cuvinte cheie: substrat, nucă de cocos, turbă, microgreens

### **INTRODUCTION**

Microgreens or as they are called "confetti vegetables", are a new class of products, which refers to species in the category of vegetables, which are produced from vegetable seeds, herbs, including wild species (Franks and Richardson, 2009; Di Gioia and Santamaria, 2015).

Their popularity is due to the bright colors, delicate textures, unique flavor enhancement properties, such as garnishes (salad, soups, desserts, etc.), but also the high content of phytonutrients and their bioactive potential. Microgreens can be distributed as fresh vegetables, but they can also be harvested by the final consumers, by delivering the product with the nutritious substrate, harvesting when needed. The most common species of microgreens belong to the following botanical families: *Brasicaceae*, *Asteraceae*, *Chenopodiaceae*, *Lamiaceae*, *Curcubitaceae*, *Apiaceae* (Delian *et al.*, 2015; Maftei *et al.*, 2018a).

<sup>&</sup>lt;sup>1</sup>University of Agricultural Sciences and Veterinary Medicine Iasi, Romania

Microgreens represent a newer category of food products, different from germs and baby products, with a height of 3 to 10 cm, with a short duration of vegetation and with a unique appearance (Kyriacou *et al.*, 2016).

The choice of the growth substrate for the production of vegetable plants of microgreens type, plays a fundamental role in determining the productivity and quality of the plants. The purpose of this paper is to establish the best type of substrate used for the production of vegetable microgreens, in order to optimize the crop. Objectives In order to achieve the proposed purpose, the following objectives were established: establishing the duration for seed germination, establishing the days needed for growth and development, establishing the average harvesting duration and assessing the uniformity of the crops.

## MATERIAL AND METHOD

The research was carried out in the Phytotron greenhouse of the Vegetable discipline, within the farm "Vasile Adamachi" lasi in 8- 22 October 2018.

Biometric observation and determination were used as a research method. The biological material used is represented by seven vegetable species namely: Pears - *Pisum sativum*, Red moon radish - *Raphanus sativus*, White summer radish - *Raphanus sativus*, Early cabbage - *Brassica oleracea*, Red cabbage - *Brassica oleracea*, Red basil cousin. violaceum, Green basil - *Ociumum basilicum* var. *viridis*. The technical material is represented by the four types of substrates: Coconut (coconut waste); Peat 80 % with 20 % coconut waste; Peat 80 % with 20 % pearlite; Peat 80 % with 20 % sand.

Beginning with seed germination, the management of microclimate factors will be based on species requirements (Ipătioaie *et al.*, 2016; Lobiuc *et al.*, 2017; Voicu *et al.*, 2017; Cărbune *et al.*, 2018; Maftei *et al.*, 2018b).

## **RESULTS AND DISCUSSIONS**

### 1. Results obtained on average days for seed germination

Results obtained regarding the number of average days for seed germination are presented in table 1 and indicate that, the best type of substrate is the one based on coconut, followed by the one based on peat with coconut, and the most poor results, being obtained on the substrate based on pearlite and peat.

Table 1

| Species                                      | Nr. days |   |   |   |   |   |   |   |   |    |    |    |  |
|--|----------|---|---|---|---|---|---|---|---|----|----|----|--|
| Species                                      | 1        | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 |  |
| Pears - <i>Pisum sativum</i>                 |          |   |   |   |   |   |   |   |   |    |    |    |  |
|  |          |   |   |   |   |   |   |   |   |    |    |    |  |
|  |          |   |   |   |   |   |   |   |   |    |    |    |  |
|  |          |   |   |   |   |   |   |   |   |    |    |    |  |
|  |          |   |   |   |   |   |   |   |   |    |    |    |  |
| Red moon radish -<br><i>Raphanus sativus</i> |          |   |   |   |   |   |   |   |   |    |    |    |  |
|  |          |   |   |   |   |   |   |   |   |    |    |    |  |
|  |          |   |   |   |   |   |   |   |   |    |    |    |  |

Results obtained on average days for seed germination

Legend: Coconut (coconut waste) 100% Peat 80 % with 20 % coconut waste Peat 80 % with 20 % pearlite Peat 80 % with 20 % sand

# 2. Results regarding the establishment of the average days of growth and development

The results regarding the establishment of the average days of plant growth and development are presented in table 2. These results highlight that for the coconut substrate, the pea needs 4-7 days average for the development of plants, early cabbage of 4-8 days, and basil for 5-9 days. Not the same can be said about the peat and pearl substrate, where the values increase by 2-3 days ie: for red cabbage a number of 7-10 days is required for the development of plants, 4-8 days for radishes and 8-12 days for basil.

Table 2

| Species                      | Nr. days |   |   |   |   |   |   |   |   |    |    |    |  |
|------------------------------|----------|---|---|---|---|---|---|---|---|----|----|----|--|
|                              | 1        | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 |  |
| Pears - <i>Pisum sativum</i> |          |   |   |   |   |   |   |   |   |    |    |    |  |
|                              |          |   |   |   |   |   |   |   |   |    |    |    |  |
|                              |          |   |   |   |   |   |   |   |   |    |    |    |  |
|                              |          |   |   |   |   |   |   |   |   |    |    |    |  |
|                              |          |   |   |   |   |   |   |   |   |    |    |    |  |
| Red moon radish -            |          |   |   |   |   |   |   |   |   |    |    |    |  |
| Raphanus sativus             |          |   |   |   |   |   |   |   |   |    |    |    |  |
|                              |          |   |   |   |   |   |   |   |   |    |    |    |  |

### Results regarding the establishment of the average days of growth and development

| White summer radish -                     |  |  |  |  |  |  |
|---|--|--|--|--|--|--|
| Raphanus sativus                          |  |  |  |  |  |  |
|   |  |  |  |  |  |  |
|   |  |  |  |  |  |  |
| Early cabbage -                           |  |  |  |  |  |  |
| Brassica oleracea                         |  |  |  |  |  |  |
| Red cabbage - <i>Brassica</i><br>oleracea |  |  |  |  |  |  |
|   |  |  |  |  |  |  |
|   |  |  |  |  |  |  |
|   |  |  |  |  |  |  |
|   |  |  |  |  |  |  |
|   |  |  |  |  |  |  |
| Red basil cousin.                         |  |  |  |  |  |  |
| violaceum                                 |  |  |  |  |  |  |
|   |  |  |  |  |  |  |
|   |  |  |  |  |  |  |
| Green basil - Ociumum                     |  |  |  |  |  |  |
| basilicum var. viridis                    |  |  |  |  |  |  |
|   |  |  |  |  |  |  |
|   |  |  |  |  |  |  |
| Legend                                    |  |  |  |  |  |  |

Coconut (coconut waste) 100% Peat 80 % with 20 % coconut waste Peat 80 % with 20 % pearlite Peat 80 % with 20 % sand

## 3. Results regarding the establishment of the average harvest days

Table 3 contains results regarding the establishment of the average harvest days, which presents us as well as the other studies on the coconut substrate the best results, followed by the substrate consisting of peat and coconut and peat and sand, and the weakest results are obtained for peat and pearl substrate.

Table 3

|                              | - |   |   |   |   |   |   |       | -    |    |    |    |    |    |
|------------------------------|---|---|---|---|---|---|---|-------|------|----|----|----|----|----|
| Species                      |   |   |   |   |   |   |   | Nr. c | lays |    |    |    |    |    |
| Species                      | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8     | 9    | 10 | 11 | 12 | 13 | 14 |
| Pears - <i>Pisum sativum</i> |   |   |   |   |   |   |   |       |      |    |    |    |    |    |
|                              |   |   |   |   |   |   |   |       |      |    |    |    |    |    |
|                              |   |   |   |   |   |   |   |       |      |    |    |    |    |    |
|                              |   |   |   |   |   |   |   |       |      |    |    |    |    |    |
|                              |   |   |   |   |   |   |   |       |      |    |    |    |    |    |
| Red moon radish -            |   |   |   |   |   |   |   |       |      |    |    |    |    |    |
| Raphanus sativus             |   |   |   |   |   |   |   |       |      |    |    |    |    |    |
|                              |   |   |   |   |   |   |   |       |      |    |    |    |    |    |
|                              |   |   |   |   |   |   |   |       |      |    |    |    |    |    |
| White summer radish -        |   |   |   |   |   |   |   |       |      |    |    |    |    |    |
| Raphanus sativus             |   |   |   |   |   |   |   |       |      |    |    |    |    |    |
|                              |   |   |   |   |   |   |   |       |      |    |    |    |    |    |

Results regarding the establishment of the average harvest days

| Early cabbage - Brassica     |  |  |  |  |  |  |  |
|------------------------------|--|--|--|--|--|--|--|
|                              |  |  |  |  |  |  |  |
| oleracea                     |  |  |  |  |  |  |  |
|                              |  |  |  |  |  |  |  |
|                              |  |  |  |  |  |  |  |
| Red cabbage - Brassica       |  |  |  |  |  |  |  |
| oleracea                     |  |  |  |  |  |  |  |
|                              |  |  |  |  |  |  |  |
|                              |  |  |  |  |  |  |  |
| Red basil cousin.            |  |  |  |  |  |  |  |
| violaceum                    |  |  |  |  |  |  |  |
|                              |  |  |  |  |  |  |  |
|                              |  |  |  |  |  |  |  |
| Green basil - <i>Ociumum</i> |  |  |  |  |  |  |  |
| basilicum var. viridis       |  |  |  |  |  |  |  |
|                              |  |  |  |  |  |  |  |
|                              |  |  |  |  |  |  |  |

Legend

Coconut (coconut waste) 100% Peat 80 % with 20 % coconut waste Peat 80 % with 20 % pearlite Peat 80 % with 20 % sand

## CONCLUSIONS

1. The best type of substrate used for growing vegetables microgreens is the one based on coconut waste, followed by 80% peat + 20% walnut waste cock.

2. The weakest type of substrate used for type culture microgreens is the 80% peat and 20% pearlite.

3. All seven species studied in the microgreens system are suitable for cultivation in this system.

4. Radish is the species that has the fastest development until harvest, followed by the opposite pole of basil, which needs the most days of growth and development.

### REFERENCES

- Cărbune R.D., Munteanu N., Stoleru C.M., Stan T., Teliban G.C., Stoleru V., 2018 - Preliminary results on germination and velocity of Kelvedon Wonder pea seeds variety. Scientific Papers, Horticulture, vol. 61, nr. 2, pp 177-182, USAMV Iasi. ISSN: 1454-7376.
- 2. Delian E., Chira A., Bădulescu L., Chira L., 2015 Insights into microgreens physiology. Scientific Papers-Series B-Horticulture, vol. 59, pp. 447-454, Bucharest.
- 3. Di Gioia F., Santamaria P., 2015 Microgreens. ECO Publishing House Logistics SRL, Bari.
- 4. Franks E., Richardson J., 2009 *Microgreens: A Guide to Groiwing Nutrient Packed Greens.* Published by Gibbs Smith, Layton, Utah USA. ISBN-13: 978-1-4236-0364-1.
- Ipătioaie D.C., Munteanu N., Stoleru V., Teliban G.C., Cojocaru Al., 2016 A study on the seed germination and plantlets spring at rhubarb species (Rheum rhabarbarum L.). Scientific Papers, Horticulture, vol. 59, nr. 2, pp. 337-342, USAMV

lasi. ISSN: 1454-7376.

- Kyriacou M., Rouphael Y., Di Gioia F., Kyratzis A., Serio F., Renna M., De Pascale S., Santamaria P., 2016 - Micro-scale vegetable production and the rice of microgreens. Trend in Food Science & Technology, v. 57, p 103-115. http://dx.doi.org/10.1016/j.tifs.2016.09.005.
- Lobiuc A., Vasilache V., Oroian M., Stoleru T., Burducea M., Pintilie O., Zamfirache M.M., 2017 - Blue and red LED illumination improves growth and bioactive compounds contents in acyanic and cyanic Ocimum basilicum L. microgreens. Molecules, 22 (12), 2111. https://doi.org/10.3390/molecules22122111.
- Maftei (Hriscu) A., Munteanu N., Stoleru V., Teliban G.C., Galea (Deleanu) F.M., 2018a - Stage of knowledge on cultivation of microgreens plants. Scientific Papers, Horticulture, vol. 61, nr. 2, pp 75-80, USAMV Iasi. ISSN: 1454-7376.
- Maftei (Hriscu) A., Munteanu N., Stoleru V., Teliban G.C., 2018b Studies of seeds germination as a technical stage in the production of microgreens. Scientific Papers, Horticulture, vol. 61, nr. 2, pp 71-74, USAMV Iasi. ISSN: 1454-7376.
- Voicu M., Munteanu N., Stoleru V., Teliban G., Stoleru C., 2017 The influence of the storage period of pea seeds on their germination capacity. Scientific Papers, Horticulture, vol. 60, no. 1, pp. 51-56, USAMV Iasi. ISSN: 1454-7376.