

THE STUDY OF THE CLIMBING POD BEANS REGARDING THE INFLUENCE OF THE CULTIVAR AND PLANT DENSITY ON SEED PRODUCTION / HECTARE UNDER THE ENVIRONMENTAL CONDITIONS OF ZONE IERNUT

STUDIU INFLUENȚEI CULTIVARULUI ȘI A DENSITĂȚII PLANTELOR/HA ASUPRA PRODUCȚIEI DE SEMINȚE/HA LA FASOLEA DE PĂSTĂI URCĂTOARE, ÎN CONDIȚIILE DE MEDIU ALE S.C.D.L. IERNUT

FARCAȘ Laura¹*, **CHIPER Lidia¹**, **RADU Aurelia¹**

*Corresponding author e-mail: minervaheitz@yahoo.com

Abstract. The paper presents the research results from a bifactorial experiment carried out at Vegetable Research and Development Station Iernut. Experimental factors were four cultivars (*Mădărășeni* cultivar and three perspective lines – *Alina*, *Ghibolească* and *Viola-2*) and three crop density determined by the number of plants in a hole (2, 3 and 4 plants / hole). The goal of the experiment was to evaluate the influence of the two factors on the seed production. The best result was obtained for all the cultivars and a density of two plants/hole.

Keywords: climbed, bean pods, cultivar, density, production

Rezumat. Lucrarea prezintă rezultatele de cercetare obținute într-o experiență de tip bifactorial organizată în condițiile de la Stațiunea de Cercetare Dezvoltare pentru Legumicultura Iernut. Factorii experimentalni luati în studiu au fost reprezentanți de patru cultivare (*Mădărășeni* și trei linii de perspectivă - *Alina*, *Ghibolească* și *Viola-2*) și trei densități de cultură determinate de numărul de plante dintr-un cuib (2, 3 și 4 plante / cuib). Scopul acestei cercetări a fost evaluarea influenței celor doi factori asupra producției de semințe. Cel mai bun rezultat a fost obținut pentru toate cultivarele la o densitate de două plante / cuib.

Cuvinte cheie: urcătoare, fasole păstăi, cultivare, densitate, producție

INTRODUCTION

The bean is an important vegetable species for both pods and grains (Avasilcăi *et al.*, 2017; Hamburdă *et al.*, 2016; Teliban *et al.*, 2014, 2015). This paper refers to the production of seeds per unit area. Local populations were studied and were created through the ameliorative selection, under the conditions from Transilvania area, valuable lines of climbing beans for pods.

¹Research and Development Station for Vegetable Growing of Iernut, Romania

MATERIAL AND METHODS

The biological material used, is part of the Măldăreșeni cultivar and of grains ameliorative lines which are valuable for Transylvania's conditions (L-Alina, L-Grasa of lernut (Ghiboleasca) and a4-L-Viola-2. The experiment has been organized in a split plot design with four replicates for the two factors: cultivar and density.

- the A-factor (the cultivar), with four graduations: A₁ - Măldăreșeni, A₂ - L-Alina, A₃ - L-Grasa of lernut (Ghiboleasca) and A₄ - L-Viola-2;

- the B-factor, the number of plants per planting hole, with three graduations: b1 - two plants on the planting hole, b2 - three plants on the planting hole and b3 - four plants on the planting hole. The distance between rows was of 80 cm and the distance between the planting holes was of 25 cm (Ruști and Munteanu, 2008; Stan *et al.*, 2003; Teodorescu *et al.*, 2012, 2014).

The influence of experimental factors was evaluated by the seed yield of each graduation of factors (Ceapoiu, 1968).

RESULTS AND DISCUSSION

The biological material used was characterized by specific features of the cultivar (tab. 1, tab. 2).

Table 1

Cultivars' description

The cultivar	Precocity	Vegetation period	Moment of crop establishment	
			For pods	For seed
Măldăreșeni	Early	77	1 – 10.05	20.04
L – Alina	Belated	115	30.04 – 05.05	25.04
L – Ghibolească	Semi-early	78	05.05 – 10.05	30.04
L – Viola-2	Semi-belated	85	03.05 – 10.05	20.04



Fig. 1 Măldăreșeni Variety - plants, pods and flowers

From the analysis of the influence of factor "A", it was found that compared to witness "A₁" cultivar was a difference: very significantly negatively for "A₂" cultivar and significantly positively for "A₃" cultivar (tab. 2).



Fig. 2 L-Alina - plant, pods and flowers

Table 2

The influence of factor "A"

The factor	Production		Diff. t/ha	Meaning
	t/ha	%		
A ₁	3.4	100	Mt	Mt
A ₂	2.76	81.17	- 0.64	000
A ₃	3.9	114.7	0.5	***
A ₄	3.66	107.6	0.26	*

LSD 5% = 0.104 x 2.26 = 0.235 t/ha

LSD 1% = 0.104 x 3.25 = 0.338 t/ha

LSD 0.1% = 0.104 x 4.78 = 0.497 t/ha

Table 3

Multiple comparisons of factor B

Clasification	Production (t/ha)	The difference between the variant from the ...place		
		III – B ₃	II – B ₂	I – B ₁
I – B ₁	4.6	2.35**	1.15	-
II – B ₂	3.45	1.2	-	
III - B ₃	2.25	-		

LSD 5% = 0.761 x 2.06 = 1.567 t/ha

LSD 1% = 0.761 x 2.8 = 2.13 t/ha

LSD 0.1% = 0.761 x 3.75 = 2.85 t/ha

The analysis of the multiple comparisons of factor "B" at the same graduation of "A" showed the significant positively differences between "B₁" and "B₃" from all cultivars, distinctly significant from the other comparisons especially for L-Ghimbolesca cultivars, L-Viola-2 or significant in the case of Mădărășeni and L - Alina (tab. 4) cultivars.

Table 4

Multiple comparisons of factor "B" at the same graduation of "A" regarding production of seed /he

Clasification	Production (t/ha)	The difference between the variant from the place		
		III	II	I
A ₁	I	4.5	2.2***	1.1*
	II	3.4	1.1*	-
	III	2.3	-	
		III	II	I
A ₂	I	3.7	1.9***	0.9*
	II	2.8	1.0*	-
	III	1.8	-	
		III	II	I
A ₃	I	5.2	2.6***	1.3**
	II	3.9	1.3**	-
	III	2.6	-	
		III	II	I
A ₄	I	5.0	2.7***	1.3**
	II	3.7	1.4**	-
	III	2.3	-	

LSD 5% = 0.416 x 2.06 = 0.857 t/ha;

LSD 1% = 0.416 x 2.8 = 1.165 t/ha

LSD 0.1% = 0.416 x 3.75 = 1.56 t/ha

The meanings of the differences between the variants (seed production per hectare) have shown very clearly and also through the analysis of comparing two “A’s” average at the same graduation of B or at different graduations of B (tab. 5).

Table 5
Comparison of two “A’s” average at the same graduation of “B” or “B’s” different graduations (the production of seed per hectare)

The factor	t/ha	The difference between the variant from the place											
		XII	XI	X	IX	VIII	VII	VI	V	IV	III	II	
I– A ₃ B ₁	5.2	3.4***	2.9***	2.9***	2.6***	2.4***	1.8***	1.5***	1.5***	1.3***	0.7**	0.2	
II– A ₄ B ₁	5.0	3.2***	2.7***	2.7***	2.4***	2.2***	1.6***	1.3***	1.3***	1.1***	0.5*	-	
III– A ₁ B ₁	4.5	2.7***	2.2***	2.2***	1.9***	1.7***	1.1***	0.8***	0.8***	0.6**	-		
IV– A ₃ B ₂	3.9	2.1***	1.6***	1.6***	1.3***	1.1***	0.5*	0.2	0.2	-			
V– A ₂ B ₁	3.7	1.9***	1.4***	1.4***	1.1***	0.9***	0.3	0	-				
VI– A ₄ B ₂	3.7	1.9***	1.4***	1.4***	1.1***	0.9***	0.3	-					
VII– A ₁ B ₂	3.4	1.6***	1.1***	1.1***	0.8***	0.6**	-						
VIII– A ₂ B ₂	2.8	1.0***	0.5*	0.5*	0.2	-							
IX– A ₃ B ₃	2.6	0.8***	0.3	0.3	-								
X– A ₁ B ₃	2.3	0.5*	0	-									
XI– A ₄ B ₃	2.3	0.5*	-										
XII– A ₂ B ₃	1.8	-											

CONCLUSIONS

1. The four cultivars are highly valued by consumers, although they are differentiated by different quantitative and qualitative features.
2. As result of the capacity of seed production, the cultivars can be classified in the following order: L-Ghimboleasca, L-Viola 2, Mădărășeni, L-Alina.

3. In all cultivars, we obtained the highest seed production for B₁ (two seed / the hole) the establishment saving biological material culture.

REFERENCES

1. Avasilcăi L., Teliban G., Morariu D.I., Stoleru V., Bibire N., Vieriu M., Panainte A.D., Munteanu N., 2017 – *Parameters of Chemical Composition of Phaseolus coccineus L. Pods Grown in Protected Areas*. Revista de chimie, vol. 68, no. 12, pp. 2955-2958, Bucharest, Romania. ISSN: 0034-7752.
2. Ceapoiu N., 1968 – *Statistical methods in agricultural and biological experiments*. Agro-Forestry Publishing House, Bucharest.
3. Hamburdă S.B., Teliban G.C., Munteanu N., Stoleru V., 2016 – *Effect of Intercropping System on the Quality and Quantity of Runner Bean (Phaseolus coccineus L.)*. Notulae Botanicae Horti Agrobotanici, Cluj-Napoca, Romania, vol. 44, Issue: 2, pp. 613-618. ISSN: 0255-965X. DOI: 10.15835/nbha44210260.
4. Ruști G., Munteanu N., 2008 – *Cultura fasolei de grădină urcătoare*. Editura "Ion Ionescu de la Brad" Iași, ISBN 978-973-147-014-6.
5. Stan N., Munteanu N., Stan T., 2003 – *Legumicultură. Volumul III*. Editura "Ion Ionescu de la Brad", Iași.
6. Teliban G. C., Munteanu N., Popa L.D., Stoleru V., Stan T., Hamburdă S.B., 2014 – *The study of the influence of the planting distance on the early production of certain runner bean cultivars (Phaseolus coccineus L.) for pods, in the environment of the polytunnel*. Lucrări științifice, seria Horticultură, vol. 57, nr. 1, pp. 105-110, USAMV Iași. ISSN: 1454-7376.
7. Teliban G.C., Munteanu N., Popa Lorena-Diana, Stoleru V., Hamburdă S.B., Stan T., Buburuz A.A., Onofrei V., Ciobanu V., 2015 – *The behavior of a runner bean assortment for pods (Phaseolus coccineus L.) in polyethylene tunnels, in a crop established by direct sowing*. Lucrări științifice, seria Horticultură, vol. 58, nr. 1, pp. 117-122, USAMV Iași. ISSN: 1454-7376.
8. Teodorescu E., Munteanu N., Vinătoru C., Stoleru V., Zamfir B., 2012 – *Phenotypical research concerning climbing bean pods belonging to the germplasm collection from Vegetable Research-Development Station Buzău*. Bulletin of University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca. Horticulture, vol. 69 (1), pp. 427-428.
9. Teodorescu E., Munteanu N., Vinătoru C., 2014 – *Drought Effects to the Dry Bean Local Population Productivity*. Bulletin UASVM Cluj-Napoca - Agriculture 71 (1), ISSN 1843-5246, Electronic ISSN 1843-5386.