



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

XXI International Grassland Congress / VIII
International Rangeland Congress

The Production and Nutritional Composition of Forage Sorghum Cultivars as Silage Crops

A. Swanepoel

Western Cape Department of Agriculture, South Africa

Philip R. Botha

Western Cape Department of Agriculture, South Africa

H. S. Gerber

Western Cape Department of Agriculture, South Africa

Robin Meeske

Western Cape Department of Agriculture, South Africa

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/15-1/22>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

The production and nutritional composition of forage sorghum cultivars as silage crops

A . Swanepoel , P .R . Botha , H .S . Gerber and R . Meeske*

Department of Agriculture Western Cape , Institute for Plant Production , * Institute for Animal Production . Outeniqua Research Farm , George , Republic of South Africa . e-mail : annelenes@elsenburg .com

Key words : forage sorghum , production , nutritional composition , silage crop

Introduction Forage sorghum (Sorghum x Sudan grass hybrid) cultivars are palatable , high producing annual summer growing grasses suited for silage production on the duplex soils of the Southern Cape of South Africa . New cultivars are released regularly and the production potential and nutritive value of these cultivars needs to be determined . The aim of the study was to quantify the dry matter production and quality of forage sorghum cultivars for silage production .

Materials and methods The trial was carried out annually during the summer over two years (2004/2005 and 2005/2006) as a small plot trial under irrigation on a Estcourt soil type . Fertiliser was applied to raise phosphorus level to 35 mg kg⁻¹ , potassium level to 80 mg kg⁻¹ and pH (KCL) to 5 .5 . Planting dates were 25th and 24th of November 2004 and 2005 respectively . Each plot received 50 kg N ha⁻¹ and 20 kg K ha⁻¹ with planting . Four weeks after emergence 85 kg N ha⁻¹ and 45 kg K ha⁻¹ were applied as top dressing and this was repeated four weeks later . When the plants reached a dry matter content of between 25 and 30% , each cultivar was cut once at a height of 150 mm , weighed and chopped with a silage chopper to obtain a representative sample . Dry matter (DM) production , crude protein (CP) , *in vitro* organic material digestibility (IVOMD) , metabolisable energy (ME) and NDF of each cultivar were determined .

Results The forage sorghum cultivars Sugargraze , SAC 710 and Hunnigreen produced more than 18 tons of DM ha⁻¹ and are recommended for silage production (Table 1) . There are a number of cultivars that have the ability to produce between 14 and 17 tons of DM ha⁻¹ and can also be considered for silage production . The CP content varied between 7% and 9% and was low . The ME of 6 .9 , 5 .9 and 6 .4 MJ kg⁻¹ DM for Hunnigreen , Sugargraze and SAC 710 respectively were lower than some of the lower producing cultivars nl . Hygro Sil BMR (8 .1 MJ kg⁻¹ DM) and Hygro Sil (7 .8 MJ kg⁻¹ DM) . The CP content (%) , ME (MJ kg⁻¹ DM) and DM production (kg ha⁻¹) of the cultivars influenced the total CP (kg ha⁻¹) and ME (MJ ha⁻¹) . Forage sorghum cultivars differ in production potential .

Table 1 The DM production , DM (%) , CP (%) , total CP , NDF (%) , ME and total ME ha⁻¹ of forage sorghum cultivars for silage production under irrigation , for one season (2005/2006) , at Outeniqua Research Farm .

Cultivars (2005/2006)	DM (ton ha ⁻¹)	DM %	CP (%)	Total CP (kg ha ⁻¹)	NDF (%)	ME (MJ kg ⁻¹ DM)	Total ME (MJ ha ⁻¹)
SAC 710	19 .0 ^a	28 .6	7 .7	1463	64	6 .4	121600
Sugargraze	18 .6 ^{ab}	29 .4	5 .3	987	68	5 .9	109740
Hunnigreen	18 .4 ^{ab}	28 .7	6 .9	1270	62	6 .9	126960
Bulkmaster	16 .6 ^{bc}	30 .6	7 .4	1228	63	6 .1	101260
Superdan 401	16 .1 ^c	29 .7	7 .0	1127	62	6 .1	98210
Super King	15 .4 ^c	28 .1	9 .1	1401	57	6 .1	93940
Revolution BMR	15 .7 ^{cd}	na	5 .7	895	71	5 .2	76440
AGR 3404	14 .7 ^{cd}	28 .5	5 .3	779	69	6 .0	88200
Everlush	14 .7 ^{cd}	28 .3	6 .0	882	73	5 .1	74970
Classic Grazer	14 .5 ^{cde}	28 .1	6 .2	899	70	6 .0	87000
NS 1	13 .0 ^{def}	27 .8	6 .0	780	70	5 .5	71500
Kow Kandy BMR	12 .7 ^{def}	29 .5	7 .9	1003	61	6 .0	76200
Hygro 2 (Wei 7)	12 .4 ^{ef}	29 .8	8 .8	1091	59	6 .8	84320
AGR 6201	12 .0 ^{7f}	28 .0	5 .6	676	56	7 .9	95353
Haymaker	11 .8 ^{3f}	28 .6	7 .9	935	66	6 .0	70800
Silage King	11 .2 ^{0fg}	28 .9	6 .7	750	63	6 .1	68320
Rambo	11 .0 ^{7fg}	28 .3	7 .9	869	68	5 .3	58830
Higro Sil BMR	10 .8 ^{3fg}	28 .9	7 .9	853	54	8 .1	87480
Higro Sil	9 .3 ^{3gh}	27 .5	8 .8	818	51	7 .8	72540
Advanta BMR	8 .3 ^{3h}	29 .3	5 .3	689	64	6 .4	53120
Hygro Graze BMR	7 .9 ^{0h}	28 .8	7 .1	561	59	7 .3	57670

^{abcd} Means with no common superscript differ significantly (P<0 .05) LSD (0 .05)=2 .21 na=not available