

Apparent climate and ecological change during MIS 3-MIS 2 in northern Alaska, Itkillik River Yedoma

Lyna Lapointe Elmrabti¹⁻²⁻³, J. Talbot¹⁻³, J. Strauss⁴, M. Kanevskiy⁵, Y. Shur⁵, B. Fréchette⁶ and D. Fortier¹⁻²⁻³⁻⁵.

¹ Département de géographie, Université de Montréal, Montréal, QC, Canada, ² Centre d'étude nordique, Université Laval, QC, Canada, ³ ArcticNet, Université Laval, QC, Canada, ⁴ Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, Germany, ⁵ Institute of Northern Engineering, University of Alaska Fairbanks, Fairbanks, AK, USA, ⁶ département des sciences de la Terre et de l'atmosphère, Université du Québec à Montréal, Montréal, QC, Canada.

1. Context

The cold-arid climate associated with the late Pleistocene environment of unglaciated Beringia was favorable to active sedimentation processes and accumulation of ground ice leading to the formation of a relic form of ice-rich syngenetic permafrost, termed yedoma. These periglacial features accumulated during Pleistocene between MIS 4 and MIS 2. Yedoma provide interesting snapshots of well-preserved paleoenvironmental data which can be linked to the framework established by continuous sequences while providing potentially older information (> 50 000 yrs BP). Also, data can be linked to terrestrial sedimentologic, cryostratigraphic and stratigraphic markers. Warmer conditions through the Holocene and current global warming might cause rapid thawing of these deposits.

2. Objectives

- A. Reconstruct vegetation dynamics since MIS 4 (pollen)
- B. Reconstruct paleoclimate since MIS 4 (Modern Analog Technique (MAT) and $\delta^{18}\text{O}$)
- C. Link sediment and ground-ice records (C and ice content) to A and B.

3. Study site




Fig 1. Overview of the main sampling profile. The pollen sampling areas are identified with black/yellow diamonds.





Fig 2. a) Regional location of Itkillik river yedoma in Alaska. b) Location of the main sampling profile on the bluff. Modified from Kanevskiy 2011.

4. Methods

- Field work**
By rappelling down from the top of the bluff.
- B. MAT :** Modern pollen data for North America (Whitmore et al. 2005, QSR 24, 1828-1848)
 $\delta^{18}\text{O}$: Finnigan MAT Delta-S mass spectrometer
- A. Pollen extraction:** sodium polytungstate density separation
Analysis: 400X – 1000X Leica microscope
- C. C :** Carbon-nitrogen-sulphur analyzer (Vario EL III, Elementar)
Ice content: wet-dry weight difference

5. Results

A. Palynological record




Fig 3. Pollen percentage diagram. Pollen zonation is given at the right side. 14C dates yrs BP represent radiocarbon ages obtained from the main sampling profile and the age-depth model is based on all the dates available from the site (Kanevskiy et al. 2011) and Strauss et al. (2012).

B. Paleoclimatic reconstruction




Fig 4. Average paleoclimatic curves from MAT and $\delta^{18}\text{O}$. Summer T °C = average of june, july and august. Seasonal T °C ranges is based on variation between summer and winter (december, january and february) temperatures.

C. Sediment and ground-ice




Fig 5. Sedimentological data from Itkillik river yedoma. TC = total carbon. IC/TC = inorganic carbon/total carbon. C/N = carbon/nitrogen.

6. Summary

Palynological records are subdivided into 4 vegetation/climate units that seem largely associated to marine isotopic stage (MIS 1 (PZ1), MIS 2 (PZ2), MIS 3 (PZ3), MIS 4 (PZ4)) that represent alternating warm and cool periods in the Earth's paleoclimate. Chronology of the site and $\delta^{18}\text{O}$ seem to support this hypothesis. Kanevskiy (2011) also concluded that the greatest part of the exposure was formed during the middle Wisconsinan/Karginsky interstadial (MIS 3) and late Wisconsinan/Sartan glaciation (MIS 2). Palynological records are consistent with other sites in Eastern Beringia (Fréchette et al. (2008)), but is more easily comparable to sites in Western Beringia due to its older age (> 45 000 yrs BP) and depositional environment .

We observed that vegetation is dominated by **Cyperaceae** during warmer periods (MIS 1 (PZ1) and MIS 3 (PZ3)). **Summer temperature are higher**, near the actual value (8,6 °C) and **correlated to $\delta^{18}\text{O}$ variations**. TC is low throughout the profile (4,8%), but reaches its highest average concentration during MIS 3 (6,7% compare to 3,7% in MIS 2). IC/TC is inversely correlated to TC. During MIS 3 and MIS 1 the C/N ratio indicates conditions more favorable to decomposition (21). The **volumetric ice content** is also higher (75%), potentially linked to climatic conditions promoting surface stabilization (indicated by the formation of peat layer) and segregation of ice during syngenetic development of the permafrost.

In contrast, during **cold and dry periods** (MIS 2 (PZ2) and MIS 4 (PZ4)), vegetation is mainly represented by **Poaceae** and a high diversity of minor herbs and forbs. TC is low and IC is dominant (84,4% during MIS 2 compare to 57,8% for MIS3). C/N ratio shows that the organic matter is not labile (35). **Volumetric ice content** values (56%) are lower, probably due to increased eolian sedimentation and faster syngenetic surface elevation, conditions less favorable to ice segregation during syngenetic permafrost development aggradation.

Seasonal temperature contrast and annual precipitation show slight variability throughout the profile, suggesting that they are not the determining factors for carbon accumulation, nor ground ice aggradation.

7. Future work

- Add pollen data from other profiles on the same bluff
- Correlate pollen data with cryostratigraphic units, ice wedge volume and particle size analysis
- Improve the accuracy of reconstructed climatic parameters

References

- Fréchette B., de Vernal A., Guiot J., Wolfe A.P., Miller G.H., Fredskild B., Kerwin M.W., Richard P.J.H. (2008). Methodological basis for quantitative reconstruction of air temperature and sunshine from pollen assemblages in Arctic Canada and Greenland. Quaternary Science Reviews, Vol. 27, p. 1197-1216.
- Kanevskiy M., Shur Y., Fortier D., Jorgenson M.T., Stephani E. (2011). Cryostratigraphy of late Pleistocene syngenetic permafrost (Yedoma) in northern Alaska, Itkillik River exposure. Quaternary research, Vol. 75, p. 584-596.
- Strauss J., Ulrich M., Buchhorn M. (2012). Expeditions to permafrost 2012: "Alaskan North Slope/Itkillik". Berichte zur Polar- und Meeresforschung , Reports on polar and marine research, Bremerhaven, Alfred Wegener Institute for Polar and Marine Research, Vol. 655, p. 3-28.
- Whitmore J., Gajewski K., Sawada M., Williams J.W., Shuman B., Bartlein P.J., Minckley T., Vieu A.E., Webb III T., Shafer S., Anderson P., Brubaker L. (2005). Modern pollen data from North America and Greenland for multiscale paleoenvironmental applications. Quaternary Science Reviews vol. 24, p. 1828-1848

