

1 **Polar Science: doi.org/10.1016/j.polar.2015.05.001**

2 **Methane excess production in oxygen-rich polar water and a model of cellular**
3 **conditions for this paradox**

4
5 E. Damm*, S. Thoms, A. Beszczynska-Möller, E.M. Nöthig and G. Kattner

6 Alfred Wegener Institute for Polar and Marine Research, P.O. Box 12061 D-27515
7 Bremerhaven, Germany

8
9 **Abstract**

10 Summer sea ice cover in the Arctic Ocean has undergone a reduction in the last decade
11 exposing the sea surface to unforeseen environmental changes. Melting sea ice increases
12 water stratification and induces nutrient limitation, which is also known to play a crucial role
13 in methane formation in oxygenated surface water. We report on an excess of methane in the
14 marginal ice zone in the western Fram Strait. Our study is based on measurements of oxygen,
15 methane, DMSP, nitrate and phosphate concentrations as well as on phytoplankton
16 composition and light transmission, conducted along the 79°N oceanographic transect, in the
17 western part of the Fram Strait and in Northeast Water Polynya region off Greenland.
18 Between the eastern Fram Strait, where Atlantic water enters from the south and the western
19 Fram Strait, where Polar water enters from the north, different nutrient limitations occurred
20 and consequently different bloom conditions were established. Ongoing sea ice melting
21 enhances the environmental differences between both water masses and initiates regenerated
22 production in the western Fram Strait. We show that in this region methane is in situ produced
23 while DMSP (dimethylsulfoniopropionate) released from sea ice may serve as a precursor for
24 the methane formation. The methane production occurs despite high oxygen concentrations in
25 this water masses. As the metabolic activity (respiration) of unicellular organisms explains the

26 presence of anaerobic conditions in the cellular environment we present a theoretical model
27 which explains the maintenance of anaerobic conditions for methane formation inside
28 bacterial cells, despite enhanced oxygen concentrations in the environment.

29

30 **1. Introduction**

31 The Arctic Ocean is one of the regions in the world where climate change is most
32 pronounced. Increased summer melting is considered to amplify biological production, due to
33 the shift from an ice-covered to an open water Arctic Ocean (Arrigo et al., 2008). However,
34 increasing water stratification during sea ice melting is likely to limit nutrient availability in
35 near-surface water, which in turn hampers the enhancement of primary production (Sakshaug,
36 2003). A characteristic feature of the Arctic Ocean is the distinct post-bloom nutrient
37 limitation found in the Atlantic-dominated and Pacific-dominated sectors. The former is
38 nitrate and phosphate co-limited while the latter is mostly nitrate-limited, which results in an
39 excess of phosphate (Yamamoto-Kawai et al., 2006). The role of nutrient limitation as a
40 possible regulator of methane production in surface water has recently been investigated (Karl
41 et al., 2008, Damm et al., 2010) while methane excess in ocean surface water relative to the
42 atmospheric equilibrium has been studied for more than three decades (Scranton and Brewer,
43 1977). Different nutrient limitations can stimulate the growth of specific members of the
44 bacterioplankton assemblage with consequences not only for the turnover of organic matter,
45 biogeochemical cycling of carbon but also for producing climate relevant traces gases
46 (Thingstad et al., 2008). Methanogenic archaea have been identified to have the ability to
47 metabolize dimethylsulfoniopropionate (DMSP) and its degradation products by producing
48 methane (Kiene et al., 1986; Oremland et al., 1989; van der Maarel and Hansen, 1997).
49 However also bacteria may be methylotroph, using a series of methylated compounds,
50 including methylated sulphur compounds such as (DMSP) and dimethylsulfide (DMS)

51 (Neufeld et al., 2008). This metabolism is referred as methylotrophic methanogenesis (Sowers
52 and Ferry, 1983). DMSP is produced by marine phytoplankton and when metabolized, is a
53 primary carbon source for heterotrophic bacteria (Kiene et al., 2000). DMSP is the precursor
54 of dimethylsulfide (DMS) or methanethiol. DMS partly escapes to the atmosphere where it is
55 the most important climate-cooling gas, counterbalancing the effect of greenhouse gases
56 (Charlson et al., 1987). Methanethiol is a key reactive intermediate utilized as sulphur and
57 carbon sources for biosynthesis or energy generation (Kiene et al., 2000). In anaerobic
58 environments methanethiol act also as precursor for methane production (Tallant and Kryzcki,
59 1997). A switch in the utilization of phosphate and DMSP degradation products in nitrate-
60 limited Pacific-derived water is also considered to produce methane in aerobic environments
61 (Damm et al., 2010). Methane excess in surface water has also been detected under multi-year
62 sea ice and in the marginal ice zone along the North-West Passage, i.e. the region from the
63 southwest edge of Greenland through the Baffin Bay to the Beaufort Sea (Kitidis et al., 2010).
64 Here we present data from Fram Strait where Atlantic water and Pacific-derived surface water
65 bodies occur adjacent to each other. We show that ongoing sea ice melting has amplified the
66 environmental differences between both water masses and we postulate that methane
67 production occurs during regenerated production in Pacific-derived water despite an apparent
68 oxygen excess. Methanogenesis in an aerobic environment is called the methane paradox as
69 this process requires strictly anaerobic conditions. However, methane concentrations above
70 the equilibrium concentration with the atmosphere are well known from the ventilated (i.e.
71 oxic) open ocean surface layer (Reeburgh, 2007). Hence we determine the maximum oxygen
72 concentration in seawater, which allows anaerobic processes to take place inside bacterial
73 cells. Since this aspect is fundamentally important we provide a detailed model description to
74 show and explain why and how it can potentially occur.

75 **2. Study area**

76

77 In the Fram Strait, the surface water (< 60 m) in general comprises two main water masses,
78 which flow in opposite directions (Rudels et al., 2000). The warm (up to 4°C) and saline (up
79 to 34.8) Atlantic water (AW) branch flows northward east of about 4°W (Fig.1). Further west,
80 colder and less saline polar surface water (PSW) occupies the upper water column. In PSW, a
81 portion is Pacific-derived water that varies inter-annually between more than 90% (Jones et
82 al., 2003) to almost zero (Falck et al. 2005). In 2008, this portion had attained just over 60%
83 (Dodd et al., 2012). The salinity of PSW was homogeneous at about 33 indicating unchanged
84 conditions since winter convection, except for some near-surface warming and freshening by
85 melt water (Fig. 2A). This distribution has been described previously for the end of the
86 summer season (Budeus et al., 1997).

87 The recurrent Northeast Water Polynya (NEWP) is localized in the region of the PSW
88 (Budeus and Schneider, 1995). Polynyas are less light-limited due to early opening of the ice
89 cover compared to adjacent regions, and primary production starts earlier in the year. In the
90 NEWP, nutrient-limited conditions occur at the end of July (Wallace et al., 1995, Kattner and
91 Budeus, 1997). In the summer of 2008, ice fields drifting from the north partly covered the
92 study area (Fig. 1). Hence stations in the middle of transect were located in partly ice-covered
93 AW and PSW and the more eastern and western stations in ice-free AW and PSW,
94 respectively.

95

96 **3. Sampling and methods**

97

98 In summer 2008, water sampling for measuring methane, oxygen, nutrients and DMSP was
99 carried out in Fram Strait during the cruise ARK-XXIII/2 with RV “Polarstern”, roughly
100 along the 79°N transect and spread on the Greenland sea shelf (Fig. 1). Further oceanographic
101 and biological data were taken in the surface water to 200 m depths. The main sampling sites
102 were along the hydrographic transect and in an opened ice lead on the Greenland shelf where

103 the sampling was repeated twice, first on July 23th (time 1) and one week later (time 2).
104 Salinity, temperature, light transmission and oxygen were measured with a Seabird SBE 911+
105 CTD and C-Star Wetlabs transmissiometer. Oxygen was measured with the SBE 43 dissolved
106 oxygen sensor SN 743 and sensor calibration was done on water samples using Winkler
107 titration.

108 Water samples for estimating the abundance of dominant phytoplankton species were
109 collected with a Niskin rosette sampling system and with an Apstein net (20 µm mesh size)
110 towed through the upper 10 m of the water column. Samples were preserved in hexamine-
111 buffered formalin (final concentration of ~1%) and dominant species or groups were counted
112 with an inverted microscope. Nutrient analyses were performed on board with a nutrient
113 analyzer (Evolution III, Alliance Instruments) according to standard methods. Methane
114 concentrations were analyzed within a few hours after sampling. The dissolved gas was
115 extracted from the water by vacuum-ultrasonic treatment and subsequently measured with a
116 gas chromatograph (Chrompack 9003 (GC) with a flame ionization detector (FID). For gas
117 chromatographic separation we used a packed column (Porapak Q 80/100 mesh). The GC
118 oven was operated isothermally (60°C) and the FID was held at 250°C. Two sets of standard
119 gas mixtures were used for calibration. The standard deviation of duplicate analyses was 5%.
120 This high overall error is almost exclusively due to the gas extraction procedure and not to
121 GC precision, which had an error of only 1%.

122 Total DMSP samples were collected directly from the Niskin sample bottles into 50 ml
123 centrifuge tubes, containing 167 µl of 50% H₂SO₄, and stored at 4°C for later analysis.
124 Dissolved DMSP samples were collected by the small volume drip filtration procedure
125 recommended by Kiene and Slezak (2006). Briefly, immediately after sampling on the rosette
126 about 50 ml of seawater was filled into a 47 mm filter tower with a Whatman GF/F glass fiber
127 filter. From the water dripping through the filter only the first 3.5 ml of filtrate were collected
128 directly into a storage tube containing 50 µl of 50% H₂SO₄. DMSP is stable for months in

129 acidic solution (Curran et al., 1999). In the home lab DMSP was analyzed as DMS after
130 alkaline cleavage. A subsample of the solution was pipetted into a 14 ml serum vial, treated
131 with 1 ml of 5 N NaOH and quickly sealed. The released DMS was purged into a cryotrap and
132 quantified with a gas chromatograph equipped with a Chromosil 330 column and a pulsed
133 flame photometric detector (PFPD). Helium was used as purge gas and carrier gas.

134

135 **4. Results and Discussion**

136 **4.1 Nutrient limitation and biological production**

137 In AW, nitrate and phosphate were abundant in the ice covered regions but depleted in open
138 waters, without changes in the Redfield ratio. In both ice-covered and ice-free PSW, nitrate
139 was undetectable in the near-surface layer (<20 m) and became limiting before phosphate
140 exhaustion (Fig. 2B and C). Hence, PSW was characterized by nitrate to phosphate ratios
141 lower than the Redfield ratio as also reported by Yamamoto-Kawai et al. (2006).

142 In addition to distinct nitrate availabilities, variations in oxygen saturation and light
143 transmissions along the E-W transect are obviously and point to bloom conditions which are
144 partly influenced by melting sea ice. As consequence, different blooms stages in ice free and
145 ice covered AW and PSW, respectively were eventually created (Fig. 2).

146 In the ice-free AW, the light transmission was reduced down to a depth of 60 m. This feature
147 is in accordance with a typical late bloom population which was observed in the non-stratified
148 water column east of 1°E (Fig. 2B and D). Besides the dominating prymnesiophyte
149 *Phaeocystis pouchetii*, many heterotrophic unicellular species were found belonging to
150 dinoflagellates and ciliates. Diatoms comprised a few *Thalassiosira* spp and very few
151 pennates. The occurrence of the two coccolithophores *Emiliana huxleyi* and *Coccolithus*
152 *pelagicus* was indicative of the minor ice influence. In comparison in the ice-free PSW an
153 impoverished phytoplankton community were found caused by the nitrate limitation and
154 reflected by the high light transmission (Fig. 2D, west of 10°W).

155 In the ice-covered regions both water masses clearly show a reduced light transmission up to
156 20 m depth. Sinking particles and ice algae released during brine drainage and ice melt may
157 create this effect (Fig. 2B, Mundy et al., 2005). Under melting ice, chlorophyll concentrations
158 are comparable in both water masses, ranging from nearly 0 to 2.6µg/L. Certainly the ice-
159 covered AW was dominated by large *Phaeocystis pouchetii* colonies, which were partly
160 covered with tiny pennate diatoms, whereas in the ice-covered PSW cold water ice-related
161 algal communities were observed. In addition the different levels of oxygen saturation despite
162 low solubility differences (by about 1.4%) were detected along the 79° transect which refers
163 to deviations in the steady state between production and respiration in both ice-covered water
164 masses (Fig. 2). Detected along the transect (Fig. 2) this observation is corroborated by the
165 relation between chlorophyll and oxygen in a new opened lead in the NEWP region a region
166 with long time ice-covered PSW (Figs. 1 and 3).

167

168 **4.2 Sea ice melting in nitrate limited sea water - Biogeochemical consequences**

169

170 The environmental differences between AW and PSW obviously create different under ice
171 bloom conditions. While in the largely ice-covered AW new production occurred and
172 favoured as nutrients are replete, a shift from new to regenerated production was evident in
173 the nitrate-limited PSW (Fig. 2). The PSW on the East Greenland shelf is generally
174 characterized by low initial nitrate concentrations (Kattner and Budéus, 1997). In summer, the
175 surface waters are widely nitrate exhausted, and ammonium uptake becomes more important
176 (Smith et al. 1997). The pronounced oxygen enhancement in the PSW combined with a highly
177 variable nitrate to phosphate ratio revealed the importance of both new and regenerated
178 production, probably dependent on the ice cover (Fig. 2E and 4). Thus, in PSW, where an
179 excess of phosphate is available, ammonium could be an alternative nitrogen source to sustain

180 primary production. In the Fram Strait region, ammonium, released from multi-year Arctic
181 sea ice, may additionally alleviate the nitrate limitation (Tovar-Sanchez et al., 2010).
182 However, during regenerated production the ability of bacteria to compete with the
183 phytoplankton community for inorganic nutrients and organic material is enhanced (Thingstad
184 et al., 2008). Hence, melting sea ice in PSW may also affect the microbial food web. An
185 important energy, carbon and sulphur source for bacterial biomass production is DMSP
186 (Kiene et al., 2000). Both water masses differed clearly with regard to their DMSP
187 concentrations. The high concentrations of DMSP in the ice-free AW were probably due to
188 DMSP release by senescing *P. pouchetii* cells, which are known to be a major producer of
189 DMSP in polar waters (Matrai and Vernet, 1997) (Fig. 2F). In comparison, low DMSP
190 concentrations in the ice-free PSW may be due to an impoverished bloom of almost non
191 DMSP producing diatoms, but perhaps also to an enhanced bacterial utilization of DMSP
192 (Fig. 2F). The correlation between DMSP and oxygen saturation suggest a coupling of DMSP
193 with the ongoing biological production in the AW ($R^2=0.600$; $p<0.001$). This correlation is
194 however not found in the PSW (Fig. 5). The enhanced DMSP concentration in PSW is
195 restricted to the upper 20 m and therefore likely induced by the DMSP release from melting
196 sea ice (Fig. 2F, 5). The production of substantial amounts of DMSP by ice algae suggested
197 by Levasseur et al. (1994) and Uzuka, (2003) corroborates this assumption. Furthermore
198 DMSP released from sea ice is reported to be partially responsible for elevated DMSP
199 concentrations in the water column at the ice edge (Trevena and Jones, 2006, Tison et al,
200 2010). A rapid microbial consumption of sea ice released DMSP (Galindo et al., 2014)
201 suggest that sea ice released DMSP may serve as an additional carbon source for the
202 microbial food web while finally the nutrient status in the water column impacts the pathway
203 of its bacterial consumption.

204

205 **4.2.1 Methane excess - a response to special environmental features?**

206

207 The methane inventories were also clearly different in both water masses. In AW, methane
208 concentrations tended to be in equilibrium or slightly under-saturated in relation to the
209 atmospheric partial pressure (3 to 3.5 nM, depending on temperature and salinity). In the ice-
210 free PSW a slight oversaturation was found, potentially generated by methane release from
211 the seafloor in the NEWP region on the shallow shelf with water depths of about 100 m. In
212 shallow polynya regions enhanced turbulence during convective mixing enhances sediment
213 resuspension and eventually methane release from the seafloor (Damm et. al., 2007).

214 In the ice-covered PSW, however, a near-surface methane excess clearly rose above the slight
215 oversaturation detected in ice-free PSW. It is striking that this methane surplus was found in
216 the region where regenerated production occurred and where nitrate was clearly depleted (Fig.
217 2). This pattern is similar to that in the central Arctic Ocean where a change in the utilization
218 of phosphate and methylated compounds is found to trigger the switch from no methane
219 production to methane production in Atlantic and Pacific surface water, i.e. in
220 nitrate/phosphate co-limited and nitrate-limited water (Damm et al., 2010).

221 Reduced turbulence in the presence of sea ice restricts the gas transfer (Rutgers van der Loeff
222 et al., 2014). Hence the partially ice-covered water tends to reduce the escape of produced
223 methane. Furthermore melting sea ice enhances the water stratification (Rabe et al., 2014).
224 Indeed the PSW has a strong stratification in the upper water column induced by the long
225 journey below melting sea ice (Fig 2A). Hence on one side, methane efflux and on the other
226 side downward mixing is hampered in PSW. Both conditions finally induce that the methane
227 excess created in sea ice-influenced water remains preserved during calm weather conditions
228 in summer. Conspicuously is that the methane excess detected under multi-year-sea-ice and in
229 the marginal ice zone along the North-West Passage (Kiditis et al., 2010) also occurs in a
230 region which receives Pacific-derived water after its journey through the Arctic Ocean (Jones
231 et al., 2003).

232 We therefore conclude that the development of a hotspot of methane production creates the
233 excess as a rapid response during regenerated production when melting sea ice supplies
234 DMSP, which may act as a potential precursor for methane formation. The microbial
235 degradation of DMSP to methane was observed in a microcosm experiment carried out with
236 seawater from the Fram Strait during this cruise (Damm et al., 2010). During the experiment
237 *Archaea* abundance remained negligible and bacteria of the clades *Rhodobacter/Roseobacter*
238 were dominant which are frequent in oligotrophic ocean surface waters and known for their
239 highly diverse and flexible metabolism. A survey of available *Roseobacter* genomes by
240 Moran et al. (2007) revealed that 50% of the genomes contained genes for DMSP
241 demethylation.

242

243 **4.3 Methane production in waters with oxygen excess - a paradox?**

244 It is conspicuous that high oxygen concentrations in surface waters did not hamper methane
245 production (Fig. 2E and G). As methanogenic activity is not favoured in an aerobic
246 environment it was assumed that this process occurs in microenvironments which are
247 sufficiently lacking in molecular oxygen (Cynar and Yayanos, 1992). The limiting conditions
248 for the maintenance of a reduced micro-niche within oxidized marine sediments were first
249 discussed by Jørgensen (1977). The question arises whether reducing conditions can exist, for
250 example, within a *Roseobacter* cell, which would allow anaerobic processes inside the
251 bacterial cell. To answer this question, we extended the model of Jørgensen (1977) by an
252 additional compartment for the cell membrane and calculated the oxygen concentration
253 profile in the interior of the cell as a function of the cell properties (cell size, rates of
254 respiratory metabolism, membrane permeability for O₂) and the external O₂ concentration.

255 We describe the bacterial cell in terms of a sphere with a radius b covered by a thin
256 membrane. The membrane is described by a homogeneous spherical shell of outer and inner
257 radii a and b (Fig. 6A). Within the interior of the sphere there is an O₂ consumption of

258 constant intensity ρ . To determine the stationary concentration profile in the interior of the
 259 sphere we have to find the solution in the region $0 \leq r \leq b$ of

$$260 \quad D \frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{dC(r)}{dr} \right) - \rho(r) = 0, \quad (1)$$

261 where D is the diffusivity of O_2 and $C(r)$ is the concentration of O_2 as a function of the radial
 262 distance r from the centre of the cell. Assuming a free floating cell, the diffusion coefficient
 263 (D) in the surrounding water is a constant and is given by the value in bulk seawater with a
 264 salinity of 33 at 0°C . Inside the cell, the salinity is probably slightly lower than in the
 265 surrounding seawater since a portion of the osmotic pressure in the cell is established by
 266 means of organic osmolytes. However, at 0°C a salinity change from 35 to about 30 (cellular
 267 interior) only has a minor impact on the diffusion coefficient for oxygen ($S=30-35$: $D =$
 268 $1.0580-1.0503 \times 10^{-5} \text{ cm}^2 \text{ s}^{-1}$, Ramsing and Gundersen, 1994). Our assumption therefore is that
 269 the diffusion coefficient of seawater ($D = D_w$) is the same for both the water in the cell and the
 270 surrounding water. Integrating and solving the equation for $\rho(r) = \rho$ and finite concentrations
 271 at $r = 0$ yields

$$272 \quad C(r) = C_b - \frac{\rho}{6D_w} (b^2 - r^2), \quad (2)$$

273 where $C_b = C(r = b)$ is the concentration in the sphere at the inner side of the membrane and
 274 D_w is the diffusion coefficient in water. To determine C_b we consider a stationary diffusion
 275 through the membrane of the thickness $h = a - b$ and with a permeability P for O_2 . It is
 276 assumed that there is no O_2 consumption in the membrane region. Hence in the region $b \leq r \leq$
 277 a , equation (1) is integrated for $\rho(r) = 0$. With the total flux of O_2 through the membrane F
 278 (units: $\text{mol } O_2 \text{ s}^{-1}$), the integration of equation (1) gives the following expression for C_b as a
 279 function of the concentration at the cell surface, $C_a = C(r = a)$,

$$280 \quad C_b = C_a - \frac{F}{4\pi D_m} \frac{a-b}{ab}, \quad (3)$$

281 where D_m is the diffusion coefficient in the membrane and F equals the total O_2 consumption
 282 in the region $0 \leq r \leq b$, which is given by the respiration rate per cell, i.e. $F = \frac{4}{3}\pi b^3 \rho$. We
 283 consider the situation where the thickness of the membrane $h = a - b$ is small in proportion to
 284 the radius b . In this case $ab \approx b^2$, and equation (3) takes the form

$$285 \quad C_b = C_a - \frac{F}{4\pi b^2} \frac{h}{D_m} = C_a - \frac{F}{4\pi b^2} \frac{1}{P} = C_a - \frac{\rho b}{3P}. \quad (4)$$

286 To determine C_a we have to find the stationary concentration profile in the cell environment
 287 by solving equation (1) for $\rho(r) = 0$ in the region $r \geq a$. With $C(r \rightarrow \infty) = C_0$, integration of
 288 equation (1) yields

$$289 \quad C(r) = C_0 - \frac{F}{4\pi D_w} \frac{1}{r} = C_0 - \frac{\rho}{3D_w} \frac{b^3}{r}, \quad (5)$$

290 where C_a follows for $r = a \approx b$. Replacing this value for C_a in equation (4) and the result for
 291 C_b in equation (2) gives

$$292 \quad C(r) = C_0 - \frac{\rho b^2}{3D_w} - \frac{\rho b}{3P} - \frac{\rho}{6D_w} (b^2 - r^2), \quad (6)$$

293 which is the equation for the concentration profile in the interior of the cell. The strongest
 294 decline of $C(r)$ occurs across the membrane of permeability P (Fig. 6B). Inside the cell the
 295 diffusion of O_2 and respiration occur on two distinct time scales. Diffusion on the micrometer
 296 scale is, as a rule, much faster compared to slow metabolic processes. This means that within
 297 the cell significant concentration gradients of O_2 do not appear. According to equation (6) the
 298 concentration profile is described by a very flat parabola with the minimum of O_2
 299 concentration at $r = 0$.

300 When $C(r)$ is zero for $r = 0$ we obtain the following equation for the maximum concentration
 301 in the environment, $C_{0,\max}$, which allows anaerobic processes to take place inside the cell:

$$302 \quad C_{0,\max} = \frac{\rho b^2}{2D_w} + \frac{\rho b}{3P}. \quad (7)$$

303 For a salinity of 33 at 0°C , the diffusion coefficient of O_2 in water is $D_w = 1.05 \times 10^{-9} \text{ m}^2 \text{ s}^{-1}$
 304 (Ramsing and Gundersen, 1994). The respiration per cell, $0.61 \text{ fmol } O_2 \text{ h}^{-1}$, was measured in
 305 laboratory experiments with *Roseobacter* cultures grown in a chemostat at 22°C (Koblížek et

306 al., 2010). The simplest correction for the temperature of the bacterial respiration is the
 307 temperature coefficient, Q_{10} , the factor by which a biological reaction changes with a
 308 temperature increase of 10°C. The model of Rivkin and Legendre (2001) predicts a Q_{10} of
 309 1.85 for bacterial respiration (Vázquez-Domínguez et al., 2007). The respiration rate
 310 measured at 22°C is thus reduced by a factor of $1/Q_{10}^{2.2} = 0.26$ for an environmental
 311 temperature of ~ 0°C. In the following a respiration rate per cell of $F = 0.16$ fmol O₂ h⁻¹ at
 312 0°C is therefore assumed. Using the cell volume of 0.53 μm³ obtained in the laboratory
 313 experiments with *Roseobacter* (Koblížek et al., 2010), it follows the radius of the sphere, $b =$
 314 0.5 μm, and the constant intensity of O₂ consumption in the interior of the sphere, $\rho = 0.084$
 315 mol m⁻³ s⁻¹. The permeability for gases of bacteria and microalgae has been determined in
 316 very few investigations. The membrane permeability for O₂ (P) follows from the permeability
 317 for CO₂ (P_{CO_2}) by the relationship (Spalding and Portis, 1985)

$$318 \quad P = P_{\text{CO}_2} \sqrt{\frac{\text{molecular weight of CO}_2}{\text{molecular weight of O}_2}}. \quad (8)$$

319 The inverse proportionality between P and the square root of molecular mass is assumed to
 320 represent a useful approximation for gases that permeate the membrane via (passive)
 321 diffusion. Using $P_{\text{CO}_2} = 3 \times 10^{-8}$ m s⁻¹, as measured for *Synechococcus* UTEX 625 (Salon et al.,
 322 1996), one obtains $P = 3.5 \times 10^{-8}$ m s⁻¹. From equation (7) it now follows that the maximum O₂
 323 concentration in the environment which allows anaerobic processes to take place inside the
 324 bacterial cell, is $C_{0,\text{max}} = 400$ μM. The latter corresponds to the O₂ concentrations observed in
 325 nitrate-limited PSW in the region where the highest methane concentrations were observed
 326 (Fig. 2E and G). Hence, in the PSW we had the situation where oxygen decreased almost to
 327 zero in the interior of the bacterial cell, but was present in the membrane region and outside
 328 the cell. The above calculations demonstrate that an oxygen excess in the surrounding
 329 medium does not exclude the establishment of anaerobic conditions within the bacterial cell.

330 Altogether our model results suggest that oxygen excess and methane production are not
331 mutually exclusive.

332 **5. Summary and conclusions**

333 A methane hotspot was detected in surface water of the western Fram Strait during summer
334 2008. We show that this methane excess is formed exclusively in Pacific-derived surface
335 water (PSW) where sea-ice melting occurred. It is not found in PSW without ice coverage
336 further west nor in Atlantic water (AW) further east. A conspicuous difference between both
337 water masses is the availability of nitrate which was clearly depleted in PSW. We show that
338 the methane excess is confined to a region where sea-ice is melting and postulate that DMSP
339 released from sea-ice act as the precursor of methane produced via methylotrophic
340 methanogenesis. Water stratification and reduced turbulence hampers the methane efflux
341 which finally induces the hot-spot in surface water. We prove by modelling that anaerobic
342 methanogenesis occurs inside a bacterial cell, despite high oxygen saturation levels in its
343 surrounding. These results support the observation that methane excess in stratified aerobic
344 seawater is coupled to an oligotrophic environment previously found in Pacific-derived water
345 in the central Arctic Ocean while a potential coupling of methanogenesis with DMSP
346 degradation processes requires further elucidation, especially the relationship between DMSP
347 turnover rates and in situ production of methane.

348 **Acknowledgements**

349 We are grateful to K.-U. Ludwichowski and M. Graeve for nutrient measurements and E.
350 Lichte for DMSP measurements on board . We thank the scientific party and crew of RV
351 Polarstern for their professional support at sea.

353

354 **References**

355 Arrigo, K.A., Dijken, G.v. and Pabi, S., 2008. Impact of a shrinking Arctic ice cover on
356 marine primary production. *Geophys. Res. Lett.*, doi 10.1029/2008GL035028.

357 Budeus, G. and Schneider, W., 1995. On the hydrography of the Northeast Water Polynya. *J.*
358 *Geophys. Res.* 100, 4287-4299.

359 Budeus, G., Schneider, W. and Kattner, G., 1997. Distribution and exchange of water masses
360 in the Northeast water Polynya (Greenland Sea). *J. Mar. Syst.* 10, 123-138.

361 Charlson, R.J., Lovelock, J.E., Andreae, M.O., and Warren, S.G., 1987. Oceanic
362 phytoplankton, atmospheric sulphure, cloud albedo and climate. *Nature.* 326, 655–661.

363 Curran, M.A.J., Jones, G.B. and Burton, H., 1999. Spatial distribution of dimethylsulfide and
364 dimethylsulfoniopropionate in the Australian sector of the Southern Ocean. *J. Geophys.*
365 *Res.* 103, 16667-16689.

366 Cynar, F.J. and Yayanos, A.A., 1992. The distribution of methane in the upper waters of the
367 southern California Bight. *J. Geophys. Res.* 97 (C7), 11.269–11.285.

368 Damm, E., Schauer, U., Rudels, B. and Hass, C., 2007. Excess of bottom-released methane in
369 an Arctic shelf sea polynya in winter. *Cont. Shelf Res.* doi:10.1016/j.-csr.2007.02.003.

370 Damm, E., Helmke, E., Thoms, S., Schauer, U., Nöthig, E., Bakker, K. and Kiene, R., 2010.
371 Methane production in aerobic oligotrophic surface water in the central Arctic Ocean.
372 *Biogeoscience.* 7, 1099-1108.

373 Dodd, P.A., Rabe, B., Hansen, E., Mackensen, A., Rohling, E., Falck, E., Reigstad, M., Jones,
374 P., Schauer, U. and Stedmon, C., 2012. The freshwater composition of the Fram Strait
375 outflow derived from a decade of tracer measurements. *J. Geophys. Res.*, 117. Doi
376 10.1029/2012JC008011.

377 Falck, E., Kattner, G., and Budeus, G., 2005. Disappearance of Pacific Water in the
378 northwestern Fram Strait. *Geophys. Res. Lett.* doi:10.1029/2005GL023400.

379 Galindo, V. Levasseur, M. Mundy, C.J., Gosselin, M. Tremblay, J.-E., Scarratt, M. Gratton,
380 Y., Papakiriakou, T., Poullin, M. and Lozotte, M., 2014. Biological and physical processes

381 influencing sea ice, under-ice algae, and dimethylsulfoniopropionate during spring in the
382 Canadian Arctic Archipelago. *J. Geophys. Res. Oceans*, 119, doi:10.1002/2013JC009497.

383 Jones, E.P., Swift, J.H., Anderson, L.G., Lipizer, M., Civitarese, G., Falkner, K.K., Kattner,
384 G. and McLaughlin, F.A., 2003. Tracing Pacific Water in the North Atlantic Ocean. *J.*
385 *Geophys. Res.* 108(C4), 3116, doi:10.1029/2001JC001141.

386 Jørgensen, B.B., 1977. Bacterial sulphate reduction within reduced microniches of oxidized
387 marine sediments. *Marine Biology*. 41, 7-17.

388 Karl, D.M., Beversdorf, L., Björkman, K., Church, M.J., Martinez, A. and DeLong, E.F.,
389 2008. Aerobic production of methane in the sea, *Nature Geosciences*. doi: 10.1038/ngeo234.

390 Kattner, G. and Budeus, G., 1997. Nutrient status of the Northeast Water Polynya. *J. Mar.*
391 *Syst.* 10, 185-197.

392 Kiene, R.P., Oremland, R.S., Catena, A., Miller, L.G., Capone, D.G., 1986. Metabolism of
393 reduced methylated sulfur-compounds in anaerobic sediments and by a pure culture of an
394 estuarine methanogen. *Applied and Environmental Microbiology* 52, 1037–1045.

395 Kiene, R.P., Linn, L.J. and Burton, J.A., 2000. New and important roles for DMSP in marine
396 microbial communities. *J. Sea Res.* 43, 209-224.

397 Kiene, R.P. and Slezak, D., 2006. Low dissolved DMSP concentrations in seawater revealed
398 by small-volume gravity filtration and dialysis sampling. *Limnology and Oceanography* 4,
399 80-95.

400 Kitidis, V., Upstill-Goddard, R.C. and Anderson, L.G., 2010. Methane and nitrous oxide in
401 surface water along the North-West Passage. *Mar. Chem.* 121, 80-86.

402 Koblížek, M., Mlčoušková, J., Kolber, Z., and Kopecký, J., 2010. On the photosynthetic
403 properties of marine bacterium COL2P belonging to *Roseobacter* clade. *Arch Microbiol.*
404 192, 41-49.

405 Levasseur, M., Gosselin, M. and Michaud, S., 1994. A new source of dimethylsulfide (DMS)
406 for the Arctic atmosphere: ice diatoms. *Mar. Biol.* 121, 181-187.

407 Matrai, P.A. and Vernet, M., 1997. Dynamics of the vernal bloom in the marginal ice zone of
408 the Barents Sea: dimethylsulfide and dimethylsulfoniopropionate budgets. *J. Geophys. Res.*
409 102, C10, 22965-22979.

410 Moran, M.A., Belas, R., Schell, M.A., Gonzalez, J.M., Sun, F., Sun, S., Binder, B.J.,
411 Edmonds, B., Ye, J.W., Orcutt, B., Howard, E.C., Meile, C., Palefsky, W., Goesmann, A.,
412 Ren, Q., Paulsen, I., Ulrich, L.E., Thompson, L.S., Saunders, E. and Buchan, A., 2007.
413 Ecological genomics of marine Roseobacters. *Appl. Environ. Microbiol.* 73, 4559–4569.

414 Mundy, C.J., Barber, D.G., and Michel, C., 2005. Variability of snow and ice thermal,
415 physical and optical properties pertinent to sea ice algae biomass during spring. *J. Mar.*
416 *Syst.* 58, 107-120.

417 Neufeld, J.D., Boden, R., Moussard, H., Schäfer, H., Murrell, J.C., 2008. Substratespecific
418 clades of active marine methylotrophs associated with a phytoplankton bloom in a temperate
419 coastal environment. *Applied and Environmental Microbiology* 74, 7321–7328.

420 Oremlad, R.S., 1979. Methanogenic activity in plankton sample and fish intestines: a
421 mechanism for in situ methanogenesis in ocean surface water. *Limnology and Oceanography*
422 24, 1136–1141.

423 Ramsing, N. and Gundersen, J., 2001. Tabulated physical parameters of interest to people
424 working with microsensors in marine systems. Techn. Rep. MPI Mar. Microbiology.
425 Bremen, 1994.

426 Reeburgh, W. S.: Oceanic methane biogeochemistry, *Chem. Rev.*, 107, 486–513,
427 doi:10.1021/cr050362v, 2007.

428 Rivkin, R.B. and Legendre, L., 2001. Biogenic carbon cycling in the upper ocean. Effects of
429 microbial respiration. *Science.* 291, 2398-2400.

430 Rudels, B., Meyer, R., Fahrbach, E., Ivanov, V., Osterhus, S., Quadfasel, D., Schauer, U.,
431 Tverberg, V. and Woodgate, R., 2000. Water mass distribution in Fram Strait and over the
432 Yermak Plateau in summer 1997. *Annales Geophysicales.* 18, 687-705.

433 Rutgers van der Loeff, M., Cassar, N., Nicolaus, M., Rabe, B., Stimac, I., 2014. The influence
434 of sea-ice cover on air-sea gas exchange estimated with radon-222 profiles. *J.Geophys.*
435 *Res.*, C – Oceans, doi: 10/1002/2013JC009321.

436 Sakshaug, E., 2003. Primary and secondary production in the Arctic Seas, in: Stein, R., and
437 McDonald, R.W. (Eds.), *The Organic Carbon Cycle in the Arctic Ocean*. pp. 57-81,
438 Springer, Berlin.

439 Salon, C., Mir, N.A. and Canvin, D.T., 1996. Influx and efflux of inorganic carbon in
440 *Synechococcus* UTEX 625. *Plant Cell Environ*, 19, 247-259.

441 Scranton, M.I. and Brewer, P.G., 1977. Occurrence of methane in the near surface waters of
442 the western subtropical North-Atlantic. *Deep-Sea Res.*, 24, 127–138.

443 Smith, Jr. W.O., Gosselin, M., Legendre, L., Wallace, D., Daly, K., Kattner, G., 1997. New
444 production in the Northeast Water Polynya 1993. *J. Mar. Syst.* 10, 199-209.

445 Sowers, K.R., Ferry, J.G., 1983. Isolation and characterization of a methylotrophic marine
446 methanogen, *Methanococcoides methylutens*, gen. nov., sp. nov. *Applied and Environmental*
447 *Microbiology* 45, 684–690.

448 Spalding, M.H. and Portis, A.R., 1985. A model of carbon dioxide assimilation in
449 *Chlamydomonas reinhardtii*. *Palta*. 164, 308-320.

450 Spreen, G., Kaleschke, L., and Heygster, G., 2008. Sea ice remote sensing using AMSR-E 89
451 GHz channels. *J. Geophys. Res.* doi:10.1029/2005JC003384.

452 Tallant, T.C. and Krycki, J.A., 1997. Methylthiol: coenzyme M methyltransferase from
453 *Methanosarcina barkeri*, an enzyme of methanogenesis from dimethylsulfide and
454 methylmercatopropionate. *J. Bacteriol.* 179, 6902–6911.

455 Thingstad, T.F., Bellerby, R.G.J., Bratbak, G., Borsheim, K.Y. Egge, J.K., Heldal, M.,
456 Larsen, A., Neill, C., Nejstgaard, J., Norland, S., Sandaa, R.A., Skoldal, E.F., Tanaka, T.,
457 Thyraug, R. and Töpper, B., 2008. Counterintuitive carbon-to-nutrient coupling in an Arctic
458 pelagic ecosystem. *Nature*. doi:10.1038/nature07235.

459 Tison, J.-L., Brabant, F., Dumont, I. and Stefels, J. 2010., High-resolution dimethyl sulphide
460 and dimethylsulfoniopropionate time series profiles in decaying summer first-year sea ice at
461 Ice Station Polarstern, western Weddell Sea, Antarctica. *J. Geophys. Res.*, 115, G04044,
462 doi10.1029/2010JG001427.

463 Tovar-Sanchez, A., Duarte, C.M., Alonso, J.C., Lacorte, S., Tauler, R. and Galban-Malagon,
464 C., 2010. Impacts of metals and nutrients released from melting multiyear Arctic sea ice. *J.*
465 *Geophys. Res.* 115, C07003, doi:10.1029/2009JC005685.

466 Trevena, A.J., and Jones, G.B. 2006. Dimethylsulphide and Dimethylsulphoniopropionate in
467 Antarctic sea ice and their release during sea ice melting, *Mar. Chem.* 98, 210-222,
468 doi10.1016/j.marchem.2005.09.005.

469 Uzuka, N., 2003. A time series observation of DMSP production in the fast ice zone near
470 Barrow (extended abstract). *Tohoku Geophys. Journ. (Sci. Rep. Tohoku Univ., Ser. 5)*, 36,
471 4, 439-442.

472 Vázquez-Domínguez, E., Vaqué, D. and Gasol, J.M., 2007. Ocean warming enhances
473 respiration and carbon demand of coastal microbial plankton. *Global Change Biology.* 13,
474 1327–1334, doi: 10.1111/j.1365-2486.2007.01377.

475 Wallace D.W.R., Minnett, P.J. and Hopkins, T.S., 1995. Nutrients, oxygen and inferred new
476 production in the Northeast Water Polynya. *J. Geophys. Res.* 100, C3, 4323-4340.

477 Yamamoto-Kawai, M., Carmack, E.C. and McLaughlin, F.A., 2006. Nitrogen balance and
478 Arctic through flow. *Nature*, 443, 43.

479

480 **Captions**

481 Fig. 1: Map of the Fram Strait with ice coverage (AMSR-E data see Spreen et al., 2008). Ice
482 coverage is shown by colors from red to blue, (100%, 0%) meaning a closed ice cover and
483 open water, respectively. Black circles indicate stations localized in AW (Atlantic water) in
484 PSW (Polar surface water) and NEWP (Northeast Water Polynya)

485 Fig. 2: Profiles (black dots) along the transect from 15°W to 4°E (ice coverage ranges from
486 7°W to 0°); the transect crosses Atlantic water (east of 4.5°W) and Polar surface water (west
487 of 4.5°W) diagrams show: potential density in sigma θ units (A), light transmission (%), (B),
488 concentrations of nitrate, oxygen and phosphate ($\mu\text{mol/l}$) (C, D, E), concentrations of DMSP
489 and methane (nM), (F and G).

490 Fig. 3: Oxygen saturation vs. chlorophyll a in Atlantic water (filled squares) oxygen is almost
491 under saturated (up to 15%) and in Polar surface water (open squares) clearly oversaturated
492 (by up to 11%) in relation to the atmospheric partial pressure. Dots, grey and black are from
493 the new opened lead at time 1 and at time 2 (one week later), respectively. In AW and in the
494 new opened lead clear different relationships are apparent while in PSW the data scatter
495 between both ratios.

496 Fig.:4 Nitrate/phosphate ratios vs. oxygen concentration in Atlantic water (red squares/green
497 triangles for ice-free and ice-covered stations) and Polar surface water (black/grey dots for
498 ice-covered and ice-free stations). In the former, Redfield ratios are almost retained, while in
499 the latter, nitrate limitation has induced increasing deviations from Redfield.

500 Fig. 5: Oxygen saturation vs. DMSP in Atlantic water (filled squares), in PSW (open squares)
501 and in the new opened lead at time 1 and at time 2 (one week later), respectively. In PSW
502 DMSP concentration higher than 20 nM are available in surface water (<20 m) and likely
503 released from melting sea ice.

504 Fig. 6: (A) Model of oxygen distribution in a bacterial cell surrounded by oxic seawater; for
505 explanation of symbols consult text.

506 (B) The oxygen concentration in the interior of the cell ($C(r)$ of equation (6)) and in the cell
507 membrane, where the concentration is described by the equation $C(r) = C_0 - \rho b^2 / (3D_w) -$
508 $\rho b^2 / (3hP)(a/r - 1)$. Shown is the crucial role of the low membrane permeability (P) for the

509 maintenance of anaerobic conditions inside the cell. The dashed lines indicate oxygen
510 concentration profiles for the n-fold increase ($n=2, 4, 8, 16$) in membrane permeability.
511