

## Cytotoxicity Activities in Local *Justicia gendarussa* Crude Extracts against Human Cancer Cell Lines

Zahidah Ayob<sup>a</sup>, Azman Abd Samad<sup>a</sup>, Siti Pauliena Mohd Bohari<sup>a\*</sup>

<sup>a</sup>Department of Biotechnology and Medical Engineering, Faculty of Biosciences and Medical Engineering, Universiti Teknologi Malaysia, 81310 UTM Johor Bahru, Johor, Malaysia

\*Corresponding author: pauliena@utm.my

### Article history

Received :29 July 2013  
Received in revised form :  
23 September 2013  
Accepted :29 September 2013

### Graphical abstract



### Abstract

The present study was designed to evaluate the cytotoxic activity of *Justicia gendarussa* methanolic leaf extracts from five different locations in Johor and also two standards flavonoids (naringenin and kaempferol) as these two compounds were detected in the chromatogram. In this study, leaf extracts and compounds were tested against various human cancer cell lines (HT-29, HeLa and BxPC-3) by using MTT assay. The results showed that methanolic leaf extracts from Mersing and kaempferol were very toxic against BxPC-3 and HeLa cells with IC<sub>50</sub> values of 16 µg/ml and 5 µg/ml, respectively. It is suggested that *J. gendarussa* leaf extracts from Mersing have potential cytotoxic activity on human cancer cell lines particularly BxPC-3 cells.

**Keywords:** Cytotoxicity; *Justicia gendarussa*; MTT assay; HT-29; HeLa; BxPC-3

### Abstrak

Kajian ini bertujuan menilai aktiviti sitotoksik ekstrak daun *Justicia gendarussa* menggunakan metanol daripada lima lokasi berbeza di Johor dan dua standard sebatian flavonoid (naringenin dan kaempferol) yang mana kompaun sebatian ini yang dikesan pada kromatogram. Dalam kajian ini, ekstrak daun dan sebatian telah diuji ke atas berbagai sel kanser manusia (HT-29, HeLa dan BxPC-3) dengan menggunakan ujian MTT. Hasil kajian menunjukkan bahawa ekstrak daun dari Mersing dan kaempferol sangat toksik kepada BxPC-3 dan HeLa dengan nilai IC<sub>50</sub> masing-masing adalah 16 µg/ml dan 5 µg/ml. Penemuan ini menunjuk ekstrak daun *J. gendarussa* dari Mersing berpotensi untuk digunakan dalam rawatan sel kanser manusia khususnya sel BxPC-3.

**Kata kunci:** Sitotoksik; *Justicia gendarussa*; ujian MTT; HT-29; HeLa; BxPC-3

© 2013 Penerbit UTM Press. All rights reserved.

### 1.0 INTRODUCTION

Cancer is the fourth leading death in Malaysia after diseases of the circulatory, respiratory systems and motor vehicle accidents that responsible for 10.3% of medically certified deaths.<sup>1</sup> Cancer causes significant morbidity, mortality and also the major health problem worldwide. The incidence of colon cancer, cervix cancer and pancreas cancer are rising in every country in the world. Although, many clinical anti-cancer drugs today are effective in killing cancer cells, the price still expensive and not affordable for some people. Furthermore, there are also undesirable side effects like excessive hair lose, loss of appetite and nausea.

Medicinal plants have provided a wide variety of phytochemical uses which have been shown to have activity across multiple signalling pathways. Medicinal plants not only disrupt aberrant signalling pathways leading to cancer (proliferation, deregulation of apoptosis, angiogenesis, invasion

and metastasis), but also synergize with chemotherapy and radiotherapy. It is not surprising many researchers are constantly turning to natural products for alternative source of medicines.<sup>1</sup>

*Justicia gendarussa* is also commonly known as Gendarussa is a member of Acanthaceae. Traditionally, *J. gendarussa* extracts have been used to treat ailments such as emetic, antipyretic, amenorrhea, stomach troubles, hemoptysis, cough and asthma.<sup>2-3</sup> Recently, *J. gendarussa* leaves and stem extracts were reported to have antiangiogenic,<sup>4</sup> hepatoprotective activity,<sup>5</sup> anti-fungal,<sup>6</sup> antioxidant,<sup>7-8</sup> anthelmintic,<sup>9</sup> anti microbial,<sup>10</sup> anti-bacterial<sup>11-12</sup> activities. Also, *J. gendarussa* leaf extracts have been used traditionally as a male contraceptive agent by several ethnic groups in central part of Papua, Indonesia and this extracts could inhibit mice spermatozoa penetrate onto mice ovum.<sup>13</sup>

Phytochemical study on the leaves of *J. gendarussa* revealed the alkaloids, amino acids, aromatics amines,<sup>14</sup> flavonoids<sup>14-15</sup> and triterpenoidal saponins<sup>7,14</sup> exist in this plant.

In addition, naringenin and kaempferol were detected in green callus and *in vitro* leaf extracts of *J. gendarussa* using GC-FID.<sup>16</sup> On the other hand, two flavonoid compounds (naringenin and kaempferol) were successfully detected and quantified in *J. gendarussa* methanolic leaf extracts (wild type) using GC-FID and RP-HPLC (unpublished reports). Naringenin and kaempferol are the natural phytoestrogen that presence in this leaf extracts and exhibit strong anti-proliferation of colon (HT-29), cervix (HeLa) and pancreas (BxPC-3) human cancer cells. The former studies reported naringenin have significant cytotoxicity and suppress apoptosis in mouse leukemia P388 cells<sup>17</sup> and cytotoxic activity on breast cancer cell, MCF-7,<sup>18</sup> HT-29,<sup>19</sup> while kaempferol were reported to inhibit cell proliferation and induce apoptosis in pancreatic cancer cells.<sup>20</sup>

To our knowledge, there is no scientific report pertaining cytotoxic activities on *J. gendarussa* methanolic leaf extracts against human cancer cell lines (HT-29, HeLa and BxPC-3). Therefore, in this study, we report for the first time on crude extracts of *J. gendarussa* against human cancer cell lines. This study was performed to screened the cytotoxic activities of methanolic leaf extracts from five different locations (Mersing, Muar, Pulai, Skudai and Batu Pahat) in Johor and two flavonoids (naringenin and kaempferol) against human cancer cell lines that might have correlation between the remain compounds in crude extracts and flavonoids compounds with the current findings. These data will provide scientific validation in the use of *J. gendarussa* leaves that have potential uses as an anti-cancer.

## ■ 2.0 EXPERIMENTAL

### 2.1 Plant Materials

*J. gendarussa* plants were collected from five different locations in Johor (Mersing, Muar, Skudai, Batu Pahat, Pulai) and maintained in a greenhouse at Faculty Biosciences and Medical Engineering, University Teknologi Malaysia (UTM). The commercial standards (kaempferol and naringenin) were purchased from Sigma Aldrich Corporation, Malaysia. Mature leaves and commercial standards (kaempferol and naringenin) were used as a starting material. The samples were used after diluted the stock solution with dimethylsulfoxide (DMSO) where DMSO at the concentrations lower than 1% had no effect on cell growth.

### 2.2 Flavonoids Detection of *J. gendarussa* Methanol Leaf Extracts

Detection of flavonoids in leaf extracts were performed using gas chromatography - flame ionization detector (GC-FID).<sup>16,21</sup>

### 2.3 Preparations of *J. gendarussa* Methanol Leaf Extracts

The matured leaves of *J. gendarussa* were air dried for 4 weeks. Then, the dried leaves were ground into small particles and approximately, 50 g of small particles of leaves were soaked into 1000 ml of methanol at room temperature for 72 hours in ratio of 1:20 (w/v).<sup>22</sup> The mixtures were filtered through sterile cotton and filtered again using Whatman No. 1 filter paper to obtain the methanol supernatants. The filtered methanol extraction was evaporated at 40 °C under reduced pressure by using rotary evaporator (EYELA N-1000, EYELA, Tokyo, Japan). The dried crude extracts were kept at 4 °C prior to use.

### 2.4 Culture of Cells

Three human cancer cell lines, HT-29 (colon adenocarcinoma), HeLa (cervix adenocarcinoma), BxPC-3 (epitheloid cervix adenocarcinoma) and one normal cell line CHO (chinese hamster ovary) were obtained from American Type Culture Collection (ATCC) and were a generous gift from Dr Salehuddin Hamdan (Animal Cell Culture Laboratory, Faculty of Biosciences and Medical Engineering, UTM). HT-29, HeLa, BxPC-3 cells were cultured in Roswell Park Memorial Institute 1640 (RPMI-1640) medium supplemented with 10% v/v fetal calf serum, 100 U/ml of penicillin and 100 µg/ml of streptomycin as a complete growth medium. Cells were maintained in 25 cm<sup>2</sup> flask at 37 °C with 5 % CO<sub>2</sub>. All of the materials were obtained from the (Gibco, Bio-Diagnostics, Petaling Jaya, Selangor, Malaysia).

### 2.5 MTT Assay

Cytotoxicity assay was determined by using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT, Sigma) based on previous study.<sup>23-24</sup> In this assay, cells were harvested when it reached 80 % confluent. Before starting the MTT assay, cells were optimized at a different seeding density that ranging from  $2.0 \times 10^3$  cell/ml to  $1.0 \times 10^6$  cell/ml in light to get the suitable seeding number for the experiment. Then, each well of the microtiter plate (96-well) was filled with 100 µl of cells suspension (HT-29 with seeding number;  $5 \times 10^4$  cell/ml), (HeLa with seeding number;  $5 \times 10^4$  cell/ml), (BxPC-3 with seeding number;  $1 \times 10^5$  cell/ml) and (CHO with seeding number;  $5 \times 10^4$  cell/ml) in RPMI- 1640 complete growth medium. After 24 hour of incubation, the cells were treated (100 µl/well) with the plant extracts in different concentration that ranging from 7.81 to 1000 µg/ml. The total volume of each well is 200 µl with technical replicates. The microtiter plates were further incubated for 72 hours with the plant extracts. After 72 hours of incubation, 20 µl of 5 mg/ml MTT was added to each well and the plates then incubated for 4 hours at 37 °C. Then, the medium in each well was carefully removed without disturbing the MTT crystals in the wells. The MTT formazan crystals were dissolved by addition of 225 µl of 100mM HCl-isopropanol to each wells. After completing the solubilization of the purple formazan the absorbance was measured using the BioRad microplate reader (BioRad, Tokyo, Japan) at a wavelength of 575 nm. The cytotoxicity was recorded as the drug concentration causing 50 % growth inhibition of cell lines (IC<sub>50</sub> value) using the formula given below:

$$\% \text{ cell viability} = \frac{A_{\text{sample}}(\text{mean})}{A_{\text{control}}(\text{mean})} \times 100 \%$$

### 2.6 Statistical Analysis

All of the samples were run in three replicates. Data obtained were analyzed using SPSS for Window software (SPSS 16.0 for Windows Evaluation Version software, SPSS Inc., USA). The normality of the data was tested using The Shapiro-Wilk test. The data were analyzed using Independence t-test for normal data and Mann-Whitney test for non-normal data.<sup>25</sup> Differences were considered to be significant if the probability  $p < 0.05$ .


### 3.0 RESULTS AND DISCUSSION

In this study, screening of cytotoxicity activities on *J. gendarussa* methanolic leaf extracts from five different locations were evaluated against human cancer cell lines using MTT assay. IC<sub>50</sub> values defined as the concentrations of the extracts to produce a 50 % reduction in viability of cells.<sup>26</sup> According to <sup>27-29</sup> studies reported that the cytotoxicity was evaluated based on IC<sub>50</sub> values i.e. IC<sub>50</sub> values below than 20 µg/ml (cytotoxicity), range between 21 - 40 µg/ml (weak cytotoxicity) and above than 41 µg/ml (not cytotoxicity). Figure 3.1, shows chromatogram that correspond to both flavonoid compounds (naringenin and kaempferol) in leaf extract from Mersing by using GC-FID. Also, *in vitro* leaf extracts and green callus extracts of *J. gendarussa* were detected and quantified the naringenin and kaempferol contents.<sup>16</sup> It shows the presence of flavonoids (naringenin and kaempferol) in *J. gendarussa* leaf extracts that could be associated to inhibit the proliferation of HT-29, HeLa and BxPC-3 cell lines.

The *J. gendarussa* leaf extracts from five different locations showed the cytotoxicity in a dose- dependent manner from lowest to highest concentration in HT-29, colon cancer cells. The similar trend of inhibitory cells was recorded in all extracts. Table 3.1 shows IC<sub>50</sub> values of leaf extracts from Mersing, Muar, Skudai, Batu Pahat and Pulai. The IC<sub>50</sub> values of leaf extracts from Muar, Skudai and Pulai showed high significantly difference (p<0.001) between different concentration of extracts. Leaf extracts from Mersing showed weak cytotoxicity with the approximately IC<sub>50</sub> values of 21 µg/ml followed by leaf extracts from Batu Pahat (36 µg/ml), Pulai (39 µg/ml), Muar (65 µg/ml) and Skudai (76 µg/ml). The results are summarized in Table 3.2. From the data obtained,

leaf extracts from Mersing was found effective inhibit the proliferation of the HT-29 cells and it is believed could be due the presence of more kaempferol contents rather than naringenin as shown in chromatogram (Figure 3.1). The results are summarized in table 3.6 showed the high cytotoxic activity on kaempferol as compared to naringenin against HT-29 cells.

Furthermore, production of flavonoids and various flavonoid compositions could be found in different regions or locations due to genetic variations and different environment conditions.<sup>30</sup> Based on the results, *J. gendarussa* leaf extracts from Mersing, Batu Pahat and Pulai were more cytotoxicity contrary on other study in *M. calabura* leaf extracts with IC<sub>50</sub> value of 46 µg/ml against HT-29 cells.<sup>22</sup>



**Figure 3.1** Chromatogram of naringenin and kaempferol contents in leaf extract from Mersing using GC-FID

**Table 3.1** Cytotoxicity study on *J. gendarussa* methanolic leaf extracts from five different locations against HT-29 cells

Concentration (µg/ml)	7.81	15.63	31.25	62.5	125	250	500	1000
<b>Leaf extracts from Mersing</b>	86.37± 9.78 ***	75.13± 9.00 ***	9.67± 1.19 **	3.97± 0.46 **	2.44± 0.77 ***	2.24± 1.58 ***	1.78± 0.40 ***	1.71± 0.36 ***
<b>Leaf extracts from Muar</b>	81.75± 0.29 ***	77.66± 3.16 ***	73.56± 0.97 **	51.92± 4.12 ***	17.76± 1.23 ***	3.86± 0.85 ***	2.10± 0.26 ***	1.74± 0.21 ***
<b>Leaf extracts from Skudai</b>	73.87± 2.52 *	71.24± 2.10 **	68.04± 2.35 *	56.83± 1.96 *	19.72± 0.42 **	3.69± 0.36 ***	2.15± 0.33 ***	2.49± 0.03 ***
<b>Leaf extracts from Batu Pahat</b>	67.43± 8.53 ***	61.04± 6.81 **	51.46± 7.57 ***	47.49± 7.71 ***	14.44± 3.16 ***	3.02± 0.70 ***	2.99± 0.84 ***	2.87± 0.31 ***
<b>Leaf extracts from Pulai</b>	73.52± 3.31 *	69.51± 4.39 **	57.12± 4.62 **	31.28± 3.25 **	4.69± 1.35 **	3.64± 1.10 **	3.62± 0.29 **	3.38± 0.11 **

Values are mean±SD for 3 replicates \*p<0.05, \*\*p<0.01, \*\*\*p<0.001 compared with control


**Table 3.2** Comparison of IC<sub>50</sub> values the various extracts of *J. gendarussa* leaf extracts in human cancer cell lines

Extracts of <i>J. gendarussa</i>	IC <sub>50</sub> values (µg/ml)		
	HT-29	HeLa	BxPC-3
Mersing	21	22	16
Muar	65	88	157
Skudai	76	39	105
Batu Pahat	36	146	410
Pulai	39	205	264

The morphological changes of the cells were observed under Nikon fluorescence microscope (Nikon, Tokyo, Japan) (10× magnification) after 72 hours treatment. The IC<sub>50</sub> leaf

extracts from various locations treated in HT-29 cancer cell lines revealed morphology changes (Figure 3.2: B,C,D,E,F) as compared to non-treated cells (Figure 3.2:A). Treated cells

showed a more prominent growth inhibition and shrinkage of the cells. On the contrary, untreated cells remained confluent throughout the incubation period (Figure 3.2:A).



**Figure 3.2** Morphology of  $IC_{50}$  in *J. gendarussa* leaf extracts from five different locations against HT-29 cancer cell lines after MTT assay (10 $\times$  magnification). (A) HT-29 cells without any treatment, control; (B) leaf extract from Mersing, 31.25  $\mu$ g/ml; (C) leaf extract from Muar, 125  $\mu$ g/ml; (D) leaf extract from Skudai, 125  $\mu$ g/ml; (E) leaf extract from Batu Pahat, 62.5  $\mu$ g/ml; (F) leaf extract from Pulai, 62.5  $\mu$ g/ml

Table 3.3 represent HeLa cancer cells viability on *J. gendarussa* leaf extracts from Mersing, Muar, Pulai, Skudai and Batu Pahat.  $IC_{50}$  values of leaf extracts from Mersing, Muar, Skudai and Batu Pahat showed high significantly difference ( $p < 0.001$ ) between different concentrations of extracts. Leaf extracts from Mersing showed weak cytotoxicity with  $IC_{50}$

values of approximately 22  $\mu$ g/ml followed by leaves extracts from Skudai (39  $\mu$ g/ml), Muar (88  $\mu$ g/ml), Batu Pahat (146  $\mu$ g/ml) and Pulai (205  $\mu$ g/ml) (Table 3.2). The cell viability of HeLa cancer cells were decreased in dose- dependent manner and the trend of inhibitory cells are same in all extracts. From the results, leaf extract from Mersing was effective against HeLa cells as compared to other extracts and this activity could be associated with the presence of more kaempferol contents as compared to naringenin contents in the extracts (Figure 3.1).

The  $IC_{50}$  methanolic leaf extracts from various locations treated in HeLa cancer cell lines revealed morphology changes (Figure 3.3: B,C,D,E,F) as compared to non-treated cells (Figure 3.3: A). Treated cells show a more prominent growth inhibition and shrinkage of the cells. On the contrary, untreated cells remained confluent throughout the incubation period (Figure 3.3:A).


The effect of *J. gendarussa* leaf extracts from Mersing, Muar, Pulai, Skudai and Batu Pahat on the cytotoxicity potential of BxPC-3 cancer cell lines are demonstrated in Table 3.4. The  $IC_{50}$  values showed high significantly difference ( $p < 0.001$ ) in leaf extracts from Skudai and Pulai, followed by leaf extract from Mersing, Muar and Batu Pahat showed moderate significant difference ( $p < 0.01$ ) between different concentration of extracts. The similar trend inhibition of cells and the cell viability of the cells were decreased in dose- dependent manner in all extracts. The result from cytotoxicity assay on BxPC-3 cancer cell lines are summarized in table 3.2, leaf extracts from Mersing showed cytotoxicity with  $IC_{50}$  values of approximately 16  $\mu$ g/ml and other extracts showed not cytotoxicity against BxPC-3.<sup>27-29</sup> The leaf extract from Mersing showed cytotoxicity and it is believed from the chromatogram indicates the presence of more kaempferol contents rather than naringenin contents that could be associated the proliferation of the cells rather than other leaf extracts (Figure 3.1).

**Table 3.3** Cytotoxicity study on *J. gendarussa* methanolic leafextracts from five different locations against HeLa cells

Concentration ( $\mu$ g/ml)	7.81	15.63	31.25	62.5	125	250	500	1000
	***	**	***	*	***	***	***	*
Leaf extracts from Mersing	82.82 $\pm$ 3.33	64.55 $\pm$ 2.73	38.13 $\pm$ 2.86	4.64 $\pm$ 0.26	1.53 $\pm$ 0.21	1.21 $\pm$ 0.26	1.54 $\pm$ 0.30	4.02 $\pm$ 0.39
	*	***	**	***	***	***	***	***
Leaf extracts from Muar	82.18 $\pm$ 5.46	69.10 $\pm$ 6.79	64.51 $\pm$ 3.43	56.66 $\pm$ 2.86	44.34 $\pm$ 1.90	32.78 $\pm$ 1.90	1.22 $\pm$ 0.89	0.89 $\pm$ 0.29
	**	***	***	***	***	***	***	***
Leaf extracts from Skudai	70.12 $\pm$ 2.78	57.44 $\pm$ 3.08	53.06 $\pm$ 1.59	43.28 $\pm$ 2.45	38.88 $\pm$ 2.91	32.49 $\pm$ 4.24	4.14 $\pm$ 0.95	5.30 $\pm$ 0.37
	*	**	***	**	***	**	**	***
Leaf extracts from Batu Pahat	85.04 $\pm$ 3.11	78.27 $\pm$ 4.92	65.48 $\pm$ 4.77	62.32 $\pm$ 4.09	52.95 $\pm$ 3.47	41.16 $\pm$ 5.37	28.74 $\pm$ 7.28	2.59 $\pm$ 1.15
	**	***	***	***	**	***	***	***
Leaf extracts from Pulai	76.21 $\pm$ 7.76	63.63 $\pm$ 7.27	63.49 $\pm$ 1.94	60.39 $\pm$ 4.96	57.33 $\pm$ 4.63	47.16 $\pm$ 5.00	32.03 $\pm$ 4.51	2.33 $\pm$ 0.58

Values are mean $\pm$ SD for 3 replicates \*, $p < 0.05$ , \*\*, $p < 0.01$ , \*\*\*, $p < 0.001$  compared with control





**Figure 3.3** Morphology of IC<sub>50</sub> in *J. gendarussa* leaf extracts from five different locations against HeLa cancer cell lines after MTT assay (10× magnification). (A) HeLa cells without any treatment, control; (B) leaf extract from Mersing, 31.25 µg/ml; (C) leaf extract from Muar, 125 µg/ml; (D) leaf extract from Skudai, 62.5 µg/ml; (E) leaf extract from Batu Pahat, 125 µg/ml; (F) leaf extract from Pulai, 250 µg/ml


**Table 3.4** Cytotoxicity study on *J. gendarussa* methanolic leaf extracts from five different locations against BxPC-3 cells

Concentration (µg/ml)	7.81	15.63	31.25	62.5	125	250	500	1000
	*	**	**	***	***	***	***	***
<b>Leaf extracts from Mersing</b>	69.44±4.25	45.79±5.25	43.16±3.95	25.76±4.54	5.28±0.19	4.42±0.82	3.94±0.87	15.21±2.39
	***	**	*	*	*	**	**	***
<b>Leaf extracts from Muar</b>	65.88±0.69	59.02±13.05	57.47±9.92	55.85±11.82	54.68±11.71	36.52±8.12	9.55±7.00	6.03±2.21
	***	***	***	***	***	***	*	***
<b>Leaf extracts from Skudai</b>	74.79±1.13	70.69±0.41	60.45±3.89	54.48±0.33	49.91±1.41	48.52±1.23	47.83±14.79	6.48±0.43
	*	*	*	*	***	***	**	***
<b>Leaf extracts from Batu Pahat</b>	90.85±5.19	87.36±5.55	87.16±1.97	82.92±2.35	75.51±1.98	70.10±1.63	41.62±7.33	4.52±0.20
	***	*	***	***	***	**	***	***
<b>Leaf extracts from Pulai</b>	79.24±2.76	73.29±17.6	69.03±6.04	64.42±5.55	58.24±6.41	52.49±3.65	14.37±1.05	6.49±0.36

Values are mean±SD for 3 replicates \*,p<0.05, \*\*,p<0.01, \*\*\*,p<0.001 compared with control

The IC<sub>50</sub> methanolic leaf extracts from various locations treated in BxPC-3 cancer cell lines revealed morphology changes (Figure 3.4: B,C,D,E,F) as compared to non-treated cells (Figure 3.4:A). Treated cells show a more prominent

growth inhibition and shrinkage of the cells. On the contrary, untreated cells remained confluent throughout the incubation period (Figure 3.4:A).



**Figure 3.4** Morphology of IC<sub>50</sub> in *J. gendarussa* leaf extracts from five different locations against BxPC-3 cancer cell lines after MTT assay (10× magnification). (A) BxPC-3 cells without any treatment, control; (B) leaf extract from Mersing, 15.63µg/ml; (C) leaf extract from Muar, 250µg/ml; (D) leaf extract from Skudai, 125µg/ml; (E) leaf extract from Batu Pahat, 500µg/ml; (F) leaf extract from Pulai, 500µg/ml

Table 3.5 demonstrated that the ability of kaempferol and naringenin to inhibit the proliferation of HT-29, HeLa and BxPC-3 cancer cell lines in this study. The results are as summarized in table 3.6, kaempferol showed highly cytotoxicity with IC<sub>50</sub> values of approximately 5 µg/ml (HeLa) and 6 µg/ml (HT-29). This followed by naringenin with IC<sub>50</sub> values of approximately 15 µg/ml (HeLa) and 19 µg/ml (HT-29) cancer cell lines. The IC<sub>50</sub> values for kaempferol shows high significantly difference (p<0.001) between difference concentrations in all compounds whereas naringenin shows high significantly difference (p<0.001) in HT-29 and BxPC-3 while weak significantly difference (P<0.05) in HeLa cancer cell lines between difference concentrations in all compounds (Table 3.5).

From the chromatogram (Figure 3.1) shows both compounds were detected in leaf extracts from Mersing and assessed in this study to determine the effectiveness against human cancer cell lines where both compounds might be contributed the proliferation of cells tested in leaf extracts. It is suggested, the combination of kaempferol and naringenin in leaf

extracts that could be associated in proliferation of the cells. Previous study also reported that combination of naringenin and kaempferol showed toxicity with IC<sub>50</sub> values of 43.73 µg/ml as compared to naringenin and kaempferol alone against HeLa.<sup>31</sup> However, in this study, kaempferol and naringenin alone showed high cytotoxicity with IC<sub>50</sub> values of approximately 5 µg/ml and 15 µg/ml, respectively against HeLa cells. The cell of HT-29, HeLa and BxPC-3 were shrinkage decreased in dose-dependent manner and the similar trends of inhibitory of cells in all compounds. On the other hand, kaempferol showed high toxicity rather than naringenin could be due to kaempferol is strong estrogenic activity as compared to naringenin. It is suggested kaempferol have benefits potential effect in preventing estrogen imbalance diseases such as breast cancer, osteoporosis, cardiovascular,<sup>32</sup> hypertension, headache, abdominal pains and rheumatism.<sup>33</sup> However, naringenin also reported induced inhibitory effects on human cancer cell lines (MDA-MB-231 and Caco-2).<sup>17</sup>

**Table 3.5** Cytotoxicity study on kaempferol and naringenin against HT-29, HeLa and BxPC-3 cells lines

Concentration (µg/ml)		3.91	7.81	15.63	31.25	62.5	125	250	500
<b>Cells</b>									
<b>HT-29</b>	<b>Kaempferol</b>	** 61.19±5.90	*** 30.09±1.86	*** 15.59±0.95	*** 3.20±0.55	*** 2.67±0.43	*** 2.62±1.38	*** 4.35±1.33	*** 4.09±0.35
	<b>Naringenin</b>	*** 70.17±5.48	** 67.06±2.63	* 56.19±1.37	*** 34.41±7.82	*** 17.04±5.08	*** 1.60±0.31	*** 1.51±1.26	*** 1.33±0.76
<b>HeLa</b>	<b>Kaempferol</b>	*** 51.22±0.67	*** 25.15±0.69	*** 13.28±2.59	*** 6.49±0.66	*** 5.71±0.86	*** 5.80±0.86	*** 4.90±1.12	*** 6.92±0.60
	<b>Naringenin</b>	* 69.73±8.40	*** 60.26±7.12	* 48.58±5.99	*** 27.94±2.66	*** 15.01±2.55	*** 4.08±1.31	*** 3.37±0.16	*** 4.22±0.96
<b>BxPC-3</b>	<b>Kaempferol</b>	** 69.55±3.76	** 67.13±10.39	*** 66.21±5.65	*** 31.54±5.90	*** 27.74±0.78	*** 27.09±1.58	*** 30.91±1.69	*** 24.51±6.45
	<b>Naringenin</b>	** 76.10±8.98	** 75.69±6.58	*** 67.96±11.57	*** 62.97±11.32	*** 46.82±0.54	*** 8.24±1.31	*** 6.37±1.12	*** 5.06±0.88


Values are mean±SD for 3 replicates \*,p<0.05, \*\*,p<0.01,\*\*\*,p<0.001 compared with control

**Table 3.6** Comparison of IC<sub>50</sub> values the compounds (kaempferol and naringenin) in human cancer cell lines

Compounds	IC <sub>50</sub> values (µg/ml)		
	HT-29	HeLa	BxPC-3
<b>Kaempferol</b>	6	5	23
<b>Naringenin</b>	19	15	57

The IC<sub>50</sub> kaempferol and naringenin in HT-29 (Figure 3.5: B,C), HeLa (Figure 3.5: E,F) and BxPC-3 (Figure 3.5: H,I) cancer cell lines revealed morphology changes as compared to non-treated cells (Figure 3.5:A,D,G). Treated cells showed a

more prominent growth inhibition and shrinkage of the cells. On the contrary, untreated cells remained confluent throughout the incubation period.



**Figure 3.5** Morphology of  $IC_{50}$  in kaempferol and naringenin against HT-29, HeLa and BxPC-3 cancer cell lines after MTT assay (10 $\times$  magnification). (A) HT-29 cells without any treatment, control; (B) HT-29 cells treated with kaempferol, 7.81  $\mu\text{g/ml}$ ; (C) HT-29 cells treated with naringenin, 31.25  $\mu\text{g/ml}$ ; (D) HeLa cells without any treatment, control; (E) HeLa cells treated with kaempferol, 7.81  $\mu\text{g/ml}$ ; (F) HeLa cells treated with naringenin, 15.63  $\mu\text{g/ml}$ ; (G) BxPC-3 cells without any treatment, control; (H) BxPC-3 cells treated with kaempferol, 31.25  $\mu\text{g/ml}$ ; (I) BxPC-3 cells treated with naringenin, 62.5  $\mu\text{g/ml}$

Table 3.7 shows the cytotoxicity study on *J. gendarussa* leaf extracts from Mersing, Muar, Pulai, Skudai and Batu Pahat and also flavonoids (kaempferol and naringenin) against CHO normal cell lines. CHO cell lines are positive control used to compare the cytotoxic activity on HT-29, HeLa and BxPC-3 human cancer cell lines. The comparisons of both compounds were performed based on  $IC_{50}$  between human cancer and normal cell lines.<sup>27-29</sup> From this finding, leaf extract from Mersing was weak cytotoxicity against CHO cells but not cytotoxicity in others leaf extracts. However,  $IC_{50}$  values of leaf extracts in Mersing against human cancer cells highly cytotoxicity as compared in CHO normal cells. It is suggested, the cytotoxic activity on leaf extracts from Mersing could be associated presence high contents of flavonoid compounds. Therefore, the active compounds particularly flavonoid contents in *J. gendarussa* leaf extracts might be contributed the cytotoxicity activity against HT-29, HeLa and BxPC-3 cancer cells lines.

**Table 3.7** Comparison of  $IC_{50}$  values the compounds and various extracts of *J. gendarussa* leaf extracts in CHO normal cell lines

Compounds	$IC_{50}$ values ( $\mu\text{g/ml}$ )
<b>Kaempferol</b>	14
<b>Naringenin</b>	21
<b>Extract of <i>J. gendarussa</i></b>	
<b>Mersing</b>	28
<b>Muar</b>	108
<b>Skudai</b>	88
<b>Batu Pahat</b>	190
<b>Pulai</b>	305

#### 4.0 CONCLUSION

In conclusion, various leaf extracts and flavonoid compounds were evaluated and exhibited a dose- dependent manner on cytotoxicity effects against human cancer cell lines. In this study, leaf extracts from Mersing demonstrated high cytotoxic against BxPC-3 pancreas cell lines. It is suggested, leaf extracts from Mersing containing active compounds particularly flavonoids that exhibit potential inhibitors of growth and act as an anti-cancer plant particularly in BxPC-3 cells. Further investigation to those researches are interested on *J. gendarussa* plant should review on positive standard drugs such as tamoxifen that could provide additional insight of this extracts uses as chemotherapeutic agents. Furthermore, this study justifies the *J. gendarussa* leaf extracts as a folklore medicinal uses and validates its potential uses as anti-cancer.

#### Acknowledgement

The authors would like to thank Universiti Teknologi Malaysia (UTM) and Ministry of High Education (MOHE) for financial support (Research University Grant, Vot 05J27) and Animal Tissue Culture, Faculty Biosciences and Medical Engineering for lab facilities.

#### References

- [1] Malek, A. S. N., Lee, G. S., I, Hong, S. L., Yaacob, H., Abdul Wahab, N., Weber, J-F, F. and Ali Shah, S. A. 2011. Phytochemical and Cytotoxic Investigations of *Curcuma mangga* Rhizomes. *Molecules*. 16: 4539-4548.
- [2] Khatijah, H. and Noraini, T. 2007. *Anatomical Atlas of Medicinal Plants*. Penerbit UKM. 1: 106.

- [3] Ayob, Z., Wagiran, A. and Abd Samad, A. 2013. Potential of Tissue Cultured Medicinal Plants in Malaysia. *Jurnal Teknologi*. 62(1):111–117.
- [4] Priyanayagam, K., Umamaheswari, B., Suseela, L., Padmini, M. and Ismail, M. 2009. Evaluation of Antiangiogenic Effect of the Leaves of *Justicia gendarussa* (Burm. f.) (Acanthaceae) by Chrio Allontic Membrane Method. *American Journal of Infectious Diseases*. 5(3): 180–182.
- [5] Krishna, K.L. Mehta, T.A. and Patel, J.A. 2010. *In-vitro* Hepatoprotective Activity of *Justicia gendarussa* Stem on Isolated Rat Hepotocytes. *Pharmacologyonline*. 2: 9–13.
- [6] Sharma, K. K., Saikia, R., Kotoky, J., Kalita, J. C. and Devi, R. 2011. Antifungal Activity of *Solanum melongena* L, *Lawsonia inermis* L. and *Justicia gendarussa* B. against Dermatophytes. *International Journal of PharmaTech Research*. 3(3): 1635–1640.
- [7] Uddin, M. R., Sinha, S., Hossain, A. S., Kaisar, M. A., Hossain, M. K. and Rashid, M. A. 2011. Chemical and Biological Investigations of *Justicia gendarussa* (Burm.f.) *Dhaka University Journal of Pharmaceutical Sciences*. 10(1): 53–57.
- [8] Fazaludeen, M. F., Manickam, C., Ashanky, I. M. A., Ahmed, M. Q. and Beg, Q. Z. 2012. Synthesis and Characterization of Gold Nanoparticles by *Justicia gendarussa* Burm. f. Leaf Extract. *Journal of Microbiology and Biotechnology Research*. 2(1): 23–34.
- [9] Saha, M. R., Debnath, P. C., Rahman, M. A. and Islam, M. A. U. 2012. Evaluation of *In Vitro* Anthelmintic Activities of Leaf and Stem Extracts of *Justicia gendarussa*. *Bangladesh Journal Pharmacological*. 7: 50–53.
- [10] Subramanian, N., Jothimanivannan, C. and Moorthy, K. 2012. Antimicrobial Activity and Preliminary Phytochemical Screening of *Justicia Gendarussa* (Burm.f.) against Human Pathogens. *Asian Journal of Pharmaceutical and Clinical Research*. 5(3): 229–233.
- [11] Sudhanandh, V. S., Arjun, J. K., Faisal, A. K., Ani, M. V., Renjini, V. S. and Narendra Babu, K. 2012. *In-vitro* Antibacterial Screening of Selected Folklore Indian Medicinal Plants with Few Clinical Pathogens. *Indian Journal of Pharmaceutical Education and Research*. 46(2): 174–178.
- [12] Kowsalya, D. and Sankaranarayanan, S. 2012. Efficacies of Bactericidal *Justicia gendarussa* Extract Inhibiting Protein Synthesis Against Methicilin Resistant *Staphylococcus aureus*. *IOSR Journal of Pharmacy and Biological Sciences*. 4(2): 32–41.
- [13] Bambang Prajogo, E. W., Farida, I., Putri Febriyanti, A. I. and Jusak, N. 2008. Effect of *Justicia gendarussa* Burm.f. Leaves Water Fraction On Male Rabbit Liver and Renal Fuction (Sub acute Toxicity Test of *Justicia gendarussa* Burm.f. Leaves Water Fraction as Male Contraceptive Agent). *Veterinaria Medika*. 1(3): 79–82.
- [14] Ratnasooriya, W. D., Deraniyagala, S. A., and Dehigaspitiya, D. C. 2007. Antinociceptive Activity and Toxicological Study of Aqueous Leaf Extract of *Justicia gendarussa* Burm. f. in Rats. *Pharmacognosy Magazine*. 3: 145–155.
- [15] Mustafa, R. A., Hamid, A. A., Mohamed, S. and Bakar, E. A. 2010. Total Phenolic Compounds, Flavonoids, and Radical Scavenging Activity of 21 Selected Tropical Plants. *Journal of Food Sciences*. 75: 28–35.
- [16] Ayob, Z., Md Saari, N.H. and Abd Samad, A. 2012. In Vitro Propagation and Flavonoid Contents in Local *Justicia gendarussa* Burm. F. *UMT 11th International Annual Symposium on Sustainability Science and Management*. 09th–11th July 2012, Terengganu, Malaysia. 403–409.
- [17] Kanno, S-I., Tomizawa, A., Hiura, T., Osanai, Y., Shouji, A., Ujibe, M., Ohtake, T., Kimura, K. and Ishikawa, M. 2005. Inhibitory Effects of Naringenin on Tumor Growth in Human Cancer Cell Lines and Sarcoma S-180-implanted Mice. *Biological Pharmaceutical Bulletin*. 28(3): 527–530.
- [18] Park, J-H., Lee, J-W., Paik, H-D., Cho, S-G., Nah, S-Y., Park, Y-S. and Han, Y.S. 2010. Cytotoxic Effects of 7-O-Butyl Naringenin on Human Breast Cancer MCF-7 Cells. *Food Science Biotechnology*. 19: 717–724.
- [19] Neal, M. D. and Jaime, A. Y. 2013. *Flavonoid Pharmacokinetics: Methods of Analysis, Preclinical and Clinical Pharmacokinetics, Safety and Toxicology*. New Jersey: John Wiley & Sons, Inc.
- [20] Zhang, Y., Chen, A. Y., Li, M., Chen, C. and Yao, Q. 2008. *Ginkgo biloba* Extract Kaempferol Inhibits Cell Proliferation and Induces Apoptosis in Pancreatic Cancer Cells. *Journal of Surgical Research*. 148: 17–23.
- [21] Sarju, N., Abd. Samad, A., Abd Ghani, M. and Ahmad, F. 2012. Detection and Quantification of Naringenin and Kaempferol in *Melastoma Decemfidum* Extracts by GC-FID and GC-MS. *Acta Chromatographica*. 24(2): 221–228.
- [22] Zakaria, Z. A., Mohamed, A. M., Mohd Jamil, N. S., Rofiee, M. S., Khairi, H. M. and Sulaiman, M. R. 2011. In Vitro Antiproliferative and Antioxidant Activities of the Extracts of *Muntingia calabura* Leaves. *The American Journal of Chinese Medicine*. 39: 183–200.
- [23] Mosmann, T. 1983. Rapid Colorimetric Assay for Cellular Growth and Survival: Application to Proliferation and Cytotoxicity Assays. *Journal of Immunological Methods*. 65: 55–63.
- [24] Deny, S., Sirat, H. M., Ahmad, F. and Ali, R. M., Aimi, N. and Kitajima, M. 2007. Antioxidant and Cytotoxic Flavonoids from the Flowers of *Melastoma malabathricum* L. *Food Chemistry*. 103 (3): 1600–1606.
- [25] Pallant, J. 2007. *SPSS Survival Manual*. Third ed. Berkshire: McGraw-Hill.
- [26] Jafari, S., Saeidnia, S., Hajimehdipoor, H., Ardekani, M.R.S., Faramarzi, M.A., Hadjiakhoondi, A. and Khanavi, M. 2013. Cytotoxic Evaluation of *Melia azedarach* in Comparison with *Azadirachta indica* and its Phytochemical Investigation. *Journal of Pharmaceutical Sciences*. 21: 1–7.
- [27] Geran, R. I., Greenberg, N. H., Macdonald, M. M., Schumacher, A. M., and Abbott, B. J. 1972. Protocols for Screening Chemical Agents and Natural Products against Animal Tumors and Other Biological Systems. *Cancer Chemotherapy Reports*. 3: 1–103.
- [28] Mohamed, S. M., Ali, A. M., Rahmani, M., Dhaliwal, J. S. and Yusoff, K. 2000. Apoptotic and Neurotic Cell Death Manifestations in Leukemic Cells treated with Methylgerambulin a Sulphone from *Glycosmis calcicola*. *Journal Biochemistry Molecular Biology and Biophysiology*. 4: 253–261.
- [29] Rohaya, A., Abdul Manaf, A., Daud, A. I., Nor Hadiani, I., Khozirah, S. and Nordin, H. L. 2005. Antioxidant, Radical - Scavenging, Anti-inflammatory, Cytotoxic and Antibacterial Activities of Methanolic Extracts of Some *Hedyotis* Species. *Life Sciences*. 76: 1953–1964.
- [30] Cristina Figueroa, A., Andres Soria2, E., Jose Cantero, J., Silvina Sanchez, M. and Ester Goleniowski, M. 2012. Cytotoxic Activity of *Thelesperma megapotamicum* Organic Fractions against MCF-7 Human Breast Cancer Cell Line. *Journal of Cancer Therapy*. 3: 103–109.
- [31] Sarju, N., Abd Ghani, M., Hamdan, S. and Abd Samad, A. 2010. Antioxidant Activity and Cytotoxicity of the Leaves of *Melastoma Decemfidum* Roxb. Ex. Jack. *National Biotechnology Seminar*. 24–26 May. PWTC, Kuala Lumpur.
- [32] Oh, S. M., Kim, Y. P. and Chung, K. H. 2006. Biphasic Effects of Kaempferol on the Estrogenicity in Human Breast Cancer Cell. *Archives of Pharmacol Research*. 29(5): 354–362.
- [33] Guo, A. J., Choi, R. C., Zheng, K. Y., Chen, V. P., Dong, T. T., Wang, Z-T., Vollmer, G., Lau, D. T. and T, W-K. 2012. Kaempferol as a Flavonoid Induces Osteoblastic Differentiation via Estrogen Receptor Signalling. *Chinese Medicine*. 7: 1–7.