

1

2

3

Structural basis of meiotic chromosome synapsis through SYCP1 self-assembly

4

5

6 **Authors**

7 James M. Dunce¹, Orla M. Dunne¹, Matthew Ratcliff^{1,4}, Claudia Millán², Suzanne Madgwick¹, Isabel

8 Usón^{2,3}, Owen R. Davies¹

9

10

11 **Affiliations**

12 ¹Institute for Cell and Molecular Biosciences, Newcastle University, Newcastle upon Tyne, UK.

13 ²Crystallographic Methods, Institute of Molecular Biology of Barcelona (IBMB-CSIC), Barcelona,
14 Spain.

15 ³ICREA, Institució Catalana de Recerca i Estudis Avançats, Barcelona, Spain.

16 ⁴Present address: Department of Biochemistry, University of Cambridge, UK.

17

18

19 **Author for Correspondence**

20 Owen R. Davies: owen.davies@newcastle.ac.uk

21

22 **Abstract**

23 Meiotic chromosomes adopt unique structures in which linear arrays of chromatin loops are bound
24 together in homologous chromosome pairs by a supramolecular protein assembly, the
25 synaptonemal complex. This three-dimensional scaffold provides the essential structural framework
26 for genetic exchange by crossing over and subsequent homologue segregation. The core
27 architecture of the synaptonemal complex is provided by SYCP1. Here, we report the structure and
28 self-assembly mechanism of human SYCP1 through X-ray crystallographic and biophysical studies.
29 SYCP1 has an obligate tetrameric structure in which an N-terminal four-helical bundle bifurcates into
30 two elongated C-terminal dimeric coiled-coils. This building-block assembles into a zipper-like lattice
31 through two self-assembly sites. N-terminal sites undergo cooperative head-to-head assembly in the
32 midline, whilst C-terminal sites interact back-to-back on the chromosome axis. Our work reveals the
33 underlying molecular structure of the synaptonemal complex in which SYCP1 self-assembly
34 generates a supramolecular lattice that mediates meiotic chromosome synapsis.

35

36

37

38

39

40

41

42

43

44 **Keywords**

45 Meiosis, chromosome structure, double-strand break, chiasmata, synaptonemal complex, SYCP1,
46 self-assembly

47

48 **Introduction**

49 The reduction in chromosome number during meiosis requires a unique programme of intricate
50 molecular processes including the synapsis of homologous chromosome pairs, their exchange of
51 genetic material by crossing over, and ultimately their segregation into haploid cells. At the centre of
52 these processes is a supramolecular protein assembly, the synaptonemal complex (SC). The SC binds
53 together homologous chromosome pairs, structured as linear arrays of chromatin loops, in a single
54 continuous synapsis along their entire length^{1,2}. SC assembly occurs in a spatiotemporal manner,
55 dependent on the prior establishment of inter-homologue recombination intermediates through
56 double-strand break induction, which act as guides to ensure the synapsis of perfectly aligned
57 homologues^{3,4}. The three-dimensional structure of the SC provides the essential architectural
58 framework for the resolution of recombination intermediates, which includes the generation of one
59 genetic crossover per chromosome arm^{5,6}. Crossovers are essential for correct segregation of
60 homologues at anaphase I, and additionally contribute to genetic diversity. The defective assembly
61 of the SC is associated with human infertility, miscarriage and aneuploidy^{7,8}. However, despite its
62 discovery more than half a century ago, the molecular structure and function of the SC have
63 remained unknown.

64

65 Electron micrographs of the SC reveal a characteristic tripartite structure that is conserved across
66 eukaryotes⁹. This consists of two lateral elements, each coating a chromosome axis, separated by a
67 100 nm central region that contains a midline 20-40 nm wide central element (Fig. 1a). The central
68 and lateral elements are connected together by a network of angled transverse filaments, which in
69 hamster have a diameter of approximately 16 Å and are spaced at a density of 50-80 per 1 µm of
70 chromosome axis¹⁰. In addition to its 100 nm width, the SC central region has a depth of up to 100
71 nm, so is a truly three-dimensional protein assembly^{11,12}.

72

73 In mammals, SC transverse filaments are formed by SYCP1¹³. This 976 amino acid protein contains a
74 central α -helical core flanked by unstructured N- and C-terminal tails (Fig. 1b). SYCP1 N- and C-
75 termini are localised within SC central and lateral elements respectively, and so is bioriented with
76 juxtaposed SYCP1 molecules providing a 150 nm separation between opposing C-termini in
77 mice^{12,14,15} (Fig. 1a). The SC contains at least two layers of SYCP1 molecules; N-termini are detected
78 in two vertically separated chains within the central element, whereas C-termini are present in a
79 single chain within the lateral element^{12,16}. SC lateral elements also contain SYCP2 and SYCP3^{17,18}, the
80 latter contributing to chromosome compaction through stabilisation of chromatin loop structures¹⁹⁻
81 ²¹. The SC central element contains initiation factors SYCE3, SYCE1 and SIX6OS1 that stabilise initial
82 tripartite structures²²⁻²⁵, and elongation complex SYCE2-TEX12 that stabilises the long-range
83 extension of the tripartite structure²⁶⁻²⁹.

84

85 SYCP1 disruption leads to a complete failure of synapsis; recombination intermediates are formed
86 but fail to resolve, crossovers fail to form, cells undergo meiotic arrest and there is a resultant
87 complete infertility⁵. Whilst SC central and lateral element components are essential for the
88 structure and function of the mature SC, SYCP1 is recruited to meiotic chromosomes in the absence
89 of other SC central and lateral element components, albeit at reduced levels, and is essential for the
90 recruitment of all SC central element proteins^{5,17,18,22,23,25-28}. Furthermore, SYCP1 in isolation has an
91 intrinsic capacity for self-assembly into rudimentary SC-like structures³⁰. Thus, SYCP1 self-assembly
92 seemingly provides the underlying architectural framework of the SC.

93

94 Here, we report the structure and self-assembly mechanism of SYCP1. The obligate unassembled
95 structure of SYCP1 is an N-terminal tetramer that bifurcates into two elongated C-terminal dimeric
96 coiled-coils. This building-block self-assembles into a supramolecular lattice that defines the SC
97 structure through sites within its N- and C-termini. Whilst N-terminal sites undergo cooperative
98 head-to-head assembly, C-terminal sites interact back-to-back in a protonation-dependent manner

99 that relies upon chromosomal recruitment by unstructured C-terminal tails. Together, our data lead
100 to a complete molecular model for the structure of SYCP1 in which recursive self-assembly at N- and
101 C-terminal sites leads to the formation of a continuous and cooperative supramolecular lattice.
102 Through this, we reveal the underlying structure of the synaptonemal complex and the molecular
103 basis of meiotic chromosome synapsis by SYCP1.

104

105

106 Results

107 The obligate structure of SYCP1

108 Human SYCP1 contains a large α -helical core (α Core) of amino acids 101-783, flanked by
109 unstructured N- and C-terminal tails (Fig. 1b and Supplementary Fig. 1a). Size-exclusion
110 chromatography multi-angle light scattering (SEC-MALS) analysis of purified recombinant SYCP1
111 α Core revealed heterogeneous 1-12 MDa species (Fig. 1c and Supplementary Fig. 1b), indicating an
112 intrinsic capacity to self-assemble *in vitro*. Self-assembly of large molecular weight species is
113 completely abrogated by deletion of the first 11 amino acids at its N-terminal tip (α N-tip), with
114 α Core- Δ Ntip (residues 112-783) forming a stable tetramer (Fig. 1c). Circular dichroism (CD)
115 spectroscopy confirms that α Core- Δ Ntip is almost entirely α -helical (Supplementary Fig. 2a,b). Size
116 exclusion chromatography small-angle X-ray scattering (SEC-SAXS) analysis reveals scattering profiles
117 and real space pair-distance distribution functions ($P(r)$ distributions) corresponding to an elongated
118 molecule of 900 Å length (Fig. 1d and Supplementary Fig. 2c). This matches its theoretical α -helical
119 coiled-coil length and is sufficient to span just over half of the inter-chromosomal distance, in
120 keeping with SYCP1 biorientation within the SC. We conclude that α Core- Δ Ntip is an extended α -
121 helical coiled-coil tetramer that represents the obligate structure of SYCP1, and self-assembly of this
122 minimum building-block into higher molecular weight species is dependent on the N-terminal tip of
123 SYCP1 α Core.

124

125 The obligate α Core- Δ Ntip is composed of two distinct structural units, an N-terminal tetramer
126 (residues 206-362) and C-terminal dimer (residues 358-783) (Fig. 1c). These boundaries were
127 identified through exhaustive screening to define clearly demarcated structural regions of maximal
128 stability; nevertheless, oligomer states and structures of these and other constructs described herein
129 are robust across a range of sequence boundaries (Supplementary Fig. 1a and Supplementary Table
130 1). The α N-tetramer and α C-dimer are almost entirely α -helical (Supplementary Fig. 2a,b); SEC-SAXS
131 analysis reveals elongated structures of respective lengths 260 Å and 645 Å (Fig. 1d and

132 Supplementary Fig. 2c-e), matching their theoretical coiled-coil lengths. The cross-sectional radius of
133 gyration (R_c) was determined as 10.3 Å and 8.9 Å for α N-tetramer and α C-dimer (Supplementary Fig.
134 2f), corresponding to the known dimensions of four-helical and dimeric coiled-coils respectively. The
135 R_c of α C-dimer (8.9 Å) indicates a diameter of 17.8 Å that closely matches the measured 16 Å
136 diameter of transverse filaments in the hamster SC¹⁰, suggesting that α C-dimers constitute the
137 individual structures visualised spanning between SC central and lateral elements.

138

139 We determined the orientation of helices within α Core- Δ Ntip, α N-tetramer and α C-dimer through
140 SEC-SAXS $P(r)$ analysis of N-terminal MBP fusion proteins, exploiting the strong scattering of globular
141 proteins in comparison to coiled-coils to identify the relative positions of globular tags. In all cases,
142 $P(r)$ distributions demonstrate strong inter-MBP peaks at short distances, compatible with their
143 parallel orientation, but lack inter-MBP peaks at long distances that would occur in anti-parallel
144 structures (Fig. 1e and Supplementary Fig. 2d-h). Similarly, an N-terminal GST fusion of α N-tetramer
145 shows only short distance inter-GST peaks (Supplementary Fig. 2 j-n). Finally, the α N-tetramer and
146 α C-dimer structures are compatible with their N-terminal fusion to a constitutive tetramer (RecE)
147 and dimer (GST) respectively (Fig. 1f and Supplementary Fig. 2i-n), confirming their parallel
148 orientation. Thus, α Core- Δ Ntip, α N-tetramer and α C-dimer are parallel coiled-coils, in keeping with
149 the biorientation of SYCP1 molecules within the SC. We conclude that the obligate structure of
150 SYCP1, which provides the minimal building block for self-assembly, is an N-terminal four-helical
151 bundle that bifurcates into C-terminal dimeric coiled-coils of sufficient length to span between SC
152 central and lateral elements (Fig. 1g).

153

154 **SYCP1 N-terminal self-assembly**

155 The α N-tip (residues 101-111) is essential for self-assembly of α Core into large molecular weight
156 species *in vitro* and is part of a short α N-end region (residues 101-206), immediately preceding the
157 α N-tetramer, that is the most highly conserved portion of SYCP1 (Fig. 1b). The X-ray crystal

158 structures of two α N-end constructs (residues 101-206 and 101-175) reveal tetrameric assemblies in
159 which two parallel dimeric coiled-coils interact head-to-head (Fig. 2a,b, Supplementary Fig. 3 and
160 Table 1). The head-to-head interface is mediated entirely by the α N-tip (Fig. 2a,b), suggesting that
161 this 'dimer of dimers' structure may be responsible for SYCP1 N-terminal self-assembly into higher
162 order structures.

163

164 The two α N-end crystal structures demonstrate a common fold in which parallel dimeric coiled-coils
165 splay apart through a wedge formed of W119 and I116 to allow the α N-tips of opposing molecules to
166 interact head-to-head (Figs. 2a,b and 3a,d). The head-to-head interface shows distinct but highly
167 related conformations in the two structures, indicating conformational plasticity. The open
168 conformation of α N-end is asymmetrical and crescent-shaped, formed of midline and lateral anti-
169 parallel coiled-coil interactions (Figs. 2a and 3a-c,g). The closed conformation of truncated α N-end is
170 a symmetrical four-helical bundle, consisting of a hydrophobic core and analogous midline and
171 lateral helical interfaces (Figs. 2b and 3d-f,h). The two conformations are formed of identical amino
172 acids undergoing largely similar coiled-coil and aromatic stacking interactions (Fig. 3b-c,e-h), and
173 likely exist in equilibrium, undergoing conformational change through a rotamer flip of central Y106
174 residues (Fig. 3g,h and Supplementary Fig. 3e). This structural plasticity may be important in
175 enforcing synapsis whilst accommodating large-scale twisting and bending of synapsed meiotic
176 chromosomes, with the open conformation permitting wider angulation between opposing SYCP1
177 molecules than the more rigid closed conformation.

178

179 SYCP1 α Core self-assembly is recapitulated by construct α N (residues 101-362) that includes both
180 α N-end and α N-tetramer (Fig. 4a and Supplementary Fig. 4a,b). Its self-assembly into large
181 molecular weight species is blocked by removal of either sequence, and is retained in the presence
182 of the unstructured N-terminal tail (Fig. 4a,b and Supplementary Fig. 4c-e). Thus, the presence of
183 α N-end and α N-tetramer is necessary and sufficient for SYCP1 N-terminal self-assembly *in vitro*.

184 Mutation of head-to-head interacting residues V105 and L109 to glutamate completely abrogates
185 α N self-assembly into large molecular weight species, leaving a stable obligate tetramer (Fig. 4a).
186 Thus, the α N-end head-to-head interaction is likely responsible for SYCP1 N-terminal self-assembly.
187 We propose that the α N-tetramer provides a structural scaffold from which two α N-end dimers
188 splay apart, with their α N-tips interacting head-to-head with opposing SYCP1 molecules. A staggered
189 configuration provides a simple model for the cooperative assembly of a continuous lattice structure
190 of potentially limitless length, which we propose defines the structural basis of midline SYCP1 N-
191 terminal self-assembly (Fig. 4c).

192

193 Isolated α N-end is monomeric (Fig. 4b), indicating that individual head-to-head interactions are
194 weak and only form when the α N-tetramer mediates lattice formation. This requirement for
195 cooperativity favours the self-assembly of a single continuous lattice between appropriately aligned
196 meiotic chromosomes rather than forming heavily branched unproductive cellular assemblies (Fig.
197 4d).

198

199 ***SYCP1 C-terminal self-assembly***

200 A highly conserved sequence at the C-terminal end of SYCP α Core caps off the α C-dimer parallel
201 coiled-coil (Fig. 1b). The X-ray crystal structure of α C-end (residues 676-770) reveals an anti-parallel
202 tetramer in which two α C-end parallel dimers interact back-to-back in an intertwined α -helical
203 assembly (Fig. 5, Supplementary Fig. 5 and Table 1). We suggest that this α C-end tetrameric
204 assembly provides the structural basis for SYCP1 C-terminal self-assembly on the chromosome axis.

205

206 In solution, α C-end is dimeric at pH 8.0 and tetrameric at pH 5.5 (Fig. 6a and Supplementary Fig.
207 6a,b). SEC-SAXS reveals that both species have similar length, but the cross-sectional radius
208 increases from 7.8 Å to 10.1 Å at pH 5.5, consistent with a transition from dimeric to four-helical
209 coiled-coil (Fig. 6b and Supplementary Fig. 6c-e). SAXS *ab initio* envelopes of the pH 8.0 and pH 5.5

210 species match the dimensions of a dimeric coiled-coil and the α C-end tetramer structure
211 respectively (Fig. 6c,d). SEC-SAXS $P(r)$ distributions of MBP fusions of α C-end at pH 8.0 show inter-
212 MBP peaks at short distances, compatible with their parallel orientation; peaks at long anti-parallel
213 distances were observed only upon MBP fusion at both termini, and for MBP- α C-end upon
214 tetrameric assembly at pH 5.5 (Fig. 6e and Supplementary Fig. 6f-j). Similarly, GST- α C-end forms a
215 stable dimer at pH 8.0 (Supplementary Fig. 6h-k). Finally, a tethered dimer of two consecutive α C-
216 end sequences joined by a flexible linker is dimeric at pH 8.0, with length 241 Å and cross-sectional
217 radius 8.8 Å, consistent with it forming two consecutive dimeric coiled-coils (Figs. 6f,g and
218 Supplementary Fig. 6c-e). It remains dimeric at pH 5.5, but becomes a compact molecule of length
219 156 Å and cross-sectional radius 10.7 Å, indicating the folding back of α C-end sequences into an anti-
220 parallel tetramer (Figs. 6f,g and Supplementary Fig. 6c-e). We conclude that α C-end is a parallel
221 dimeric coiled-coil that undergoes pH-induced back-to-back assembly into the anti-parallel tetramer
222 observed in the crystal structure.

223

224 The α C-end crystal structure has a highly conserved central tetrameric interface in which H717 and
225 Y721 residues (invariant throughout vertebrates) form a hydrophobic core and engage in hydrogen
226 bonding networks with Q720 residues (Figs. 1b and 5a,b, Supplementary Figs. 1a and 5g). The
227 position of H717 residues suggested that their protonation may mediate pH-induced assembly. We
228 introduced mutation H717W Y721F, designed to stabilise the core whilst eliminating pH-sensitivity
229 (Supplementary Fig. 5h), into an extended α C-end construct (residues 676-783) that accentuates the
230 elution difference between dimer and tetramer (Supplementary Fig. 7). H717W Y721F is tetrameric
231 at pH 8.0 (Fig. 6h and Supplementary Fig. 7j), suggesting that pH-induced assembly in wild type
232 involves stabilisation of the core through H717 protonation. Accordingly, mutation H717E blocked
233 pH-induced tetrameric assembly (Supplementary Fig. 7k).

234

235 The central interface leads to pinch points, where N-terminal parallel coiled-coil dimers are flanked
236 by angulated C-terminal chains (Fig. 5a,c,d). The coiled-coil includes C703 heptad interactions that
237 are disulphide and non-disulphide at the respective smoothly and sharply angulated ends of the
238 molecule. An alternative α C-end crystal form contains symmetry-related pinch points with C703
239 partial disulphide bonds and smoothly angulated flanking chains (Supplementary Fig. 5). Whilst
240 disulphide bond formation may be a crystallisation artefact, it may also provide an intriguing means
241 for stabilising assembly *in vivo*; notably, the α N-end head-to-head assembly includes similar heptad
242 interactions between pairs of C183 and C190 residues.

243

244 The ends of the tetrameric structure are formed of four-helical bundles, consisting of a hydrophobic
245 core and anti-parallel coiled-coil interfaces (Fig. 5a,e-g). Hydrophobic core residues outline heptad
246 repeats within N- and C-terminal chains, with the latter constituting a three-heptad leucine zipper
247 (Fig. 5g). These residues likely also mediate parallel coiled-coil interactions in the dimeric
248 conformation. Amino acids L679 and I688 mediate anti-parallel interactions but lie outwith the
249 hydrophobic core heptads, so may be specific for the tetramer. The mutation L679A I688A
250 eliminated tetramer assembly but retained dimer formation (Fig. 6h and Supplementary Fig. 7l). We
251 conclude that heptad residues of the α C-end termini are bifunctional in mediating parallel dimeric
252 and anti-parallel tetrameric interactions, with the conformational change triggered by structural
253 alteration of the protonation-sensitive central interface.

254

255 In the cell, back-to-back assembly of α C-end may be triggered by its concentration on the
256 chromosome axis, through local protonation induced by the high proton density in the close
257 proximity of DNA³¹ or by specific interactions with chromosome axis proteins. Thus, protonation-
258 dependent conformational change of α C-end provides an elegant mechanism for triggering SYCP1 C-
259 terminal self-assembly upon chromosomal recruitment.

260

261 ***DNA binding by SYCP1***

262 The α C-end tetrameric structure contains a series of surface basic patches separated by ~ 30 Å (Fig.
263 7a), suggesting a direct interaction with the DNA backbone. Analysis by electrophoretic mobility shift
264 assay (EMSA) revealed strong double-stranded DNA binding of tetrameric α C-end at pH 5.5, but not
265 of the dimer at pH 8.0 (Fig. 7b). The presence of DNA-binding interfaces on both surfaces of the α C-
266 end tetramer could mediate the formation of large protein-DNA assemblies, possibly accounting for
267 the range of species observed. The α C-end tetrameric conformation is likely stabilised by interaction
268 with DNA, and so SYCP1 C-terminal self-assembly and DNA-binding may be mutually reinforcing.

269

270 How is the SYCP1 C-terminus first recruited to chromosomes? The SYCP1 C-terminal tail contains
271 basic patches that could be obligate DNA-binding sites. A C-terminal construct including both α C-end
272 and Ctail (residues 640-976) interacts with DNA at neutral pH; DNA-binding is dependent on the Ctail
273 and is diminished upon deletion of α C-end (Fig. 7c and Supplementary Fig. 8a-d). We observe slightly
274 enhanced DNA-binding at neutral pH by SYCP1 constructs extended N-terminally to include the α C-
275 dimer and α Core (residues 358-976 and 101-976) (Fig. 7c and Supplementary Fig. 8e-h). Electron
276 microscopy reveals the formation of ~ 10 nm wide protein-DNA complexes by α C-end~Ctail, which
277 develop a wider frayed appearance upon inclusion of the α C-dimer, consistent with transverse
278 filaments emanating from a core protein-DNA structure (Fig. 7d). Finally, we tested DNA-binding of
279 full length SYCP1 using refolded protein that demonstrates α -helical structure and higher order
280 assembly consistent with our findings for α Core (Supplementary Fig. 9). Full length SYCP1 interacts
281 with DNA at neutral pH, and the interaction is disrupted by deletion of the Ctail (Fig. 7e). Together,
282 these data demonstrate that SYCP1 binds DNA through its Ctail and the interaction is enhanced by
283 the α C-end and wider SYCP1 structure.

284

285 We propose that SYCP1 molecules are recruited to meiotic chromosomes through sites within Ctails,
286 leading to their concentration on chromatin. The close proximity of DNA and/or interactions with

287 axis proteins then triggers protonation-induced assembly of α C-end into tetramers that bind DNA
288 and strengthen axis associations. The anti-parallel α C-end tetramers also mediate back-to-back
289 interactions between SYCP1 molecules, which given their known orientation within the SC, likely
290 result in looped U-shaped linkages between adjacent α C-dimer transverse filaments (Fig. 7f). Thus,
291 SYCP1 C-terminal self-assembly integrates DNA-binding and interactions between adjacent
292 transverse filaments to achieve SYCP1 coating of chromosome axes.

293

294 **Discussion**

295 We integrate our crystallographic and biophysical findings into a molecular model for meiotic
296 chromosome synapsis by SYCP1. The SYCP1 core consists of an α N-tetramer that bifurcates into two
297 α C-dimers (Fig. 8a). This tetrameric building-block self-assembles into a supramolecular lattice
298 through its N- and C-terminal ends. In the midline, α N-end dimers splay from α N-tetramer scaffolds
299 and interact head-to-head in a highly cooperative lattice. In the lateral element, α C-end dimers
300 assemble back-to-back as discrete intertwined tetramers that tether together adjacent α C-dimer
301 transverse filaments and reinforce chromosomal associations of C-terminal tails. Together, N- and C-
302 terminal self-assembly collaborate to generate a cooperative zipper-like supramolecular lattice of
303 SYCP1 molecules capable of mediating continuous synapsis between homologous chromosomes (Fig.
304 8b). During SC assembly, midline lattice formation and chromosomal recruitment likely occur
305 concomitantly in one dynamic process of progressive chromosome synapsis. Whilst we cannot
306 exclude additional roles for N- and C-terminal tails in SYCP1 assembly *in vivo*, these regions are
307 largely unstructured and have no effect on oligomer states *in vitro*. Conserved amino acid sequences
308 within C-terminal tails may mediate currently unidentified interactions with chromosome axis
309 proteins, which act in concert with direct DNA-binding to achieve meiotic chromosome recruitment
310 of SYCP1.

311

312 Our model for SYCP1 self-assembly is consistent with the dimensions of the native SC. The SYCP1
313 tetrameric core has a length of 900 Å, sufficient to span just over half of the inter-chromosomal
314 distance. The α C-dimer has an 8.9 Å cross-sectional radius and 645 Å length, matching the
315 dimensions of individual transverse filaments measured by electron microscopy in the hamster SC¹⁰.

316 We propose that α C-dimers constitute the transverse filaments visualised spanning between central
317 and lateral elements, with α N-tetramers buried within the central element. Importantly, anti-parallel
318 tetramer formation by α C-end explains how parallel SYCP1 molecules interact back-to-back to
319 achieve the well-established biorientation of SYCP1 N- and C-termini within the SC^{12,14,15}. A recent

320 study reported that a region similar to α C-end is an anti-parallel dimer³², incompatible with
321 established localisation patterns. Examination of their structural data (pdb 4YTO) reveals the
322 presence of an anti-parallel tetramer within the crystal lattice, indicating that the anti-parallel dimer
323 of the asymmetric unit was incorrectly attributed as the biological molecule (Supplementary Figure
324 5i).

325

326 The three-dimensional SC assembly contains at least two layers of transverse filament proteins^{11,12,16},
327 which is compatible with the SYCP1 supramolecular assembly that we describe. We propose that
328 two parallel head-to-head SYCP1 lattices are connected by vertically (or obliquely) orientated back-
329 to-back assemblies within lateral elements (Supplementary Fig. 10a). This model is consistent with
330 the observed vertical separation of SYCP1 N-termini by up to 100 nm, and the presence of single
331 tracks of SYCP1 C-termini within lateral elements^{12,16}.

332

333 How is SYCP1 self-assembly directed to occur predominantly between aligned chromosomes? Whilst
334 SYCP1 can form chromatin-free polycomplexes in meiotic tissue³³, assembly into SCs is heavily
335 favoured. Two distinct mechanisms cooperate to favour timely SYCP1 self-assembly between aligned
336 chromosomes. Firstly, α N-end head-to-head interactions are individually weak and thus the prior
337 accumulation of juxtaposed SYCP1 molecules between aligned chromosomes may nucleate its
338 cooperative supramolecular assembly. Secondly, α C-end self-assembly occurs through a
339 protonation-induced conformational change triggered by the proton density in the immediate
340 vicinity of DNA³¹ and/or axis protein interactions, thereby coupling assembly to chromosomal
341 recruitment.

342

343 The nascent synapsis generated by SYCP1 self-assembly is stabilised and matured into a full SC
344 through assembly of central element proteins SYCE3, SYCE1, SIX6OS1 and SYCE2-TEX12^{16,22,23,25-29}.

345 Their recruitment is dependent on SYCP1 and is essential for the tripartite structure and meiotic

346 function of the SC^{22,23,25-28}. Central element assembly likely occurs concomitantly with SYCP1 self-
347 assembly, rapidly converting the underpinning SYCP1 structural framework into a mature SC. Initial
348 SYCP1 assemblies recruit central element proteins to stabilise the nascent lattice, enabling its
349 growth, and providing a mutually reinforcing cycle that results in full synapsis (Fig. 8c). Central
350 element proteins may provide vertical and longitudinal supports between α N-tetramers that rigidify
351 SYCP1 hemi-lattices and orientate α N-end sites for long-range cooperative head-to-head assembly
352 (Fig. 8d and Supplementary Fig. 10b). They may further act as transverse bridges that connect hemi-
353 lattices across the midline to directly reinforce α N-end head-to-head interactions. Initiation factors
354 SYCE3, SYCE1 and SIX6OS1 may act as transverse bridges and vertical supports²²⁻²⁵, whilst SYCE2-
355 TEX12 may provide longitudinal supports that enable SC elongation²⁶⁻²⁹. This results in a mature SC in
356 which an underlying SYCP1 lattice is structurally supported by the central element. The true
357 molecular roles of SC central element proteins will be revealed upon their structure elucidation, and
358 it will be intriguing to see whether they simply dock onto the SYCP1 lattice or induce structural
359 remodelling upon recruitment.

360

361 Whilst SYCP1 chromosome axis recruitment is retained upon disruption of SC lateral element
362 proteins, synapsis is discontinuous, indicating that chromosome axis structure facilitates the efficient
363 loading of SYCP1 necessary for continuous synapsis^{17,18,20}. This may occur through positioning
364 chromatin loops to achieve a regular spacing of SYCP1 molecules that is compatible with long-range
365 lattice formation. SYCP1 loading may similarly be regulated by the underlying chromatin structure.
366 For example, if both surfaces of α C-end tetramers interact with DNA, they may sit between adjacent
367 nucleosomes and would be spaced apart by the 11 nm nucleosome diameter.

368

369 How is the SC supramolecular structure efficiently disassembled following its function in meiosis?
370 SYCP1 self-assembly is intrinsic to the protein sequence and hence independent of post-translational
371 modifications, but phosphorylation has been implicated in SC disassembly³⁴. Whilst there are no

372 clear candidate sites within SYCP1 α N-end or α C-end, phosphorylation of the numerous predicted
373 sites within the C-terminal tail could destabilise axis assembly. Similarly, central element protein
374 phosphorylation could destabilise SYCP1 midline lattice assembly. The molecular features of α N-end
375 and α C-end that achieve cooperative assembly may facilitate the continuous turnover of SYCP1
376 molecules within the SC. Whilst dynamic interchange will normally lead to continual renewal of the
377 SYCP1 lattice, phosphorylation-induced destabilisation of self-assembly sites would shift the balance
378 towards a net loss of molecules and ultimately disassembly.

379

380 SYCP1 fulfils the classic functions of coiled-coil proteins in acting as molecular spacers that scaffold
381 supramolecular assemblies and separate functional units³⁵. SYCP1 imposes a 100 nm synapsis
382 between homologous chromosomes, raising the question of why it is necessary to impose an
383 evolutionarily conserved separation between homologues? This distance may be optimal for the
384 maintenance and resolution of meiotic recombination intermediates, and so an answer may lie in
385 differences in recombination in the few meiotic organisms that lack an SC¹. Interestingly, the SC
386 central region and central element are approximately 10% narrower in female mice than in males³⁶.
387 This variation can be accommodated by the SYCP1 lattice that we propose through alteration in
388 angulation of α N-end assemblies and α C-dimers (Supplementary Fig. 10c). Furthermore, midline
389 angulation and SC central region width are determined by the frequency of α C-end binding to
390 chromosomes; thus, midline variation could originate from differences in chromosomal axis
391 structure between sexes.

392

393 Despite evolutionary conservation of the SC ultrastructure, its constituent protein sequences are
394 divergent between vertebrates and lower eukaryotes². Nevertheless, yeast transverse filament
395 protein Zip1 is approximately the same size as SYCP1 and displays similar patterns of conservation
396 and structure prediction. Thus, it is possible that Zip1 adopts a similar structure and self-assembly
397 mechanism through underlying structural conservation.

398

399 The molecular functions of the SC in recombination, crossover formation and interference remain
400 unknown. Nevertheless, we speculate that its three-dimensional structure may direct these
401 processes by regulating enzymatic access to recombination sites. The ability of coiled-coil proteins to
402 transmit conformational changes recursively may further enable the SC to communicate signals
403 along synapsed chromosomes. As our understanding of the SC structure deepens, its molecular
404 functions will gradually be uncovered, ultimately leading to a complete mechanistic understanding
405 of recombination and crossover formation within the functional architecture of the SC.

406

407 **Accession codes**

408 Crystallographic structure factors and atomic co-ordinates have been deposited in the Protein Data
409 Bank (PDB) under accession numbers 6F5X, 6F62, 6F63 and 6F64.

410

411 **Acknowledgements**

412 O.R.D. thanks D.W. Sinclair for scientific inspiration and support. We thank Diamond Light Source
413 and the staff of beamlines I02, I04-1 and B21 (proposals mx9948, mx13587, sm14435, sm15580,
414 sm15897 and sm15836). We thank A. Baslé and H. Waller for assistance with X-ray crystallographic
415 and CD data collection, and L. J. Salmon and V. A. Jatikusumo for work in the early stages of this
416 project. I.U. is funded by grants BIO2015-64216-P and MDM2014-0435-01 (MINECO, Spanish
417 Ministry of Economy and Competitiveness). C.M. is supported by a MINECO BES-2015-071397
418 scholarship associated to the Structural Biology Maria de Maeztu Unit of Excellence. S.M. is
419 supported by a Wellcome Trust Career Re-entry Fellowship (062376). O.R.D. is a Sir Henry Dale
420 Fellow jointly funded by the Wellcome Trust and Royal Society (Grant Number 104158/Z/14/Z).

421

422 **Author contributions**

423 J.M.D. performed the majority of biochemical, biophysical and crystallographic experiments. O.M.D.
424 performed SAXS experiments and analyses. M.R. crystallised truncated α N-end and analysed α N
425 constructs. C.M. and I.U. solved the α C-end crystal structures. O.R.D. solved the α N-end crystal
426 structures and built/refined all structures. S.M. assisted with initial experiments. J.M.D. and O.R.D.
427 designed experiments; O.R.D. wrote the manuscript.

428

429 **Competing financial interests**

430 The authors declare no competing interests.

431

432 **References**

- 433 1. Zickler, D. & Kleckner, N. Recombination, Pairing, and Synapsis of Homologs during Meiosis.
434 *Cold Spring Harb Perspect Biol* **7**(2015).
- 435 2. Page, S.L. & Hawley, R.S. The genetics and molecular biology of the synaptonemal complex.
436 *Annu Rev Cell Dev Biol* **20**, 525-58 (2004).
- 437 3. Hunter, N. Meiotic Recombination: The Essence of Heredity. *Cold Spring Harb Perspect Biol*
438 **7**(2015).
- 439 4. Baudat, F., Imai, Y. & de Massy, B. Meiotic recombination in mammals: localization and
440 regulation. *Nat Rev Genet* **14**, 794-806 (2013).
- 441 5. de Vries, F.A. et al. Mouse Sycp1 functions in synaptonemal complex assembly, meiotic
442 recombination, and XY body formation. *Genes Dev* **19**, 1376-89 (2005).
- 443 6. Kouznetsova, A., Benavente, R., Pastink, A. & Hoog, C. Meiosis in mice without a
444 synaptonemal complex. *PLoS One* **6**, e28255 (2011).
- 445 7. Handel, M.A. & Schimenti, J.C. Genetics of mammalian meiosis: regulation, dynamics and
446 impact on fertility. *Nat Rev Genet* **11**, 124-36 (2010).
- 447 8. Nagaoka, S.I., Hassold, T.J. & Hunt, P.A. Human aneuploidy: mechanisms and new insights
448 into an age-old problem. *Nat Rev Genet* **13**, 493-504 (2012).
- 449 9. Westergaard, M. & von Wettstein, D. The synaptonemal complex. *Annu Rev Genet* **6**, 71-110
450 (1972).
- 451 10. Solari, A.J. & Moses, M.J. The structure of the central region in the synaptonemal complexes
452 of hamster and cricket spermatocytes. *J Cell Biol* **56**, 145-52 (1973).
- 453 11. Schmekel, K., Skoglund, U. & Daneholt, B. The three-dimensional structure of the central
454 region in a synaptonemal complex: a comparison between rat and two insect species,
455 *Drosophila melanogaster* and *Blaps cribrosa*. *Chromosoma* **102**, 682-92 (1993).
- 456 12. Schucker, K., Holm, T., Franke, C., Sauer, M. & Benavente, R. Elucidation of synaptonemal
457 complex organization by super-resolution imaging with isotropic resolution. *Proc Natl Acad*
458 *Sci U S A* **112**, 2029-33 (2015).
- 459 13. Meuwissen, R.L. et al. A coiled-coil related protein specific for synapsed regions of meiotic
460 prophase chromosomes. *EMBO J* **11**, 5091-100 (1992).
- 461 14. Liu, J.G. et al. Localization of the N-terminus of SCP1 to the central element of the
462 synaptonemal complex and evidence for direct interactions between the N-termini of SCP1
463 molecules organized head-to-head. *Exp Cell Res* **226**, 11-9 (1996).
- 464 15. Schmekel, K. et al. Organization of SCP1 protein molecules within synaptonemal complexes
465 of the rat. *Exp Cell Res* **226**, 20-30 (1996).
- 466 16. Hernandez-Hernandez, A. et al. The central element of the synaptonemal complex in mice is
467 organized as a bilayered junction structure. *J Cell Sci* **129**, 2239-49 (2016).
- 468 17. Yang, F. et al. Mouse SYCP2 is required for synaptonemal complex assembly and
469 chromosomal synapsis during male meiosis. *J Cell Biol* **173**, 497-507 (2006).
- 470 18. Yuan, L. et al. The murine SCP3 gene is required for synaptonemal complex assembly,
471 chromosome synapsis, and male fertility. *Mol Cell* **5**, 73-83 (2000).
- 472 19. Syrjanen, J.L., Pellegrini, L. & Davies, O.R. A molecular model for the role of SYCP3 in meiotic
473 chromosome organisation. *Elife* **3**(2014).
- 474 20. Yuan, L. et al. Female germ cell aneuploidy and embryo death in mice lacking the meiosis-
475 specific protein SCP3. *Science* **296**, 1115-8 (2002).
- 476 21. Syrjanen, J.L. et al. Single-molecule observation of DNA compaction by meiotic protein
477 SYCP3. *Elife* **6**(2017).
- 478 22. Schramm, S. et al. A novel mouse synaptonemal complex protein is essential for loading of
479 central element proteins, recombination, and fertility. *PLoS Genet* **7**, e1002088 (2011).

- 480 23. Bolcun-Filas, E. et al. Mutation of the mouse Syce1 gene disrupts synapsis and suggests a link
481 between synaptonemal complex structural components and DNA repair. *PLoS Genet* **5**,
482 e1000393 (2009).
- 483 24. Costa, Y. et al. Two novel proteins recruited by synaptonemal complex protein 1 (SYCP1) are
484 at the centre of meiosis. *J Cell Sci* **118**, 2755-62 (2005).
- 485 25. Gomez, H.L. et al. C14ORF39/SIX6OS1 is a constituent of the synaptonemal complex and is
486 essential for mouse fertility. *Nat Commun* **7**, 13298 (2016).
- 487 26. Bolcun-Filas, E. et al. SYCE2 is required for synaptonemal complex assembly, double strand
488 break repair, and homologous recombination. *Journal of Cell Biology* **176**, 741-747 (2007).
- 489 27. Hamer, G. et al. Characterization of a novel meiosis-specific protein within the central
490 element of the synaptonemal complex. *Journal of Cell Science* **119**, 4025-4032 (2006).
- 491 28. Hamer, G. et al. Progression of meiotic recombination requires structural maturation of the
492 central element of the synaptonemal complex. *J Cell Sci* **121**, 2445-51 (2008).
- 493 29. Davies, O.R., Maman, J.D. & Pellegrini, L. Structural analysis of the human SYCE2–TEX12
494 complex provides molecular insights into synaptonemal complex assembly. *Open Biology* **2**,
495 120099 (2012).
- 496 30. Ollinger, R., Alsheimer, M. & Benavente, R. Mammalian protein SCP1 forms synaptonemal
497 complex-like structures in the absence of meiotic chromosomes. *Mol Biol Cell* **16**, 212-7
498 (2005).
- 499 31. Hanlon, S., Wong, L. & Pack, G.R. Proton equilibria in the minor groove of DNA. *Biophys J* **72**,
500 291-300 (1997).
- 501 32. Seo, E.K., Choi, J.Y., Jeong, J.H., Kim, Y.G. & Park, H.H. Crystal Structure of C-Terminal Coiled-
502 Coil Domain of SYCP1 Reveals Non-Canonical Anti-Parallel Dimeric Structure of Transverse
503 Filament at the Synaptonemal Complex. *PLoS One* **11**, e0161379 (2016).
- 504 33. Goldstein, P. Multiple synaptonemal complexes (polycomplexes): origin, structure and
505 function. *Cell Biol Int Rep* **11**, 759-96 (1987).
- 506 34. Cahoon, C.K. & Hawley, R.S. Regulating the construction and demolition of the
507 synaptonemal complex. *Nat Struct Mol Biol* **23**, 369-77 (2016).
- 508 35. Truebestein, L. & Leonard, T.A. Coiled-coils: The long and short of it. *Bioessays* **38**, 903-16
509 (2016).
- 510 36. Agostinho, A. et al. Sexual dimorphism in the width of the mouse synaptonemal complex. *J*
511 *Cell Sci* **131**(2018).
- 512

513 **Figure Legends**

514 **Figure 1**

515 **The obligate tetrameric structure of SYCP1.**

516 (a) SYCP1 molecules are bioriented within the synaptonemal complex (SC), with midline N-termini
517 and chromosome-bound C-termini, providing a 100 nm separation between chromosome axes. (b)
518 Sequence analysis of SYCP1 demonstrating the presence of an α -helical core (amino acids 101-783)
519 that is highly conserved at both ends, flanked by unstructured N- and C-terminal tails. Amino acid
520 conservation was calculated amongst vertebrate sequences. The principal protein constructs
521 analysed in this study are indicated along with their amino acid boundaries. An extensive summary
522 of SYCP1 constructs is provided in Supplementary Fig. 1a and biophysical data are compiled in
523 Supplementary Table 1. (c) SEC-MALS analysis; light scattering (LS) and differential refractive index
524 (dRI) are shown as solid and dashed lines respectively, with fitted molecular weights (Mw) plotted as
525 diamonds across elution peaks. SYCP1 α Core (101-783) forms large molecular species of 1-12 MDa,
526 whereas α Core- Δ Ntip (112-783) is a 306 kDa tetramer (theoretical tetramer – 320 kDa) consisting of
527 a 68 kDa α N-tetramer (theoretical tetramer – 76 kDa) and 97 kDa α C-dimer (theoretical dimer – 101
528 kDa). (d) SEC-SAXS $P(r)$ distributions of α Core- Δ Ntip, α N-tetramer and α C-dimer; maximum
529 dimensions (D_{max}) and cross-sectional radii (R_c) are indicated. (e) SEC-SAXS $P(r)$ distributions of
530 MBP- α Core- Δ Ntip, MBP- α N-tetramer, MBP- α C-dimer and MBP; intra-MBP and inter-MBP peaks are
531 indicated. (f) SEC-MALS analysis showing that RecE- α N-tetramer is a 208 kDa tetramer (theoretical
532 tetramer- 214 kDa) and GST- α C-dimer is a 157 kDa dimer (theoretical dimer - 160 kDa). (g) Model of
533 the SYCP1 obligate unassembled structure. The SYCP1 α -helical core has a parallel organisation and
534 consists of a 260 Å α N-tetramer that bifurcates into two 645 Å α C-dimer coiled-coils.

535

536 **Figure 2**

537 **Crystal structures of the SYCP1 α N-end head-to-head assembly in open and closed conformations.**

538 (a) Crystal structure of SYCP1 α N-end (101-206) demonstrating head-to-head 'dimer of dimers'
539 assembly of two eleven heptad parallel coiled-coils, spanning a total length of 288 Å. The long
540 dimeric coiled-coils are interrupted by a wedge-like structure that splays apart the two α -helices to
541 enable their α N-tip sites to mediate midline head-to-head assembly in an open conformation. The
542 head-to-head interface provides 1,990 Å² buried surface area in addition to 4,520 Å² for each coiled-
543 coil dimer alone. (b) Crystal structure of truncated SYCP1 α N-end (101-175) demonstrating a similar
544 head-to-head 'dimer of dimers' assembly of two seven heptad parallel coiled-coils, spanning 194 Å,
545 with α N-tips undergoing head-to-head assembly in a closed conformation. The head-to-head
546 interface provides 2,950 Å² buried surface area in addition to the 2,210 Å² for each coiled-coil dimer
547 alone. C-terminal interactions of α N-end-truncated chains within the crystal lattice were determined
548 to be artefactual owing to their absence in the α N-end structure and through *in vitro* mutagenesis
549 experiments (M.R. and O.R.D., unpublished data).

550

551 **Figure 3**

552 **Head-to-head assembly interfaces of SYCP1 α N-end.**

553 (a-c) Crystal structure of SYCP1 α N-end (101-206). (a) A wedge structure formed of residues I116 and
554 W119 splays apart coiled-coil α -helices to enable their head-to-head assembly. (b-c) The open
555 assembly is formed of one midline and two lateral interfaces. (b) The midline interface (open) is an
556 anti-parallel coiled-coil between symmetry-related chain A copies, with heptad residues L102, V105,
557 L109 and E112. (c) The lateral interface (open) is an anti-parallel association of unique chains A and
558 B, formed of coiled-coil and aromatic stacking interactions of residues L102, L109, Y106 and Y110.
559 (d-f) Crystal structure of truncated SYCP1 α N-end (101-175). (d) Similar to α N-end, a wedge
560 structure of residues I116 and W119 splays apart α -helices to enable their head-to-head assembly.
561 (e-f) The closed assembly is formed of a hydrophobic core and interfaces that are analogous to the
562 midline and lateral interfaces of the open conformation. (e) The midline interface (closed) is an anti-
563 parallel coiled-coil of heptad residues G101, V105, K108 and E112. (f) The lateral interface (closed) is

564 anti-parallel, consisting of coiled-coil and interlaced aromatic stacking interactions of residues L102,
565 L109, Y106 and Y110. **(g-h)** Cross-sections through the α N-end head-to-head open and closed
566 conformations. **(g)** The open conformation contains no hydrophobic core and is asymmetrical in
567 nature, with midline chain A copies flanked by two copies of chain B. **(h)** The closed conformation is
568 formed of symmetry-related chains and contains a hydrophobic core of residues L102, L109 and
569 I116.

570

571 **Figure 4**

572 **SYCP1 N-terminal self-assembly into higher order structures is mediated by α N-end head-to-head**
573 **interactions.**

574 **(a-b)** SEC-MALS analysis. **(a)** SYCP1 α N (101-362) (black, left) forms large molecular species of 3-25
575 MDa, whereas α N- Δ Ntip (112-362) (grey) and α N (101-362) V105E L109E (black, right) form
576 tetramers of 118 kDa and 121 kDa respectively (theoretical tetramers – 121 kDa and 126 kDa). **(b)**
577 SYCP1 α N-end, truncated α N-end and α N-end~Ntail are monomeric species of 15 kDa, 10 kDa and
578 23 kDa respectively (theoretical monomers - 13 kDa, 9 kDa and 20 kDa). **(c)** Model of SYCP1 N-
579 terminal self-assembly. SYCP1 α N-ends splay from α N-tetramers and interact head-to-head in the
580 midline to create a continuous lattice-like assembly. **(d)** SYCP1 N-terminal self-assembly is predicted
581 to be highly cooperative, enabling stable structure formation through a series of individually weak
582 head-to-head associations. This allows transient chromosome associations to be formed and
583 remodelled rapidly, ultimately favouring a single continuous assembly between aligned chromosome
584 axes.

585

586 **Figure 5**

587 **Crystal structure of the SYCP1 C-terminal tetrameric assembly.**

588 **(a)** Crystal structure of SYCP1 α C-end (676-770) in crystal form 1, demonstrating an anti-parallel
589 tetrameric assembly of length 142 Å. The structure includes a central tetrameric interface flanked by

590 C703 pinch points that lead to lateral four-helical bundles. N- and C-termini are coloured in green
591 and red respectively. **(b)** The central tetrameric interface consists of two stacked layers each
592 containing a hydrogen bonding network of pairs of H717, Q720 and Y721 residues. **(c-d)** The C703
593 pinch point consists of a parallel dimeric coiled-coil (containing C703) flanked by surrounding anti-
594 parallel chains. **(c)** The parallel dimeric coiled-coil is formed of heptad residues D700, C703, I707,
595 M710 and M714 **(d)** The flanking chains have a distinct angulation at E731 and provide pseudo-cores
596 of loose anti-parallel interactions. **(e-g)** The lateral four-helical bundle (4HB) is formed of a
597 hydrophobic core and anti-parallel interfaces. **(e)** The lateral 4HB type 1 interface is an anti-parallel
598 coiled-coil of heptad residues L679, V682 and K686, L753, K757, L760 and K764. **(f)** The lateral 4HB
599 type 2 interface is an anti-parallel coiled-coil of heptad residues L678, E681, A685, I688 and A692,
600 L745, L749, E752, L756 and E759. **(g)** Cross-section through the lateral 4HB assembly. A hydrophobic
601 core is formed from residues that also contribute to 4HB anti-parallel interfaces and are predicted to
602 mediate the formation of N- and C-terminal parallel dimeric coiled-coils in the non-assembled
603 conformation. L679 and I688 are the only hydrophobic 4HB residues not also implicated in the
604 putative parallel dimeric coiled-coil structure.

605

606 **Figure 6**

607 **SYCP1 α C-end undergoes pH-induced assembly into an anti-parallel tetramer.**

608 **(a)** SEC-MALS analysis. SYCP1 α C-end (676-770) is a 22 kDa dimer at pH 8.0 (grey) (theoretical dimer
609 – 23 kDa) and a 43 kDa tetramer at pH 5.5 (black) (theoretical tetramer – 46 kDa). **(b)** SEC-SAXS P(r)
610 distributions of α C-end at pH 5.5 (black) and pH 8.0 (grey). **(c-d)** SAXS *ab initio* models of the
611 tetrameric and dimeric conformations of SYCP1 α C-end (676-770) at **(c)** pH 5.5 and **(d)** pH 8.0.
612 Averaged models were generated from 20 independent DAMMIF runs with NSD values 0.527 (\pm
613 0.014) and 0.513 (\pm 0.014), and reference model χ^2 values 1.81 and 1.49. The α C-end tetrameric
614 crystal structure and a theoretical dimeric coiled-coil were docked into the respective envelopes. **(e)**
615 SEC-SAXS P(r) distributions of N-terminal, C-terminal and both N- and C-terminal MBP fusions of α C-

616 end at pH 8.0, alongside MBP- α C-end at pH 5.5. (f) SEC-MALS analysis reveals that α C-end tethered
617 dimer forms dimers of 44 kDa and 43 kDa (theoretical dimer of dimers – 47 kDa) at pH 5.5 (black,
618 right) and pH 8.0 (black, left), with an increase in elution volume at pH 5.5. The α C-end (single chain)
619 tetramer at pH 5.5 is shown in grey. (g) SEC-SAXS P(r) distributions of the α C-end tethered dimer at
620 pH 5.5 (black) and pH 8.0 (grey). (h) SEC-MALS analysis of SYCP1 α C-end extended (676-783) point
621 mutants. H717W Y721F (black) forms 40 kDa tetramers at pH 8.0 (theoretical tetramer – 52 kDa).
622 L679A I688A (grey) fails to undergo pH-induced assembly and remains mostly as a 26 kDa dimer
623 (theoretical dimer – 26 kDa) at pH 4.6.

624

625 **Figure 7**

626 **DNA binding by SYCP1.**

627 (a) Surface electrostatic potential of the SYCP1 α C-end crystal structure (red – electronegative; blue
628 – electropositive). The flat surface of the α C-end structure contains five demarcated basic patches
629 that are separated by approximately 30 Å. (b) EMSA analysing the ability of SYCP1 α C-end (676-770)
630 to interact with linear double-stranded DNA (dsDNA) at pH 8.0 (top) and pH 5.5 (bottom).
631 Uncropped gel images are shown in Supplementary Data Set 1 (c) EMSA of MBP fusions of α C-
632 end~Ctail (640-976), Ctail (784-976), α C-dimer~Ctail (358-976) and α Core~Ctail (101-976) with linear
633 dsDNA at pH 8.0. (d) Electron microscopy (EM) analysis of MBP fusions of α C-dimer~Ctail (358-976)
634 and α C-end~Ctail (640-976) in complex with plasmid dsDNA. Scale bars, 50 nm. (e) EMSA of refolded
635 full length SYCP1 (1-976) and Δ Ctail (1-783) with linear dsDNA at pH 8.0. (f) Model of SYCP1
636 chromosomal axis assembly. SYCP1 molecules are initially recruited to chromosomes through Ctail
637 DNA-binding sites. The close proximity of DNA and/or interactions with chromosome axis proteins
638 then triggers protonation-induced assembly of α C-ends into anti-parallel tetramers that bind DNA
639 and thereby reinforce Ctail interactions. This results in the complete coating of the chromosome axis
640 with SYCP1 molecules linked together through U-shaped assemblies that are anchored to
641 chromosomal DNA.

642

643 **Figure 8**

644 **Meiotic chromosome synapsis through SYCP1 self-assembly.**

645 (a) Model of the SYCP1 obligate unassembled structure. The α Core consists of a parallel α N-tetramer
646 that splays into two α C-dimers. The α N-tetramer splays at its N-terminus into α N-end self-assembly
647 sites that lead to unstructured Ntails. The α C-dimers terminate as α C-end self-assembly sites,
648 leading to unstructured Ctails that contain DNA-binding sequences. (b) Model of chromosome
649 synapsis by SYCP1. The bifurcating SYCP1 α Core presents pairs of α N-end and α C-end self-assembly
650 sites in the midline and chromosome axis respectively. α N-end sites undergo head-to-head assembly
651 through their α N-tips to provide zipper-like associations that mediate synapsis of SYCP1-coated
652 homologous chromosomes. α C-end sites undergo back-to-back assembly into tetrameric structures
653 that bind directly to DNA within the lateral element and reinforce axis associations of Ctails.
654 Together, these distinct mechanisms of SYCP1 self-assembly generate a supramolecular lattice
655 between meiotic chromosome pairs. (c) Concomitant and mutually reinforcing assembly of SYCP1
656 and central element proteins in SC formation. Initial SYCP1 contacts trigger central element
657 recruitment, enabling growth of the SYCP1 assembly, extending the central element and thereby
658 enabling further SYCP1 growth. (d) Model of the mature SC. The central element may provide
659 vertical and longitudinal supports between SYCP1 α N-tetramers to rigidify SYCP1 hemi-lattices and
660 orientate α N-ends for long range cooperative head-to-head assembly. They may also act as
661 transverse bridges that provide direct connections across the midline to reinforce SYCP1 head-to-
662 head interactions. This leads to the formation of a mature SC in which an underlying SYCP1 lattice is
663 stabilised and extended across long distances through central element assembly.

664

665

Table 1 Data collection and refinement statistics

| | SYCP1 α N-end 101-206 Open conformation (PDB 6F62) | SYCP1 α N-end truncated 101-175 Closed conformation (PDB 6F5X) | SYCP1 α C-end 676-770 Crystal form 1 (PDB 6F63) | SYCP1 α C-end 676-770 Crystal form 2 (PDB 6F64) |
|---|--|--|---|---|
| Data collection | | | | |
| Space group | I2 | I222 | C2 | I4 ₁ 22 |
| Cell dimensions | | | | |
| <i>a</i> , <i>b</i> , <i>c</i> (Å) | 65.67, 37.31, 108.52 | 28.64, 39.38, 165.77 | 233.42, 42.85, 43.69 | 43.38, 43.38, 292.18 |
| α , β , γ (°) | 90.00, 106.66, 90.00 | 90.00, 90.00, 90.00 | 90.00, 93.61, 90.00 | 90.00, 90.00, 90.00 |
| Wavelength (Å) | 0.9282 | 1.7712 | 0.9795 | 0.9795 |
| Resolution (Å) | 34.87–2.06 (2.12–2.06) ^a | 41.44–1.91 (1.95–1.91) ^a | 116.48–2.15 (2.27–2.15) ^a | 42.91–2.48 (2.58–2.48) ^a |
| <i>R</i> _{merge} | 0.071 (0.919) | 0.028 (0.678) | 0.052 (0.695) | 0.080 (2.567) |
| <i>R</i> _{pim} | 0.023 (0.286) | 0.017 (0.541) | 0.032 (0.429) | 0.023 (0.727) |
| <i>I</i> / σ (<i>I</i>) | 15.0 (1.8) | 27.9 (1.8) | 12.4 (1.9) | 14.8 (1.5) |
| <i>CC</i> _{1/2} | 0.999 (0.969) | 1.000 (0.839) | 0.998 (0.872) | 1.000 (0.935) |
| Completeness (%) | 99.9 (100.0) | 99.3 (92.1) | 97.4 (88.0) | 99.8 (99.7) |
| Redundancy | 11.0 (11.4) | 5.9 (3.7) | 3.6 (3.5) | 13.2 (13.2) |
| Refinement | | | | |
| Resolution (Å) | 27.23–2.07 | 41.44–1.91 | 58.26–2.15 | 39.63–2.49 |
| UCLA anisotropy (Å) | 2.1, 2.1, 2.6 | 1.9, 2.0, 2.1 | 2.2, 2.3, 2.2 | 2.9, 2.9, 2.5 |
| No. reflections | 12467 | 6754 | 21416 | 4138 |
| <i>R</i> _{work} / <i>R</i> _{free} | 0.2264/0.2441 | 0.2272/0.2392 | 0.2186/0.2526 | 0.2251/0.2517 |
| No. atoms | 1866 | 677 | 3318 | 806 |
| Protein | 1744 | 633 | 3143 | 786 |
| Ligand/ion | 18 | 12 | 0 | 4 |
| Water | 104 | 32 | 175 | 16 |
| <i>B</i> -factors | 42.79 | 58.4 | 46.97 | 60.86 |
| Protein | 42.51 | 57.7 | 47.30 | 60.80 |
| Ligand/ion | 62.91 | 83.9 | N/A | 81.59 |
| Water | 44.02 | 61.7 | 41.20 | 58.92 |
| R.m.s. deviations | | | | |
| Bond lengths (Å) | 0.002 | 0.009 | 0.004 | 0.004 |
| Bond angles (°) | 0.334 | 1.020 | 0.511 | 0.575 |

^aValues in parentheses are for highest-resolution shell.

666 **Online Methods**

667 **Recombinant protein expression and purification**

668 Sequences corresponding to regions of human SYCP1 were cloned into pHAT4, pGAT3 or pMAT11
669 vectors for expression as TEV-cleavable N-terminal His₆-, His₆-GST or His₆-MBP fusion proteins
670 respectively. A list of protein constructs, including sequence boundaries, is provided in
671 Supplementary Fig. 1a and Supplementary Table 1. Constructs were expressed in BL21 (DE3) cells
672 (Novagen®), in 2xYT media, induced with 0.5 mM IPTG for 16 hours at 25°C. Cells were lysed by
673 sonication in 20 mM Tris pH 8.0, 500 mM KCl, and fusion proteins were purified from clarified lysate
674 through consecutive Ni-NTA (Qiagen), amylose (NEB) or glutathione sepharose (GE Healthcare), and
675 HiTrap Q HP (GE Healthcare) ion exchange chromatography. Affinity tags were removed by
676 incubation with TEV protease and cleaved samples were purified by HiTrap Q HP ion exchange
677 chromatography and size exclusion chromatography (HiLoad™ 16/600 Superdex 200, GE Healthcare)
678 in 20 mM Tris pH 8.0, 150 mM KCl, 2 mM DTT. Protein samples were concentrated using Amicon
679 Ultra® 10,000 MWCO centrifugal filter units (Millipore), and were stored at -80°C following flash-
680 freezing in liquid nitrogen. Protein samples were analysed by SDS-PAGE with Coomassie staining,
681 and concentrations were determined by UV spectroscopy using a Cary 60 UV spectrophotometer
682 (Agilent) with extinction coefficients and molecular weights calculated by ProtParam
683 (<http://web.expasy.org/protparam/>).

684

685 **Purification of refolded full length SYCP1**

686 Full-length human SYCP1 (amino acids 1-976) was expressed using a pHAT4 vector in Rosetta (DE3)
687 cells, grown in 2xYT media and induced with 0.5 mM IPTG for 3 hours at 37°C. Cells were lysed by
688 sonication in 20 mM Tris pH 8.0, 500 mM NaCl and the insoluble fraction pelleted through
689 centrifugation at 40,000 g for 30 minutes. The resultant pellet was washed in 20 mM Tris pH 8.0, 500
690 mM NaCl prior to solubilisation in 20 mM Tris pH 8.0, 500 mM NaCl, 8M urea pH 8.0. DNA-containing
691 hydrogels were formed by consecutive dialysis into 20 mM Tris pH 8.0, 500 mM NaCl, 500 mM L-

692 arginine pH 8.0, followed by 20 mM Tris pH 8.0, 500 mM NaCl. Soluble SYCP1 was produced by
693 removal of DNA from the denatured material through ion exchange chromatography, prior to the
694 refolding protocol through dialysis, as described above.

695

696 **Circular dichroism (CD) spectroscopy**

697 Far UV circular dichroism (CD) spectroscopy data were collected on a Jasco J-810 spectropolarimeter
698 (Institute for Cell and Molecular Biosciences, Newcastle University). CD spectra were recorded in
699 10mM Na₂HPO₄/ NaH₂PO₄ pH 7.5, at protein concentrations between 0.1-0.5 mg/ml, using a 0.2 mm
700 pathlength quartz cuvette (Hellma), at 0.2 nm intervals between 260 and 185 nm at 4°C. Spectra
701 were averaged across nine accumulations, corrected for buffer signal, smoothed and converted to
702 mean residue ellipticity ([θ]) (x1000 deg.cm².dmol⁻¹.residue⁻¹). Deconvolution was performed using
703 the CDSSTR algorithm of the Dichroweb server (<http://dichroweb.cryst.bbk.ac.uk>)³⁷. CD thermal
704 denaturation was performed in 20 mM Tris pH 8.0, 150 mM KCl, 2 mM DTT, at protein
705 concentrations between 0.1-0.4 mg/ml, using a 1 mm pathlength quartz cuvette (Hellma). Data were
706 recorded at 222 nm, between 5°C and 95°C, at 0.5°C intervals with ramping rate of 2°C per minute,
707 and were converted to mean residue ellipticity ([θ]₂₂₂) and plotted as % unfolded ($[\theta]_{222,x} -$
708 $[\theta]_{222,5} / ([\theta]_{222,95} - [\theta]_{222,5})$). Melting temperatures (T_m) were estimated as the points at which samples
709 are 50% unfolded. SYCP1 αC-end constructs were also analysed in 50 mM NaOAc pH 5.5 or 4.6, 150
710 mM KCl.

711

712 **Size-exclusion chromatography multi-angle light scattering (SEC-MALS)**

713 The absolute molar masses of SYCP1 constructs were determined by size-exclusion chromatography
714 multi-angle light scattering (SEC-MALS). Protein samples at >1 mg/ml were loaded onto a Superdex™
715 200 Increase 10/300 GL size exclusion chromatography column (GE Healthcare) in 20 mM Tris pH
716 8.0, 150 mM KCl, 2 mM DTT, at 0.5 ml/min using an ÄKTA™ Pure (GE Healthcare). SYCP1 αC-end
717 constructs were also analysed in 50 mM NaOAc pH 5.5 or 4.6, 150 mM KCl, 2 mM DTT. The column

718 outlet was fed into a DAWN® HELEOS™ II MALS detector (Wyatt Technology), followed by an
719 Optilab® T-rEX™ differential refractometer (Wyatt Technology). Light scattering and differential
720 refractive index data were collected and analysed using ASTRA® 6 software (Wyatt Technology).
721 Molecular weights and estimated errors were calculated across eluted peaks by extrapolation from
722 Zimm plots using a dn/dc value of 0.1850 ml/g. SEC-MALS data are presented with light scattering
723 (LS) and differential refractive index (dRI) profiles, with fitted molecular weights (M_w) plotted across
724 elution peaks.

725

726 **Electrophoretic mobility shift assay (EMSA)**

727 SYCP1 protein constructs were incubated with 25 or 32 μ M (per base pair) 470 or 75 bp linear dsDNA
728 substrate at concentrations indicated, in 20 mM Tris pH 8.0, 150 mM KCl or 50 mM NaOAc pH 5.5,
729 150 mM KCl, for 5 minutes at 4°C. Glycerol was added at a final concentration of 3% and samples
730 were analysed by electrophoresis on a 0.5% (w/v) agarose gel in 0.5x TBE pH 8.0 or 25 mM GABA pH
731 5.5 at 20V for 4 hours at 4°C. DNA was detected by SYBR™ safe (ThermoFisher).

732

733 **Size-exclusion chromatography small-angle X-ray scattering (SEC-SAXS)**

734 SEC-SAXS experiments were performed at beamline B21 of the Diamond Light Source synchrotron
735 facility (Oxfordshire, UK). Protein samples at concentrations >10 mg/ml were loaded onto a
736 Superdex™ 200 Increase 10/300 GL size exclusion chromatography column (GE Healthcare) in 20
737 mM Tris pH 8.0, 150 mM KCl at 0.5 ml/min using an Agilent 1200 HPLC system. SYCP1 α C-end
738 constructs were also analysed in 50 mM NaOAc pH 5.5 or 4.6, 150 mM KCl. The column outlet was
739 fed into the experimental cell, and SAXS data were recorded at 12.4 keV, detector distance 4.014 m,
740 in 3.0 s frames. Data were subtracted and averaged, and analysed for Guinier region R_g and cross-
741 sectional R_g (R_c) using ScÅtter 3.0 (<http://www.bioisis.net>). Approximate parameters for real space
742 analysis were determined using the server www.bayesapp.org, and $P(r)$ distributions fitted using
743 PRIMUS³⁸. *Ab initio* modelling was performed using DAMMIF³⁹ run in interactive mode with random

744 chain selected as expected shape. 10-20 independent runs were performed and averaged. Crystal
745 structures and models were docked into DAMAVER molecular envelopes using SUPCOMB⁴⁰.

746

747 **Electron Microscopy**

748 Electron microscopy (EM) was performed using an FEI Philips CM100 transmission electron
749 microscope at the Electron Microscopy Research Services, Newcastle University. MBP fusion SYCP1
750 samples at 10 μ M were incubated with 100 μ M (per base pair) plasmid double-stranded DNA in 20
751 mM Tris pH 8.0, 250 mM KCl for 10 minutes, and applied to carbon-coated EM grids. Negative
752 staining was performed using 2% (wt/vol) uranyl acetate.

753

754 **Protein crystallisation and X-ray structure solution of SYCP1 α N-end (101-206)**

755 SYCP1 α N-end (101-206) protein crystals were obtained through vapour diffusion in hanging drops,
756 by mixing 200 nl of protein at 10 mg/ml with 100 nl of crystallisation solution (100 mM MES pH 6.0,
757 40% (v/v) MPD) and equilibrating at 4°C for 4-9 days. Crystals were flash frozen in liquid nitrogen. X-
758 ray diffraction data were collected at 0.9282 Å, 100 K, as 2000 consecutive 0.10° frames of 0.050 s
759 exposure on a Pilatus 6M detector at beamline I04-1 of the Diamond Light Source synchrotron
760 facility (Oxfordshire, UK). Data were indexed and integrated in XDS⁴¹ using AutoPROC⁴²; datasets
761 from three crystals were scaled together using XSCALE⁴³ and then merged in Aimless⁴⁴. Crystals
762 belong to monoclinic spacegroup I2 (cell dimensions $a = 65.67$ Å, $b = 37.31$ Å, $c = 108.52$ Å, $\alpha = 90^\circ$, β
763 $= 106.66^\circ$, $\gamma = 90^\circ$), with two SYCP1 chains per asymmetric unit. Data were corrected for anisotropy
764 using the UCLA diffraction anisotropy server (<https://services.mbi.ucla.edu/anisyscale/>)⁴⁵, imposing
765 anisotropic limits of 2.1 Å, 2.1 Å, 2.6 Å, with principal components of 24.09 Å², 6.01 Å² and -20.19 Å².
766 Structure solution was achieved using AMPLE⁴⁶ on the CCP4 online web server
767 (<https://www.ccp4.ac.uk/ccp4online/>), through molecular replacement of Quark *ab initio* model
768 decoys⁴⁷, with auto-tracing and rebuilding in SHELX E and ARP/wARP. Phase improvement was
769 achieved through iterative re-building by PHENIX Autobuild⁴⁸. The structure was completed through

770 manual model building in Coot and refinement using PHENIX refine⁴⁸, with the addition of two 2-
771 methyl-2,4-pentanediol (MPD) ligands and two chloride ions. Refinement was performed using
772 isotropic atomic displacement parameters with riding hydrogens. The structure was refined against
773 anisotropy corrected 2.07 Å data to R and R_{free} values of 0.2264 and 0.2441 respectively, with 100%
774 of residues within the favoured regions of the Ramachandran plot, clashscore of 3.05 and overall
775 MolProbity score of 1.10.

776

777 **Protein crystallisation and X-ray structure solution of truncated SYCP1-αN-end (101-175)**

778 SYCP1 αN-end-tr (101-175) protein crystals were obtained through vapour diffusion in hanging
779 drops, by mixing 1 µl of protein at 10 mg/ml with 1 µl of crystallisation solution (140 mM NaCl, 70
780 mM Na/K phosphate pH 6.2, 35% (v/v) PEG200) and equilibrating at 20°C for 4-9 days. Crystals were
781 soaked for 30 minutes in crystallisation solution containing 40% (v/v) PEG200 and 100 mM NaI, prior
782 to flash freezing in liquid nitrogen. X-ray diffraction data were collected at 1.7712 Å, 100 K, as 2000
783 consecutive 0.10° frames of 0.050 s exposure on a Pilatus 6M detector at beamline I02 of the
784 Diamond Light Source synchrotron facility (Oxfordshire, UK). Data were indexed, integrated and
785 scaled in XDS⁴¹ and XSCALE⁴³, and merged in Aimless⁴⁴. Crystals belong to orthorhombic spacegroup
786 I222 (cell dimensions a = 28.64 Å, b = 39.38 Å, c = 165.77 Å, α = 90°, β = 90°, γ = 90°), with one SYCP1
787 chain per asymmetric unit. SAD structure solution was achieved through identification of five
788 putative iodide sites and secondary structure auto-tracing by SHELX C/D/E, utilising the HKL2MAP
789 interface⁴⁹. Phase improvement was achieved through iterative re-building by PHENIX Autobuild⁴⁸.
790 Data were corrected for anisotropy using the UCLA diffraction anisotropy server
791 (<https://services.mbi.ucla.edu/anisoscale/>)⁴⁵, imposing anisotropic limits of 1.9 Å, 2.0 Å, 2.1 Å, with
792 principal components of 13.25 Å², 0.78 Å² and -14.08 Å². The structure was completed through
793 manual model building in Coot and refinement using PHENIX refine⁴⁸, with the truncation to two
794 iodide sites (based on anomalous difference map peaks) and the addition of a triethylene glycol
795 ligand (PGE). Refinement was performed using isotropic atomic displacement parameters with five

796 TLS groups. The structure was refined against anisotropy corrected 1.91 Å data to R and R_{free} values
797 of 0.2272 and 0.2392 respectively, with 100% of residues within the favoured regions of the
798 Ramachandran plot, clashscore of 6.77 and overall MolProbity score of 1.37.

799

800 **Protein crystallisation and X-ray structure solution of SYCP1 α C-end (676-770) crystal form 1**

801 SYCP1 α C-end (676-770) protein crystals were obtained through vapour diffusion in hanging drops,
802 by mixing 100 nl of protein at 31 mg/ml with 100 nl of crystallisation solution (3.5 M sodium formate
803 pH 7.0) and equilibrating at 20°C for 2 months. Crystals were soaked in a cryoprotectant solution of 6
804 M sodium formate pH 7.0 and flash frozen in liquid nitrogen. X-ray diffraction data were collected at
805 0.9795 Å, 100 K, as 2000 consecutive 0.10° frames of 0.080 s exposure on a Pilatus 6M detector at
806 beamline I02 of the Diamond Light Source synchrotron facility (Oxfordshire, UK). Data were indexed
807 and integrated in XDS⁴¹, and scaled and merged in Aimless⁴⁴, using AutoPROC⁴². Crystals belong to
808 monoclinic spacegroup C2 (cell dimensions $a = 233.42$ Å, $b = 42.85$ Å, $c = 43.69$ Å, $\alpha = 90^\circ$, $\beta = 93.61^\circ$,
809 $\gamma = 90^\circ$), with four SYCP1 chains per asymmetric unit. Structure solution was achieved through
810 fragment-based molecular replacement using ARCIMBOLDO_SHREDDER⁵⁰, a program that derives
811 small models from distant homologs, decomposes and refines the fragments against PHASER's⁵¹ gyre
812 and gimble functions⁵², and combines partial solutions⁵³ for expansion through density modification
813 and main chain tracing with SHELXE⁵⁴ to generate the full structure. The SYCP1- α C-end I4₁22
814 structure (crystal form 2) was used as a starting template for generating 74 models containing 99
815 amino acids each. A phase set combining 25 partial solutions was expanded into a full solution,
816 recognisable by a correlation coefficient of 48.2%. Phase improvement was achieved through
817 iterative re-building by PHENIX Autobuild⁴⁸. Data were corrected for anisotropy using the UCLA
818 diffraction anisotropy server (<https://services.mbi.ucla.edu/anisoscale/>)⁴⁵, imposing anisotropic
819 limits of 2.2 Å, 2.3 Å, 2.2 Å, with principal components of 18.46 Å², 3.44 Å² and -21.90 Å². The
820 structure was completed through manual model building in Coot and refinement using PHENIX
821 refine⁴⁸. Refinement was performed using isotropic atomic displacement parameters with seven TLS

822 groups per chain. The structure was refined against anisotropy corrected 2.15 Å data to R and R_{free}
823 values of 0.2186 and 0.2526 respectively, with 100% of residues within the favoured regions of the
824 Ramachandran plot, clashscore of 6.86 and overall MolProbity score of 1.38.

825

826 **Protein crystallisation and X-ray structure solution of SYCP1 α C-end (676-770) crystal form 2**

827 SYCP1 α C-end (676-770) protein crystals were obtained through vapour diffusion in hanging drops,
828 by mixing 100 nl of protein at 15 mg/ml with 100 nl of crystallisation solution (0.1 M sodium
829 cacodylate pH 6.5, 1.4M sodium acetate) and equilibrating at 20°C for 2 months. Crystals were
830 soaked in a cryoprotectant solution of 0.1 M sodium cacodylate pH 6.5, 1.4 M sodium acetate, 20%
831 PEG400 and flash frozen in liquid nitrogen. X-ray diffraction data were collected at 0.9795 Å, 100 K,
832 as 2000 consecutive 0.10° frames of 0.080 s exposure on a Pilatus 6M detector at beamline I02 of
833 the Diamond Light Source synchrotron facility (Oxfordshire, UK). Data were indexed and integrated
834 in XDS⁴¹ using AutoPROC⁴², scaled in XSCALE⁴³ and scaled merged in Aimless⁴⁴. Crystals belong to
835 tetragonal spacegroup I4₁22 (cell dimensions a = 43.38 Å, b = 43.38 Å, c = 292.18 Å, $\alpha = 90^\circ$, $\beta = 90^\circ$,
836 $\gamma = 90^\circ$), with one SYCP1 chain per asymmetric unit. Structure solution was achieved through
837 fragment-based molecular replacement using ARCIMBOLDO_LITE⁵⁵. Substructures made up of two
838 ideal polyalanine helices of 30 residues each were located with PHASER, profiting from its features
839 for small fragments, and were extended with SHELXE within the Arcimboldo mode for coiled coil
840 structures⁵⁶. A correct solution was identified by a SHELXE Correlation Coefficient of 40.5%. Phase
841 improvement was achieved through iterative re-building by PHENIX Autobuild⁴⁸. Data were
842 corrected for anisotropy using the UCLA diffraction anisotropy server
843 (<https://services.mbi.ucla.edu/anisoscale/>)⁴⁵, imposing anisotropic limits of 2.9 Å, 2.9 Å, 2.5 Å, with
844 principal components of 16.05 Å², 16.05 Å² and -32.09 Å². The structure was completed through
845 manual model building in Coot and refinement using PHENIX refine⁴⁸, with the addition of one
846 acetate ligand. Refinement was performed using isotropic atomic displacement parameters with
847 riding hydrogens. The structure was refined against anisotropy corrected 2.49 Å data to R and R_{free}

848 values of 0.2251 and 0.2517 respectively, with 100% of residues within the favoured regions of the
849 Ramachandran plot, clashscore of 1.24 and overall MolProbity score of 0.84.

850

851 **Protein sequence and structure analysis**

852 Conservation of SYCP1 amongst vertebrate sequences was calculated as per residue scores for the
853 full SYCP1 sequence and the α C-end structure by ConSurf (<http://consurf.tau.ac.il/>), and secondary
854 structure prediction was performed by JNet (<http://www.compbio.dundee.ac.uk/www-jpred/>).
855 Protein structures were superposed and rmsd values calculated by PHENIX superpose⁴⁸. Structural
856 assemblies were analysed by PISA. Molecular structure images were generated using the PyMOL
857 Molecular Graphics System, Version 1.3 Schrödinger, LLC.

858

859 **Data availability**

860 Crystallographic structure factors and atomic co-ordinates have been deposited in the Protein Data
861 Bank (PDB) under accession numbers 6F5X, 6F62, 6F63 and 6F64. Uncropped gel images are shown
862 in Supplementary Data Set 1. All other data are available from the corresponding author upon
863 reasonable request. A Life Sciences Reporting Summary for this article is available.

864

865 **Methods-only references**

- 866 37. Whitmore, L. & Wallace, B.A. Protein secondary structure analyses from circular dichroism
867 spectroscopy: methods and reference databases. *Biopolymers* **89**, 392-400 (2008).
- 868 38. P.V.Konarev, V.V.V., A.V.Sokolova, M.H.J.Koch, D. I. Svergun. PRIMUS - a Windows-PC based
869 system for small-angle scattering data analysis. *J Appl Cryst.* **36**, 1277-1282 (2003).
- 870 39. Franke, D. & Svergun, D.I. DAMMIF, a program for rapid ab-initio shape determination in
871 small-angle scattering. *J Appl Crystallogr* **42**, 342-346 (2009).
- 872 40. Kozin, M.B. & Svergun, D.I. Automated matching of high- and low-resolution structural
873 models. *Journal of Applied Crystallography* **34**, 33-41 (2001).
- 874 41. Kabsch, W. Xds. *Acta Crystallogr D Biol Crystallogr* **66**, 125-32 (2010).
- 875 42. Vonrhein, C. et al. Data processing and analysis with the autoPROC toolbox. *Acta Crystallogr*
876 *D Biol Crystallogr* **67**, 293-302 (2011).
- 877 43. Diederichs, K., McSweeney, S. & Ravelli, R.B. Zero-dose extrapolation as part of
878 macromolecular synchrotron data reduction. *Acta Crystallogr D Biol Crystallogr* **59**, 903-9
879 (2003).
- 880 44. Evans, P.R. An introduction to data reduction: space-group determination, scaling and
881 intensity statistics. *Acta Crystallogr D Biol Crystallogr* **67**, 282-92 (2011).
- 882 45. Strong, M. et al. Toward the structural genomics of complexes: crystal structure of a PE/PPE
883 protein complex from *Mycobacterium tuberculosis*. *Proc Natl Acad Sci U S A* **103**, 8060-5
884 (2006).
- 885 46. Bibby, J., Keegan, R.M., Mayans, O., Winn, M.D. & Rigden, D.J. AMPLE: a cluster-and-
886 truncate approach to solve the crystal structures of small proteins using rapidly computed
887 ab initio models. *Acta Crystallogr D Biol Crystallogr* **68**, 1622-31 (2012).
- 888 47. Xu, D. & Zhang, Y. Ab initio protein structure assembly using continuous structure fragments
889 and optimized knowledge-based force field. *Proteins* **80**, 1715-35 (2012).
- 890 48. Adams, P.D. et al. PHENIX: a comprehensive Python-based system for macromolecular
891 structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-21 (2010).
- 892 49. Pape, T. & Schneider, T.R. HKL2MAP: a graphical user interface for macromolecular phasing
893 with SHELX programs. *Journal of Applied Crystallography* **37**, 843-844 (2004).
- 894 50. Millan, C. et al. Exploiting distant homologues for phasing through the generation of
895 compact fragments, local fold refinement and partial solution combination. *Acta Crystallogr*
896 *D Struct Biol* **74**, 290-304 (2018).
- 897 51. McCoy, A.J. et al. Phaser crystallographic software. *Journal of Applied Crystallography* **40**,
898 658-674 (2007).
- 899 52. McCoy, A.J. et al. Gyre and gimble: a maximum-likelihood replacement for Patterson
900 correlation refinement. *Acta Crystallogr D Struct Biol* **74**, 279-289 (2018).
- 901 53. Millan, C. et al. Combining phase information in reciprocal space for molecular replacement
902 with partial models. *Acta Crystallogr D Biol Crystallogr* **71**, 1931-45 (2015).
- 903 54. Uson, I. & Sheldrick, G.M. An introduction to experimental phasing of macromolecules
904 illustrated by SHELX; new autotracing features. *Acta Crystallogr D Struct Biol* **74**, 106-116
905 (2018).
- 906 55. Rodriguez, D.D. et al. Crystallographic ab initio protein structure solution below atomic
907 resolution. *Nature Methods* **6**, 651-U39 (2009).
- 908 56. Caballero, I. et al. ARCIMBOLDO on coiled coils. *Acta Crystallogr D Struct Biol* **74**, 194-204
909 (2018).

910















