

# Modulation of *Pleurodeles waltl* DNA Polymerase $\mu$ Expression by Extreme Conditions Encountered during Spaceflight

Véronique Schenten<sup>1</sup>✉, Nathan Guéguinou<sup>1</sup>✉, Sarah Baatout<sup>2,3</sup>, Jean-Pol Fripiat<sup>1\*</sup>

**1** Stress Immunity Pathogens Laboratory, EA7300, Lorraine University, Vandœuvre-lès-Nancy, France, **2** Radiobiology Unit, Belgian Nuclear Research Centre, SCK-CEN, Mol, Belgium, **3** Department of Molecular Biotechnology, Ghent University, Ghent, Belgium

## Abstract

DNA polymerase  $\mu$  is involved in DNA repair, V(D)J recombination and likely somatic hypermutation of immunoglobulin genes. Our previous studies demonstrated that spaceflight conditions affect immunoglobulin gene expression and somatic hypermutation frequency. Consequently, we questioned whether Pol $\mu$  expression could also be affected. To address this question, we characterized Pol $\mu$  of the Iberian ribbed newt *Pleurodeles waltl* and exposed embryos of that species to spaceflight conditions or to environmental modifications corresponding to those encountered in the International Space Station. We noted a robust expression of Pol $\mu$  mRNA during early ontogenesis and in the testis, suggesting that Pol $\mu$  is involved in genomic stability. Full-length Pol $\mu$  transcripts are 8–9 times more abundant in *P. waltl* than in humans and mice, thereby providing an explanation for the somatic hypermutation predilection of G and C bases in amphibians. Pol $\mu$  transcription decreases after 10 days of development in space and radiation seem primarily involved in this down-regulation. However, space radiation, alone or in combination with a perturbation of the circadian rhythm, did not affect Pol $\mu$  protein levels and did not induce protein oxidation, showing the limited impact of radiation encountered during a 10-day stay in the International Space Station.

**Citation:** Schenten V, Guéguinou N, Baatout S, Fripiat J-P (2013) Modulation of *Pleurodeles waltl* DNA Polymerase  $\mu$  Expression by Extreme Conditions Encountered during Spaceflight. PLoS ONE 8(7): e69647. doi:10.1371/journal.pone.0069647

**Editor:** John R. Battista, Louisiana State University and A & M College, United States of America

**Received:** April 23, 2013; **Accepted:** June 12, 2013; **Published:** July 31, 2013

**Copyright:** © 2013 Schenten et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This work was supported by the French National Space Agency (CNES) (www.cnes.fr), the National Research Agency (grant ANR NT09-632666)(www.agence-nationale-recherche.fr), the French Ministry of Higher Education and Research (www.enseignementsup-recherche.gouv.fr), the Lorraine University (www.univ-lorraine.fr/), the Lorraine area (www.lorraine.eu/) and BELSPO/PRODEX/ESA (contract 42-000-90-380) (www.belspo.be). N.G. was supported by a PhD fellowship from the National Research Fund of Luxembourg (www.fnr.lu) and V.S. by a CNES post-doctoral fellowship. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: jean-pol.fripiat@univ-lorraine.fr

✉ These authors contributed equally to this work.

## Introduction

The immune system is affected by environmental changes encountered during spaceflight such as confinement, microgravity, biomechanical stresses, radiation and disruption of the circadian rhythm. Indeed, astronauts have experienced altered immune function and increased vulnerability to infections during spaceflights dating back to the Apollo and Skylab missions [1]. The effects of spaceflights on the immune system should therefore be considered more thoroughly before undertaking prolonged space missions.

Until now, studies regarding spaceflight-induced immune dysfunctions focused mostly on innate immunity and T-cell responses while humoral immunity and antibodies, despite their important functions, have rarely been investigated [2]. To improve our knowledge concerning the effects of spaceflight on humoral immunity, we used the Iberian ribbed newt *Pleurodeles waltl* as animal model. This amphibian species is a good model for studies applicable for human health because it was shown that the cardinal elements of the adaptive immune system are shared by all gnathostomes [3]. This observation was confirmed in *P. waltl*. Indeed, it was shown that this animal possesses at least 10

subgroups of VH genes, 8 JH and 8 D segments in its genome and that it uses these genes and segments in the same way as humans and rodents to create its functional antibody heavy chain genes [4]. It was also demonstrated that *P. waltl* possesses three isotypes of antibodies, IgM, IgY and IgP, which have their human physiological counterparts [5]. And finally, several immunologically important genes and transcripts, such as those encoding AID, Ikaros and the C3 complement component, were shown to be conserved in *P. waltl* and have expression patterns similar to those observed in mammals ([6,7] and unpublished data).

Using that species, we previously showed that spaceflight affects antibody production in response to an antigenic stimulation. Indeed, the expression of IgY heavy chains (the physiological counterpart of mammalian IgA molecules) was increased in flown *P. waltl* [8] as previously observed in cosmonauts [9]. Furthermore, the use of the different VH gene subgroups [8] and the expression of individual VH genes [10] were observed to be modified under spaceflight conditions. In addition, we recently demonstrated that somatic hypermutation, that diversify antibody binding-sites to improve their affinity, occur in space following immunization but at a lower frequency [11] and that gravity changes during animal development affect IgM heavy chain transcription [12].

To create functional immunoglobulin (Ig) genes, B lymphocytes from jawed vertebrates associate V<sub>H</sub>-D, D-J<sub>H</sub>, V <sub>$\kappa$</sub> -J <sub>$\kappa$</sub>  and V <sub>$\lambda$</sub> -J <sub>$\lambda$</sub>  gene and segments during their development in primary lymphoid organs. This process, called V(D)J recombination, involves DNA double-strand breaks inflicted by the products of the RAG1 and RAG2 recombination-activating genes [13]. The processing of V(D)J coding joints is then sustained by core NHEJ components (Ku-Artemis-DNA-PK and XLF-XRCC4-DNA ligase IV) [14] but DNA polymerases of the X family further contribute, particularly when the generated nicks leave 3' overhangs, which are good substrates for the polymerases [15–17]. The polymerases of the X family include the DNA polymerases beta (Pol $\beta$ ), lambda (Pol $\lambda$ ), mu (Pol $\mu$ ) and terminal deoxynucleotidyl transferase (TdT), which are structurally related enzymes specialized in repair pathways involving double-strand breaks (DSBs) and gaps [18]. Pol $\mu$  is more closely related to TdT than to other polymerase X members and shares the same exon-intron organization. However, the roles of TdT and Pol $\mu$  are distinct [19].

In 2003, Bertocci et al. reported that Ig light chain gene rearrangement is altered in Pol $\mu$ -deficient mice, with an extensive reduction in  $\kappa$  light chain CDR3 length [20]. This observation highlighted the contribution of this protein in Ig light gene rearrangement, in contrast to TdT, which is mainly involved in Ig heavy chain gene rearrangement [21,22]. This difference is reinforced by the fact that the Pol $\mu$  and TdT expression patterns are different. In adults, Pol $\mu$  is expressed in a wide range of tissues [23], while TdT expression is limited to hematopoietic organs.

Several studies suggested that Pol $\mu$  could also be involved in somatic hypermutation (SHM) of Ig genes that diversify antibody binding sites [24–27].

Finally, it was shown that Pol $\mu$  is involved in the repair of radiation-induced DNA damage. Indeed, exposure of cells to ionizing radiation (IR) induced a reduction in properly spliced Pol $\mu$  mRNAs [23] and an increase in Pol $\mu$  protein that forms discrete nuclear foci coinciding with IR-induced foci of  $\gamma$ H2AX, a marker of sites of DNA DSBs [28]. Another study showed that a lack of Pol $\mu$  results in delayed DSB repair kinetics and in persistent DNA damage [29].

As these data indicate that Pol $\mu$  is involved in V(D)J recombination, DNA repair and likely somatic hypermutation, we characterized that polymerase in *P. waltl* and determined how spaceflight-associated environmental modifications affect Pol $\mu$  expression during *P. waltl* development in the International Space Station. This study is important because several studies showed that spaceflight has a negative impact on the immune system (see above) and exposure of astronauts to low doses of cosmic radiation is considered as a threat for long-term space missions [30].

## Materials and Methods

### Ethic Statement

Animals were treated in accordance with the National Legislation and the Council Directive of the European Communities on the Protection of Animals Used for Experimental and Other Scientific Purposes (86/609/EEC). Animal husbandry conditions were approved by the French Ministry of Agriculture and Fisheries (agreement DDSV54/SPA/07/130) and the protocol was approved by the regional ethic committee: Comité régional d'éthique pour l'expérimentation animale, Nancy - Lorraine - Nord-Est (Permit Number: 04112004).

### Animals and RNA Extractions

*P. waltl* adults and embryos were reared in our animal facilities in running tap water at 18°C. Embryonic and larval develop-

mental stages were defined according to Gallien and Durocher [31]. Ten embryos were randomly chosen for each developmental stage studied and homogenized in TRIzol reagent (Invitrogen, Carlsbad, CA, USA) for RNA extraction. Total RNA was also isolated from the thymus of stage 44 (36-day-old) larvae and from various adult tissues.

### AMPHIBODY Space Experiment

This experiment was performed in the International Space Station (ISS) from March 30 to April 09, 2006 using miniaquaria developed by EADS-Astrium and the German Space Agency [12]. Six miniaquaria were mounted on the stationary positions of the Kubik incubator designed for space experiments (Comat Aerospace, Toulouse, France) for microgravity exposure, 6 on incubator's 1 g-centrifuge to perform an in-flight 1 g control and 12 in an incubator kept on Earth as ground controls. Embryos developed during 10 days in the ISS. Living larvae were recovered and fixed 14 h after landing. Note that the Kubik's centrifuge was unexpectedly shut off 30 h before landing, thereby exposing larvae to several hours of microgravity and invalidating our in-flight 1 g control.

### Other Animal Treatments

*P. waltl* embryos exposures to *i*) hypergravity in a centrifuge running at 3G, *ii*) simulated microgravity in a random positioning machine (RPM) that randomize the direction of the gravity force resulting in an average net force approaching zero, *iii*) altered circadian rhythm and *iv*) simulated ISS radiation environment using a combination of neutrons (total dose of 2.0 mGy) and  $\gamma$  rays (total dose of 1.9 mGy) as representative of low-linear energy transfer (LET) and high-LET particles received during 10 days in the space station, were performed as previously described [12]. Note that a full reproducibility of space radiation on Earth is not possible this is why we use the word 'simulation'. Additionally, to induce a mechanical stress, *P. waltl* larvae at stage 36–37 of development were placed in 50 mL of water at 20°C and subjected to vibrations for 5 h at 15 Hz. In each case, larvae of the same developmental stage reared under standard conditions were used as controls.

### Cloning and Sequencing of *P. waltl* Polymerase $\mu$ cDNA

One microgram of spleen RNA was reverse transcribed using random primers and M-MLV reverse transcriptase (Invitrogen, Carlsbad, CA, USA). Primers were designed in conserved regions identified by aligning Pol $\mu$  mRNA sequences from *Homo sapiens*, *Mus musculus*, *Danio rerio* and *Xenopus tropicalis*. A first PCR reaction was performed with the Pol $\mu$  For and Pol $\mu$  Rev primers (Table 1) on spleen cDNA using Goldstar Taq polymerase (Eurogentec, Seraing, Belgium). Two PCR products of 917 and 814 bp were obtained, cloned and sequenced. Next, specific primers (Polmu GSP1 and Polmu GSP2) and nested primers (Polmu NGSP1 and Polmu NGSP2, Table 1) were designed from the obtained sequence to perform 3'- and 5'-RACE using the BD SMART RACE cDNA Amplification kit (Clontech, Mountain View, USA). RACE products were cloned and sequenced to obtain full-length *P. waltl* Pol $\mu$  cDNA.

### Real-time PCR

qPCR primers were designed using the GenScript software. Pol $\mu$  primers (Polmu long For and Polmu long Rev) were defined in exons 7 and 8 to quantify only the longest isoform (Table 1). DNA-PKc-, Gadd45-, GAPDH-,  $\alpha$ -actin-, TAFII- and mtRNA16S-specific primers were described previously [12].

**Table 1.** Primers used in this study.

Primer name	Primer sequence (5' to 3')	Purpose	Annealing temp. (°C)
Polmu For	CCAGTATATGCCTGCCAGAGA	PCR	63
Polmu Rev	TGCTTTGATCCAGTCCACCC	PCR	63
Polmu GSP1	ACTGTTGCCGAACCTGCCTCTCT	5'-RACE	63
Polmu NGSP1	GCCAGTATCTCCAAGCATCCGTGA	5'-RACE	63
Polmu GSP2	AGAGCTCCAGCACAGCAGCAGAAAA	3'-RACE	64
Polmu NGSP2	CATTTGCTCTGCTTGGTGACTG	3'-RACE	64
Polmu long For	TGGAAGCTGACAAGATCCAG	qPCR & PCR	59
Polmu long Rev	CTGCTTCCTCTTTGAAACC	qPCR	59

doi:10.1371/journal.pone.0069647.t001

qPCRs were performed using the Mesa Green qPCR Master Mix (Eurogentec, Seraing, Belgium) in a Mastercycler® realplex2 real-time PCR engine (Eppendorf, Hamburg, Germany). The cycling protocol was as follows: 3 min at 95°C, then 40 cycles of 15 s at 95°C and 30 s at 59°C. Standard curves were generated to ensure that the amplification efficiencies were similar and in the range of 95%–105%. Each qPCR was performed in triplicate and repeated at least 2 times. The data were analyzed using the relative Pfaffl model [32]. Relative expressions, expressed in arbitrary units (A.U.), were calculated using 4 housekeeping genes ( $\alpha$ -actin, mtRNA16S, TafII and GAPDH), the GeNorm software and the Vandesompele method [33]. Primers Polmu long For and Polmu Rev (Table 1) were used to amplify, using conventional PCR, Polμ variants in the spleen and testis of *P. waltl*.

### Production of Anti-polymerase μ Polyclonal Antibodies

To produce polyclonal antibodies specific for *P. waltl* Polμ, a 16 amino acid peptide was designed, produced and coupled to KLH by Eurogentec (Seraing, Belgium). BALB/c mice were immunized with this immunogen. The first dose consisted of 100 μg of the immunogen emulsified with TiterMax® adjuvant (Sigma-Aldrich, Saint-Quentin Fallavier, France) in a 1:1 ratio. Three weeks later, a second dose of 100 μg of immunogen mixed with TiterMax® in a 1:0.5 ratio was injected. One week after this second injection, mice antisera were collected, purified with the Rapid Antibody Purification Kit (Cell Biolabs, San Diego, CA, USA) and concentrated using a Centricon Plus-70 device (Millipore, Billerica, MA, USA).

### Western Blotting

Proteins were prepared by lysing a pool of 3 larvae in lysis buffer containing Triton X-100. Samples of 50 μg of protein were heated at 97°C for 5 minutes, run on a 12% Bis-Tris gel and electrotransferred to a PVDF membrane (Amersham, Buckinghamshire, UK). Immunodetection was performed using homemade polyclonal antibodies against Polμ (see above) and a horseradish peroxidase (HRP)-conjugated rat anti-mouse secondary antibody. Immunodetection was performed using the Super-Signal West Pico Chemiluminescent Substrate (Pierce, Erembodegem, Belgium).

### Protein Oxidation

The detection of oxidatively damaged proteins was performed with an OxiSelect™ Protein Carbonyl Immunoblot Kit (Cell Biolabs, Inc., San Diego, CA, USA) according to the manufacturer's recommendations. Briefly, proteins were separated by SDS-PAGE and blotted on PVDF membranes that were immuno-

stained to detect carbonyls. Carbonyls are primarily aldehydes resulting from oxidation of various amino acids. This staining procedure involves a derivatization of carbonyl groups with 2,4-dinitrophenylhydrazine (DNPH), followed by the detection of DNPH by a specific antibody. Following the oxiblot, membranes were Coomassie Blue stained to ensure equal protein loading in each lane. All measurements of oxidation were normalized to the amount of protein in each sample. To ensure that this approach recognizes only carbonylated proteins, the molecular weight marker, made of recombinant and thus unmodified proteins, was used as a negative control. In all cases, the marker was detected by Coomassie blue staining but not in the oxiblots thereby demonstrating that this protocol reveals only derivatized carbonyl groups and not native proteins.

### Statistics

Statistical analyses were performed using SPSS v13.0 (SPSS Inc., USA). Outlier values were determined by a boxplot of each studied group. When normality and homogeneity of variances were ascertained, as determined by Kolmogorov-Smirnov and Levene tests, respectively, Student's *t*-test analyses were performed to establish two group comparisons. Otherwise, Mann-Whitney tests were used for two group comparisons. *p*-values <0.05 and <0.10 were selected to indicate significance and trends, respectively.


## Results

### *P. waltl* Polymerase μ

To isolate the Polμ cDNA of *P. waltl*, we designed primers in conserved regions revealed by the alignment of Polμ sequences from different species. These primers allowed the amplification of two products that presented strong homologies with the corresponding regions of human, mouse, *Xenopus* and zebrafish Polμ mRNAs. Specific primers were then designed to perform RACE-PCR and recover both extremities of *P. waltl* Polμ cDNAs. The sequence of the full-length cDNA has been submitted to the EMBL database (accession number HE583591).

We predicted the *P. waltl* Polμ exon limits by alignment with Polμ cDNA sequences from species whose exon borders are known. This analysis revealed that, as in other species [23,25,34], *P. waltl* Polμ is encoded by 11 exons, of which exon 8 could be alternatively spliced (Figure 1).

To confirm that we isolated the Polμ cDNA of *P. waltl*, we built a cladogram (Figure 1C). This cladogram shows that the sequence identified in *P. waltl* clusters with Polμ sequences from other species, thereby demonstrating that we isolated the cDNA coding



**Figure 1. *Pleurodeles waltl* DNA polymerase  $\mu$ .** Schematic representation of *P. waltl* Pol $\mu$  mRNAs (A). The short variant (Pol $\mu$   $\Delta$  Ex 8) presents a 103 bp deletion corresponding to exon 8. This deletion removes one part of the active site of the protein. The BRCT (breast cancer suppressor protein carboxy-terminal) and the Pol X domain, which contains the active site, are required for Pol $\mu$  function. Pol $\mu$  variants were amplified using primers indicated by horizontal arrows (B). Quantification of the obtained PCR products using a Gel Doc 2000 and the Quantity One v.4.3.1 software (Bio-Rad, Hercules, CA, USA) indicates that the short isoform represents 16% and 6% of the Pol $\mu$  transcripts in the spleen and testis of *P. waltl*, respectively. Ctl<sup>-</sup> = negative control. MW = 100 bp DNA ladder. A cladogram was then constructed by neighbor joining (C) supported with 1000 bootstrap replications using the MEGA4 software (<http://www.megasoftware.net/>) and the following sequences: Terminal deoxynucleotidyl transferase (TdT); *Bos taurus* (DAA14763), *Homo sapiens* (BAB72001), *Mus musculus* (NP\_001036693), *Gallus gallus* (NP\_990720), *Ambystoma mexicanum* (AA092254), *Xenopus laevis* (NP\_001079251), *Danio rerio* (AAS89780). Polymerase beta; *Mus musculus* (NP\_035260), *Homo sapiens* (NP\_002681), *Xenopus laevis* (NP\_001081643), *Xenopus tropicalis* (AAH74537), *Xiphophorus maculatus* (AAU11319). Polymerase lambda; *Danio rerio* (NP\_998408), *Xenopus tropicalis* (NP\_001093716), *Mus musculus* (NP\_064416), *Homo sapiens* (NP\_001167555). Polymerase mu; *Homo sapiens* (NP\_037416), *Mus musculus* (NP\_059097), *Danio rerio* (NP\_956542), *Xenopus tropicalis* (NP\_001164987), *Pleurodeles waltl* (HE583591). The scale bar corresponds to the evolutionary distance. doi:10.1371/journal.pone.0069647.g001

for Pol $\mu$  and not for another polymerase. Moreover, this tree confirms that Pol $\mu$  is more closely related to TdT than to other polymerase X members and that the divergence of Pol $\mu$  and TdT is likely much more recent than the divergence of Pol $\beta$  and Pol $\lambda$ .

Moreover, the two regions known to be necessary for Pol $\mu$  function are conserved (data not shown): BRCT required for the interaction with core NHEJ factors Ku and XRCC4-Ligase IV [17,28] and the Pol X domain, which contains the active site

(DNA, NTP and metal binding sites) responsible for polymerase activity. Thus, *P. waltl* Pol $\mu$  has the domains shown to be required for its activity in mammals.


### Expression during Early Development and in Adult Tissues

To determine when Pol $\mu$  mRNA is expressed during early *P. waltl* development and situate its expression relative to the first expression of AID, RAG1 and IgM heavy chain transcripts [4,6], we performed real-time PCR experiments on cDNAs prepared with total RNA extracted from embryos from stage 2 (four cells, 7.5 h after laying) to stage 33a (213 h after laying) of development. Our results (Figure 2A) indicate that the amount of Pol $\mu$  transcripts is important during the first stages of development, is weak during neurulation and increases again during the tail-bud stages when the first IgM heavy chain mRNAs appear. We also

quantified Pol $\mu$  transcripts in adult tissues, in which strong signals were observed in the testis, the thymus and the spleen (Figure 2B).

### Pol $\mu$ Expression in Larvae Submitted to Spaceflight and Spaceflight-associated Environmental Conditions

Because our previous studies suggested that spaceflight conditions could alter the V(D)J recombination process [8], we quantified Pol $\mu$  transcripts in larvae of the AMPHIBODY space experiment. Our results show a tendency for a decrease in Pol $\mu$  transcripts ( $p=0.089$ ) in larvae that developed in the ISS compared to ground controls (Figure 3A). To identify the cause of decreased Pol $\mu$  transcription, we recreated in the laboratory, to the best of our abilities, the various environmental changes encountered during spaceflight and quantified Pol $\mu$  mRNAs in larvae subjected to each of these changes (Figures 3B–F). Our results demonstrate that neither hypergravity nor simulated microgravity in the RPM affected the expression of Pol $\mu$



**Figure 2. *P. waltl* polymerase  $\mu$  mRNA expression profiles.** Pol $\mu$  transcripts were quantified by real-time PCR during early stages of development (A) and in various tissues (B). Arrows indicate the expression of AID, IgM heavy chains and RAG1 transcripts, as reported in previous studies [4,6]. Developmental stages, as well as the time elapsed since laying (between brackets), are indicated. Ten embryos were studied for each developmental stage. The relative value obtained for the thymus of stage 44 larvae was set to 1 as a reference for tissue expression. Values represent means  $\pm$  SEM.

doi:10.1371/journal.pone.0069647.g002


transcripts, indicating that an alteration in gravity is not responsible for the decrease in Pol $\mu$  transcription. Perturbation of the circadian rhythm (Figure 3D) and mechanical stress (Figure 3E) also did not affect Pol $\mu$  transcription, while exposure to simulated space radiation led to a significant decrease in Pol $\mu$  mRNAs ( $p = 0.006$ ) (Figure 3F).

Consequently, we wondered whether the simulation of space radiation could also impact the expression of another gene involved in NHEJ, DNA-PKc, a kinase involved in the repair of DNA breaks generated by radiation and Ig gene rearrangement. Our results indicate that space radiation does not affect the transcription of DNA-PKc (Figure 3G). To determine whether the stress created by the irradiation protocol was high, we quantified Gadd45, a canonical stress response gene. As shown in Figure 3H, the transcription of Gadd45 is also unaffected by the simulation of space radiation. Finally, we quantified Pol $\mu$  protein in control and irradiated larvae. Our results show that space radiation has no

impact on Pol $\mu$  protein levels (Figures 3I and 3J). We attempted to quantify DNA-PKc and Gadd45 protein levels, but the commercial antibodies tested did not recognize these proteins in *P. waltl* samples.

### Cumulative Effects of Space Radiation and Circadian Rhythm Perturbation

During spaceflight, astronauts are submitted to a combination of stresses and not to a single stress, as tested above. Therefore, we analyzed the effects of a combination of two environmental modifications on Pol $\mu$ , DNA-PKc and Gadd45 mRNA levels and on Pol $\mu$  protein levels. Because it is not possible to test all possible combinations, we opted for the combination of space radiation with the perturbation of the circadian rhythm, as the latter is known to modulate immunity [35]. Exposure of larvae to darkness and simulated space radiation led to significantly increased Pol $\mu$  mRNA levels (2 fold increase) (Figure 4A) and to the almost tripled



**Figure 3. Effects of spaceflight conditions.** Pol $\mu$  mRNAs were quantified in larvae that developed in microgravity in the ISS (A), in hypergravity (B), in the RPM to simulate microgravity (C) or in the dark (D). The same study was performed with larvae subjected to a mechanical stress (E) or the ISS radiation environment (F). The effects of space radiation on DNA-PKc (G) and Gadd45 (H) mRNAs and on the amount of Pol $\mu$  protein (I and J) were also investigated. Western blot results were analyzed and quantified by densitometry using a Fusion Fx7 imaging system (Vilber Lourmat, Marne-la-Vallée, France) and Pol $\mu$  expression was normalized to  $\beta$ -actin (J). Each lane of the western blot contains proteins from 3 larvae. Results are expressed as the mean  $\pm$  SEM. "A.U." stands for "arbitrary units". doi:10.1371/journal.pone.0069647.g003

expression of both DNA-PKc and Gadd45 transcripts (Figures 4B and 4C). However, the amount of Pol $\mu$  protein was unaffected by this combination of environmental changes (Figures 4D and 4E). Thus, it appears that space radiation combined with darkness induces a greater response in larvae compared to either condition alone, as shown by increased DNA-PKc, Gadd45 and Pol $\mu$  mRNA levels, even though the induction of Pol $\mu$  transcription is not sufficient to alter the amount of Pol $\mu$  protein.


### Oxidative Stress in Irradiated Larvae and in Larvae Irradiated in the Dark

It has previously been shown that radiation can induce oxidative stress and that some markers, such as Gadd45, are sensitive to an oxidative environment. Therefore, we determined whether the variations in mRNA expression found in this study could be due to oxidative stress. To test this hypothesis, we measured the carbonylation of proteins in control and irradiated larvae and in larvae irradiated in the dark. Our results show that neither radiation (Figure 5A) nor the combination of space radiation and darkness (Figure 5B) induced oxidative stress. Thus, the variations of Pol $\mu$ , DNA-PKc and Gadd45 transcripts described in this study (Figure 4) are not due to increased oxidative stress.

## Discussion


### Overexpression of Full-length Pol $\mu$ mRNA in *P. waltl*

In this study, we report the first sequencing of Pol $\mu$  in a urodele amphibian. We also describe a splice variant with a 103 bp deletion, corresponding to exon 8 (Figure 1A), which represents, respectively, 16% and 6% of the 11-exon isoform expressed in the spleen and testis of adults (Figure 1B). No other splice variant was revealed by our RACE-PCRs. Numerous human and murine Pol $\mu$  splice variants have been described representing approximately 90% of Pol $\mu$  mRNA species, in which splicing affected exons 5 to 11, encoding the active site of the polymerase. In most cases, these splicing events induced a premature stop codon and truncated proteins [23,27], which could function as regulators of the activity of this enzyme [36]. Similarly, the splice variant detected in this study lacks one part of its active site and will consequently likely be nonfunctional. However, in *P. waltl*, the full-length Pol $\mu$  isoform represents 84–94% of Pol $\mu$  mRNAs. This corresponds to an 8–9 fold overexpression compared to humans and mice, in which the full-length isoform represents only ~10% of the total Pol $\mu$  mRNAs. Given that the overexpression of full-length Pol $\mu$  in the RAMOS cell line increased the frequency of somatic hypermutations at G/C residues and decreased those at A/T residues by a factor of 6 [24], this overexpression of full-length Pol $\mu$  in *P. waltl* could explain why SHM affects mainly G/C residues in Ig VH regions of this animal [11].



**Figure 4. Effects of the combination of space radiation and darkness.** Pol $\mu$  (A), DNA-PKc (B) and Gadd45 (C) mRNAs were quantified by qPCR in larvae that developed in the dark and under the simulation of ISS radiation. Pol $\mu$  proteins were quantified by western blotting (D). Each lane contains proteins from 3 larvae. Western blot results were analyzed and quantified by densitometry using a Fusion Fx7 imaging system (Vilber Lourmat, Marne-la-Vallée, France) and Pol $\mu$  expression was normalized to  $\beta$ -actin (E). Results are expressed as the mean  $\pm$  SEM. "A.U." stands for "arbitrary units".

doi:10.1371/journal.pone.0069647.g004



**Figure 5. Protein oxidation evaluated by measuring the carbonylation of amino acid side chains.** Representative oxiblots of carbonylated proteins from larvae submitted to space radiation (**A**) or larvae that developed in the dark and under the simulation of ISS radiation (**B**) are shown. Each lane contains proteins from 3 larvae. Equal protein loading was assessed by Coomassie staining. Western blot results were analyzed and quantified by densitometry using a Fusion Fx7 imaging system (Vilber Lourmat, Marne-la-Vallée, France) and carbonylation was normalized to the total amount of protein. Results are expressed as the mean  $\pm$  SEM. doi:10.1371/journal.pone.0069647.g005

### Pol $\mu$ Expression Profiles

Using real-time PCR, we detected high quantities of Pol $\mu$  transcripts at early developmental stages (segmentation and gastrulation) (Figure 2A). Expression of Pol $\mu$  was approximately 4 times lower during neurulation and increased again when Ig heavy chain mRNAs appeared. These results suggest that Pol $\mu$  has a role during early embryonic development that is distinct from antibody V(D)J recombination. During early ontogeny, somatic cells proliferate actively and are sensitive to genotoxic stresses. Consequently, robust machineries are required to preserve cell viability and genome stability [37,38]. Pol $\mu$ , which is a component of the cellular response to DSBs [28,39], is well-equipped to connect embryonic DSBs and prevent genomic damage, thereby providing an explanation to its very early developmental expression. The fact that Pol $\mu$  expression increases when Ig heavy chain gene rearrangement starts was expected because it was shown that Pol $\mu$  adds nontemplated nucleotides at D-J<sub>H</sub> joints in mouse embryos when TdT is not yet expressed [40].

As reported in humans [23], Pol $\mu$  is expressed in a wide range of *P. waltil* tissues, with greater representation in lymphoid organs (Figure 2B). Expression in these tissues was expected because we previously detected a high level of RAG1 mRNA in the thymus of 40-day-old larvae and lower levels in the spleens of 2-year-old adults [4], and Pol $\mu$  repairs RAG-induced DSBs [40]. Figure 2B also shows that the expression of Pol $\mu$  in the spleen decreases with age. Some authors have suggested that due to its mutagenic potential, its preferential expression in secondary lymphoid tissues and its expression in the nucleus of centroblasts from human tonsils, Pol $\mu$  could play a role in somatic hypermutation of Ig genes [24–27]. However, an analysis of Pol $\mu$ -deficient mice did not support this hypothesis [36,41]. This discrepancy could be due to

redundancy between DNA polymerases involved in the hypermutation process. Nevertheless, if Pol $\mu$  is involved in the SHM machinery, the decrease in Pol $\mu$  expression with age suggests that the efficiency of that machinery, which is required for Ab affinity maturation, could decrease with age.

We also detected a high level of Pol $\mu$  mRNAs in the testis, a tissue in which many repair or translesional DNA polymerases are strongly expressed. A similar observation was made in the ovary and testis of zebrafish [34]. This expression pattern could be related to early embryonic expression. Indeed, as suggested above, Pol $\mu$  could be involved in the protection of the genome of germinal cells against DSBs and in limiting the transmission of mutations between generations. This hypothesis is reinforced by evidence that suggests a role for Pol $\lambda$ , another PolX family member, in DNA repair synthesis associated with meiosis, as high expression of this protein was observed in pachytene spermatocytes [42].

### Modulation of Pol $\mu$ Expression by Spaceflight-encountered Environmental Conditions

Because Pol $\mu$  plays a role in hematopoiesis [40], V(D)J recombination [20] and potentially SHM, and given our previous studies that demonstrated that the space environment affects Ig gene expression and decreases SHM [2,8,10–12,43], we investigated the effects of spaceflight conditions on Pol $\mu$  expression. Our results show that Pol $\mu$  transcription tends to decrease after 10 days of development under spaceflight conditions (Figure 3A) and that radiation, and not the other environmental modifications encountered during the flight, seem primarily involved in this down-regulation. The fact that radiation caused a decrease in Pol $\mu$  mRNA is not surprising because Aoufouchi et al. [23] showed that



ionizing radiation downregulates Pol $\mu$  transcription in RAMOS cells. The decrease in Pol $\mu$  mRNA in response to space radiation was however not confirmed at the protein level (Figures 3I and 3J). In HEK cells, the amount of Pol $\mu$  protein was shown to be increased by radiation exposure [28]. However, in that study the cells were exposed to 12 Gy of radiation, while in our study *P. walli* larvae were exposed to total doses of 1.9 mGy of  $\gamma$  rays and 2.0 mGy of neutrons. Differences in the doses of radiation likely explain why in this study no increase in Pol $\mu$  protein could be detected. Recently, Wyrobek et al. [44] evaluated transcriptional changes across 5 doses of radiation (1–10 cGy, considered to be low doses) and found Gadd45 to be a low dose response gene. However, even the doses considered to be low in that study were higher than the doses used here to mimic space radiation, thereby explaining why no change in Gadd45 mRNA was detected (Figure 3H).

High doses of radiation (>2 Gy) are known to induce the formation of reactive oxygen species (ROS) [45], which cause damage to biological molecules, including DNA and proteins. Recently, Smith et al. [46] showed that a low dose of radiation (417  $\mu$ Gy/h) produced too few ROS to affect cells. This paper supports our data showing the absence of oxidative stress in larvae that received neutrons and  $\gamma$  rays at a rate of 30  $\mu$ Gy/h to mimic radiations received during 10 days in the ISS (Figure 5).

Because astronauts are exposed to a multitude of environmental modifications during spaceflight, we investigated the cumulative effect of space radiation and another important stress encountered during spaceflight, namely, the disruption of the circadian rhythm. Our results show that the quantity of Pol $\mu$  mRNA doubled in larvae exposed to radiation and darkness (Figure 4A), while radiation alone decreased the production of Pol $\mu$  transcripts (Figure 3F), and darkness had no influence (Figure 3D). These observations show that cumulating environmental modifications can induce different effects, likely because a combination of two modifications causes greater stress, as revealed by the amounts of DNA-PKc and Gadd45 mRNAs, which were tripled in larvae irradiated in the dark (Figures 4B and 4C). However, once again, changes in Pol $\mu$  mRNA levels were not confirmed at the protein

level (Figures 4D and 4E). This observation, combined with the absence of protein oxidation (Figure 5), indicates that radiation encountered during a 10 day stay in the ISS has a limited impact, even though Pol $\mu$  transcription appears to be highly radiation-sensitive given its ability to be modified by the very low doses of radiation used in this study. Nevertheless, this conclusion does not exclude the possibilities that *i*) ROS could be produced under these conditions, as melatonin, whose synthesis is induced by darkness, can act as a free radical scavenger [47] and *ii*) exposure of astronauts to low doses of radiation during very long space mission, such as to Mars, could represent a threat.

## Conclusion

In conclusion, this paper examines polymerase  $\mu$  of a urodele amphibian and the expression of this enzyme in larvae subjected to spaceflight-induced environmental changes. We noted robust expression of Pol $\mu$  mRNA during early ontogenesis and in the testis, likely to protect rapidly dividing cells and germinal cells from DNA damage. Pol $\mu$  expression is also high when Ig gene rearrangements occur, confirming its involvement in V(D)J recombination. Pol $\mu$  transcripts are less abundant under spaceflight conditions because they are sensitive to space radiation. However, space radiation, alone or in combination with darkness, did not affect the synthesis of the corresponding Pol $\mu$  protein and did not induce oxidative stress, showing the limited impact of radiation encountered during a 10 day stay in the ISS.

## Acknowledgments

We thank D. Chapat for his support during the space mission, C. Huin-Schohn and M. Bascove for their assistance and the European Space Agency for the spaceflight opportunity.

## Author Contributions

Conceived and designed the experiments: SB JPF. Performed the experiments: VS NG. Analyzed the data: VS NG JPF. Wrote the paper: VS NG SB JPF.

## References

- Kimzey SL (1977) Hematology and immunology studies. In: Johnson RS, Dietlein LF, editors. Biomedical results from Skylab. Washington, DC: NASA. 248–282.
- Guéguinou N, Huin-Schohn C, Bascove M, Bueb JL, Tschirhart E, et al. (2009) Could spaceflight-associated immune system weakening preclude the expansion of human presence beyond Earth's orbit? *J Leukoc Biol* 86: 1027–1038. (doi: 10.1189/jlb.0309167).
- Cooper AD, Alder MN (2006) The evolution of adaptive immune systems. *Cell* 124: 815–822. (doi: 10.1016/j.cell.2006.02.001).
- Frippiat C, Kremarik P, Ropars A, Dournon C, Frippiat JP (2001) The recombination-activating gene 1 of *Pleurodeles waltl* (urodele amphibian) is transcribed in lymphoid tissues and in the central nervous system. *Immunogenetics* 52: 264–275. (doi: 10.1007/s002510000275).
- Schaerlinger B, Bascove M, Frippiat JP (2008) A new isotype of immunoglobulin heavy chain in the urodele amphibian *Pleurodeles waltl* predominantly expressed in larvae. *Mol Immunol* 45: 776–786. (doi: 10.1016/j.molimm.2007.06.356).
- Bascove M, Frippiat JP (2010) Molecular characterization of *Pleurodeles waltl* activation-induced cytidine deaminase. *Mol Immunol* 47: 1640–1649. (doi: 10.1016/j.molimm.2010.01.005).
- Boudarra N, Frippiat C, Dournon C, Frippiat JP (2002) An alternative internal splicing site defines new *Ikaros* isoforms in *Pleurodeles waltl*. *Dev Comp Immunol* 26: 659–673. (doi: 10.1016/S0145-305X(02)00014-9).
- Boxio R, Dournon C, Frippiat JP (2005) Effects of a long-term spaceflight on immunoglobulin heavy chains of the urodele amphibian *Pleurodeles waltl*. *J Appl Physiol* 98: 905–910. (doi: 10.1152/jappphysiol.00957.2004).
- Konstantinova IV, Rykova MP, Lesnyak AT, Antropova EA (1993) Immune changes during long-duration missions. *J Leuk Biol* 54: 189–201.
- Bascove M, Huin-Schohn C, Guéguinou N, Tschirhart E, Frippiat JP (2009) Spaceflight-associated changes in immunoglobulin VH gene expression in the amphibian *Pleurodeles waltl*. *FASEB J* 23: 1607–1615. (doi: 10.1096/fj.08-121327).
- Bascove M, Guéguinou N, Schaerlinger B, Gauquelin-Koch G, Frippiat JP (2011) Decrease in antibody somatic hypermutation frequency under extreme, extended spaceflight conditions. *FASEB J* 25: 2947–2955. (doi: 10.1096/fj.11-185215).
- Huin-Schohn C, Guéguinou N, Schenten V, Bascove M, Gauquelin-Koch G, et al. (2013) Gravity changes during animal development affect IgM heavy-chain transcription and probably lymphopoiesis. *FASEB J* 27: 333–341. (doi: 10.1096/fj.12-217547).
- Bassing CH, Swat W, Alt FW (2002) The mechanism and regulation of chromosomal V(D)J recombination. *Cell* 109: S45–55. (doi: 10.1016/S0092-8674(02)00675-X).
- Gu J, Lu H, Tippin B, Shimazaki N, Goodman MF et al. (2007) XRCC4:DNA ligase IV can ligate incompatible DNA ends and can ligate across gaps. *EMBO J* 26: 1010–1023. (doi: 10.1038/sj.emboj.7601559).
- Lieber MR, Lu H, Gu J, Schwarz K (2008) Flexibility in the order of action and in the enzymology of the nuclease, polymerases, and ligase of vertebrate non-homologous DNA end joining: relevance to cancer, aging, and the immune system. *Cell Res* 18: 125–133. (doi: 10.1038/cr.2007.108).
- Ma Y, Lu H, Tippin B, Goodman MF, Shimazaki N, et al. (2004) A biochemically defined system for mammalian nonhomologous DNA end joining. *Mol Cell* 16: 701–713. (doi: 10.1016/j.molcel.2004.11.017).
- Nick McElhinny SA, Havener JM, Garcia-Diaz M, Juárez R, Bebenek K, et al. (2005) A gradient of template dependence defines distinct biological roles for family X polymerases in nonhomologous end joining. *Mol Cell* 19: 357–366. (doi: 10.1016/j.molcel.2005.06.012).
- Moon A F, Garcia-Diaz M, Batra VK, Beard WA, Bebenek K, et al. (2007) The X family portrait: Structural insights into biological functions of X family polymerases. *DNA Repair (Amst)* 12: 1709–1725. (doi: 10.1016/j.dnarep.2007.05.009).
- Bertocci B, De Smet A, Weill JC, Reynaud CA (2006) Nonoverlapping functions of DNA polymerases  $\mu$ ,  $\lambda$ , and terminal deoxynucleotidyltransferase

- during immunoglobulin V(D)J recombination *in vivo*. *Immunity* 25: 31–41. (doi: 10.1016/j.immuni.2006.04.013).
20. Bertocci B, De Smet A, Berek C, Weill JC, Reynaud CA (2003) Immunoglobulin kappa light chain gene rearrangement is impaired in mice deficient for DNA polymerase mu. *Immunity* 19: 203–211. (doi: 10.1016/S1074-7613(03)00203-6).
  21. Gilfillan S, Dierich A, Lemeur M, Benoist C, Mathis D (1993) Mice lacking TdT: mature animals with an immature lymphocyte repertoire. *Science* 261: 1175–1178. (doi: 10.1126/science.8356452).
  22. Komori T, Okada A, Stewart V, Alt FW (1993) Lack of N regions in antigen receptor variable region genes of TdT-deficient lymphocytes. *Science* 261: 1171–1175. (doi: 10.1126/science.8356451).
  23. Aoufouchi S, Flatter E, Dahan A, Faili A, Bertocci B, et al. (2000) Two novel human and mouse DNA polymerases of the polX family. *Nucleic Acids Res* 28: 3684–3693. (doi: 10.1093/nar/28.18.3684).
  24. Ruiz JF, Lucas D, Garcia-Palmero E, Saez AI, González MA, et al. (2004) Overexpression of human DNA polymerase mu (Pol mu) in a Burkitt's lymphoma cell line affects the somatic hypermutation rate. *Nucleic Acids Res* 32: 5861–5873. (doi: 10.1093/nar/gkh929).
  25. Domínguez O, Ruiz JF, Laín de Lera T, García-Díaz M, González MA, et al. (2000) DNA polymerase mu (Pol mu), homologous to TdT, could act as a DNA mutator in eukaryotic cells. *EMBO J* 19: 1731–1742. (doi: 10.1093/emboj/19.7.1731).
  26. Ruiz JF, Domínguez O, Laín de Lera T, Garcia-Díaz M, Bernad A, et al. (2001) DNA polymerase mu, a candidate hypermutase? *Philos Trans R Soc Lond B Biol Sci* 356: 99–109. (doi: 10.1098/rstb.2000.0754).
  27. Reynaud CA, Frey S, Aoufouchi S, Faili A, Bertocci B, et al. (2001) Transcription, beta-like DNA polymerases and hypermutation. *Philos Trans R Soc Lond B Biol Sci* 356: 91–97. (doi: 10.1098/rstb.2000.0753).
  28. Mahajan KN, McElhinny SAN, Mitchell BS, Ramsden DA (2002) Association of DNA polymerase mu (pol mu) with Ku and ligase IV: role for pol mu in end-joining double-strand break repair. *Mol Cell Biol* 22: 5194–5202. (doi: 10.1128/MCB.22.14.5194-5202.2002).
  29. Chayot R, Danckaert A, Montagne B, Ricchetti M (2010) Lack of DNA polymerase  $\mu$  affects the kinetics of DNA double-strand break repair and impacts on cellular senescence. *DNA Repair (Amst)* 9: 1187–1199. (doi: 10.1016/j.dnarep.2010.09.001).
  30. Space Radiation Expert Group Report (2012) Towards Human Exploration of Space: a European Strategy. European Science Foundation publication, Strasbourg, France.
  31. Gallien L, Durocher M (1957) Table chronologique du développement chez *Pleurodeles waltl Michah*. *Bull Biol Fr Belg* 91: 97–114.
  32. Pfaffl MW (2001) A new mathematical model for relative quantification in real-time RTPCR. *Nucleic Acids Res* 29: e45.
  33. Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, et al. (2002) Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol* 3:research0034.
  34. Beetz S, Diekhoff D, Steiner LA (2007) Characterization of terminal deoxynucleotidyl transferase and polymerase mu in zebrafish. *Immunogenetics* 59: 735–744. (doi: 10.1007/s00251-007-0241-7).
  35. Scheiermann C, Kunisaki Y, Frenette PS (2013) Circadian control of the immune system. *Nat Rev Immunol* 3: 190–198. (doi: 10.1038/nri3386).
  36. Lucas D, Laín de Lera T, González MA, Ruiz JF, Domínguez O, et al. (2005) Polymerase mu is up-regulated during the T cell-dependent immune response and its deficiency alters developmental dynamics of spleen centroblasts. *Eur J Immunol* 35: 1601–1611. (doi: 10.1002/eji.200526015).
  37. Heyer BS, MacAuley A, Behrendtsen O, Werb Z (2000) Hypersensitivity to DNA damage leads to increased apoptosis during early mouse development. *Genes Dev* 14: 2072–2084. (doi: 10.1101/gad.14.16.2072).
  38. Reddy YV, Perkins EJ, Ramsden DA (2006) Genomic instability due to V(D)J recombination-associated transposition. *Genes Dev* 20: 1575–1582. (doi: 10.1101/gad.1432706).
  39. Capp JP, Boudsocq F, Bernard AG, Lopez BS, Cazaux C, et al. (2007) Involvement of DNA polymerase mu in the repair of a specific subset of DNA double-strand breaks in mammalian cells. *Nucleic Acids Res* 35: 3551–3560. (doi: 10.1093/nar/gkm243).
  40. Gozalbo-López B, Andrade P, Terrados G, Andrés B, Serrano N, et al. (2009) A role for DNA polymerase mu in the emerging DJH rearrangements of the postgastrulation mouse embryo. *Mol Cell Biol* 29: 1266–1275. (doi: 10.1128/MCB.01518-08).
  41. Bertocci B, De Smet A, Flatter E, Dahan A, Bories JC, et al. (2002) Cutting edge: DNA polymerases mu and lambda are dispensable for Ig gene hypermutation. *J Immunol* 168: 3702–3706.
  42. García-Díaz M, Domínguez O, López-Fernández LA, de Lera LT, Saniger ML, et al. (2000) DNA polymerase lambda (Pol lambda), a novel eukaryotic DNA polymerase with a potential role in meiosis. *J Mol Biol* 301: 851–867. (doi: 10.1006/jmbi.2000.4005).
  43. Guéguinou N, Bojados M, Jamon M, Derradji H, Baatout S, et al. (2012) Stress response and humoral immune system alterations related to chronic hypergravity in mice. *Psychoneuroendocrinology* 37: 137–147. (doi: 10.1016/j.psycuen.2011.05.015).
  44. Wyrobek AJ, Manohar CF, Krishnan VV, Nelson DO, Furtado MR, et al. (2011) Low dose radiation response curves, networks and pathways in human lymphoblastoid cells exposed from 1 to 10cGy of acute gamma radiation. *Mutat Res* 722: 119–130. (doi: 10.1016/j.mrgentox.2011.03.002).
  45. Leach K, Van Tuyle G, Lin PS, Schmidt-Ullrich R, Mikkelsen RB (2001) Ionizing radiation-induced, mitochondria-dependent generation of reactive oxygen/nitrogen. *Cancer Res* 61: 3894–3901.
  46. Smith JT, Willey NJ, Hancock JT (2012) Low dose ionizing radiation produces too few reactive oxygen species to directly affect antioxidant concentrations in cells. *Biol Lett* 8: 594–597. (doi: 10.1098/rsbl.2012.0150).
  47. Galano A, Tan DX, Reiter RJ (2012) On the free radical scavenging activities of melatonin's metabolites, AFMK and AMK. *J Pineal Res* 54: 245–57. (doi: 10.1111/jpi.12010).