



Dear Author

Here are the proofs of your article.

- You can submit your corrections **online** or by **fax**.
- For **online** submission please insert your corrections in the online correction form. Always indicate the line number to which the correction refers.
- For **fax** submission, please ensure that your corrections are clearly legible. Use a fine black pen and write the correction in the margin, not too close to the edge of the page.
- Together with the proof please return the cover sheet (including the *Copyright Transfer Statement*) and the *Offprint Order Form*. They can either be scanned and sent electronically or sent by fax.
- Remember to note the journal title, article number, and your name when sending your response via e-mail, fax or regular mail.
- **Check** the metadata sheet to make sure that the header information, especially author names and the corresponding affiliations are correctly shown.
- **Check** the questions that may have arisen during copy editing and insert your answers/corrections.
- **Check** that the text is complete and that all figures, tables and their legends are included. Also check the accuracy of special characters, equations, and electronic supplementary material if applicable. If necessary refer to the *Edited manuscript*.
- The publication of inaccurate data such as dosages and units can have serious consequences. Please take particular care that all such details are correct.
- Please **do not** make changes that involve only matters of style. We have generally introduced forms that follow the journal's style. Substantial changes in content, e.g., new results, corrected values, title and authorship are not allowed without the approval of the responsible editor. In such a case, please contact the Editorial Office and return his/her consent together with the proof.
- If we do not receive your corrections **within 48 hours**, we will send you a reminder.

Please note

Your article will be published **Online First** approximately one week after receipt of your corrected proofs. This is the **official first publication** citable with the DOI. **Further changes are, therefore, not possible.**

After online publication, subscribers (personal/institutional) to this journal will have access to the complete article via the DOI using the URL:

<http://dx.doi.org/10.1007/s12031-014-0249-z>

If you would like to know when your article has been published online, take advantage of our free alert service. For registration and further information, go to:

<http://www.link.springer.com>.

Due to the electronic nature of the procedure, the manuscript and the original figures will only be returned to you on special request. When you return your corrections, please inform us, if you would like to have these documents returned.

The **printed version** will follow in a forthcoming issue.

Metadata of the article that will be visualized in OnlineFirst

1	Article Title	Molecular Mechanisms Underlying the Nephroprotective Effects of PACAP in Diabetes		
2	Article Sub- Title			
3	Article Copyright - Year	Springer Science+Business Media New York 2014 (This will be the copyright line in the final PDF)		
4	Journal Name	Journal of Molecular Neuroscience		
5	Corresponding Author	Family Name	Reglodi	
6		Particle		
7		Given Name	Dora	
8		Suffix		
9		Organization	University of Pécs	
10		Division	Department of Anatomy, PTE-MTA "Lendulet" PACAP Research Team	
11		Address	Szigeti u 12, Pécs 7624, Hungary	
12		e-mail	dora.reglodi@aok.pte.hu	
13	Author	Family Name	Banki	
14		Particle		
15		Given Name	Eszter	
16		Suffix		
17		Organization	University of Pécs	
18		Division	Department of Anatomy, PTE-MTA "Lendulet" PACAP Research Team	
19		Address	Szigeti u 12, Pécs 7624, Hungary	
20		e-mail		
21	Author	Family Name	Kovacs	
22		Particle		
23		Given Name	Krisztina	
24		Suffix		
25		Organization	University of Pécs	
26		Division	Department of Biochemistry and Medical Chemistry	
27		Address	Pécs, Hungary	
28		e-mail		

29		Family Name	Nagy
30		Particle	
31		Given Name	Daniel
32		Suffix	
33		Organization	University of Pécs
34	Author	Division	Department of Anatomy, PTE-MTA "Lendulet" PACAP Research Team
35		Address	Szigeti u 12, Pécs 7624, Hungary
36		Organization	University of Pécs
37		Division	Department of Pharmacology and Pharmacotherapy
38		Address	Pécs, Hungary
39		e-mail	
<hr/>			
40		Family Name	Juhasz
41		Particle	
42		Given Name	Tamas
43		Suffix	
44	Author	Organization	University of Debrecen
45		Division	Department of Anatomy, Histology and Embryology, Medical and Health Science Center
46		Address	Debrecen, Hungary
47		e-mail	
<hr/>			
48		Family Name	Degrell
49		Particle	
50		Given Name	Peter
51		Suffix	
52	Author	Organization	University of Pécs
53		Division	Internal Medicine 2/ Nephrology Center
54		Address	Pécs, Hungary
55		e-mail	
<hr/>			
56		Family Name	Csanaky
57		Particle	
58		Given Name	Katalin
59	Author	Suffix	
60		Organization	University of Pécs
61		Division	Department of Anatomy, PTE-MTA "Lendulet" PACAP Research Team
62		Address	Szigeti u 12, Pécs 7624, Hungary

63		e-mail	
64		Family Name	Kiss
65		Particle	
66		Given Name	Peter
67		Suffix	
68	Author	Organization	University of Pécs
69		Division	Department of Anatomy, PTE-MTA "Lendulet" PACAP Research Team
70		Address	Szigeti u 12, Pécs 7624, Hungary
71		e-mail	
72		Family Name	Jancso
73		Particle	
74		Given Name	Gabor
75		Suffix	
76	Author	Organization	University of Pécs
77		Division	Surgical Research and Techniques
78		Address	Pécs, Hungary
79		e-mail	
80		Family Name	Toth
81		Particle	
82		Given Name	Gabor
83		Suffix	
84	Author	Organization	University of Szeged
85		Division	Department of Medical Chemistry
86		Address	Szeged, Hungary
87		e-mail	
88		Family Name	Tamas
89		Particle	
90		Given Name	Andrea
91		Suffix	
92	Author	Organization	University of Pécs
93		Division	Department of Anatomy, PTE-MTA "Lendulet" PACAP Research Team
94		Address	Szigeti u 12, Pécs 7624, Hungary
95		e-mail	
96		Received	15 November 2013
97	Schedule	Revised	
98		Accepted	22 January 2014

99 Abstract

Diabetic nephropathy is the leading cause of end-stage renal failure and accounts for 30–40 % of patients entering renal transplant programmes. The nephroprotective effects of the neuropeptide pituitary adenylate cyclase-activating polypeptide (PACAP38) against diabetes have been shown previously, but the molecular mechanisms responsible for these effects remain unknown. In the present study, we showed that PACAP treatment counteracted the diabetes-induced increase in the level of the proapoptotic pp38MAPK and cleaved caspase-3 and also decreased the p60 subunit of NFκB. The examined antiapoptotic factors, including pAkt and pERK1/2, showed a slight increase in the diabetic kidneys, while PACAP treatment resulted in a notable elevation of these proteins. PCR and Western blot revealed the downregulation of fibrotic markers, like collagen IV and TGF-β1 in the kidney. PACAP treatment resulted in increased expression of the antioxidant glutathione. We conclude that the nephroprotective effect of PACAP in diabetes is, at least partly, due to its antiapoptotic, antifibrotic and antioxidative effect in addition to the previously described antiinflammatory effect.

100 Keywords
separated by ' - '

PACAP - Diabetes - Kidney - Oxidative stress - Apoptosis

101 Foot note
information

4 **Molecular Mechanisms Underlying the Nephroprotective**
5 **Effects of PACAP in Diabetes**7 **Eszter Banki · Krisztina Kovacs · Daniel Nagy · Tamas Juhasz ·**
8 **Peter Degrell · Katalin Csanaky · Peter Kiss · Gabor Jancso · Gabor Toth ·**
9 **Andrea Tamas · Dora Reglodi**12 Received: 15 November 2013 / Accepted: 22 January 2014
13 © Springer Science+Business Media New York 201414 **Abstract** Diabetic nephropathy is the leading cause of end-
15 stage renal failure and accounts for 30–40 % of patients enter-
16 ing renal transplant programmes. The nephroprotective effects
17 of the neuropeptide pituitary adenylate cyclase-activating poly-
18 peptide (PACAP38) against diabetes have been shown previ-
19 ously, but the molecular mechanisms responsible for these
20 effects remain unknown. In the present study, we showed that
21 PACAP treatment counteracted the diabetes-induced increase
22 in the level of the proapoptotic pp38MAPK and cleaved
23 caspase-3 and also decreased the p60 subunit of NFκB. The
24 examined antiapoptotic factors, including pAkt and pERK1/2,showed a slight increase in the diabetic kidneys, while PACAP 25
treatment resulted in a notable elevation of these proteins. PCR 26
and Western blot revealed the downregulation of fibrotic 27
markers, like collagen IV and TGF-β1 in the kidney. PACAP 28
treatment resulted in increased expression of the antioxidant 29
glutathione. We conclude that the nephroprotective effect of 30
PACAP in diabetes is, at least partly, due to its antiapoptotic, 31
antifibrotic and antioxidative effect in addition to the previously 32
described antiinflammatory effect. 33**Keywords** PACAP · Diabetes · Kidney · Oxidative stress · 34
Apoptosis 35E. Banki · D. Nagy · K. Csanaky · P. Kiss · A. Tamas ·
D. Reglodi (✉)
Department of Anatomy, PTE-MTA “Lendulet” PACAP Research
Team, University of Pécs, 7624 PécsSzigeti u 12, Hungary
e-mail: dora.reglodi@aok.pte.huK. Kovacs
Department of Biochemistry and Medical Chemistry,
University of Pécs, Pécs, HungaryD. Nagy
Department of Pharmacology and Pharmacotherapy,
University of Pécs, Pécs, HungaryT. Juhasz
Department of Anatomy, Histology and Embryology, Medical and
Health Science Center, University of Debrecen, Debrecen, HungaryP. Degrell
Internal Medicine 2/ Nephrology Center, University of Pécs, Pécs,
HungaryG. Jancso
Surgical Research and Techniques, University of Pécs, Pécs,
HungaryG. Toth
Department of Medical Chemistry, University of Szeged, Szeged,
Hungary**Introduction** 36In the present study, we aimed at investigating the nephro- 37
protective mechanisms of the neuropeptide pituitary adenylate 38
cyclase-activating polypeptide (PACAP) in an experimental 39
diabetic nephropathy. Diabetic nephropathy (DN) is a common 40
microvascular complication of diabetes. The development as 41
well as the progression of the disease is multifactorial due to its 42
profound genetic background besides the well-known environ- 43
mental impacts, explaining why 30–40 % of all diabetic patients 44
suffer from this diabetic complication (Klein et al. 1984). 45
Glucose uptake is insulin independent in neuronal, retinal, 46
glomerular and endothelial cells, leading to excessive glucose 47
uptake in extracellular hyperglycemia (Di Mario and Pugliese 48
2001). Although a few decades ago diabetic nephropathy was 49
considered a primarily glomerular disease, nowadays increasing 50
emphasis is put on the tubulointerstitial alterations. Indeed, the 51
stage of tubulointerstitial injury is considered to be a reliable 52
prognostic factor in progressive kidney diseases, including di- 53
abetic nephropathy (Nath 1998). Key factors that are involved in 54
diabetic kidney damage are as follows: (1) oxidative stress, (2) 55
overproduction of advanced glycation end products (AGE), (3) 56

57	apoptosis, (4) excessive production of pro-sclerotic growth factors and (5) inflammation due to the overproduction of pro-inflammatory cytokines (Gnudi 2012; Sun et al. 2013). Since the neuropeptide PACAP is known to exert anti-inflammatory, antiapoptotic and antioxidant effects, it seems to be a suitable candidate to prevent the development or delay the progression of DN.	110
58		111
59		112
60		113
61		114
62		115
63		116
64	PACAP is a member of the vasoactive intestinal polypeptide (VIP)/secretin/glucagon family and exists in two biologically active forms, PACAP1-27 and PACAP1-38. PACAP38 has been shown to have more prolonged effects compared to PACAP27 in most studies (Araki and Takagi 1992; Lindén et al. 1999). PACAP acts via G-protein-coupled receptors: PAC1, specific for PACAP, and VPAC1 and VPAC2 which also bind VIP with the same affinity.	117
65		
66		
67		
68		
69		
70		
71		
72	In addition to its very first known effect, namely adenylate cyclase activation in the hypophysis, PACAP exerts numerous effects in the endocrine, respiratory, gastrointestinal and urogenital systems (Miyata et al. 1989; Girard et al. 2012; Koppan et al. 2012; Moody et al. 2012; Nedvig et al. 2012; Syed et al. 2012; Wada et al. 2013). Moreover, the neuropeptide has been shown to be involved in neuroprotection and general cytoprotection. PACAP is also involved in the regulation of carbohydrate metabolism, although its exact role seems to be complex. Its ability to protect β cells and enhance insulin secretion glucose-dependently has been shown in several studies (Sakurai et al. 2011). Recently, PACAP has been proven to upregulate selenoprotein T in pancreatic β cells, leading to increased insulin secretion (Prevost et al. 2013). However, PACAP also effectively stimulates the release of adrenalin and glucagon. As a result, publications seem to be rather contradictory in the effect of PACAP treatment on blood glucose levels (Sekiguchi et al. 1994; Filipsson et al. 1998; Yada et al. 2000).	118
73		
74		
75		
76		
77		
78		
79		
80		
81		
82		
83		
84		
85		
86		
87		
88		
89		
90		
91	PACAP exerts renoprotective effects against several nephrotoxic agents, like hydrogen peroxide-induced oxidative stress in vitro and in vivo or ischemia/reperfusion injury (Szakaly et al. 2008; Horvath et al. 2011; Khan et al. 2012; Reglodi et al. 2012). PACAP has also been shown to attenuate kidney injuries induced by multiple myeloma, cyclosporine A, gentamicin and short-term diabetes (Arimura et al. 2006; Li et al. 2007, 2008). Recently, we have provided evidence for similar protection against 8-week diabetes-induced kidney damage (Banki et al. 2013).	119
92		
93		
94		
95		
96		
97		
98		
99		
100		
101	The protective effects of PACAP are mediated through its anti-inflammatory, antiapoptotic and antioxidative effects. The anti-inflammatory effects involve the inhibition of pro-inflammatory cytokine (i.e. tumor necrosis factor α (TNF α) and interleukin-6 (IL-6) production and NF κ B activation through PAC1- and VPAC1 receptor-mediated signaling (Arimura et al. 2006). We have already proven the importance of the anti-inflammatory effect in long-term diabetic nephropathy, resulting in marked downregulation of several cytokines, like	120
102		
103		
104		
105		
106		
107		
108		
109		
	cytokine-induced neutrophil chemokine (CINC-1), tissue inhibitor of metalloproteinase 1 (TIMP-1), lipopolysaccharide-induced CXC chemokine (LIX) and monokine induced by gamma interferon (MIG) (Banki et al. 2013). However, the further mechanisms leading to the significant nephroprotective effect in diabetic nephropathy remain unknown.	121
		122
		123
		124
		125
		126
		127
		128
		129
		130
		131
		132
		133
		134
		135
		136
		137
		138
		139
		140
		141
		142
		143
		144
		145
		146
		147
		148
		149
		150
		151
		152
		153
		154
		155
		156

157	NM_007393, amplicon size 462 bp). Amplifications were	Western blot signals was measured by using ImageJ 1.40 g	208
158	performed in a thermal cycler (Labnet MultiGene™ 96-well	freeware and the results were normalised to actin.	209
159	Gradient Thermal Cycler; Labnet International, Edison, NJ,		
160	USA) in a final volume of 25 µL (containing 1 µL forward	Biochemical Assay of Oxidative Stress Markers	210
161	and reverse primers [0.4 µM], 0.5 µL dNTP [200 µM] and 5		
162	U of Promega GoTaq® DNA polymerase in 1× reaction	Malondialdehyde (MDA) was measured as previously de-	211
163	buffer) as follows: 95 °C, 2 min, followed by 35 cycles	scribed (Placer et al. 1966). Briefly, MDA was measured in	212
164	(denaturation, 94 °C, 1 min; annealing at 52 °C for Col4a1	kidney homogenates with the addition of TBA (saturated	213
165	or 53 °C for Actb for 1 min; extension, 72 °C, 90 s) and then	thiobarbituric acid in 10 % perchloric acid)-TCA (20 % tri-	214
166	72 °C, 10 min. PCR products were analysed by electrophore-	chloroacetic acid) reagent. After incubation at 100 °C for	215
167	sis in 1.2 % agarose gel containing ethidium bromide. Actin	20 min, samples were placed in ice-cold water and were	216
168	was used as the internal control. Optical density of signals was	centrifuged for 15 min at 4,000 rpm. MDA concentration	217
169	measured by using ImageJ 1.40 g freeware and results were	was determined spectrophotometrically by measuring the ab-	218
170	normalised to actin.	sorbance at 532 nm against TBA+TCA reagent. From the	219
		concentration vs. extinction curve, the MDA value of the	220
171	Western Blot	tissue sample could be calculated in micromole per gram	221
		tissue weight.	222
172	The specimens were put into 100 µL of ice-cold homogeni-	Reduced glutathione (GSH) was quantified as described by	223
173	zation buffer containing 50 mM Tris-HCl buffer (pH 7.0),	Sedlak and Lindsay (1968). After adding 10 % TCA, kidney	224
174	10 µg/mL Gordox, 10 µg/mL leupeptine, 1 mM	homogenates were centrifuged for 15 min at 4,000 rpm. TRIS-	225
175	phenylmethylsulphonyl-fluoride (PMSF), 5 mM benzamidine	buffer was added to the supernatant and samples were mea-	226
176	and 10 µg/mL trypsin inhibitor as protease inhibitors. Samples	sured at 412 nm after adding DTNB to the mixture. Values of	227
177	were stored at -70 °C. Tissue samples were sonicated by	glutathione were expressed in micromole per gram tissue	228
178	pulsing burst for 30 s at 40 A (Cole-Parmer, IL, USA).	weight.	229
179	Samples for SDS-PAGE were prepared by the addition of	Kidney homogenates were centrifuged for 20 min at	230
180	fivefold concentrated electrophoresis sample buffer (20 mM	16,000 rpm, and the supernatant was used to measure the	231
181	Tris-HCl pH 7.4, 0.01 % bromophenol blue dissolved in 10 %	concentration of superoxide dismutase (SOD) as described	232
182	SDS, 100 mM β-mercaptoethanol) to kidney homogenates to	previously (Misra and Firdovich 1972). SOD inhibited the	233
183	set equal protein concentration of samples and boiled for	transformation of adrenaline to adrenochrome, which	234
184	10 min. About 40 µg of protein was separated by 7.5 %	absorbed maximally at 480 nm. Quantification of SOD is	235
185	SDS-PAGE gel for detection of tAkt, pAkt, pERK1/2,	based on the degree of inhibition. The value of SOD was	236
186	p38MAPK, pp38MAPK, cleaved caspase-3, TGF-β1, colla-	given in international units per gram tissue weight.	237
187	gen type IV, β-actin and NFκB. Proteins were transferred		
188	electrophoretically to nitrocellulose membranes. After	Electron Microscopy	238
189	blocking with 5 % non-fat dry milk in phosphate-buffered		
190	saline with 0.1 % Tween 20 (PBST), membranes were washed	Kidney samples were cut into maximum 1 mm ³ pieces and	239
191	and exposed to the following primary antibodies overnight at	were fixed in 5 % glutaraldehyde. Post-fixation was per-	240
192	4 °C: monoclonal anti-Akt, monoclonal phospho-specific	formed with 1 % osmium tetroxide. After dehydration in	241
193	anti-Akt-1 Ser473, monoclonal phospho-specific anti-ERK1/	ascending alcohol and subsequent transfer to propylene oxide,	242
194	2 Thr202/Tyr204, monoclonal anti-p38 MAPK, monoclonal	samples were embedded in Araldite resin. Semithin sections	243
195	phospho-specific anti-p38 MAPK (1:500; Cell Signaling	were cut by a ultramicrotome (Leica Ultracut R) and stained	244
196	Technology, USA), monoclonal anti-NFκB, monoclonal	by toluidine blue. Ultrathin sections were prepared from the	245
197	anti-caspase-3 (1:500; Santa Cruz, USA), polyclonal anti-	area of interest and were contrasted by uranyl-acetic acid and	246
198	TGF-β1 antibody (1:400; Abcam, Cambridge, UK), mono-	lead citrate. Slides were eventually examined using JEOL	247
199	clonal anti-collagen type IV antibody (1:400; Chemicon/	1200 EX-II electron microscope.	248
200	Millipore, USA) and monoclonal anti-actin antibody		
201	(1:10,000; Sigma, Hungary) were used. After washing for	Statistical Analysis	249
202	40 min in PBST, membranes were incubated with anti-		
203	mouse IgG (1:1500; Bio-Rad Laboratories, USA) or anti-	Statistical analysis was performed by Microsoft Office Excel	250
204	rabbit IgG (1:3,000; Bio-Rad Laboratories, USA). Signals	and GraphPad software. Analysis of variance (ANOVA) with	251
205	were detected by enhanced chemiluminescence (Millipore,	Bonferroni correction was used to detect significant differ-	252
206	USA) according to the instructions of the manufacturer.	ences between groups. <i>p</i> value less than 0.05 was considered	253
207	Actin was used as the internal control. Optical density of	to be statistically significant.	254

255 **Results**

256 RT-PCR analysis revealed marked elevation in the collagen IV
 257 mRNA, a collagen uniquely present in the basement mem-
 258 brane. PACAP treatment successfully counteracted this in-
 259 crease (Fig. 1). We then measured the protein expression of
 260 two key factors involved in the fibrotic processes of diabetic
 261 nephropathy, namely collagen IV and TGF-β1. Diabetes result-
 262 ed in excessive expression of collagen IV and TGF-β1, which
 263 was attenuated by PACAP38, reaching the level of normal
 264 kidneys in the case of collagen IV (Figs. 2 and 3a, b).

265 Next we investigated the levels of anti- and proapoptotic
 266 proteins by Western blot. PACAP alone caused increased
 267 expression of the phosphorylated form of Akt. Diabetic neph-
 268 ropathy is accompanied by excessive apoptosis, shown by
 269 the upregulation of the phosphorylated forms of the proapoptotic
 270 pp38MAPK. However, the antiapoptotic Akt and ERK1/2 were also
 271 activated. PACAP treatment in diabetic animals led to a remark-
 272 able increase in the activation of the antiapoptotic factors, like
 273 pAkt and pERK1/2, and decreased the level of pp38MAPK. PACAP
 274 treatment was effective in decreasing the elevated cleaved cas-
 275 pase-3 levels observed in diabetic animals. We then aimed at
 276 measuring the level of p60 NFκB, a protein known to control both
 277 cytokine production and cell survival. We found that p60 NFκB
 278 was upregulated in the diabetic samples. PACAP treatment in
 279 diabetic kidneys resulted in a remarkable decrease in the p60
 280 subunit of NFκB (Figs. 2 and 3c-h).

282 Biochemical assay of the oxidative stress markers revealed
 283 a significant elevation in the kidney GSH concentration of the
 284 PACAP-treated diabetic group compared to the untreated
 285 diabetic one. No changes were observed in the kidney SOD
 286 or MDA concentration of the diabetic groups compared to that
 287 of the intact animals; however, PACAP caused a significant
 288 increase in the SOD level of the control animals (Fig. 4).

289 Electron microscopy revealed segmental thickening of the
 290 glomerular basement membrane (GBM) in several parts of the
 291 untreated diabetic glomeruli. The thickness of these parts of
 292 the GBM was significantly greater than the GBM in control,




Fig. 1 mRNA expression pattern of collagen type IV in control, PACAP-treated control, diabetic and PACAP-treated diabetic kidneys. For RT-PCR reactions, actin was used as the control. *** $p < 0.001$ vs. control; #### $p < 0.001$ vs. control + PACAP; &&& $p < 0.001$ vs. diabetes




Fig. 2 Western blots of collagen type IV, TGF-β1, pERK1/2, pAkt, pp38MAPK, cleaved caspase-3 and p60 NFκB in untreated or PACAP-treated control and diabetic animal groups. For Western blot analysis, β-actin was used as a control

PACAP-treated control and PACAP-treated diabetic kidneys. 293
 However, we could not detect any changes between the non- 294
 thickened part of the GBM in diabetic animals and the thick- 295
 ness of the GBM in the control. Most importantly, PACAP- 296
 treated diabetic animals did not show this focal segmental 297
 thickening; there was no difference between PACAP-treated 298
 or untreated control and PACAP-treated diabetic animals. 299
 Podocytes in the PACAP-treated diabetic kidneys did not 300
 show any morphological alterations compared to the control 301
 groups, although severe podocyte injury was present in the 302
 diabetic glomeruli with marked foot process broadening and 303
 extensive flattening (Fig. 5). 304

Discussion

In the present study, we demonstrated that in vivo PACAP 306
 treatment exhibits protective effect through inhibiting apoptotic 307
 fibrotic and oxidative pathways, key mediators in the 308
 development and progression of diabetic nephropathy and 309
 preventing diabetes-induced podocyte injury in 8-week dia- 310
 betes. The present experiment was based on our previous 311
 finding, showing that PACAP38 effectively counteracted the 312




Fig. 3 Effect of 8-week PACAP treatment and diabetes on the protein expression of collagen type IV (a), TGF-β1 (b), pERK1 (c), pERK2 (d), pAkt (e), pp38MAPK (f), cleaved caspase-3 (g) and p60 NFκB (h).

p*<0.05; *p*<0.01; ****p*<0.001 vs. control; #*p*<0.05; ###*p*<0.01; ####*p*<0.001 vs. control+PACAP; &*p*<0.05; &&*p*<0.01; &&&*p*<0.001 vs. diabetes

313 histological alterations of 8-week diabetic nephropathy.
 314 Decreased PAS-positive area expansion, glycogen deposits
 315 in tubular epithelial cells and significantly diminished vascu-
 316 lar hyalinosis proved the ameliorative effect of PACAP.
 317 Moreover, we provided evidence that this effect is at least
 318 partially mediated through its antiinflammatory effect (Banki
 319 et al. 2013).

320 As we mentioned earlier, the effect of exogenous PACAP
 321 on glucose homeostasis is not fully elucidated. However, the
 322 pivotal role of endogenous PACAP on regulating blood sugar
 323 level seems to be more evident. Chronic administration of the
 324 antagonist PACAP6-27 was found to deteriorate insulin sen-
 325 sitivity and glucose tolerance in mice (Green et al. 2006).
 326 Newborn PACAP knockout mice showed decreased glucose
 327 and intrahepatic glucagon levels, but significantly higher

328 insulin levels compared to the control PACAP^{+/+} mice (Gray 328
 329 et al. 2001). Although PACAP protects pancreatic β cells 329
 330 against streptozotocin-induced apoptosis, under our experi- 330
 331 mental circumstances, we could not find significant changes 331
 332 between the blood sugar levels of the PACAP-treated and 332
 333 untreated diabetic rats after 8 weeks of survival, meaning that 333
 334 this factor cannot be responsible for the ameliorative effect in 334
 335 diabetic nephropathy (Onoue et al. 2008; Banki et al. 2013). 335

336 In the present study, we found that diabetes resulted in 336
 337 upregulated NFκB levels compared to the controls, while 337
 338 PACAP treatment effectively diminished the renal NFκB 338
 339 expression. Several previous studies revealed that PACAP, which 339
 340 is structurally similar to the related VIP peptide, prevents NFκB 340
 341 translocation to the nucleus via inhibition of IκB phosphoryla- 341
 342 tion both in vitro and in vivo (Leceta et al. 2000; Delgado and 342




Fig. 4 Changes in renal expression of glutathione (GSH; **a**), superoxide dismutase (SOD; **b**) and malondialdehyde (MDA; **c**). Values are given as mean micromoles per gram, international units per gram and micromoles per gram \pm SEM, respectively. * $p < 0.05$ vs. diabetes + PACAP; # $p < 0.05$ vs. control

343 Ganea 2001; Delgado et al. 2002; Ganea and Delgado 2003).
 344 Antioxidants and inhibitors of the renin-angiotensin-
 345 aldosterone system, which exert their nephroprotective effect
 346 partially through controlling the action of NF κ B, are extensively
 347 used in the treatment of diabetic nephropathy (Liu et al.
 348 2009). In diabetic nephropathy, NF κ B can be activated by
 349 mesangial cells, hyperglycemia-induced ROS production,
 350 renin-angiotensin-aldosterone system (RAAS) activation,
 351 TGF- β 1, TNF α and interleukins (Iwamoto et al. 2005; Gnudi
 352 2012). NF κ B plays an important role in several renal diseases,
 353 such as nephritis, proteinuria and tubulointerstitial disorders,
 354 and it is the key transcriptional regulator of diabetic kidney
 355 disease (Sakurai et al. 1996; Iwamoto et al. 2005). It activates
 356 the transcription of a variety of factors involved in cell prolifer-
 357 ation and inflammation contributing to the progression of
 358 diabetic nephropathy (Navarro-González et al. 2011).

Mesangial cell proliferation, tubular cell damage and excessive
 secretion of proinflammatory cytokines, chemokines and adhe-
 sion molecules occur as a result of NF κ B activation (Mezzano
 et al. 2004; Chen et al. 2008).

Activation of NF κ B under diabetic conditions is assumed
 to mediate ROS-induced apoptotic changes (Aoki et al. 2011).
 Extracellular signal-regulated kinase (ERK) 1/2, p38 mitogen
 activated protein kinase (MAPK) and Jun kinase-mediated
 pathways are stimulated upon NF κ B activation. In this study,
 we showed that PACAP increased the expression of
 antiapoptotic factors, like pAkt and pERK1/2, while down-
 regulated the proapoptotic pp38MAPK. The markedly in-
 creased expression of the antiapoptotic proteins in untreated
 diabetic animals may result from the induction of the protec-
 tive compensatory mechanisms. The antiapoptotic effect of
 PACAP was associated with decreased cleaved caspase-3.
 Caspase-3, as an effector caspase, plays a critical role in
 receptor-mediated, mitochondria-dependent and endoplasmic
 reticulum stress-induced apoptotic mechanisms, which are
 involved in hyperglycemia-induced podocyte loss (Susztak
 et al. 2006; Tunçdemir and Oztürk 2011; Gui et al. 2012).
 Apoptosis is a rare event in the normal kidney; however, it is
 present in human diabetic kidney biopsies, similar to other
 kidney disorders. Not only proximal and distal tubular cells
 but also endothelial and mesangial cells are affected by apo-
 ptosis in diabetes (Woo 1995; Kumar et al. 2004).

Similar to the pathomechanism of the tubulointerstitial injury
 in myeloma kidney, tubular epithelial cells start to produce
 excessive amount of proinflammatory cytokines in diabetic
 nephropathy as well. TGF- β 1 is a prosclerotic cytokine pro-
 duced by mesangial and proximal tubular cells (Gilbert et al.
 1998). It is the key mediator of hyperglycemia-induced changes
 in the kidney, accelerating the production of extracellular matrix
 in excess through epithelial-to-mesenchymal cell transformation
 (di Paolo et al. 1996; Hills Paul and Squires 2010). The accu-
 mulation of fibronectin, collagen IV and laminin is caused by
 simultaneous overproduction and decreased breakdown of pro-
 teins in mesangial matrix, glomerular and tubular basal mem-
 brane and interstitium, resulting in a severely damaged renal
 morphology and function (Mauer et al. 1984; Steffes et al.
 1989). Similar to our results in rat kidney, PACAP was found
 to significantly lower the TGF- β 1 production in stimulated
 macrophages (Sun et al. 2000). Li et al. reported that PACAP
 provided a protective effect against early diabetic nephropathy.
 They found that this effect is mediated via inhibiting TGF- β 1
 and TNF α pathways, resulting in remarkably attenuated histo-
 logical changes in the PACAP-treated animals. Similar to Li
 et al., in the present study, we showed decreased TGF- β 1 and
 collagen IV levels in the PACAP-treated diabetic compared to
 the untreated diabetic kidneys (Li et al. 2008). Therefore, these
 findings provide explanation for the suppressed extracellular
 matrix expansion observed in the histological sections of
 PACAP-treated animals. Antioxidants and AGE inhibitors,




Fig. 5 Representative electron microscopic images of diabetic (a) and PACAP-treated diabetic kidneys (b). Focal segmental thickening of the glomerular basement membrane (GBM) (*paired arrows*) was observed in the diabetic kidney, while other parts of the GBM remained unchanged (*paired arrowheads*). Fusion of the foot processes of the podocytes (*encircled*) was also remarkable. PACAP treatment effectively counteracted the diabetes-induced GBM thickening and podocyte injury (b). Scale bar= 2 μ m. *Inserts* show *enlarged encircled parts* of pictures a and b to visualize the severe injury of the podocyte foot processes (fused foot processes

marked by *asterisks*, c), while PACAP treatment resulted in intact podocyte foot processes (foot processes marked by *asterisks*, d). Thickness of GBM (e). Basement membrane thickness measurements were performed in the entire basement membrane in control, control + PACAP and in diabetes + PACAP-treated groups. Data are given separately for segmental thickenings in diabetic animals (*bar*: diabetes segmental) and for areas showing no segmental thickening (*bar*: diabetes nonsegmental). Data show mean \pm SEM. *** p <0.001 vs. all other groups

412 which are already commonly used in clinical practice, also
 413 effectively reduce TGF- β 1, fibronectin and collagen IV levels
 414 (Ha et al. 1999; Kelly et al. 2001). The results of anti-TGF- β 1
 415 treatments are promising; therefore, any candidate, which
 416 downregulates the TGF- β 1 pathway, could be effective in the
 417 treatment of diabetic nephropathy (Sharma et al. 1996; Ziyadeh
 418 et al. 2000).

419 The importance of oxidative stress in the development of
 420 diabetes and diabetic complications is well known (Usuki et al.
 421 2011). The diabetogenic effect of streptozotocin in pancreatic β
 422 cells also involves the overproduction of nitric oxide (NO) and
 423 superoxide (O_2^-) (Nukatsuka et al. 1988; Kaneto et al. 1995).
 424 The mitochondrial free radicals were shown to stimulate the
 425 polyol-PKC, hexosamine and AGE pathways, contributing to
 426 the progression of DN. The vital role of glutathione is based on
 427 its ability to neutralize electrophils produced by metabolic
 428 processes or external stimuli in order to prevent the organism
 429 against their harmful effects. Conditions associated with high
 430 levels of free radicals, like diabetes, probably inactivate the
 431 mitochondrial GSH carriers, resulting in depressed antioxidant
 432 mechanisms. Non-enzymatic glycation of renal mitochondrial
 433 proteins was shown to be at least partially responsible for these

434 changes (Lash 2006). Moreover, increased activity of NADPH
 435 oxidase results in severely lowered NADPH levels, leading to
 436 impaired glutathione recovery. Therefore, not only the in-
 437 creased oxidative stress but also the inactivated defence mech-
 438 anisms contribute to the oxidative damage of the renal cells,
 439 leading to more severe consequences of the oxidative agents
 440 (Stanton 2011; Gnudi 2012). Changes in the concentration and
 441 redox status of GSH lead to mitochondrial DNA damage and
 442 induction of apoptotic pathways (Marchetti et al. 1997; Davis
 443 et al. 2001). We found that PACAP is capable of normalising
 444 the decreased GSH levels in the diabetic kidney, suggesting an
 445 antioxidative effect of PACAP. This finding is similar to our
 446 previous studies, where PACAP was found to exert antioxidative
 447 effects in oxidative stress-induced renal and hepatic cell
 448 damage, intestinal cold preservation and warm ischemic injury
 449 (Ferencz et al. 2010a, 2010b; Horvath et al. 2011). Usually
 450 diabetes is also associated with decreased superoxide dismutase
 451 (SOD) and elevated malondialdehyde (MDA) levels; however,
 452 under our experimental circumstances, we could not detect
 453 diabetes-induced changes in these factors.

454 Based on these data, PACAP seems to be a promising
 455 candidate in treating diabetic nephropathy. Similar to other

456 studies showing that PACAP is a strong cytoprotective agent,
 457 we have also provided evidence that PACAP is highly effective
 458 in diabetic nephropathy. This efficacy is most probably due to
 459 PACAP acting at several levels and directly and/or indirectly
 460 (via antiinflammatory, antiapoptotic and antioxidant mecha-
 461 nisms) affecting more cell types in the kidney. However, several
 462 aspects have to be examined before the clinical application of
 463 PACAP. In rats, no side effects were observed after systemic
 464 administration of PACAP. Indeed, even a human study proved
 465 that systemic infusion of the peptide causes no changes in the
 466 physiological parameters apart from transient flushing (Li et al.
 467 2007). The other drawback of systemic PACAP administration
 468 is the poor bioavailability, since the half life of the peptide in the
 469 circulation is only 5–10 min due to its rapid degradation by
 470 dipeptidyl peptidase IV (DPP-IV) (Banks et al. 1993;
 471 Bourgault et al. 2008). However, nowadays, several studies
 472 aim at finding an easy and reliable way of PACAP treatment
 473 (Onoue et al. 2011).

474 In summary, our present study demonstrated the molecular
 475 mechanisms involved in the protective effect of PACAP.
 476 Besides the contribution of the previously proven
 477 antiinflammatory effect, antiapoptotic, antioxidative and
 478 antifibrotic mechanisms are responsible for the protective
 479 effect of PACAP in 8-week diabetic nephropathy in rats.

480 **Acknowledgments** This study was supported by OTKA K104984,
 481 108596, PD109644, TAMOP 4.2.2.A-11/1/KONV-2012-0024, Arimura
 482 Foundation and PTE-MTA “Lendület” Program. This research was realized
 483 in the frames of TAMOP 4.2.4. A/2-11-1-2012-0001 “National Excellence
 484 Program—Elaborating and operating an inland student and researcher
 485 personal support system convergence program”, Bolyai Scholarship. The
 486 project was subsidized by the European Union and co-financed by the
 487 European Social Fund.

488 **References**

490 Aoki M, Nata T, Morishita R et al (2011) Endothelial apoptosis induced
 491 by oxidative stress through activation of NF-kappaB: antiapoptotic
 492 effect of antioxidant agents on endothelial cells. *Hypertension* 38:
 493 48–55
 494 Araki N, Takagi K (1992) Relaxant effect of pituitary adenylate cyclase-
 495 activating polypeptide on guinea-pig tracheal smooth muscle. *Eur J*
 496 *Pharmacol* 216:113–117
 497 Arimura A, Li M, Batuman V (2006) Potential protective action of
 498 pituitary adenylate cyclase-activating polypeptide (PACAP38) on
 499 in vitro and in vivo models of myeloma kidney injury. *Blood* 107:
 500 661–668
 501 Banki E, Degrell P, Kiss P et al (2013) Effect of PACAP treatment on
 502 kidney morphology and cytokine expression in rat diabetic nephrop-
 503 athy. *Peptides* 42:125–130
 504 Banks WA, Kastin AJ, Komaki G, Arimura A (1993) Passage of pituitary
 505 adenylate cyclase activating polypeptide1-27 and pituitary adenylate
 506 cyclase activating polypeptide1-38 across the blood–brain barrier. *J*
 507 *Pharmacol Exp Ther* 267:690–696

Bourgault S, Vaudry D, Botia B et al (2008) Novel stable PACAP analogs 508
 with potent activity towards the PAC1 receptor. *Peptides* 29:919– 509
 932 510
 Chen L, Zhang J, Zhang Y, Wang Y, Wang B (2008) Improvement of 511
 inflammatory responses associated with NF-kappa B pathway in 512
 kidneys from diabetic rats. *Inflamm Res* 57:199–204 513
 Davis W Jr, Ronai Z, Tew KD (2001) Cellular thiols and reactive oxygen 514
 species in drug-induced apoptosis. *J Pharmacol Exp Ther* 296:1–6 515
 Delgado M, Ganea D (2001) Inhibition of endotoxin-induced macro- 516
 phage chemokine production by VIP and PACAP in vitro and 517
 in vivo. *Arch Physiol Biochem* 109:377–382 518
 Delgado M, Jonakait GM, Ganea D (2002) Vasoactive intestinal peptide 519
 and pituitary adenylate cyclase-activating polypeptide inhibit che- 520
 mokine production in activated microglia. *Glia* 39:148–161 521
 Di Mario U, Pugliese G (2001) 15th Golgi lecture: from hyperglycaemia 522
 to the dysregulation of vascular remodelling in diabetes. 523
Diabetologia 44:674–692 524
 Di Paolo S, Gesualdo L, Ranieri E, Grandaliano G, Schena FP (1996) 525
 High glucose concentration induces the overexpression of 526
 transforming growth factor-beta through the activation of a 527
 platelet-derived growth factor loop in human mesangial cells. *Am* 528
J Pathol 149:2095–2106 529
 Ferencz A, Kiss P, Weber G et al (2010a) Comparison of intestinal warm 530
 ischemic injury in PACAP knockout and wild-type mice. *J Mol* 531
Neurosci 42:435–442 532
 Ferencz A, Weber G, Helyes Z, Hashimoto H, Baba A, Reglodi D 533
 (2010b) Presence of endogenous PACAP-38 ameliorated intestinal 534
 cold preservation tissue injury. *J Mol Neurosci* 42:428–434 535
 Filipsson K, Pacini G, Scheurink AJ, Ahren B (1998) PACAP stimulates 536
 insulin secretion but inhibits insulin sensitivity in mice. *Am J* 537
Physiol 274:834–842 538
 Ganea D, Delgado M (2003) The neuropeptides VIP/PACAP and T cells: 539
 inhibitors or activators? *Curr Pharm Des* 9:997–1004 540
 Gilbert RE, Cox A, Wu LL et al (1998) Expression of transforming 541
 growth factor-beta1 and type IV collagen in the renal 542
 tubulointerstitium in experimental diabetes: effects of ACE inhibition. 543
Diabetes 47:414–422 544
 Girard BM, Tompkins JD, Parsons RL, May V, Vizzard MA (2012) 545
 Effects of CYP-induced cystitis on PACAP/VIP and receptor ex- 546
 pression in micturition pathways and bladder function in mice with 547
 overexpression of NGF in urothelium. *J Mol Neurosci* 48:730–743 548
 Gnudi L (2012) Cellular and molecular mechanisms of diabetic glomer- 549
 ulopathy. *Nephrol Dial Transplant* 27:2642–2649 550
 Gray SL, Cummings KJ, Jirik FR, Sherwood NM (2001) Targeted 551
 disruption of the pituitary adenylate cyclase-activating polypeptide 552
 gene results in early postnatal death associated with dysfunction of 553
 lipid and carbohydrate metabolism. *Mol Endocrinol* 15:1739–1747 554
 Gui D, Guo Y, Wang F et al (2012) Astragaloside IV, a novel antioxidant, 555
 prevents glucose-induced podocyte apoptosis in vitro and in vivo. 556
PLoS ONE 7:e39824 557
 Ha H, Yu MR, Kim KH (1999) Melatonin and taurine reduce early 558
 glomerulopathy in diabetic rats. *Free Radic Biol Med* 26:944–950 559
 Hills Paul E, Squires PE (2010) TGF-beta1-induced epithelial-to- 560
 mesenchymal transition and therapeutic intervention in diabetic 561
 nephropathy. *Am J Nephrol* 31:68–74 562
 Horvath G, Brubel R, Kovacs K et al (2011) Effects of PACAP on 563
 oxidative stress-induced cell death in rat kidney and human hepatoc- 564
 yte cells. *J Mol Neurosci* 43:67–75 565
 Iwamoto M, Mizuiri S, Arita M, Hemmi H (2005) Nuclear factor-kB 566
 activation in diabetic rat kidney: evidence for involvement of P- 567
 selectin in diabetic nephropathy. *Tohoku J Exp Med* 206:163–171 568
 Kaneto H, Fujii J, Seo HG et al (1995) Apoptotic cell death triggered by 569
 nitric oxide in pancreatic beta-cells. *Diabetes* 44:733–738 570

- 571 Kelly DJ, Gilbert RE, Cox AJ, Soulis T, Jerums G, Cooper ME (2001) Aminoguanidine ameliorates overexpression of pro-sclerotic growth factors and collagen deposition in experimental diabetic nephropathy. *J Am Soc Nephrol* 12:2098–2107
- 572
573
574
- 575 Khan AM, Li M, Abdunour-Nakhoul S, Maderdrut JL, Simon EE, Batuman V (2012) Delayed administration of pituitary adenylate cyclase-activating polypeptide 38 ameliorates renal ischemia/reperfusion injury in mice by modulating Toll-like receptors. *Peptides* 38:395–403
- 576
577
578
579
- 580 Klein R, Klein BE, Moss SE, Davis MD, DeMets DL (1984) The Wisconsin epidemiologic study of diabetic retinopathy. IV. Diabetic macular edema. *Ophthalmology* 91:1464–1474
- 581
582
- 583 Koppan M, Varnagy A, Reglodi D et al (2012) Correlation between oocyte number and follicular fluid concentration of pituitary adenylate cyclase-activating polypeptide (PACAP) in women after superovulation treatment. *J Mol Neurosci* 48:617–622
- 584
585
586
587
- 588 Kumar D, Robertson S, Burns KD (2004) Evidence of apoptosis in human diabetic kidney. *Mol Cell Biochem* 259:67–70
- 589
- 590 Lash LH (2006) Mitochondrial glutathione transport: physiological, pathological and toxicological implications. *Chem Biol Interact* 163:54–67
- 591
592
- 593 Leceta J, Gomariz RP, Martinez C et al (2000) Receptors and transcriptional factors involved in the anti-inflammatory activity of VIP and PACAP. *Ann N Y Acad Sci* 921:92–102
- 594
595
- 596 Li M, Maderdrut JL, Lertora JJ, Batuman V (2007) Intravenous infusion of pituitary adenylate cyclase-activating polypeptide (PACAP) in a patient with multiple myeloma and myeloma kidney: a case study. *Peptides* 28:1891–1895
- 597
598
599
- 600 Li M, Maderdrut JL, Lertora JJ, Arimura A, Batuman V (2008) Renoprotection by pituitary adenylate cyclase activating polypeptide in multiple myeloma and other kidney diseases. *Regul Pept* 145:24–32
- 601
602
603
- 604 Lindén A, Cardell LO, Yoshihara S, Nadel JA (1999) Bronchodilation by pituitary adenylate cyclase-activating peptide and related peptides. *Eur Respir J* 14:443–451
- 605
606
- 607 Liu XP, Pang YJ, Zhu WW et al (2009) Benazepril, an angiotensin-converting enzyme inhibitor, alleviates renal injury in spontaneously hypertensive rats by inhibiting advanced glycation end-product-mediated pathways. *Clin Exp Pharmacol Physiol* 36:287–296
- 608
609
610
611
- 612 Marchetti P, Decaudin D, Macho A et al (1997) Redox regulation of apoptosis: impact of thiol oxidation status on mitochondrial function. *Eur J Immunol* 27:289–296
- 613
614
- 615 Mauer SM, Steffes MW, Ellis EN, Sutherland DE, Brown DM, Goetz FC (1984) Structural-functional relationships in diabetic nephropathy. *J Clin Invest* 74:1143–1155
- 616
617
- 618 Mezzano S, Aros C, Droguett A et al (2004) NF-kappaB activation and overexpression of regulated genes in human diabetic nephropathy. *Nephrol Dial Transplant* 19:2505–2512
- 619
620
- 621 Misra HP, Firdovich I (1972) The role of superoxide anion in the oxidation of epinephrine and a simple assay for superoxide dismutase. *J Biol Chem* 247:3170–3175
- 622
623
- 624 Miyata A, Arimura A, Dahl RR et al (1989) Isolation of a novel 38 residue-hypothalamic polypeptide which stimulates adenylate cyclase in pituitary cells. *Biochem Biophys Res Commun* 164:567–574
- 625
626
627
- 628 Moody TW, Di Florio A, Jensen RT (2012) PYK-2 is tyrosine phosphorylated after activation of pituitary adenylate cyclase activating polypeptide receptors in lung cancer cells. *J Mol Neurosci* 48:660–666
- 629
630
631
- 632 Nath KA (1998) The tubulointerstitium in progressive renal disease. *Kidney Int* 54:992–994
- 633
- Navarro-González JF, Mora-Fernández C, Muros de Fuentes M, García-Pérez J (2011) Inflammatory molecules and pathways in the pathogenesis of diabetic nephropathy. *Nat Rev Nephrol* 7:327–340
- 634
635
636
637
- Nedvig K, Weber G, Nemeth J et al (2012) Changes of PACAP immunoreactivities and cytokine levels after PACAP-38 containing intestinal preservation and autotransplantation. *J Mol Neurosci* 48:788–794
- 638
639
640
641
- Nukatsuka M, Sakurai H, Yoshimura Y, Nishida M, Kawada J (1988) Enhancement by streptozotocin of O₂⁻ radical generation by the xanthine oxidase system of pancreatic beta-cells. *FEBS Lett* 239:295–298
- 642
643
644
645
- Onoue S, Hanato J, Yamada S (2008) Pituitary adenylate cyclase activating polypeptide attenuates streptozotocin-induced apoptotic death of RIN-m5F cells through regulation of Bcl-2 family protein mRNA expression. *FEBS J* 275:5542–5551
- 646
647
648
649
- Onoue S, Hanato J, Kuriyama K, Mizumoto T, Yamada S (2011) Development of PACAP38 analogue with improved stability: physicochemical and in vitro/in vivo pharmacological characterization. *J Mol Neurosci* 43:85–93
- 650
651
652
653
- Placer ZA, Cushman LL, Johnson BC (1966) Estimation of product of lipid peroxidation (malondialdehyde) in biochemical systems. *Anal Biochem* 16:359–364
- 654
655
656
- Prevost G, Arabo A, Jian L et al (2013) The PACAP-regulated gene Selenoprotein T is abundantly expressed in mouse and human β -cells and its targeted inactivation impairs glucose tolerance. *Endocrinology* 154:3796–3806
- 657
658
659
660
- Reglodi D, Kiss P, Horvath G et al (2012) Effects of pituitary adenylate cyclase activating polypeptide in the urinary system, with special emphasis on its protective effects in the kidney. *Neuropeptides* 46:61–70
- 661
662
663
664
- Sakurai H, Hisada Y, Ueno M, Sugiura M, Kawashima K, Sugita T (1996) Activation of transcription factor NF-kB in experimental glomerulonephritis in rats. *Biochim Biophys Acta* 1316:132–138
- 665
666
667
- Sakurai Y, Shintani N, Hayata A, Hashimoto H, Baba A (2011) Trophic effects of PACAP on pancreatic islets: a mini-review. *J Mol Neurosci* 43:3–7
- 668
669
670
- Sedlak J, Lindsay R (1968) Estimation of total protein-bound and non-protein sulphhydryl groups in tissue with Ellman's reagent. *Anal Biochem* 25:192–205
- 671
672
673
- Sekiguchi Y, Kasai K, Hasegawa K, Suzuki Y, Shimoda S (1994) Glycogenolytic activity of pituitary adenylate cyclase activating polypeptide (PACAP) in vivo and in vitro. *Life Sci* 55:1219–1228
- 674
675
676
677
- Sharma K, Jin Y, Guo J, Ziyadeh FN (1996) Neutralization of TGF-beta by anti-TGF-beta antibody attenuates kidney hypertrophy and the enhanced extracellular matrix gene expression in STZ-induced diabetic mice. *Diabetes* 45:522–530
- 678
679
680
681
- Stanton RC (2011) Oxidative stress and diabetic kidney disease. *Curr Diab Rep* 11:330–336
- 682Q4
683
- Steffes MW, Osterby R, Chavers B, Mauer SM (1989) Mesangial expansion as a central mechanism for loss of kidney function in diabetic patients. *Diabetes* 38:1077–1081
- 684
685
686
- Sun W, Tadmori I, Yang L, Delgado M, Ganea D (2000) Vasoactive intestinal peptide (VIP) inhibits TGF-beta1 production in murine macrophages. *J Neuroimmunol* 107:88–99
- 687
688
689
- Sun YM, Su Y, Li J, Wang LF (2013) Recent advances in understanding the biochemical and molecular mechanism of diabetic nephropathy. *Biochem Biophys Res Commun* 433:359–361
- 690
691
692
- Susztak K, Raff AC, Schiffer M, Böttinger EP (2006) Glucose-induced reactive oxygen species cause apoptosis of podocytes and podocyte depletion at the onset of diabetic nephropathy. *Diabetes* 55:225–233
- 693
694
695
696

- 697 Syed AU, Koide M, Braas KM, May V, Wellman GC (2012) Pituitary
698 adenylate cyclase-activating polypeptide (PACAP) potently dilates
699 middle meningeal arteries: implications for migraine. *J Mol*
700 *Neurosci* 48:574–583
- 701 Szakaly P, Kiss P, Lubics A et al (2008) Effects of PACAP on survival and
702 renal morphology in rats subjected to renal ischemia/reperfusion. *J*
703 *Mol Neurosci* 36:89–96
- 704 Tunçdemir M, Oztürk M (2011) The effects of angiotensin-II receptor
705 blockers on podocyte damage and glomerular apoptosis in a rat
706 model of experimental streptozotocin-induced diabetic nephropathy.
707 *Acta Histochem* 113:826–832
- 708 Usuki S, Tsai YY, Morikawa K et al (2011) IGF-1 induction by acylated
709 steryl β -glucosides found in a pre-germinated brown rice diet reduces
710 oxidative stress in streptozotocin-induced diabetes. *PLoS ONE* 6:
711 e28693
- 712 Wada Y, Nakamachi T, Endo K et al (2013) PACAP attenuates NMDA-
713 induced retinal damage in association with modulation of the
714 microglia/macrophage status into an acquired deactivation subtype.
715 *J Mol Neurosci* 51:493–502
- 716 Woo D (1995) Apoptosis and loss of renal tissue in polycystic kidney
717 diseases. *N Engl J Med* 333:18–25
- 718 Yada T, Sakurada M, Filipsson K, Kikuchi M, Ahren B (2000)
719 Intraperitoneal PACAP administration decreases blood glucose in
720 GK rats, and in normal and high fat diet mice. *Ann N Y Acad Sci*
721 921:259–263
- 722 Ziyadeh FN, Hoffman BB, Han DC et al (2000) Long term prevention of
723 renal insufficiency, excess matrix gene expression and glomerular
724 mesangial matrix expression by treatment with monoclonal anti-
725 TGF-beta antibody in db/db diabetic mice. *Proc Natl Acad Sci U*
726 *S A* 97:8015–8020

UNCORRECTED PROOF

AUTHOR QUERIES

AUTHOR PLEASE ANSWER ALL QUERIES.

- Q1. Please check corresponding author if captured and presented correctly.
- Q2. Figures 1,3 & 5 contains poor quality text. Please provide replacement. Otherwise, please advise if okay to proceed with the figure/s as is.
- Q3. “Green et al. 2006” is cited in text but not given in the reference list. Please provide details in the list or delete the citation from the text.
- Q4. Please advise if the captured abbreviated journal title is correct.

UNCORRECTED PROOF