

Endocrine factors in the hypothalamic regulation of food intake in females: a review of the physiological roles and interactions of ghrelin, leptin, thyroid hormones, oestrogen and insulin

V. Somogyi, A. Gyorffy, T. J. Scalise, D. S. Kiss, G. Goszleth, T. Bartha, V. L. Frenyo and A. Zsarnovszky*

Department of Physiology and Biochemistry, Szent Istvan University Faculty of Veterinary Sciences, 1078 Istvan u. 2, Budapest, Hungary

Abstract

Controlling energy homeostasis involves modulating the desire to eat and regulating energy expenditure. The controlling machinery includes a complex interplay of hormones secreted at various peripheral endocrine endpoints, such as the gastrointestinal tract, the adipose tissue, thyroid gland and thyroid hormone-exporting organs, the ovary and the pancreas, and, last but not least, the brain itself. The peripheral hormones that are the focus of the present review (ghrelin, leptin, thyroid hormones, oestrogen and insulin) play integrated regulatory roles in and provide feedback information on the nutritional and energetic status of the body. As peripheral signals, these hormones modulate central pathways in the brain, including the hypothalamus, to influence food intake, energy expenditure and to maintain energy homeostasis. Since the growth of the literature on the role of various hormones in the regulation of energy homeostasis shows a remarkable and dynamic expansion, it is now becoming increasingly difficult to understand the individual and interactive roles of hormonal mechanisms in their true complexity. Therefore, our goal is to review, in the context of general physiology, the roles of the five best-known peripheral trophic hormones (ghrelin, leptin, thyroid hormones, oestrogen and insulin, respectively) and discuss their interactions in the hypothalamic regulation of food intake.

Key words: Hypothalamic regulation: Food intake: Women: Hormones

Introduction

The goal of the present review is to present an overview of the most studied hormonal factors that contribute to the central regulation of food intake and energy homeostasis which simultaneously provide peripheral feedback signals to the relevant hypothalamic feedback mechanisms. The most important hormones playing such dual roles are ghrelin, leptin, thyroid hormones, insulin and oestrogen. Of course, in addition to the listed hormones there are several other humoral factors, such as catecholamines, etc, contributing directly or indirectly to the regulation of energy metabolism; however, attempts to cover all contributions are beyond the scope of the present overview. The listed factors highly influence the appetitive trigger that stimulates food-seeking behaviour, ingestion and energy expenditure, thereby establishing a dynamic and plastic information channel between central, mostly hypothalamic, circuits and peripheral tissues. In the present search

for the blueprint of the complex regulatory machinery, the hypothesis is that the functions of the hormones discussed in the present review cannot be understood and may be misleading if examined in isolated environments (i.e. singled out from the complex pool of the regulatory circuit). They can only function appropriately when acting together and adequate hormone interactions take place. Thus, we will compare and review the literature on the individual roles of the selected hormones with those observed under the influence of one or more of these hormones present in the actual context examined or discussed. We believe that comparison of individual and combined hormone effects will highlight the shortcomings of classical experimental protocols operating on the basis of only one variable and then interpreting the resulting data as lone-standing constituents of a more complex mechanism. In fact, we believe that this comprehensive review underscores that physiologically relevant data can only be obtained from complex biological systems,

Abbreviations: AgRP, agouti-related protein; D1, type I deiodinase; D2, type II deiodinase; D3, type III deiodinase; E2, 17 β -oestradiol; ER α , oestrogen receptor α ; ER β , oestrogen receptor β ; ERE, oestrogen response element; MSH, melanocyte-stimulating hormone; NPY, neuropeptide Y; POMC, pro-opiomelanocortin; STAT3, signal transducer and activator of transcription 3; T₃, triiodothyronine; T₄, thyroxine; TR α , thyroid hormone receptor α ; TR β , thyroid hormone receptor β ; TSH, thyroid-stimulating hormone; UCP, uncoupling protein.

* **Corresponding author:** Attila Zsarnovszky, fax +36 1 478 4165, email zsarnovszky.attila@aotk.szie.hu

and clinical application of experimental results should be designed on the basis of the complex interplay of the biological factors involved.

To best illustrate their individual and combined roles, each of the factors will be discussed separately, then in combination based on the available literature. Considering that we are reviewing an enormous field of research, we will keep the main stream of discussion within the boundaries of general physiology. However, we will use comparative animal physiology or specify sex differences where the examples provide clarity and reveal their complexity for the discussion of feedback loops or mechanisms in the regulation of metabolic homeostasis.

General physiological roles of ghrelin, leptin, thyroid hormones, oestrogen and insulin

Ghrelin

Ghrelin is an acylated twenty-eight-amino acid orexigenic peptide hormone first isolated from rat stomach by Kojima *et al.*⁽¹⁾. Ghrelin serum concentrations are below detectable levels during fetal life but rise sharply during the second to third postnatal weeks⁽²⁾. Significant ghrelin production has been observed in the human and rat placenta⁽³⁾. The X/A-like secretory cells in the oxyntic glands of the stomach are responsible for most of the circulating levels of ghrelin⁽¹⁾, although the hypothalamus, hypophysis, kidney and the intestines also contribute to ghrelin production. A high degree of ghrelin expression was detected in several types of tumours, particularly the adrenocorticotroph hormone-secreting thymus tumours, gastrinomas, insulinomas, and in human pituitary carcinomas^(4,5). Plasma levels of ghrelin are reduced by about 80% after gastrectomy or fundectomy⁽⁶⁾, indicating that most ghrelin is produced by the oxyntic glands of the stomach.

Besides its role in the regulation of neural circuits that underly appetitive behaviour, feeding⁽⁷⁻⁹⁾, digestion⁽¹⁰⁻¹²⁾ and thermogenesis⁽¹³⁾, ghrelin modulates a number of additional physiological processes that, overall, may affect energy homeostasis, such as growth hormone production^(7,10,14), cell proliferation^(9,15,16), apoptosis⁽⁹⁾, sleep-wake behaviour⁽¹⁷⁾ and cardiovascular functions^(12,18,19). In addition, ghrelin receptors are found on neurons in the arcuate nucleus of the hypothalamus where appetite-stimulating neuropeptide Y (NPY) and agouti-related protein (AgRP) are co-expressed^(13,20), within the hypothalamus, pituitary, small and large intestines, placenta, kidney, testes, ovary and lymphocytes and on insulin-secreting β -cells in a dose-dependent manner^(11,12,21).

Intracerebroventricular or peripheral (intravenous) application of ghrelin stimulates food intake (appetite)^(22,23). Studies by several groups have demonstrated that ghrelin supports lipogenesis and suppresses fat pad

degradation^(24,25), increasing body weight and adiposity, and leading to obesity. However, skeletal mass or bone growth remains unaffected^(24,25). Thus, ghrelin accelerates the maturation and growth of adipocytes *in vitro*, but may not augment adipocyte differentiation *in vivo*^(26,27).

Although ghrelin is an energy-deficit signal, ghrelin-null or growth hormone secretagogue receptor (GHS-R)-null mice do not display deficiencies in skeletal growth, body composition and food intake^(28,29). On high-fat diets, these knock-out animals usually do not become obese and/or insulin resistant^(30,31).

The major stimulus in ghrelin secretion is fasting⁽³²⁾. All parameters, including the pulse amplitude of ghrelin release, are augmented by fasting⁽³³⁾. Circulating levels of ghrelin increase after fasting and revert after feeding. However, while a relatively short, 24 h fasting increases serum ghrelin levels, a longer period of food restriction does not affect postprandial ghrelin concentrations^(34,35), suggesting that ghrelin levels reflect short-term energy deficits and corroborate a positive energy balance. The preprandial rise of ghrelin's plasma level in obese animals is slower and delayed compared with those animals of normal weight^(36,37). While programmed feeding allows for the observation of marked changes in plasma ghrelin levels, under *ad libitum* feeding, ghrelin (as well as leptin) secretion shows a pulsatile rhythm or pattern, with a peak every 30 min on average. Ghrelin release is not influenced by simple expansion (mechanical distention) of the stomach, but depends on the ingredients of the meal⁽³⁷⁻³⁹⁾, thus indicating that the energy content of the meal may be a determining factor⁽⁴⁰⁾.

In rodents and humans, ghrelin plasma concentrations follow changes in body weight and adiposity, but show no alterations in circulating levels in ageing rats and mice. Likewise, no alterations were found in ghrelin receptor expression in the pituitary gland and brain of ageing mice^(41,42). Ghrelin binds to HDL, decreasing their ability to bind and carry lipid molecules⁽⁴³⁾, inhibiting lipid export from responsive tissue. A circadian rhythm in the fluctuation of plasma ghrelin levels exists in rodents, with the highest concentrations of circulating ghrelin occurring before the dark period⁽⁴⁴⁾ and after waking/just before breakfast in humans⁽⁴⁵⁾. Gastric ghrelin secretion increases in the dark period. In contrast, daytime hypophyseal ghrelin release seems to be higher compared with night-time values. The biodegradation of ghrelin has an important role in the adjustment of plasma ghrelin levels. Tissue esterases and proteases are the key enzymes in the inactivation and biodegradation of ghrelin in the plasma.

The hypothalamic arcuate nucleus is a principal target of ghrelin. Ghrelin receptors are found on neurons in the arcuate nucleus of the hypothalamus where NPY and AgRP are co-expressed^(13,20). Distinct populations of neurons were identified in the arcuate nucleus as major players in the regulation of food intake, such as NPY- and AgRP-containing neurons⁽⁴⁶⁾ responsible for orexigenic effects

(hunger cells), and the melanocyte-stimulating hormone/pro-opiomelanocortin (α -MSH/POMC) satiety cells and cocaine- and amphetamine-regulated transcript (CART)-producing neurons responsible for anorexigenic mechanisms. Hewson *et al.*⁽⁴⁷⁾ suggest that ghrelin exerts its stimulatory effect on appetite and food intake/feeding activity by stimulating NPY release. Intracerebroventricular administration of anti-NPY IgG suppresses ghrelin's (appetite-stimulating) action, although anti-AgRP IgG had the same effect. Nakazato *et al.*⁽²⁴⁾ also describe ghrelin's enhancing effect on AgRP and NPY mRNA levels but found no modification of the influence of NPY/AgRP neurons on energy expenditure.

In concert with these findings, immunohistochemical analyses show a direct connection between NPY/AgRP neurons and ghrelin-containing fibres⁽⁴⁸⁾. According to Andrews *et al.*⁽⁴⁹⁾, ghrelin seems to act on NPY/AgRP neurons by inducing mitochondrial respiration of these cells, most probably by influencing mitochondrial proliferation and activity of NPY/AgRP neurons via affecting uncoupling protein (UCP) 2.

Some diseases and disorders are marked by altered ghrelin secretion. Prader–Willi syndrome (a genetic disorder characterised by mental retardation, hyperphagia, short stature and muscular hypotonia) presents with higher plasma ghrelin concentrations^(50–52), a possible factor for the hyperphagic symptom of the disease. Anorexia nervosa is also characterised by a high ghrelin plasma concentration^(53,54) which returns to a normal level after recovery and weight gain⁽⁵⁵⁾. Patients suffering from short-bowel syndrome have lower plasma ghrelin levels, supposedly due to the missing ghrelin-producing tissues⁽⁵⁶⁾. The same is observed after gastric bypass surgery⁽⁵⁷⁾, since ghrelin is predominantly produced in the stomach.

Since more than 90% of ghrelin is produced by the stomach and duodenum, parts of the gastrointestinal tract affected by gastric bypass surgery, a significant decrease in ghrelin level was observed in most patients who underwent this operation^(58–60). Indeed, one of the mechanisms by which gastric bypass reduces body weight is suppression of ghrelin production. However, the effects of bypass surgery are much more complex and are not fully understood. There are several other sources reporting that there were no alterations in ghrelin levels after gastric bypass or gastric banding⁽⁶¹⁾. This discrepancy in post-surgical ghrelin levels may be due to the difference in surgical techniques involving, for example, treatment of the vagus nerve⁽⁶²⁾.

Ghrelin appears to be evolutionarily conserved, as it has been found in birds, reptiles, amphibians and fish, although its effects may have served different functions in specific species. In mammals, it is the only gut-derived hormone that stimulates food intake, while in chickens, it has the contradistinct effect of inhibition of feeding^(63,64). Ghrelin's functions in other bird species are still unknown, albeit energy balance and thermogenesis may be regulated

by ghrelin there as well. Nevertheless, the mechanism is different from that of mammals and is yet to be determined.

Leptin

Leptin is anorexigenic, inhibits excessive deposition of lipids in various tissues and enables the organism to adapt to periods of food deprivation or undernutrition⁽⁶⁵⁾. The name 'leptin' is derived from the Greek word *leptos*, meaning 'thin'. Leptin is a member of the cytokine family of peptide hormones first identified as a 16kDa protein secreted by the adipose tissue to regulate feeding behaviour and energy balance^(66,67). The gene encoding for leptin is identical to the *ob* gene, discovered in 1995 by Geffroy *et al.*⁽⁶⁸⁾ as the factor responsible for obesity when mutated. Hence, the gene was called '*ob*', abbreviated from the word 'obesity'. Geffroy *et al.* localised the leptin's *ob* gene to the 7th chromosome. Initial evidence correlated levels of serum leptin with adipose tissue mass, suggesting that it was produced by the adipose tissue itself⁽⁶⁹⁾. Higher levels of *ob* mRNA are expressed in the fat of obese population samples compared with thin controls, and leptin release correlates with the size of adipocytes, i.e. larger cells produce proportionally higher quantities of leptin due to accumulation of TAG^(70,71). Leptin is produced predominantly, but not exclusively, by white adipose tissue⁽⁷²⁾. It is also released by brown adipose tissue, placenta (syncytiotrophoblasts), ovary, skeletal muscle, stomach (epithelial cells in the lower part of fundic glands), mammary epithelial cells, heart, bone marrow, pituitary and the liver (especially in birds)⁽⁷³⁾. In these organs and tissues, leptin acts as a paracrine or autocrine hormone^(65,74,75).

In an early experiment, mouse *ob* protein (leptin) was purified and administered in small doses both centrally and peripherally to genetically obese (*ob/ob*) mice, diet-induced obese mice and diabetic (*db/db*) obese mice. Leptin reduced food intake and body weight in both genetically obese and diet-induced obese mice, but had no effect on *db/db* obese animals. The behavioural effects after central (intracerebral) administration suggested that leptin acted directly on the neuronal networks that control feeding and energy balance⁽⁷⁶⁾. To summarise, it was proposed that the *ob* gene encoded the *ob* protein (leptin) that regulated food intake and energy balance, and that the extreme obesity of the genetically obese (*ob/ob*) mice was attributable to mutations in the *ob* gene. It was also suggested that the *db* gene encoded the receptors for this factor and the extreme obesity of diabetic (*db/db*) mice was attributed to mutations in the *db* gene⁽⁷⁷⁾. In another human study on two severely obese children, serum leptin levels were very low despite the markedly elevated fat mass. In both children, a mutation in the leptin gene was found, providing further evidence to support the role of the *ob* gene in the regulation of leptin levels⁽⁷⁸⁾.

Leptin receptor (Ob-R) is coded by the *db* gene. Cloning of the mouse *db* gene revealed that it encodes five alternative isoforms (a–e). All isoforms possess identical ligand-binding domains (extracellular), but differ in their C-terminus (intracellular), with Ob-Rb as the only isoform able to activate the correct signal transduction pathway⁽⁷⁹⁾. A further experiment showed that the mutant genotype *db/db* lacks Ob-Rb, and the phenotype of the affected mice was identical to that of genetically obese (*ob/ob*) mice, characterised by obesity and diabetes (from which the name of the gene, *db*, is derived)⁽⁸⁰⁾. Expression of the other receptor isoforms (Ob-Ra, Ob-Rc, Ob-Rd and Ob-Re) remained unaffected, leading to the conclusion that, of the known receptor isoforms, only the Ob-Rb receptor is essential for leptin's weight-regulating effects⁽⁸¹⁾.

Using *in situ* hybridisation, Ob-Rb receptors were detected in high levels in hypothalamic neurons and hypothalamic cells, suggesting that these anatomical regions are crucial sites of leptin's actions⁽⁸²⁾. This is supported by evidence that a single intracerebroventricular injection of leptin reduces food intake at doses with no measurable effect when delivered peripherally⁽⁸³⁾. In addition to the central nervous system, relatively high Ob-Rb mRNA expression levels were detected in the adipose tissue, skeletal muscle and liver, suggesting that leptin is likely to play a role in peripheral lipid metabolism as well^(84,85).

The transport mechanisms of leptin through the blood–brain barrier is still undefined, but it has been hypothesised that the receptor Ob-Ra could be involved in an active transport mechanism in certain Ob-Rb-containing neurons that actually project to neurohaemal brain regions (i.e. circumventricular organs) and thus avoid the blood–brain barrier⁽⁸⁶⁾.

In the hypothalamus, the lateral hypothalamic and ventromedial nuclei project both within and outside the hypothalamus and modulate the activity of the parasympathetic and sympathetic nervous systems, two overlapping and redundant mechanisms of the central nervous system that depend upon fight-or-flight or homeostatic demands of the organism^(87–89). The dorsomedial hypothalamic nucleus, via its direct anatomical connections with the parasympathetic nervous system, may be involved in integrating information between the lateral hypothalamic, ventromedial and paraventricular nuclei. The paraventricular nucleus controls secretion of peptides from both the posterior and anterior pituitary and projects to nuclei with sympathetic or parasympathetic efferents^(86,90,91).

Leptin's central effects on the regulation of food intake and energy metabolism are not confined to a single mechanism of action. Besides its direct effects on relevant neural circuits, indirect mechanisms of leptin actions are also known. In general, leptin down-regulates orexigenic peptides, and up-regulates anorexigenic peptides, leading to a reduction in food intake. However, there is a dynamic and interactive effect between orexigenic and anorexigenic factors. For example, α -MSH agonists decrease food intake,

while α -MSH antagonists act in the opposite manner to reduce the anorexigenic effect of leptin⁽⁹²⁾. On the other hand, peripheral and central effects of leptin can merge to shape the final biological outcome: by the facilitation of lipolysis in the adipose tissue, leptin increases the plasma level of NEFA, the latter which influence the function of relevant hypothalamic neurons, while leptin also acts directly on hypothalamic orexigenic neurons. Additionally, leptin up-regulates uncoupling in UCP-expressing hypothalamic neurons and probably other cells as well, thereby propelling energy metabolism/utilisation/use of energy. Therefore, along with its stimulatory effect on sympathetic tone, leptin's overall effects also include increased heat production⁽⁹³⁾. In addition to these central actions, leptin exerts considerable peripheral effects that can be considered upstream signals of extended hypothalamic leptin effects. In short, leptin decreases the levels of insulin and glucocorticoids, enhances the secretion of growth hormone, catecholamines and the thyroid hormones, thus stimulating energy expenditure and hepatic lipogenesis⁽⁶⁵⁾. Leptin augments mitochondrial fatty acid oxidation especially in hepatocytes and skeletal muscle cells^(94,95) and increases insulin sensitivity and glucose metabolism in myocytes^(65,96). NPY/AgRP neurons are major targets of leptin in the hypothalamus^(97,98). In these cells, the two antagonistic hormones, ghrelin and leptin, act in opposite ways to modulate neuronal activity and influence food intake, as discussed below.

Thyroid hormones

Traditionally, the hormone products of the thyroid gland have been known as 'thyroid hormones'. Today, the source and function of these hormones comprise a large body of knowledge, allowing for a remarkably complex view on the physiological roles of thyroid hormones. This brief summary is confined to those salient thyroid hormone effects that, in one way or another, are involved in the hypothalamic regulation of food intake.

There are various types of thyroid hormones: thyroxine (T_4), containing four iodine atoms, is generally considered to be a prohormone, while triiodothyronine (T_3), containing three iodine atoms, is biologically more active than T_4 . These two hormones are inactivated by conversion to reverse triiodothyronine (rT_3) or diiodothyronine (T_2).

Thyroid hormones are crucial for the maintenance of energy equilibrium of the body: thyroid hormones regulate the rate of metabolic processes (including the BMR), increase O_2 consumption and heat production. Consequently, thyroid hormones are equally necessary for normal reproduction, growth and development⁽⁹⁹⁾.

In humans, most (80%) of the circulating T_3 originates from the periphery (i.e. not from the thyroid gland), where T_4 is converted into T_3 by the corresponding deiodinase enzymes. The most important peripheral

T₃-producing organs are the liver, kidney and the skeletal muscle⁽¹⁰⁰⁾. In rats, more T₃ is produced in the thyroid gland (55%) than in other peripheral tissues⁽¹⁰¹⁾, while in birds almost all (99%) T₃ is synthesised in peripheral tissues, outside of the thyroid gland⁽¹⁰²⁾.

The function of the thyroid gland is regulated by a negative feedback mechanism. The hypothalamic nuclei produce thyrotropin-releasing hormone that stimulates the adenohypophysis to produce and secrete thyroid-stimulating hormone (TSH). TSH increases T₄ production in the thyroid gland. An increase of plasma T₄ concentration, in turn, down-regulates the production of thyrotropin-releasing hormone, TSH, and T₄ as well⁽¹⁰³⁾.

The peripheral T₄-to-T₃ conversion is catalysed by deiodinase enzymes⁽¹⁰⁴⁾. There are three types of deiodinases in different tissues: type I deiodinase (D1), type II deiodinase (D2) and type III deiodinase (D3). D1 is expressed in the central nervous system, especially the pituitary, in rats but not in humans^(105–107). Of the lower vertebrates, the fish brain was found to contain D1⁽¹⁰⁸⁾.

D2 expression was described in the brain (including the pineal gland) and pituitary of rats⁽¹⁰⁶⁾ mainly as a glial enzyme most abundant in astrocytes and tanocytes (near the mediobasal hypothalamus)^(109,110). In humans, D2 is the only thyroid hormone-activating 5' deiodinase in the brain⁽¹⁰⁵⁾.

D3 is ubiquitously expressed in the adult rat brain, especially in the forebrain, cerebral cortex, pyramidal cells of the hippocampus, granule cells of the dentate gyrus, and in the piriform cortex⁽¹¹¹⁾. These brain regions are the richest in thyroid hormone receptors⁽¹¹²⁾. In neonatal rat brain, D3 expression is age dependent⁽¹¹³⁾. In contrast to D2, D3 can be found mostly in neural cells⁽¹¹¹⁾.

D1 and D2 activate (when pH < 7, D1 may also inactivate), while D3 inactivates, thyroid hormone. In mammals, the bulk production of T₃ is mediated by D1. D1 is expressed in the liver, kidneys, thyroid gland, skeletal muscles, lung, and in the hypophysis^(114–118). In mammals, hepatic and kidney-derived D1 is the most important source of T₃. These organs produce T₃ and release it into the bloodstream. Therefore, these are called hormone-exporting organs^(116,117,119). D2 is also expressed in the liver in birds but not in mammals⁽¹⁰²⁾.

Peripheral tissues can adjust the ratio of active and inactive thyroid hormones, according to their actual energy needs and energy supply through their deiodinase activity. Therefore, the peripheral tissues can become more or less independent from the central regulatory processes. In the case of appropriate energy supply, the body primarily needs T₃, thereby turning on thyroid hormone-activating deiodination^(117,119,120).

Thyroid hormones act through binding to their cognate receptors (thyroid hormone receptors). Thyroid hormone receptors are nuclear receptors that can stimulate ligand-dependent transcription of certain genes^(121,122). In mammals, there are four thyroid hormone receptor isoforms,

thyroid hormone receptor α (TR α) 1–2 and thyroid hormone receptor β (TR β) 1–2, encoded by two highly conserved genes (*Thra* and *Thrb*, respectively)^(123,124). TR α 2 has not been found in non-mammalian species⁽¹²⁵⁾. Unlike the other thyroid hormone receptors, TR α 2 acts as a weak repressor instead of functioning as a receptor of T₃⁽¹²⁶⁾. Thyroid hormone receptors have both T₃-dependent and -independent functions. Interestingly, T₃ can convert a T₃-independent repressor into a T₃-dependent activator⁽¹²⁷⁾.

The distribution pattern of *Thra* and *Thrb* is age and tissue dependent. While TR α 1 is widely expressed, TR β 1–2 mRNA expression is restricted to specific ontogenetic states and is highly tissue specific⁽¹²⁸⁾. Based on knock-out and knock-in experiments, *Thra* was found to influence cardiac functions, thermogenesis, haemopoiesis, and the maturation of intestines and bones^(128–131). *Thrb* is crucial for the proper endocrine and sensory functions such as the hypothalamic–pituitary–thyroid axis, hepatic reactions to T₃, behaviour, audition, colour sensation and tactile senses^(132–138). TR α 1 and TR β 1 isoforms can be co-expressed in the same tissues and can substitute for each other's function to a certain extent^(138,139).

Nutrients entering the small intestines elicit enteroglucagon production in the intestinal wall that reaches the liver before the absorption of the nutrients⁽¹⁴⁰⁾. Absorbed nutrients quickly enter the liver through the portal vein. Carbohydrates, and to a lesser extent proteins, induce hepatic D1 gene expression^(141–143) and down-regulate D3 activity. T₃, in turn, increases hepatic and muscular GLUT (GLUT-1 and GLUT-5) gene expression^(101,144). The latter also enhances hepatic thyroid hormone activation⁽¹⁴⁵⁾. Increased energy intake, therefore, induces increased plasma T₃ concentration that increases BMR⁽¹¹⁷⁾. Increased, but still physiological plasma T₃ concentrations, in turn, trigger appetite in the hypothalamic ventromedial nucleus. This is a self-strengthening mechanism^(146,147).

In piglets, thyroid hormone plasma concentration (mostly T₃) increases sharply after food intake, reaching its peak after 60 min. The rise in thyroid hormone concentration mainly depends on the energy content of the food and is relatively independent from the glucose concentration of the blood⁽¹⁰¹⁾. In contrast to feeding, starvation elicits a decrease in plasma T₃ concentration in mammals. T₄ response to starvation, however, depends on the species. For example, starvation leaves plasma T₄ concentration unaffected in humans⁽¹⁴⁸⁾, but in rats^(149,150) and swine^(151–153) plasma T₃ concentrations drop after starvation. In chickens, starvation causes decreased plasma T₃ and increased plasma T₄ concentration^(115,154–157) due to increased hepatic D3⁽¹¹¹⁾ and decreased hepatic D2 activity and gene expression⁽¹⁵⁸⁾. In cows, effects of starvation are difficult to examine due to the characteristics of rumen function. Therefore, the effect of fluctuations in feeding on plasma thyroid hormone concentration was

tested and it was found that reduced food intake resulted in decreased plasma thyroid hormone concentration⁽¹⁵⁹⁾.

Dairy cows represent a unique category with regard to thyroid hormone physiology. These animals can enter a physiologically starvation-like period during lactation when energy metabolism turns towards so-called negative energy balance. Under such conditions, plasma T₄ concentration reaches its nadir and T₃ plasma concentration is low. Later, plasma thyroid hormone concentrations increase and reach their peak during the dry period⁽¹⁵⁹⁾. Romo *et al.*⁽¹⁶⁰⁾ and Pezzi *et al.*⁽¹¹⁷⁾ found that hepatic D1 activity is the lowest during early lactation and suggested a possible role of fatty liver in the decreased enzyme activity and the consequently decreased hepatic thyroid hormone activation. However, decreased feed (and consequently carbohydrate) intake can also be acting in the background of decreased thyroid hormone activation since T₄-to-T₃ conversion is a carbohydrate-dependent process⁽¹⁴¹⁾ and glucose directly triggers hepatic D1 expression⁽¹⁴³⁾.

As mentioned above, hypothalamic NPY/AgRP neurons are in a special position to trigger food intake, and both leptin and ghrelin are able to influence the activity of these cells. Coppola *et al.*⁽¹⁶¹⁾ clearly demonstrated that arcuate nucleus NPY/AgRP neurons are supplied with T₃ by hypothalamic tanycytes, and that T₃, imported from tanycytes, can activate intracellular mechanisms in NPY/AgRP cells to increase neuronal firing and, consequentially, to initiate food intake. These observations imply a possible interaction between the three hormones discussed so far, in the regulation of NPY/AgRP neuronal activity. However, according to Ishii *et al.*⁽¹⁶²⁾, subcutaneously injected T₃ results in hyperphagia via hypothalamic adenosine monophosphate activated protein kinase (AMPK) phosphorylation, without affecting local (hypothalamic) NPY or POMC mRNA expression levels, suggesting that additional mechanisms may also be involved in the hypothalamic regulation of food intake.

Oestrogen

Traditionally, oestrogens (mostly 17β-oestradiol; E2) have been regarded exclusively as sex hormones, and, indeed, many of their direct actions are exerted on reproductive organs and tissues. Oestrogens are primarily synthesised within the developing follicles of the ovaries and exert an influence on the female reproductive system. However, other sites of biosynthesis are present throughout the body, including the adipose tissue and the brain itself. Adipose tissue contributes significantly to the maintenance of plasma concentration of E2. Besides the adipose tissue, aromatase (oestrogen synthase) expression is also found in muscle, skin fibroblasts, osteoblasts and chondrocytes of bones, vascular endothelial and aortic smooth muscle cells, Leydig cells, as well as a number of sites in the brain including the medial preoptic/anterior hypothalamus,

the mediobasal hypothalamus and the amygdala^(163,164). Oestrogens exert an effect on almost every cell type, regulating an eclectic assortment of functions.

In males, regulation of E2 mainly occurs through the conversion of testosterone using aromatase enzyme throughout the body, including the adipose tissue that represents a major source of ovary-independent production⁽¹⁶⁴⁾. A positive relationship was described between BMI and plasma E2 levels in men and postmenopausal women, and circulating levels of E2 decreased in response to fasting or energy restriction in fertile women⁽¹⁶⁵⁾. Rats lacking oestrogen receptor α (ERα) or aromatase have increased fat mass and hyperlipidaemia⁽¹⁶⁶⁾. This suggests that E2 has a role in signalling the central nervous system about the status of peripheral energy stores.

It has long been known that E2 modulates hunger^(166–168). The question, however, of how much is the share of E2 alone, besides other relevant hormones, in the regulation of food intake is not fully known. Apart from the actual initiation of food intake, effects on energy expenditure highly influence evolving hunger signals. Therefore, E2 can modulate feedback hunger signals through its central and peripheral regulatory role in the initiation of heat production⁽¹⁶⁹⁾. The homeostatic functions of E2, such as reproduction, stress responses, feeding, sleep cycles and temperature regulation were originally thought to be mediated through E2's classical intracellular/nuclear receptors (oestrogen receptors) ERα and oestrogen receptor β (Erβ), which act as transcription factors, binding to the oestrogen-responsive elements (ERE)^(170,171) of respective genes. Keeping with the general context of the present review, a recent study reported that elimination of ERα (ERαKO) from the hypothalamic ventromedial nucleus in rats leads to the development of the metabolic syndrome, with a significant deficit in energy expenditure⁽¹⁷²⁾. In addition to nuclear oestrogen receptors, there are other extranuclear pathways activated after ligands bind to the oestrogen receptor in the plasma membrane^(173,174), or activate intracellular proteins directly responsible for the regulation of cellular energy levels, such as UCP⁽¹⁷⁵⁾ or ectonucleotidases⁽¹⁷⁶⁾. Oestrogens can also initiate cellular actions through membrane-bound E2-sensitive molecular complexes, such as G-protein-coupled receptor 30, ion channels and receptors for other ligands⁽¹⁷⁷⁾. An interesting mechanism of oestrogenic effects is the synaptogenic influence of E2 on hypothalamic neuronal connectivity. It is now well established that E2 induces synaptic reorganisation in the hypothalamus to regulate the secretion and release of gonadotropin-releasing hormone and, consequentially, pituitary luteinising hormone release⁽¹⁷⁸⁾. Synaptic plasticity plays a crucial role in the hypothalamic mediation of leptin and ghrelin effects as well⁽¹⁷⁹⁾, and overall synaptic reorganisation in the hypothalamus seems to be shaped by all three hormones (E2, leptin and ghrelin). However, E2 appears to exert its own synaptogenic influence in the hypothalamus,

since apart from the ovarian cycle, hunger is also modulated by the actual levels of E2^(167,168). It is now well established that oestrogens (E2 above all) are trophic hormones. As such, a plethora of studies reported that E2 interacts with other trophic hormones, including the ones discussed in the present review, to regulate developmental events and energy metabolism (food intake and energy expenditure)^(180–183). A relevant selection of these interactions will be discussed below. At this point, we only note that E2, while targeting hypothalamic neurons, is able to influence the function of NPY/AgRP-containing neurons of the arcuate nucleus, the major mediators–initiators of food intake, and thus play a significant role in shaping the overall outcome of NPY/AgRP neuronal activity^(179,182). The latter role includes a blunting effect after fasting-induced activation of these neurons⁽¹⁸³⁾. Interestingly, however, E2 not only attenuates hunger by down-regulating relevant hypothalamic ‘hunger cell’ activity (NPY/AgRP neurons), but also modulates neurotransmission on POMC neurons (satiety cells) to attenuate the cannabinoid-induced changes in appetite⁽¹⁸⁴⁾.

Insulin

Insulin is secreted by pancreatic islet cells (β -cells) and is known as a major regulator of blood glucose levels. To date, a large body of evidence demonstrates actions of insulin in the hypothalamic feedback loop of energy balance regulation⁽¹⁸⁵⁾. The actual concentration of plasma insulin is proportional to body adiposity, although it also depends on food consumption and type of diet. Secretion and synthesis of insulin are affected by a number of genetic and environmental/epigenetic factors. Food intake, diet, fasting and re-feeding influence insulin entry to the brain or its ability to get into the brain tissue⁽¹⁸⁶⁾. Eating increases the concentration of insulin in the brain whereas fasting decreases the concentration. Plasma insulin levels directly correlate with those measured in the cerebrospinal fluid. Insulin enters the brain, crossing the blood–brain barrier by a receptor-mediated transport, and stimulates anorexigenic pathways as an adiposity signal leading to reduced food intake. Insulin receptors are expressed in hypothalamic neurons, predominantly in the arcuate nucleus⁽¹⁸⁷⁾, a nucleus associated with the regulation of food intake, energy expenditure and food reward and the site of several orexigenic and anorexigenic neural signals^(180,182,188,189). In rats, hyperphagia caused by insulin-deficient type 1 diabetes can be reversed by intracerebroventricular administration of insulin. Thus, insulin interacts with both orexigenic and anorexigenic neuromodulators^(189,190).

Reports from the past three decades show that insulin exerts multiple effects in the brain, and its role has been validated by the identification of receptors and transport machinery in the central nervous system. Insulin acts on hypothalamic circuits that regulate food intake, metabolism and other crucial functions, and insulin-responsive neurons

include the hypothalamic NPY/AgRP neurons⁽¹⁹¹⁾, the latter which may represent a special cell population allowing for a cross-talk between the major hormonal signals discussed in the present review. The effects of insulin leading to decreased food intake are, at least in part, due to the inhibition of NPY expression of neurons of the arcuate nucleus⁽¹⁹²⁾. An interesting hypothalamic effect of insulin was observed by Lin *et al.*⁽¹⁹³⁾, who found that hypothalamic NPY/AgRP and POMC neurons mediate insulin’s central effects to regulate hepatic glucose production.

Part of the endocrine cross-talk appears to manifest in the hormonal regulation of hypothalamic synaptic reorganisation, where the interaction of insulin-like growth factors and their receptors are a well-known episode in the regulation of synaptic plasticity⁽¹⁷⁸⁾.

Despite the abundance of relevant data, a clearcut borderline between the effects of insulin alone and those exerted through the interactions of insulin with other peripheral trophic hormones in the hypothalamic regulation of food intake is not yet known. Therefore, further effects of insulin on food intake will be discussed below in the context of hormone interactions.

Hormone interactions and combined hormone effects

Interactions of ghrelin and leptin

In mammals, ghrelin stimulates food intake and appears to act as the functional antagonist of leptin⁽¹⁹⁴⁾. There is a fine balance between the two physiologically antagonistic hormones as demonstrated by Giovambattista *et al.*⁽¹⁹⁵⁾, who detected enhanced leptin secretion after prolonged ghrelin exposure. In turn, moderate hyperleptinaemia/leptin administration prevents the increase of plasma ghrelin during short-term energy restriction⁽¹⁹⁶⁾. Ghrelin’s hunger signal stimulates the hypothalamic regions responsible for food intake in the lateral hypothalamic nuclei and arcuate nucleus. In contrast to leptin, plasma levels of ghrelin fluctuate in relation to food intake. The low ghrelin levels in obese individuals are associated with low growth hormone concentration and high leptin levels^(32,197). Rigamonti *et al.*⁽¹⁹⁸⁾ found higher concentrations of plasma ghrelin in young patients compared with elderly patients. Thus, it is possible that a high concentration of leptin suppresses ghrelin’s plasma level. This idea is supported by observations that plasma leptin concentration correlates with the body fat mass which is high in obesity, decreased in fasting and low in anorexia nervosa.

An interesting addition for ghrelin and leptin regulation is the linkage to sleep–wake behaviour⁽¹⁷⁾. It has been demonstrated that inadequate or bad-quality sleep is associated with changed levels of ghrelin and leptin, and is in connection with energy balance disorders leading to weight gain. Normally, ghrelin levels rise at night. In sleep deprivation or insomnia, however, night-time ghrelin

levels fall significantly and are followed by increased daytime ghrelin levels. Administration of ghrelin increases the ratio of non-rapid eye movement (REM) sleep in human subjects. The REM phase is thought to be the stage with the highest metabolic rate. Lack of it leads to hormonal changes that increase appetite. It is noteworthy that insufficient sleep can decrease daytime levels of leptin; however, its nocturnal levels do not show significant change between healthy individuals and insomniacs.

The anatomy and synaptology of the hypothalamic leptin/ghrelin-responsive neurons provide an explanation for the aforementioned interactions. In a review, Horvath⁽¹⁷⁹⁾ pointed to the ability of hypothalamic neuronal circuits to rearrange themselves in response to peripheral and central hormone effects. Earlier, this phenomenon had not been considered as part of the mechanism responsible for the regulation of daily energy metabolism. However, as stressed in that review, experiments on *ob/ob* mice and transgenic animals provided clearcut evidence for the ability of leptin and ghrelin to induce synaptic reorganisation on hypothalamic neurons that are in a key position (NPY/AgRP and POMC cells) to regulate food intake and other determinants of energy homeostasis. Hypocretin/orexin-containing neurons of the lateral hypothalamus are major targets of gastric ghrelin and, at the same time, these neurons directly innervate raphe serotonergic neurons to regulate sleep–wake behaviour. Elevation of serum ghrelin levels increases the number of excitatory synapses and the ratio of stimulatory/inhibitory synapses on lateral hypothalamic hypocretin/orexin neurons. Since such a ratio in cellular synaptic status is a key factor in the shaping of neuronal activity, the consequence of such a synaptic reorganisation is that the neurons in question, as major sources of the stimulatory innervation of arcuate nucleus NPY/AgRP cells, increase the activity of the latter cell population, thereby initiating/increasing food intake. Such synaptic plasticity occurs simultaneously on POMC neurons as well, where an inhibitory dominance occurs at the time of increased NPY/AgRP cell activity. The described synaptological basis of the mechanism includes the anti-ghrelin-like (antagonistic) effects of leptin, but is also modulated by the synaptogenic effects of E2 and other hormones discussed in the present review.

Interactions of ghrelin and insulin

As mentioned above, ghrelin release is not influenced by simple expansion (mechanical distention) of the stomach; instead, it is highly dependent on the ingredients of the meal. Elevated blood glucose level, after oral or intravenous application, leads to decreased plasma ghrelin^(197,199,200). In turn, ghrelin seems to enhance the degradation/catabolism of carbohydrates and results in an increase in the RQ. According to Ariyasu *et al.*⁽³⁶⁾, insulin-induced hypoglycaemia facilitates ghrelin secretion in rodents and this can be reversed by glucose injections.

Ghrelin levels are increased under lipid-rich diets, but not under low-protein diets^(201,202).

As emphasised in the present review, the arcuate nucleus is a key brain region regulating energy balance. Peripheral hormonal signals converge to influence arcuate nucleus circuits and shape overall energy metabolism. Among these signals are ghrelin and insulin, whose common targets are the arcuate nucleus NPY/AgRP neurons, the 'hunger cells' of the hypothalamus. As described above, ghrelin stimulates the activity, while insulin down-regulates NPY secretion of these cells; thus these two hormones antagonise in the regulation of 'hunger cell' function. Apart from their hypothalamic effects, parallel brain circuits are also modulated by the two hormones. For example, there is evidence that brainstem regions, such as the ventral tegmental area, are sensitive to leptin, insulin and ghrelin and that the activity of dopaminergic cells of the ventral tegmental area are modulated by the above hormones^(203,204). In turn, the ventral tegmental area dopaminergic cells innervate a number of forebrain structures, forming circuits that are commonly associated with the expectation of feeding-associated reward^(205–207).

It is noteworthy that besides their central effects, there are certain peripheral interactions of ghrelin and insulin that also influence hypothalamic neural mechanisms. For example, a wide range of findings supports roles of ghrelin in the regulation of insulin release. Ghrelin receptors are present on insulin-secreting β -cells⁽²¹⁾ and ghrelin immunoreactivity is reported in pancreatic islets partly overlapped with glucagon-immunoreactive cells. Data suggest that ghrelin is synthesised and secreted in pancreatic α -cells and feeds back on the same α -cells in an autocrine and/or paracrine manner. Ghrelin concentration in pancreatic veins is higher than in pancreatic arteries, suggesting that pancreatic islets produce ghrelin as well^(208,209). In support of the latter, in their review, Dezaki *et al.*⁽²¹⁰⁾ described that ghrelin is present in pancreatic islet cells and regulates β -cell insulin secretion.

It was recently established that (exogenous) ghrelin administration increases plasma glucose and reduces insulin levels⁽²¹¹⁾. These results are supported by the finding that ghrelin knock-out mice show elevated insulin production in pancreatic islets, which leads to decreased blood glucose levels^(212,213). The main regulator of insulin release seems to be the glucose-induced Ca influx into the β -cells. Dezaki⁽²¹³⁾ described the Ca-mediated insulin release that could be attenuated by ghrelin administration. The mechanism underlying this observation involves ghrelin's activating effect on β -cell growth hormone secretagogue receptors (GHS-R) to enhance K^+ currents through delayed rectifier K-channels via G-protein activation; this, in turn, attenuates the glucose-induced Ca influx and thereby decreases insulin secretion⁽²¹⁰⁾.

Yet another finding to consider is that ghrelin attenuates the expression of UCP2 in *ob/ob* mice, which, under normal circumstances, down-regulates glucose-induced

ATP production, reducing insulin release from β -cells. In summary, ghrelin inhibits insulin secretion in islet β -cells in two ways: (1) by acutely enhancing Kv channels and suppressing Ca signalling; and (2) by chronically up-regulating UCP2. Consequently, ghrelin inhibits insulin release, thereby increasing glycaemia, and affecting feeding behaviour and adiposity^(210,211).

Interactions of leptin and thyroid hormones

Both leptin and the thyroid hormones are key regulators of body weight by affecting food intake and energy expenditure⁽²¹⁴⁾. Leptin decreases food intake and increases energy expenditure, while the thyroid hormones T_4 and T_3 increase both food intake and energy expenditure. This indicates that thyroid hormones and leptin could play a crucial role in the regulation of the balance between energy intake and energy expenditure, and, consequently, in the regulation of body weight and body fat mass. The correlation between serum leptin and T_3 concentrations has been studied extensively. Results so far are conflicting, particularly those from human studies⁽²¹⁵⁾.

Study on cultured rat adipocytes demonstrated that leptin mRNA production and thus leptin secretion was inhibited by Triac (triiodothyroacetic acid, a metabolite of T_3) and T_3 in a dose-dependent manner and with similar potency⁽²¹⁶⁾. This was confirmed in an *in vivo* rat study, in which intraperitoneal administration of different doses of T_3 (and consequently increased serum T_3) led to a decrease in white adipose leptin mRNA levels and a decrease in serum leptin concentration. This observation indicates that T_3 regulates serum leptin through the modulation of leptin gene (*ob*) expression⁽²⁰⁵⁾.

Conversely, administration of leptin has been shown to affect serum thyroid hormone levels. When administered to fasted rats, leptin prevented the reduction in thyroid hormone levels by maintaining normal pro-thyrotropin-releasing hormone (precursor to thyrotropin-releasing hormone) mRNA activity in the hypothalamus⁽²¹⁷⁾. Likewise, leptin enhanced T_4 -to- T_3 conversion⁽²¹⁷⁾. Leptin also prevents the suppression of TSH in the anterior pituitary after food restriction, thereby maintaining serum T_3 and T_4 levels. However, this anti-inhibitory effect of TSH also depends on thyroid gland activity, as it was not observed in hypothyroid rats⁽²¹⁸⁾.

In the brain, T_4 -to- T_3 conversion is catalysed by D2. According to Coppola *et al.*⁽¹⁶¹⁾, in the arcuate nucleus, D2 activity takes place in tanycytes, and resulting T_3 is then transferred to NPY/AgRP neurons. T_3 in the latter cell population triggers a series of events including the up-regulation of UCP2 and NPY, eventually leading to mitochondrial proliferation and increased neuronal activity. The physiological outcome of these cellular events manifests in the initiation of or in an increase in food intake. As previously mentioned, leptin up-regulates uncoupling in UCP2-expressing hypothalamic neurons.

This might be, at least in part, due to leptin's ability to increase T_3 levels by acting centrally on D2 activity⁽²¹⁹⁾. Interestingly, however, as an anorexigenic hormone, leptin down-regulates NPY/AgRP neuronal functions. Therefore, in terms of food intake, it seems likely that T_3 's and leptin's opposing physiological effects on NPY neurons are mediated by different signalling modalities with possible cross-talk at the level of UCP2. UCP2, and other uncoupling enzymes elsewhere in the body, may be common targets of T_3 and leptin in their regulatory actions in heat production. The latter regulatory process probably involves parallel neural circuits. For example, the expression level of UCP1 in brown adipose tissue is a marker of sympathetic nervous system activity. It has been shown that D2 activity, in brown adipose tissue in particular, can be stimulated by the sympathetic nervous system⁽²¹⁹⁾ and, not surprisingly, central leptin administration increases UCP1 expression in brown adipose tissue.

In sum, T_3 regulates serum leptin levels by inhibiting leptin gene expression, but leptin maintains normal thyroid hormone levels in food-restricted animals by preventing the suppression of thyrotropin-releasing hormone and TSH⁽²²⁰⁾. Leptin also increases D1 activity peripherally, and D2 activity centrally in food-restricted animals, resulting in an increase in T_3 as long as sufficient T_4 is available.

Interactions of oestrogen and leptin

The interactions of E2 and leptin are particularly interesting and deserve more attention, since the two hormones have similar effects on food intake and energy homeostasis. For example, leptin limits food intake, i.e. decreased leptin levels result in hyperphagia. Oestrogen also mimics this effect: ovariectomy leads to increased food intake that can be reversed by E2 replacement.

Noteworthy among their interactions are the two hormone's effects on the regulation of each other's plasma levels. Such effects are exerted through indirect and direct pathways. Leptin influences E2 levels indirectly by modulating the regulation of the reproductive axis. It is well known that undernutrition inhibits female reproductive functions and lowers ovarian E2 in favour of self-maintenance and thermoregulation^(221–223). Low levels of ovarian E2 during fasting can be increased by eliminating leptin deficiency. Similarly, low E2 levels in *ob/ob* mice can also be increased by leptin⁽¹⁶⁵⁾. Furthermore, leptin administration increases basal luteinising hormone levels, ovarian uterine weights and restores fertility in *ob/ob* mice⁽²²¹⁾. This suggests that leptin could be the adiposity signal to the reproductive axis signalling the body's current nutritional status, allowing reproduction to continue if significant adipose tissue is available and inhibiting it if body fat stores are depleted⁽²²⁴⁾. The inhibiting effect of leptin deficiency on reproduction is mediated through the down-regulation of gonadotropin-releasing hormone^(165,225).

Although gonadotropin-releasing hormone secretion appears as an important platform of leptin influence on oestrogenic effects, leptin may also influence cellular E2 availability directly. Leptin stimulates aromatase activity in adipose tissue and the presence of leptin receptors (Ob-R) in the ovary implies that leptin may have direct influence on ovarian E2 production and/or oestrogenic effects⁽¹⁶⁵⁾.

There is evidence suggesting that leptin production is directly affected by E2. Ovariectomy or deletion of ER α increases serum leptin levels and leads to an increase of body fat mass⁽¹⁶⁶⁾. Plasma leptin levels are higher in women than in men with an equivalent fat mass and leptin secretion is higher in isolated female omental tissue than male donors, suggesting that sex steroids exert effects on leptin production. Women have higher subcutaneous tissue fat and less intra-abdominal fat than men. Therefore, this sex-related difference in fat distribution may account for these sex differences in leptin levels. Conversely, administration of E2 to cultured rat adipocytes induces leptin secretion and E2 receptors have been found in adipose tissue, suggesting that E2 exerts its effects over leptin secretion directly on adipocytes⁽¹⁶⁵⁾.

In ovariectomised rats, the minimum effective leptin dose resulting in weight loss is greater than in controls, implying that E2 deficiency may attenuate leptin's effects⁽¹⁶⁵⁾. A number of further studies confirm this attenuating effect of E2 over leptin and provide more insight into the potential mechanism. Meli *et al.*⁽¹⁶⁶⁾ found that changes in body weight and fat content caused by ovariectomy are accompanied by alterations in serum leptin concentration. Serum leptin levels are higher in ovariectomised than sham controls, and are reversed by E2 treatment. Increased serum leptin levels after E2 withdrawal are secondary to the increase in fat mass. Mid- or long-term hypo-oestrogenism causes a lower expression of Ob-R in adipose tissue and the hypothalamus compared with sham-operated rats. Lack of E2 leads to hypothalamic NPY overproduction and impairs central leptin sensitivity. The decrease in Ob-R and the associated leptinaemia support central leptin insensitivity. Oestrogen status, therefore, modulates serum leptin levels, as well as hypothalamic and white adipose Ob-Rb expression, thereby implying close cross-talk between central and peripheral leptin sensors in the regulation of body fat mass.

The above phenomenon was observed outside of the hypothalamus as well: Chen *et al.*⁽²²⁶⁾ found a relationship between E2 levels and Ob-R expression in the dorsal root ganglion. Oestrogen was found to increase the expression of Ob-R protein and mRNA in ovariectomised rats. As a result of co-labelling for Ob-R and ER α /ER β , 100% co-localisation of ER α with Ob-R occurred, while ER β -Ob-R co-localisation occurred in only 15% of dorsal root ganglion neurons. The authors concluded that E2 may up-regulate the expression of Ob-R protein and mRNA in an ER α -dependent manner.

A study by Ainslie *et al.*⁽²²⁷⁾ confirmed that E2-deficient rats may be leptin insensitive. The authors showed that the effect of central leptin administration on food intake and physical activity in ovariectomised rats is significantly less than in sham controls. However, they did not find differences in Ob-Rb mRNA expression between sham and ovariectomised rats, suggesting that central leptin insensitivity may occur due to a defect at, or downstream to, the hypothalamic Ob-Rb itself.

Once leptin binds to Ob-Rb, the associated Janus kinase 2 tyrosine kinase is activated, followed by the activation of signal transducer and activator of transcription 3 protein (STAT3), STAT5, phosphatidylinositol 3'-kinase and MAPK pathways. STAT3 is a crucial member of these pathways, as ablation of Ob-Rb-STAT3 signalling/neuronal STAT3/STAT3 in Ob-Rb neurons mimics the obesity, hyperphagia and decreased energy expenditure of leptin-deficient *db/db* mice. Following phosphorylation of STAT3 by Janus kinase 2 and the subsequent dimerisation and nuclear transport, STAT3 alters cell transcription and function. It was proposed that E2 may interact with the leptin pathway at the level of STAT3. STAT3 has been reported as a target of E2 signalling in cells. In a neural STAT3 knock-out murine model, mice exhibited obesity and infertility which E2 administration failed to correct⁽¹⁸²⁾.

Oestrogen receptors are functionally linked to STAT3: they have a direct physical coupling and involve co-regulators such as cAMP-response element binding protein. STAT3 also enhances the transactivational function of oestrogen receptors, which suggests that oestrogen receptors and STAT3 act mutually as co-activators to facilitate signalling⁽¹⁸²⁾.

Conversely, Chen & Heiman⁽²²⁸⁾ sought to determine whether ovariectomy leads to leptin resistance. Ovariectomy caused hyperphagia and increased fat mass. Leptin treatment increased fat utilisation and prevented reduction in energy expenditure. Withdrawal of leptin stimulated hyperphagia. The authors concluded that loss of ovarian function in rats is not associated with a change in leptin sensitivity.

It is well established that the hypothalamus plays an important role in the regulation of food intake and body weight since chemical or physical lesions of these regions cause hyperphagia and obesity^(165,182). It seems clear at this point that the combined effects of leptin and E2 are exerted by their actions on the same population of hypothalamic neurons⁽²²⁹⁾. Evidence indicating that E2 and leptin may cooperate in the regulation of energy metabolism and reproduction is provided by the co-localisation of E2 and leptin receptors in hypothalamic neurons. Diano *et al.*⁽²³⁰⁾ reported co-localisation of leptin receptors with ER α in the medial preoptic area, arcuate nucleus, ventromedial nucleus and paraventricular nuclei in female rats. The authors suggested that this extensive co-localisation of receptors in neuronal perikarya of the hypothalamus indicates a functional link between these hormones and the

regulation of a variety of behavioural and neuroendocrine mechanisms. In a previous study, Meli *et al.*⁽¹⁶⁶⁾ also found considerable co-localisation of oestrogen receptors and Ob-R in hypothalamic nuclei.

The possible mechanisms through which E2 and leptin may cooperate in the brain include interrelated modifications of hypothalamic orexigenic and anorexigenic neuropeptides. In the hypothalamus, anorexigenic arcuate nucleus POMC neurons inhibit food intake through the release of the cleavage product melanocyte-stimulating hormone, while the orexigenic NPY neurons act against this to promote feeding⁽¹⁶⁵⁾. When leptin signalling is diminished, or during fasting, POMC expression is reduced, while increasing levels of leptin stimulate POMC transcription through STAT3 activation^(165,182). Mystkowski & Schwartz⁽¹⁶⁵⁾ found that ovariectomy in rats leads to increased POMC expression in the arcuate nucleus, implying an adaptive response of POMC neurons to attenuate increases in body weight. Furthermore, decreased body weight following the administration of supraphysiological amounts of E2 in male rats is followed by the induction of arcuate nucleus POMC gene expression. Oestrogen-induced weight loss elicits a compensatory reduction of POMC gene expression. Therefore, increased melanocortin signalling does not mediate E2 effects on energy homeostasis.

Gao & Horvath⁽¹⁸²⁾ claim, however, that E2 exhibits molecular and cellular effects on POMC similar to leptin in the arcuate nucleus. They state that both leptin and E2 alter the synaptic input on hypothalamic melanocortin neurons, inducing an increased POMC tone and decreased food intake and adiposity, allowing for the assumption that E2 and leptin regulate POMC expression by the induction of synaptic remodelling.

According to Hirose *et al.*⁽²³¹⁾, the increased serum leptin levels did not up-regulate POMC expression in the hypothalamus of ER α -null mice, while ablation of E2 by ovariectomy and the accompanying higher serum leptin levels up-regulated POMC expression, indicating that ER α is essential for the up-regulation of POMC and that this pathway is probably mediated in an E2-independent manner. ER α can be transcriptionally activated by ligand-independent activation. The authors suggest that leptin may activate ER α through the MAPK pathway via Janus kinase 2. Others, however, suggest that E2 influences the distribution of body fat and adipose tissue metabolism via both ER β and ER α : according to Shin *et al.*⁽²³²⁾, obesity and blood leptin levels are influenced by the ER β :ER α ratio.

It may be possible that E2 exerts its effects over POMC via a putative membrane-associated oestrogen receptor that is distinct from ER α and ER β . Qiu *et al.*⁽¹⁷¹⁾ identified this receptor in dopamine-containing and POMC hypothalamic neurons; its activation leads to desensitisation of γ -aminobutyric acid (GABA_B) receptors in POMC neurons of the arcuate nucleus by the phospholipase C, protein

kinase C, protein kinase A pathway, facilitating the formation of synaptic contacts with hypothalamic areas controlling appetite by POMC neurons.

Interactions between insulin and leptin, oestrogen or thyroid hormones

Leptin, like insulin, plays an important and key role in food reward. The mechanisms are unclear but are being intensely investigated⁽²³³⁾. Insulin is a major modulator of leptin levels^(234–239), in a dose-dependent manner^(240–242). In diabetic rats, insulin administration reverses the previously reduced leptin level⁽²⁴³⁾. Hypothalamic neurons (predominantly in the arcuate nucleus) that express leptin receptors⁽⁸²⁾ and insulin receptors^(244–246) widely overlap. In these neurons, the signalling pathways of leptin and insulin converge on PI3 kinase whose product, PIP3, regulates excitability of ATP-dependent K (K_{ATP}) channels, thereby controlling food intake, energy expenditure and glucose metabolism⁽²⁴⁷⁾. The activation of K_{ATP} channels results in hyperpolarisation and thereby inactivation of glucose-responsive/insulin-sensitive neurons. It has been shown that the inhibition of PI3 kinase blocks the anorexigenic effects of both leptin and insulin in rats⁽¹⁸⁸⁾. Insulin and leptin both target arcuate nucleus NPY neurons, where PI3 kinase-mediated signals seem to be shared and are common regulators in the two hormones' activating effects⁽²⁴⁸⁾. Insulin, in interaction with serotonin and leptin, affects the activity of POMC neurons, inducing α -MSH release, which leads to a negative regulation of energy balance.

It has been reported that both ER α knock-out and aromatase knock-out mice are obese^(249,250), but ovariectomised ER α knock-out mice display decreases in body weight, fat-pad weight, adipocyte size and improved insulin and glucose metabolism indicative of ER β -mediated activity on adipose tissue which are opposite to those of ER α ⁽²⁵¹⁾. ER α plays a role in the development of hepatic insulin sensitivity^(252,253) since the absence of ER α in the ventromedial nucleus leads to weight gain and insulin resistance, adiposity, hyperphagia and hyperglycaemia⁽¹⁷²⁾. It is well known that the insulin-activated GLUT-4 transporter is the main carrier for glucose into cells. In ER α knock-out mice, the activity of these transporters is substantially reduced; ER β knock-out mice do not show this deficiency⁽²⁵⁴⁾. Insulin–E2 interactions are also obvious in certain physiological or pathological conditions, such as pregnancy, menopause or diabetes. In pregnant rats, sexual steroid levels are increased, resulting in elevated insulin levels. Likewise, E2 administration in menopausal women corrects insulin secretion and decreases insulin resistance^(255–257). Cellular investigations demonstrate that ER α is also present in pancreatic β -cells, which is responsible for the short-term insulinotropic effect of E2 via inhibition of K_{ATP} channels.

A hypothesis has arisen that partially explains the synergistic interrelationship of insulin and T_3 underlying mechanisms involved in glucose and lipid metabolism. According to this, T_3 plays a more significant role in glucose metabolism and regulation than previously thought. T_3 regulates glucose uptake, storage, glycolysis, as well as lipid oxidation and uptake in insulin-sensitive cells. Disturbances in T_3 levels cause insulin resistance and hyperglycaemia, which is indicative of its contribution to deficiencies/pathogenesis leading to type 2 diabetes. T_3 increases glucose uptake directly. Expression of (regulatory) glucose transport proteins, such as GLUT-1, GLUT-2 and GLUT-4 are regulated by both insulin and T_3 . A family of membrane-associated proteins regulated by T_3 , named β_3 -adrenergic receptors, plays a role in NEFA oxidation and insulin secretion. T_3 and TAG in fat cells regulate insulin secretion and insulin sensitivity by an as yet unknown mechanism^(258–266). Additionally, T_3 regulates the expression of transmembrane enzymes K^+/Na^+ and Ca^{2+} ATP-ase, involved in driving membrane-bound protein pumps, which transport molecules such as glucose and amino acids into the cells⁽²⁶⁷⁾. Thyroid hormones reduce β -cell mass, the main source of insulin production, although several studies have demonstrated that thyroid hormones have cytotoxic effects on β -cells *in vivo* and *in vitro*. In fact, thyroid hormones induce apoptotic cell death in insulin-producing cells^(268–272).

Interactions between oestrogen and thyroid hormones

Although clinical reports point to a relationship between thyroid disorders and insulin sensitivity particularly prevalent in older populations of females more than males, the mechanisms which link sex steroids, thyroid hormones and metabolic homeostasis in both sexes are not well known⁽²⁷³⁾. In postmenopausal women receiving E2 treatment, lower levels of free T_3 and free T_4 have been observed although there was no rise in serum TSH⁽²⁷⁴⁾. Although E2 treatment did not alter the serum concentrations of TSH in euthyroid or untreated hypothyroid rats, it increased the T_3 effect in serum TSH in hypothyroid rats. Oestrogen treatment increased thyroid hormone receptor expression in the pituitary, suggesting that an increase in thyroid receptors may be induced by E2 in rats⁽²⁷⁵⁾. During development, treatment with testosterone and E2 had similar effects in both sexes, producing lower levels of α and β subunit mRNA in exogenous sex steroid-treated hypothyroid rats compared with hypothyroids not receiving exogenous sex-steroid treatment. These data suggest that gonadal steroids render the pituitary thyrotroph cells more sensitive to thyroid hormones⁽²⁷⁶⁾. This mechanism between oestrogen receptor and thyroid hormone receptor interaction will be explored further, keeping in mind the relationship as it pertains to feeding, appetite, metabolism and energy balance modulated by sex hormones, particularly E2 in women.

Thyroid hormones affect reproductive behaviours of both males and females in addition to playing a permissive role in the photoperiodicity of both sexes^(277–279). Both thyroid hormones and sex steroids modulate metabolic homeostasis and oxidative metabolism at the tissue level via their respective non-genomic membrane and genomic nuclear receptors, sometimes acting synergistically between genomic and non-genomic mechanisms^(280–286) as described above. These mechanisms, in turn, feed back into the regulation of the hypothalamo–pituitary–adrenal axis, hypothalamo–pituitary–gonad axis, as well as the ‘hypothalamus–gut’ circuits^(287–289). Additionally, thyroid hormones contribute to the maintenance of non-reproductive conditions, homeostasis in both male and female birds and mammals, for example, intracellular sites of activity including mitochondria, endoplasmic reticulum, plasma membrane, synapse^(124,290), lipogenesis, lipolysis, thermogenesis, growth, development, differentiation^(124,291,292), myelination of axons⁽²⁹³⁾ via nuclear recruitment of co-activators and co-repressors by ligand-bound nuclear receptors⁽²⁹⁴⁾. Mood and cognition in women are also affected by E2 and thyroid hormones – signals that underlie appetitive behaviour at converging pathways in the hypothalamic nuclei, brain stem and target organs of humoral and efferent signals^(283–285,287,295–297). Such mechanisms conduct cross-talk among isoforms of nuclear oestrogen receptors and thyroid hormone receptors in diverse tissues and in terms of genomic expression^(280,290,298–300). While it is tempting to envision an overlapping function between oestrogen receptor and thyroid hormone receptor subtypes and isoforms, different co-localisation studies indicate distinct non-overlapping functions⁽²⁹⁰⁾. In support of this idea, the phenotypes of the specific knock-out models suggest functions unique to specific isoforms⁽²⁹⁴⁾. On the other hand, the modulation of the signals arising from E2 and thyroid hormones converging must influence underlying behaviour, such as appetite, feeding, homeostasis and metabolism, concentrated in hypothalamic nuclei.

Two genes, *Thra* and *Thrb*, encode the mammalian nuclear receptors for T_3 , confirming that T_3 acts directly on transcription. TR α 1 (and TR α 2)^(291,300) is encoded by the *Thra* gene, and TR β 1 and TR β 2 are encoded by the *Thrb* gene; another TR β 3 isoform that may be rat specific⁽³⁰¹⁾. These receptors have a similar, if not identical, structure, containing a central DNA-binding domain and a C-terminus domain, which activates transcription upon ligand binding⁽³⁰²⁾. Both oestrogen receptors and thyroid hormone receptors are ligand-dependent nuclear transcription factors that can bind to an identical half-site, AGGTCA, of their cognate hormone response elements, although generalisations of receptor cross-activity are not easily made. For example, co-transfection of a mutated TR α 1 also inhibited the E2 effect without binding to the AGGTCA sequence⁽³⁰³⁾. Studies of inhibitory effects of liganded TR α 1 on E2 induction suggest

that oestrogen receptors and thyroid hormone receptors may interact to modulate E2-sensitive gene expression, in a ligand-dependent and isoform-specific manner such as for the hypothalamic preproenkephalin promoter⁽³⁰³⁾. However, in contrast to TR α , the isoforms TR β 1 and TR β 2 failed to inhibit E2-induced activation of the consensus ERE but, rather, attenuated E2-mediated transcriptional activation⁽²⁹⁷⁾. Thus, oestrogen receptor and thyroid hormone receptor isoforms have modular protein structures with high homology in the central DNA-binding domain where they bind common enhancer elements (ERE) within the promoter sequence modulating gene transcription^(280,297,304–309).

Thyroid hormone receptors and oestrogen receptors are members of the nuclear receptor superfamily, composed of forty-eight ligand-activated Zn-finger transcription factors involved in important physiological functions such as metabolism and energy homeostasis, carbohydrate and lipid metabolism in addition to development, reproduction, etc^(304,305,310,311). Briefly, members of the superfamily of nuclear receptors reside in either the cytoplasm or nucleus of the cell. When ligands or small molecules bind to the cytoplasmic nuclear receptor complex, it undergoes conformational change, dimerisation and translocation to the nucleus where it binds to a sequence of DNA called the hormone response element. There the nuclear receptor complex then co-recruits or co-represses other proteins downstream at the promoter site where transcription factors are activated and changes in cell function ensue.

Specifically, half of the T₃ consensus response element DNA sequence is identical to the oestrogen receptor response element (ERE); therefore thyroid hormone receptors bind to a consensus ERE^(297,300,306,307,312) on several receptor genes in the hypothalamus. Thyroid hormone receptors and oestrogen receptors are both present in rat hypothalamic nuclear extracts and bind to the promoter regions in the hypothalamic gene preproenkephalin, causing interactions between liganded thyroid hormone receptors and oestrogen receptors that affect preproenkephalin transcription^(296,312). Likewise, the oxytocin gene promoter has a composite hormone response element to which several members of this superfamily of nuclear receptors can bind to whereby thyroid hormones interfere with E2-stimulated oxytocin mRNA reducing E2-stimulated behaviour^(300,307). The ERE identified in the promoter region of the progesterone receptor is also bound by thyroid hormone receptors⁽²⁹⁷⁾. Progesterone, important in reproductive behaviour, correlates with a rise in appetite in cycling women⁽³¹³⁾. Thus, since T₃ may reduce E2-dependent sexual behaviour through interactions between thyroid hormone receptors and oestrogen receptors in the nuclei of hypothalamic neurons which underlie sexual behaviour⁽³⁰³⁾, their interaction may modulate eating behaviour in women throughout the cycle by acting at the nuclear receptor superfamily

consensus ERE as outlined above. Indeed, cursory searches for feeding and oestrogen receptors, thyroid hormone receptors, progesterone receptor, oxytocin and preproenkephalin return hundreds of papers altogether on feeding and each of the factors referred to above.

Over the past 30 years much has been explored on the interaction of thyroid hormones and E2 at regions of DNA co-activating and co-repressing genomic expression⁽³¹⁴⁾ and the non-genomic membrane-initiated molecular mechanisms of hormone signalling⁽³⁰⁹⁾, i.e. the cross-talk that produces a synergy between activation and repression of genomic and non-genomic mechanisms⁽³¹⁰⁾. Using a reproductive behavioural assay of rats, i.e. lordosis, it has been shown that sex hormone-dependent hypothalamic gene expression and behaviour are influenced by synaptic inputs which interfere with oestrogen receptor-dependent neuroendocrine function and behaviour⁽³¹⁴⁾. For example, the two different genes that code for TR α 1 and TR β have opposite effects on female sexual behaviours^(290,308), whereas hypothalamic neurons that express E2-sensitive genes, for example, the preproenkephalin promoter, are inhibited by TR α 1 in the ventromedial hypothalamus (VMH) but not the amygdala or caudate/putamen^(303,307,310). High doses of thyroid hormones reduce lordosis behaviour of ovariectomised E2-treated female mice⁽³¹⁵⁾ but in total pituitary RNA levels, dependent upon new protein synthesis, E2 induced increases while T₃ inhibited E2 induction⁽³¹⁶⁾. Studies of sex specificity between E2 and thyroid hormones in the regulation of pituitary function found that pituitary weight and total cellular RNA levels were significantly decreased following ovariectomy in females but increased following gonadectomy in males without changes in pituitary DNA levels compared with intact rats. Oestrogen treatment produced a significant increase in pituitary RNA in ovariectomised females but not gonadectomised males, although concomitant administration of T₃ blocked the increase in pituitary RNA in females⁽³¹⁴⁾. In contrast, thyroid hormone inhibited ER α proteolysis in pituitary lactotropes, did not prevent induction of prolactin gene expression or proliferation and inhibited the ERE reporter gene, suggesting that other endocrine factors can indirectly modulate ER α activity⁽³¹⁶⁾.

Thyroid hormone receptor is known to bind an ERE which inhibits E2-dependent transactivation⁽³¹⁷⁾. Areas of the brain involved in reproductive functions and behaviour may play a role in other appetitive behaviours such as appetite and food seeking, namely the olfactory bulb, hippocampus and granular layer of the cerebellar cortex, where TR α 1 and TR α 2 transcript levels are highest. Feedback mechanisms common to reproduction and feeding/appetitive/energy homeostasis occur in hypothalamic nuclei. For example, such mechanisms exist in the paraventricular nuclei with hypophysiotrophic neurons projecting to the median eminence, and pre-autonomic neurons controlling autonomic projections to

the liver^(287,293,318,319), arcuate nucleus^(283,297), supraoptic nucleus particularly the TR α ^(317,320), and adrenal afferents to the brainstem and suprachiasmatic nucleus via separate sympathetic and parasympathetic neuron populations⁽³¹⁹⁾. In addition, cross-talk between E2 and thyroid hormones modulates the activity in the hippocampus and amygdala contributing to cognition and mood, triggers for emotional eating in women⁽²⁸²⁾.

Nuclear receptors such as oestrogen receptors and thyroid hormone receptors bind low-molecular-weight ligands of E2, selective oestrogen response modulators, T₃ and T₄^(278,321), where they co-activate and co-repress transcription factors at targeted genomic regions^(286,304,305,321–323). However, E2 and thyroid hormones also bind to plasma membrane receptor sites initiating non-genomic, ion and intracellular Ca²⁺ channels, and G-protein-coupled secondary messenger cascades discussed above^(177,281,282,309,310,324). Future investigation into the regulation of metabolic homeostasis between the central nervous system and peripheral tissue may incorporate genomic analysis of the superfamily of nuclear receptors (oestrogen receptors and thyroid hormone receptors) in concert with the neuromodulatory factors outlined above.

Conclusion

The goal of the present review was to highlight the role of the complex hormonal interplay in the regulation of food intake and energy homeostasis. The hormones discussed in the present review are crucial in the regulation of food intake and energy homeostasis in that they play the most significant roles in the function of relevant central, mostly hypothalamic, circuits. However, they also act as the peripheral signals/mediators of peripheral tissues to inform hypothalamic circuits of the real-time status of the body's constituent energy level. Thus, it appears that ghrelin, leptin, thyroid hormones, insulin and E2 form a 'second zone' in the modulation of energy homeostasis (Fig. 1). This second zone lies between the central 'hotspot' of hypothalamic circuits and the peripheral tissues. The peripheral tissues, as appears from our review, represent a third, 'outer ring/zone' in this respect, where local interactions take place between the discussed hormones, on the one hand, and between the respective tissues and the trophic hormones, on the other hand. The apparent integrity of the discussed regulatory mechanisms suggests that the interpretation of hormonal effects alone or in simple combinations is not sufficient to understand their true nature, as both individual (feedback signals) and combined effects are necessary in the complex orchestration of energy homeostasis, including the regulation of food intake and energy expenditure. In other words, it seems most likely that the discussed hormones act as feedback signals, functioning as individual hormone signals, while their interactive effects propel and shape the bidirectional




Fig. 1. Simplified scheme of interactions between leptin (L), ghrelin (G), oestrogen (E2), insulin (I) and thyroid hormones (THs). The hypothalamus (●) is in the centre of the neuroendocrine regulation of food intake, while functionally relevant brain regions (■) influence central hypothalamic functions. The listed hormones, as feedback signals, affect the hypothalamic regulation of their own and each other's secretion, while the secreted hormones regulate the functions of relevant peripheral organs (□) directly and through local hormone interactions as well. Peripheral organs, in their interactions with the inner and outer environment (outermost circle), modulate the humoral feedback factors to tune to adequate hypothalamic regulation. All together, the 'pool' of neuromodulators acts to invoke feeding behaviour or to inhibit it, thus maintaining the energy homeostasis of the organism.

message between the central nervous system and the periphery. Our summary, therefore, points to the need for studies involving thorough, multifactorial investigation of the topic, including the analysis of simultaneous changes in the aforementioned variables from the central nervous system, as well as peripheral tissues of the experimental subjects.

Acknowledgements

Work on this project was greatly supported by Országos Tudományos Kutatási Alapprogramok (OTKA; Hungarian Scientific Research Fund) OTKA 72186 and OTKA 81745, Gazdaság és Versenyképesség Operatív Programjának (GVOP; Hungarian Economy and Competitiveness Programme) GVOP-3.2.1.-2004-0225/3.0., and the János Bolyai Research Scholarship of the Hungarian Academy of Sciences.

The authors' contributions were as follows: V. S. reviewed leptin; A. G. reviewed thyroid hormones; T. J. S. reviewed the interactions of oestrogens and thyroid hormones and edited the revised manuscript; D. S. K. reviewed insulin; G. G. reviewed ghrelin; T. B. reviewed the interactions of leptin and thyroid hormones; V. L. F. reviewed the interactions of ghrelin and leptin; A. Z. reviewed oestrogen and the interactions of oestrogen and

leptin, and wrote the abstract. The introduction and conclusion were a joint effort by all authors.

The authors declare that they have no conflict of interest with regard to the present article.

References

- Kojima M, Hosoda H, Date Y, *et al.* (1999) Ghrelin is a growth-hormone-releasing acylated peptide from stomach. *Nature* **402**, 656–660.
- Lee HM, Wang G, Englander EW, *et al.* (2002) Ghrelin, a new gastrointestinal endocrine peptide that stimulates insulin secretion: enteric distribution, ontogeny, influence of endocrine, and dietary manipulations. *Endocrinology* **143**, 185–190.
- Gualillo O, Caminos J, Blanco M, *et al.* (2001) Ghrelin, a novel placental-derived hormone. *Endocrinology* **142**, 788–794.
- Korbonits M, Bustin SA, Kojima M, *et al.* (2001) The expression of the growth hormone secretagogue receptor ligand ghrelin in normal and abnormal human pituitary and other neuroendocrine tumors. *J Clin Endocrinol Metab* **86**, 881–887.
- Kanamoto N, Akamizu T, Hosoda H, *et al.* (2001) Substantial production of ghrelin by a human medullary thyroid carcinoma cell line. *J Clin Endocrinol Metab* **86**, 4984–4990.
- Dornonville de la Cour C, Bjorkqvist M, Sandvik AK, *et al.* (2001) A-like cells in the rat stomach contain ghrelin and do not operate under gastrin control. *Regul Pept* **99**, 141–150.
- Sun Y, Asnicar M & Smith RG (2007) Central and peripheral roles of ghrelin on glucose homeostasis. *Neuroendocrinology* **86**, 215–228.
- Grouselle D, Chaillou E, Caraty A, *et al.* (2008) Pulsatile cerebrospinal fluid and plasma ghrelin in relation to growth hormone secretion and food intake in the in the sheep. *J Neuroendocrinol* **20**, 1138–1146.
- Campa D, Pardini B, Naccarati A, *et al.* (2010) Polymorphisms of genes coding for ghrelin and its receptor in relation to colorectal cancer risk: a two-step gene-wide case-control study. *BMC Gastroenterol* **10**, 112–117.
- Dass NB, Munonyara M, Bassil AK, *et al.* (2003) Growth hormone secretagogue receptors in rat and human gastrointestinal tract and the effects of ghrelin. *Neuroscience* **120**, 443–453.
- Ghelardoni S, Carnicelli V, Frascarelli S, *et al.* (2006) Ghrelin tissue distribution: comparison between gene and protein expression. *J Endocrinol Invest* **29**, 115–121.
- Tesauro M, Schinzari F, Caramanti M, *et al.* (2010) Metabolic and cardiovascular effects of ghrelin. *Int J Pept* (publication 16 March 2010).
- Holst B, Holliday ND, Bach A, *et al.* (2004) Common structural basis for constitutive activity of the ghrelin receptor family. *J Biol Chem* **279**, 53806–53817.
- Smith RG (2005) Development of growth hormone secretagogues. *Endocr Rev* **26**, 346–360.
- Korbonits M, Bustin SA, Kojima M, *et al.* (2001) The expression of the growth hormone secretagogue receptor ligand ghrelin in normal and abnormal human pituitary and other neuroendocrine tumors. *J Clin Endocrinol Metab* **86**, 881–887.
- Volante M, Allia E, Gugliotta P, *et al.* (2002) Expression of ghrelin and of the GH secretagogue receptor by pancreatic islet cells and related endocrine tumors. *J Clin Endocrinol Metab* **87**, 1300–1308.
- Motivala SJ, Tomiyama AJ, Ziegler M, *et al.* (2009) Nocturnal levels of ghrelin and leptin and sleep in chronic insomnia. *Psychoneuroendocrinology* **34**, 540–545.
- Broglio F, Prodam F, Me E, *et al.* (2005) Ghrelin: endocrine, metabolic and cardiovascular actions. *J Endocrinol Invest* **28**, 23–25.
- Dimitrova DZ, Dimitrov SD, Iliev I, *et al.* (2010) Ghrelin signaling in human mesenteric arteries. *J Physiol Pharmacol* **61**, 383–390.
- Holst B & Schwartz TW (2004) Constitutive ghrelin receptor activity as a signaling set-point in appetite regulation. *Trends Pharmacol Sci* **25**, 113–117.
- Reimer MK, Pacini G & Ahren B (2003) Dose-dependent inhibition by ghrelin of insulin secretion in the mouse. *Endocrinology* **144**, 916–921.
- Tschop M, Smiley DL & Heiman ML (2000) Ghrelin induces adiposity in rodents. *Nature* **407**, 908–913.
- Wren AM, Small CJ, Ward HL, *et al.* (2000) The novel hypothalamic peptide ghrelin stimulates food intake and growth hormone secretion. *Endocrinology* **141**, 4325–4328.
- Nakazato M, Murakami N, Date Y, *et al.* (2001) A role for ghrelin in the central regulation of feeding. *Nature* **409**, 194–198.
- De Ambrogio M, Volpe S & Tamanini C (2003) Ghrelin: central and peripheral effects of a novel peptidyl hormone. *Med Sci Monit* **9**, RA217–RA224.
- Choi K, Roh SG, Hong YH, *et al.* (2003) The role of ghrelin and growth hormone secretagogues receptor on rat adipogenesis. *Endocrinology* **144**, 754–759.
- Davies JS, Kotokorpi P, Eccles SR, *et al.* (2009) Ghrelin induces abdominal obesity via GHS-R-dependent lipid retention. *Mol Endocrinol* **23**, 914–924.
- Sun Y, Ahmed S & Smith RG (2003) Deletion of ghrelin impairs neither growth nor appetite. *Mol Cell Biol* **23**, 7973–7981.
- Wortley KE, Anderson KD, Garcia K, *et al.* (2004) Genetic deletion of ghrelin does not decrease food intake, but influences metabolic fuel preference. *Proc Natl Acad Sci U S A* **101**, 8227–8232.
- Wortley KE, del Rincon JP, Murray JD, *et al.* (2005) Absence of ghrelin protects against early-onset obesity. *J Clin Invest* **115**, 3573–3578.
- Longo KA, Charoenthongtrakul S, Giuliana DJ, *et al.* (2008) Improved insulin sensitivity and metabolic flexibility in ghrelin receptor knockout mice. *Regul Pept* **150**, 55–61.
- Tschop M, Wawarta R, Riepl RL, *et al.* (2001) Post-prandial decrease of circulating human ghrelin levels. *J Endocrinol Invest* **24**, RC19–RC21.
- Bagnasco M, Kalra PS & Kalra SP (2002) Ghrelin and leptin pulse discharge in fed and fasted rats. *Endocrinology* **143**, 726–729.
- Natalucci G, Reidl S, Gleiss A, *et al.* (2005) Spontaneous 24-h ghrelin secretion pattern in fasting subjects: maintenance of a meal-related pattern. *Eur J Endocrinol* **152**, 845–850.
- Liu J, Prudom CE, Nass R, *et al.* (2008) Novel ghrelin assays provide evidence for independent regulation of ghrelin acylation and secretion in healthy young men. *J Clin Endocrinol Metab* **93**, 1980–1987.
- Ariyasu H, Takaya K, Hosoda H, *et al.* (2002) Delayed short-term secretory regulation of ghrelin in obese animals: evidenced by a specific RIA for the active form of ghrelin. *Endocrinology* **143**, 3341–3350.
- le Roux CW, Patterson M, Vincent RP, *et al.* (2005) Postprandial plasma ghrelin is suppressed proportional to meal

- calorie content in normal-weight but not obese subjects. *J Clin Endocrinol Metab* **90**, 1068–1071.
38. Yin J, Li X, Li D, *et al.* (2009) Dietary supplementation with zinc oxide stimulates ghrelin secretion from the stomach of young pigs. *J Nutr Biochem* **20**, 783–790.
 39. Zhang H, Yin J, Li D, *et al.* (2007) Tryptophan enhances ghrelin expression and secretion associated with increased food intake and weight gain in weanling pigs. *Domest Anim Endocrinol* **33**, 47–61.
 40. Wertz-Lutz AE, Jennings JS & Clapper JA (2010) Plasma ghrelin concentrations of beef cattle consuming a similar amount of dietary energy supplied by different ingredients. *J Anim Sci* **88**, 2289–2299.
 41. Sun Y, Garcia JM & Smith RG (2007) Ghrelin and growth hormone secretagogue receptor expression in mice during aging. *Endocrinology* **148**, 1323–1329.
 42. Hsu YW, Pan YJ, Cho YM, *et al.* (2010) Aging effects on exercise-induced alternations in plasma acylated ghrelin and leptin in male rats. *Eur J Appl Physiol* (epublication ahead of print version 3 November 2010).
 43. Beaumont NJ, Skinner VO, Tan TM, *et al.* (2003) Ghrelin can bind to a species of high density lipoprotein associated with paraoxonase. *J Biol Chem* **278**, 8877–8880.
 44. Bodosi B, Gardi J, Hajdu I, *et al.* (2004) Rhythms of ghrelin, leptin and sleep in rats: effects of the normal diurnal cycle, restricted feeding, and sleep deprivation. *Am J Physiol Regul Integr Comp Physiol* **287**, R1071–R1079.
 45. Yildiz BO, Suchard MA, Wong ML, *et al.* (2004) Alterations in the dynamics of circulating ghrelin, adiponectin, and leptin in human obesity. *Proc Natl Acad Sci U S A* **101**, 10434–10439.
 46. Hahn TM, Breininger JF, Baskin DG, *et al.* (1998) Coexpression of Agrp and NPY in fasting-activated hypothalamic neurons. *Nat Neurosci* **1**, 271–272.
 47. Hewson AK & Dickson SL (2000) Systemic administration of ghrelin induces Fos and Egr-1 proteins in the hypothalamic arcuate nucleus of fasted and fed rats. *J Neuroendocrinol* **12**, 1047–1049.
 48. Cowley MA, Smith RG, Diano S, *et al.* (2003) The distribution and mechanism of action of ghrelin in the CNS demonstrates a novel hypothalamic circuit regulating energy homeostasis. *Neuron* **37**, 649–661.
 49. Andrews ZB, Liu ZW, Wallingford N, *et al.* (2008) UCP2 mediates ghrelin's action on NPY/AgRP neurons by lowering free radicals. *Nature* **454**, 846–851.
 50. DelParigi A, Tschop M, Heiman ML, *et al.* (2002) High circulating ghrelin: a potential cause for hyperphagia and obesity in Prader-Willi syndrome. *J Clin Endocrinol Metab* **87**, 5461–5464.
 51. Cummings DE, Clement K, Purnell JQ, *et al.* (2002) Elevated plasma ghrelin levels in Prader Willi syndrome. *Nat Med* **8**, 643–644.
 52. Haqq AM, Farooqi IS, O'Rahilly S, *et al.* (2003) Serum ghrelin levels are inversely correlated with body mass index, age, and insulin concentrations in normal children and are markedly increased in Prader-Willi syndrome. *J Clin Endocrinol Metab* **88**, 174–178.
 53. Cuntz U, Fruhauf E, Wawarta R, *et al.* (2002) A role for the novel weight-regulating hormone ghrelin in anorexia nervosa. *Am Clin Lab* **21**, 22–23.
 54. Soriano-Guillen L, Barrios V, Campos-Barros A, *et al.* (2004) Ghrelin levels in obesity and anorexia nervosa: effect of weight reduction or recuperation. *J Pediatr* **144**, 36–42.
 55. Otto B, Cuntz U, Fruhauf E, *et al.* (2001) Weight gain decreases elevated plasma ghrelin concentrations of patients with anorexia nervosa. *Eur J Endocrinol* **145**, 669–673.
 56. Krsek M, Rosicka M, Papezova H, *et al.* (2003) Plasma ghrelin levels and malnutrition: a comparison of two etiologies. *Eat Weight Disord* **8**, 207–211.
 57. Geloneze B, Tambascia MA, Pilla VF, *et al.* (2003) Ghrelin: a gut–brain hormone: effect of gastric bypass surgery. *Obes Surg* **13**, 17–22.
 58. Cummings DE, Weigle DS, Frayo RS, *et al.* (2002) Plasma ghrelin levels after diet-induced weight loss or gastric bypass surgery. *N Engl J Med* **346**, 1623–1630.
 59. Trostler N, Mann A, Zilberbush N, *et al.* (1995) Weight loss and food intake 18 months following vertical banded gastroplasty or gastric bypass for severe obesity. *Obes Surg* **5**, 39–51.
 60. Cummings DE, Foster-Schubert KE, Carlson MJ, *et al.* (2007) Possible hormonal mechanisms mediating the effects of bariatric surgery. In *Obesity Surgery: Principle and Practice*, pp. 137–147 [C Pitombo, editor]. New York: McGraw-Hill.
 61. Rodieux F, Giusti V, D'Alessio DA, *et al.* (2008) Effects of gastric bypass and gastric banding on glucose kinetics and gut hormone release. *Obesity* **16**, 298–305.
 62. Williams DL, Grill HJ, Cummings DE, *et al.* (2003) Vagotomy dissociates short- and long-term controls of circulating ghrelin. *Endocrinology* **144**, 5184–5187.
 63. Furuse M, Tachibana T, Ohgushi A, *et al.* (2001) Intracerebroventricular injection of ghrelin and growth hormone releasing factor inhibits food intake in neonatal chicks. *Neurosci Lett* **301**, 123–126.
 64. Kaiya H, Furuse M, Miyazato M, *et al.* (2009) Current knowledge of the roles of ghrelin in regulating food intake and energy balance in birds. *Gen Comp Endocrinol* **163**, 33–38.
 65. Chilliard Y, Delavaud C & Bonnet M (2005) Leptin expression in ruminants: nutritional and physiological regulations in relation with energy metabolism. *Domest Anim Endocrinol* **29**, 3–22.
 66. Zhang Y, Proenca R, Maffei M, *et al.* (1994) Positional cloning of the mouse obese gene and its human homologue. *Nature* **372**, 425–432.
 67. Kelesidis T, Kelesidis I, Chou S, *et al.* (2010) Narrative review: the role of leptin in human physiology: emerging clinical applications. *Ann Intern Med* **152**, 93–100.
 68. Geffroy S, De Vos P, Staels B, *et al.* (1995) Localization of the human *ob* gene to chromosome 7q32 by fluorescence *in situ* hybridization. *Genomics* **28**, 603–604.
 69. Attele AS, Shi ZQ & Yuan CS (2002) Leptin, gut, and food intake. *Biochem Pharmacol* **63**, 1579–1583.
 70. Maffei M, Halaas J, Ravussin E, *et al.* (1995) Leptin levels in human and rodent: measurement of plasma leptin and *ob* RNA in obese and weight-reduced subjects. *Nat Med* **1**, 1155–1161.
 71. Friedman JM & Halaas JL (1998) Leptin and the regulation of body weight in mammals. *Nature* **395**, 763–770.
 72. Soukas A, Cohen P, Succi ND, *et al.* (2000) Leptin-specific patterns of gene expression in white adipose tissue. *Genes Dev* **14**, 963–980.
 73. Taouis M, Chen JW, Daviaud C, *et al.* (1998) Cloning the chicken leptin gene. *Gene* **208**, 239–242.
 74. Trayhurn P (2005) Endocrine and signalling role of adipose tissue: new perspectives on fat. *Acta Physiol Scand* **184**, 285–293.
 75. Zieba DA, Amstalden M & Williams GL (2005) Regulatory roles of leptin in reproduction and metabolism: a comparative review. *Domest Anim Endocrinol* **29**, 166–185.
 76. Campfield LA, Smith FJ, Guisez Y, *et al.* (1995) Recombinant mouse *OB* protein: evidence for a peripheral signal linking adiposity and central neural networks. *Science* **269**, 546–549.

77. Coleman DL (1978) Obese and diabetes: two mutant genes causing diabetes-obesity syndromes in mice. *Diabetologia* **14**, 141–148.
78. Montague CT, Farooqi IS, Whitehead JP, *et al.* (1997) Congenital leptin deficiency is associated with severe early-onset obesity in humans. *Nature* **387**, 903–908.
79. Tartaglia LA, Dembski M, Weng X, *et al.* (1995) Identification and expression cloning of a leptin receptor, OB-R. *Cell* **83**, 1263–1271.
80. Brann DW, Wade MF, Dhandapani KM, *et al.* (2002) Leptin and reproduction. *Steroids* **67**, 95–104.
81. Chen H, Charlat O, Tartaglia LA, *et al.* (1996) Evidence that the diabetes gene encodes the leptin receptor: identification of a mutation in the leptin receptor gene in *db/db* mice. *Cell* **84**, 491–495.
82. Fei H, Okano HJ, Li C, *et al.* (1997) Anatomic localization of alternatively spliced leptin receptors (Ob-R) in mouse brain and other tissues. *Proc Natl Acad Sci U S A* **94**, 7001–7005.
83. Halaas JL, Boozer C, Blair-West J, *et al.* (1997) Physiological response to long-term peripheral and central leptin infusion in lean and obese mice. *Proc Natl Acad Sci U S A* **94**, 8878–8883.
84. Fruhbeck GA (2001) A heliocentric view of leptin. *Proc Nutr Soc* **60**, 301–318.
85. Chelikani PK, Glimm DR & Kennelly JJ (2003) Short Communication: tissue distribution of leptin and leptin receptor mRNA in the bovine. *J Dairy Sci* **86**, 2369–2372.
86. Ondicova K & Mravec B (2010) Multilevel interactions between the sympathetic and parasympathetic nervous systems: a minireview. *Endocr Regul* **44**, 69–75.
87. Yi CX, Serlie MJ, Ackermans MT, *et al.* (2009) A major role for perifornical orexin neurons in the control of glucose metabolism in rats. *Diabetes* **58**, 1998–2005.
88. Lam CK, Chari M, Su BB, *et al.* (2010) Activation of *N*-methyl-D-aspartate (NMDA) receptors in the dorsal vagal complex lowers glucose production. *J Biol Chem* **285**, 21913–21921.
89. Kalsbeek A, Bruinstroop E, Yi CX, *et al.* (2010) Hypothalamic control of energy metabolism via the autonomic nervous system. *Ann N Y Acad Sci* **2121**, 114–129.
90. Hayes MR, Skibicka KP & Grill HJ (2008) Caudal brainstem processing is sufficient for behavioral, sympathetic, and parasympathetic responses driven by peripheral and hind-brain glucagon-like-peptide-1 receptor stimulation. *Endocrinology* **149**, 4059–4068.
91. Rinaman L (2010) Ascending projections from the caudal visceral nucleus of the solitary tract to brain regions involved in food intake and energy expenditure. *Brain Res* **1350**, 18–34.
92. Satoh N, Ogawa Y, Katsuura G, *et al.* (1998) Satiety effect and sympathetic activation of leptin are mediated by hypothalamic melanocortin system. *Neurosci Lett* **249**, 107–110.
93. Scarpace PJ, Matheny M, Pollock BH, *et al.* (1997) Leptin increases uncoupling protein expression and energy expenditure. *Am J Physiol* **273**, E226–E230.
94. Chilliard Y, Bonnet M, Delavaud C, *et al.* (2001) Leptin in ruminants. Gene expression in adipose tissue and mammary gland, and regulation of plasma concentration. *Domest Anim Endocrinol* **21**, 271–295.
95. Gallardo N, Bonzon-Kulichenko E, Fernandez-Agullo T, *et al.* (2007) Tissue-specific effects of central leptin on the expression of genes involved in lipid metabolism in liver and white adipose tissue. *Endocrinology* **148**, 5604–5610.
96. Bobe G, Young JW & Beitz DC (2004) Invited review: pathology, etiology, prevention, and treatment of fatty liver in dairy cows. *J Dairy Sci* **87**, 3105–3124.
97. Mizuno TM & Mobbs CV (1999) Hypothalamic agouti-related protein messenger ribonucleic acid is inhibited by leptin and stimulated by fasting. *Endocrinology* **140**, 814–817.
98. Baskin DG, Hahn TM & Schwartz MW (1999) Leptin sensitive neurons in the hypothalamus. *Horm Metab Res* **31**, 345–350.
99. Connor EE, Laiakis EC, Fernandes VM, *et al.* (2005) Molecular cloning, expression and radiation hybrid mapping of the bovine deiodinase type II (DIO2) and deiodinase type III (DIO3) genes. *Anim Genet* **36**, 240–243.
100. Larsen PR (1997) Update on the human iodothyronine selenodeiodinases, the enzymes regulating the activation and inactivation of thyroid hormone. *Biochem Soc Trans* **25**, 588–592.
101. Hulbert AJ (2000) Thyroid hormones and their effects: a new perspective. *Biol Rev Camb Philos Soc* **75**, 519–631.
102. Darras VM, Verhoelst CH, Reyns GE, *et al.* (2006) Thyroid hormone deiodination in birds. *Thyroid* **16**, 25–35.
103. Tveit B & Larsen F (1983) Suppression and stimulation of TSH and thyroid hormones in bulls during starvation and refeeding. *Acta Endocrinol (Copenh)* **103**, 223–226.
104. St Germain DL & Galton VA (1997) The deiodinase family of selenoproteins. *Thyroid* **7**, 655–668.
105. Campos-Barros A, Hoell T, Musa A, *et al.* (1996) Phenolic and tyrosyl ring iodothyronine deiodination and thyroid hormone concentrations in the human central nervous system. *J Clin Endocrinol Metab* **81**, 2179–2185.
106. Bianco AC, Salvatore D, Gereben B, *et al.* (2002) Biochemistry, cellular and molecular biology, and physiological roles of the iodothyronine selenodeiodinases. *Endocr Rev* **23**, 38–89.
107. Gereben B, Zeold A, Dentice M, *et al.* (2008) Activation and inactivation of thyroid hormone by deiodinases: local action with general consequences. *Cell Mol Life Sci* **65**, 570–590.
108. Valverde C, Orozco A, Becerra A, *et al.* (2004) Halometabolites and cellular dehalogenase systems: an evolutionary perspective. *Int Rev Cytol* **234**, 143–199.
109. Guadano-Ferraz A, Obregon MJ, St Germain DL, *et al.* (1997) The type 2 iodothyronine deiodinase is expressed primarily in glial cells in the neonatal rat brain. *Proc Natl Acad Sci U S A* **94**, 10391–10396.
110. Tu HM, Kim SW, Salvatore D, *et al.* (1997) Regional distribution of type 2 thyroxine deiodinase messenger ribonucleic acid in rat hypothalamus and pituitary and its regulation by thyroid hormone. *Endocrinology* **138**, 3359–3368.
111. Tu HM, Legradi G, Bartha T, *et al.* (1999) Regional expression of the type 3 iodothyronine deiodinase messenger ribonucleic acid in the rat central nervous system and its regulation by thyroid hormone. *Endocrinology* **140**, 784–790.
112. Puymirat J, L'Héroult S & Dussault JH (1992) Expression of thyroid hormone receptors mRNAs in rat cerebral hemisphere neuronal cultures. *Brain Res Dev Brain Res* **69**, 173–177.
113. Escamez MJ, Guadano-Ferraz A, Cuadrado A, *et al.* (1999) Type 3 iodothyronine deiodinase is selectively expressed in areas related to sexual differentiation in the newborn rat brain. *Endocrinology* **140**, 5443–5446.
114. Rudas P, Bartha T & Frenyo LV (1993) Thyroid hormone deiodination in the brain of young chickens acutely adapts to changes in thyroid status. *Acta Vet Hung* **41**, 381–393.
115. Van der Geyten S, Van Rompaey E, Sanders JP, *et al.* (1999) Regulation of thyroid hormone metabolism during fasting

- and refeeding in chicken. *Gen Comp Endocrinol* **116**, 272–280.
116. Kohrle J (1999) Local activation and inactivation of thyroid hormones: the deiodinase family. *Mol Cell Endocrinol* **151**, 103–119.
 117. Pezzi C, Accorsi PA, Vigo D, *et al.* (2003) 5'-Deiodinase activity and circulating thyronines in lactating cows. *J Dairy Sci* **86**, 152–158.
 118. Bianco AC & Larsen PR (2005) Intracellular pathways of iodothyronine metabolism. In *Werner and Ingbar's The Thyroid: A Fundamental and Clinical Text*, 9th ed., pp. 109–125 [LE Braverman and RD Utiger, editors]. Philadelphia: Lippincott Williams & Wilkins.
 119. Larsen PR, Silva JE & Kaplan MM (1981) Relationships between circulating and intracellular thyroid hormones: physiological and clinical implications. *Endocr Rev* **2**, 87–102.
 120. Rudas P & Pethes G (1986) Acute changes of the conversion of thyroxine to triiodothyronine in hypophysectomized and thyroidectomized chickens exposed to mild cold (10 degrees). *Gen Comp Endocrinol* **63**, 408–413.
 121. Sap J, de Magistris L, Stunnenberg H, *et al.* (1990) A major thyroid hormone response element in the third intron of the rat growth hormone gene. *EMBO J* **9**, 887–896.
 122. Brent GA, Moore DD & Larsen PR (1991) Thyroid hormone regulation of gene expression. *Annu Rev Physiol* **53**, 17–35.
 123. Shi YB, Yaoita Y & Brown DD (1992) Genomic organization and alternative promoter usage of the two thyroid hormone receptor β genes in *Xenopus laevis*. *J Biol Chem* **267**, 733–738.
 124. Zhang J & Lazar MA (2000) The mechanism of action of thyroid hormones. *Annu Rev Physiol* **62**, 439–466.
 125. Koenig RJ, Lazar MA, Hodin RA, *et al.* (1989) Inhibition of thyroid hormone action by a non-hormone binding c-erbA protein generated by alternative mRNA splicing. *Nature* **337**, 659–661.
 126. Salto C, Kindblom JM, Johansson C, *et al.* (2001) Ablation of TR α 2 and a concomitant overexpression of α 1 yields a mixed hypo- and hyperthyroid phenotype in mice. *Mol Endocrinol* **15**, 2115–2128.
 127. Chen JD & Evans RM (1995) A transcriptional co-repressor that interacts with nuclear hormone receptors. *Nature* **377**, 454–457.
 128. Bradley DJ, Towle HC & Young WS III (1994) α and β Thyroid hormone receptor (TR) gene expression during auditory neurogenesis: evidence for TR isoform-specific transcriptional regulation *in vivo*. *Proc Natl Acad Sci U S A* **91**, 439–443.
 129. Angelin-Duclos C, Domenget C, Kolbus A, *et al.* (2005) Thyroid hormone T₃ acting through the thyroid hormone α receptor is necessary for implementation of erythropoiesis in the neonatal spleen environment in the mouse. *Development* **132**, 925–934.
 130. O'Shea PJ, Bassett JH, Sriskantharajah S, *et al.* (2005) Contrasting skeletal phenotypes in mice with an identical mutation targeted to thyroid hormone receptor α 1 or β . *Mol Endocrinol* **19**, 3045–3059.
 131. Plateroti M, Kress E, Mori JI, *et al.* (2006) Thyroid hormone receptor α 1 directly controls transcription of the β -catenin gene in intestinal epithelial cells. *Mol Cell Biol* **26**, 3204–3214.
 132. Amma LL, Campos-Barros A, Wang Z, *et al.* (2001) Distinct tissue-specific roles for thyroid hormone receptors β and α 1 in regulation of type 1 deiodinase expression. *Mol Endocrinol* **15**, 467–475.
 133. Ng L, Pedraza PE, Faris JS, *et al.* (2001) Audiogenic seizure susceptibility in thyroid hormone receptor β -deficient mice. *Neuroreport* **12**, 2359–2362.
 134. Flores-Morales A, Gullberg H, Fernandez L, *et al.* (2002) Patterns of liver gene expression governed by TR β . *Mol Endocrinol* **16**, 1257–1268.
 135. Abel ED, Moura EG, Ahima RS, *et al.* (2003) Dominant inhibition of thyroid hormone action selectively in the pituitary of thyroid hormone receptor- β null mice abolishes the regulation of thyrotropin by thyroid hormone. *Mol Endocrinol* **17**, 1767–1776.
 136. Esaki T, Suzuki H, Cook M, *et al.* (2003) Functional activation of cerebral metabolism in mice with mutated thyroid hormone nuclear receptors. *Endocrinology* **144**, 4117–4122.
 137. Siesser WB, Cheng SY & McDonald MP (2005) Hyperactivity, impaired learning on a vigilance task, and a differential response to methylphenidate in the TR β PV knock-in mouse. *Psychopharmacology (Berl)* **181**, 653–663.
 138. Forrest D, Erway LC, Ng L, *et al.* (1996) Thyroid hormone receptor β is essential for development of auditory function. *Nat Genet* **13**, 354–357.
 139. Gothe S, Wang Z, Ng L, *et al.* (1999) Mice devoid of all known thyroid hormone receptors are viable but exhibit disorders of the pituitary–thyroid axis, growth, and bone maturation. *Genes Dev* **13**, 1329–1341.
 140. Drucker DJ (2002) Gut adaptation and the glucagon-like peptides. *Gut* **50**, 428–435.
 141. Danforth E Jr & Burger AG (1989) The impact of nutrition on thyroid hormone physiology and action. *Annu Rev Nutr* **9**, 201–227.
 142. Bartha T (1993) Thyroid hormone metabolism in broiler chickens. PhD Thesis. Catholic University of Leuven.
 143. Brown SB, Maloney M & Kinlaw WB (1997) 'Spot 14' protein functions at the pretranslational level in the regulation of hepatic metabolism by thyroid hormone and glucose. *J Biol Chem* **272**, 2163–2166.
 144. Matosin-Matekalo M, Mesonero JE, Laroche TJ, *et al.* (1999) Glucose and thyroid hormone co-regulate the expression of the intestinal fructose transporter GLUT5. *Biochem J* **339**, 233–239.
 145. Sato K & Robbins J (1981) Thyroid hormone metabolism in primary cultured rat hepatocytes. Effects of glucose, glucagon, and insulin. *J Clin Invest* **68**, 475–483.
 146. Kong WM, Martin NM, Smith KL, *et al.* (2004) Triiodothyronine stimulates food intake via the hypothalamic ventromedial nucleus independent of changes in energy expenditure. *Endocrinology* **145**, 5252–5258.
 147. Györfi A, Rónai ZS, Áprily SZ, *et al.* (2008) Metabolic and hormonal aspects of fatty liver production in liver- and meat-type goose hybrids (article in Hungarian). *Magy Allatorv Lapja* **130**, 156–164.
 148. Hugues JN, Burger AG, Pekary AE, *et al.* (1984) Rapid adaptations of serum thyrotrophin, triiodothyronine and reverse triiodothyronine levels to short-term starvation and refeeding. *Acta Endocrinol (Copenh)* **105**, 194–199.
 149. Kaplan MM & Utiger RD (1978) Iodothyronine metabolism in liver and kidney homogenates from hyperthyroid and hypothyroid rats. *Endocrinology* **103**, 156–161.
 150. Boelen A, Wiersinga WM & Fliers E (2008) Fasting-induced changes in the hypothalamus–pituitary–thyroid axis. *Thyroid* **18**, 123–129.
 151. Slebodzinski AB, Brzezinska-Slebodzinska E & Drews R (1982) Reciprocal changes in serum 3, 3', 5'-tri-iodothyronine concentration and the peripheral thyroxine inner ring monodeiodination during food restriction in the young pig. *J Endocrinol* **95**, 349–355.
 152. Macari M & Baraldi SM (1983) Effects of fasting, noradrenaline and glucose tolerance test on free fatty acids and blood glucose in thyroid-deficient pigs. *Comp Biochem Physiol B* **74**, 743–747.

153. Houpt KA, Reimers TJ & Boyd RD (1986) Changes in free fatty acids and triiodothyronine in response to feeding in pigs. *Physiol Behav* **37**, 573–576.
154. King DB, King CR & Eshleman JR (1977) Serum triiodothyronine levels in the embryonic and post-hatching chicken, with particular reference to feeding-induced changes. *Gen Comp Endocrinol* **31**, 216–233.
155. Scanes CG & Griminger P (1990) Endocrine–nutrition interactions in birds. *J Exp Zool Suppl* **4**, 98–105.
156. Buyse J, Janssens K, Van der Geyten S, *et al.* (2002) Pre- and postprandial changes in plasma hormone and metabolite levels and hepatic deiodinase activities in meal-fed broiler chickens. *Br J Nutr* **88**, 641–653.
157. Reyns GE, Janssens KA, Buyse J, *et al.* (2002) Changes in thyroid hormone levels in chicken liver during fasting and refeeding. *Comp Biochem Physiol B Biochem Mol Biol* **132**, 239–245.
158. Györfly A, Sayed-Ahmed A, Zsarnovszky A, *et al.* (2009) Effects of energy restriction on thyroid hormone metabolism in chickens. *Acta Vet Hung* **57**, 319–330.
159. Pethes G, Bokori J, Rudas P, *et al.* (1985) Thyroxine, triiodothyronine, reverse-triiodothyronine, and other physiological characteristics of periparturient cows fed restricted energy. *J Dairy Sci* **68**, 1148–1154.
160. Romo GA, Elsasser TH, Kahl S, *et al.* (1997) Dietary fatty acids modulate hormone responses in lactating cows: mechanistic role for 5'-deiodinase activity in tissue. *Domest Anim Endocrinol* **14**, 409–420.
161. Coppola A, Liu WZ, Andrews Z, *et al.* (2007) A central thermogenic-like mechanism in feeding regulation: an interplay between arcuate nucleus T₃ and UCP2. *Cell Metab* **5**, 21–33.
162. Ishii S, Kamegai J, Tamura H, *et al.* (2008) Triiodothyronine (T₃) stimulates food intake via enhanced hypothalamic AMP-activated kinase activity. *Regul Pept* **151**, 164–169.
163. Nelson LR & Bulun SE (2001) Estrogen production and action. *J Am Acad Dermatol* **45**, S116–S124.
164. Sharpe RM (1998) The roles of oestrogen in the male. *Trends Endocrinol Metab* **9**, 371–377.
165. Mystkowski P & Schwartz MW (2000) Gonadal steroids and energy homeostasis in the leptin era. *Nutrition* **16**, 937–946.
166. Meli R, Pacilio M, Raso GM, *et al.* (2004) Estrogen and raloxifene modulate leptin and its receptor in hypothalamus and adipose tissue from ovariectomized rats. *Endocrinology* **145**, 3115–3121.
167. Wade GN, Gray JM & Bartness TJ (1985) Gonadal influences on adiposity. *Int J Obes* **9**, Suppl. 1, 83–92.
168. Shimomura Y, Shimizu H, Kobayashi I, *et al.* (1989) Importance of feeding time in pair-fed, ovariectomized rats. *Physiol Behav* **45**, 1197–1200.
169. Uchida Y, Kano M, Yasuhara S, *et al.* (2010) Estrogen modulates central and peripheral responses to cold in female rats. *J Physiol Sci* **60**, 151–160.
170. Qiu J, Bosch MA, Tobias SC, *et al.* (2003) Rapid signaling of estrogen in hypothalamic neurons involves a novel G-protein-coupled estrogen receptor that activates protein kinase C. *J Neurosci* **23**, 9529–9540.
171. Qiu J, Bosch MA, Tobias SC, *et al.* (2006) A G-protein-coupled estrogen receptor is involved in hypothalamic control of energy homeostasis. *J Neurosci* **26**, 5649–5655.
172. Musatov S, Chen W, Pfaff DW, *et al.* (2007) Silencing of estrogen receptor α in the ventromedial nucleus of hypothalamus leads to metabolic syndrome. *Proc Natl Acad Sci U S A* **104**, 2501–2506.
173. Wong JK, Le HH, Zsarnovszky A, *et al.* (2003) Estrogens and ICI182,780 (Faslodex) modulate mitosis and cell death in immature cerebellar neurons via rapid activation of p44/p42 mitogen-activated protein kinase. *J Neurosci* **23**, 4984–4995.
174. Belcher SM & Zsarnovszky A (2001) Estrogenic actions in the brain: estrogen, phytoestrogens, and rapid intracellular signaling mechanisms. *J Pharmacol Exp Ther* **299**, 408–414.
175. Horvath TL, Warden CH, Hajos M, *et al.* (1999) Brain uncoupling protein 2: uncoupled neuronal mitochondria predict thermal synapses in homeostatic centers. *J Neurosci* **19**, 10417–10427.
176. Kiss DS, Zsarnovszky A, Horvath K, *et al.* (2009) Ecto-nucleoside triphosphate diphosphohydrolase 3 in the ventral and lateral hypothalamic area of female rats: morphological characterization and functional implications. *Reprod Biol Endocrinol* **7**, 31–43.
177. Ropero AB, Alonso-Magdalena P, Quesada I, *et al.* (2008) The role of estrogen receptors in the control of energy and glucose homeostasis. *Steroids* **73**, 874–879.
178. Naftolin F, Garcia-Segura LM, Horvath TL, *et al.* (2007) Estrogen-induced hypothalamic synaptic plasticity and pituitary sensitization in the control of the estrogen-induced gonadotrophin surge. *Reprod Sci* **14**, 101–116.
179. Horvath TL (2005) The hardship of obesity: a soft-wired hypothalamus. *Nat Neurosci* **8**, 561–565.
180. Kalsbeek A, Fliers E, Hofman MA, *et al.* (2006) *Hypothalamic Integration of Energy Metabolism: Progress in Brain Research*. Oxford: Elsevier.
181. Gao Q & Horvath TL (2007) Neurobiology of feeding and energy expenditure. *Ann Rev Neurosci* **30**, 367–398.
182. Gao Q & Horvath TL (2008) Cross talk between estrogen and leptin signalling in the hypothalamus. *Am J Physiol Endocrinol Metab* **294**, E817–E826.
183. Olofsson LE, Pierce AA & Xu AW (2009) Functional requirement of AgRP and NPY neurons in ovarian cycle-dependent regulation of food intake. *Proc Natl Acad Sci U S A* **106**, 15932–15937.
184. Kellert BA, Nguyen MC, Nguyen C, *et al.* (2009) Estrogen rapidly attenuates cannabinoid-induced changes in energy homeostasis. *Eur J Pharmacol* **622**, 15–24.
185. Schwartz MW, Figlewicz DP, Baskin DG, *et al.* (1992) Insulin in the brain: a hormonal regulator of energy balance. *Endocr Rev* **13**, 387–414.
186. Kaiyala KJ, Prigeon RL, Kahn SE, *et al.* (2000) Obesity induced by a high-fat diet is associated with reduced brain insulin transport in dogs. *Diabetes* **49**, 1525–1533.
187. Pardini AW, Nguyen HT, Figlewicz DP, *et al.* (2006) Distribution of insulin receptor substrate-2 in brain areas involved in energy homeostasis. *Brain Res* **1112**, 169–178.
188. Gerozissis K (2008) Brain insulin, energy and glucose homeostasis; genes, environment and metabolic pathologies. *Eur J Pharmacol* **585**, 38–49.
189. Figlewicz DP (2003) Adiposity signals and food reward: expanding the CNS roles of insulin and leptin. *Am J Physiol Regul Integr Comp Physiol* **284**, R882–R892.
190. Air EL, Benoit SC, Clegg DJ, *et al.* (2002) Insulin and leptin combine additively to reduce food intake and body weight in rats. *Endocrinology* **143**, 2449–2452.
191. Pralong FP (2010) Insulin and NPY pathways and the control of GnRH function and puberty onset. *Mol Cell Endocrinol* **324**, 82–86.
192. Frankish HM, Dryden S, Hopkins D, *et al.* (1995) Neuropeptide Y, the hypothalamus, and diabetes: insights into the central control of metabolism. *Peptides* **16**, 757–771.
193. Lin HV, Plum L, Ono H, *et al.* (2010) Divergent regulation of energy expenditure and hepatic glucose production by

- insulin receptor in agouti-related protein and POMC neurons. *Diabetes* **59**, 337–346.
194. Zigman JM & Elmquist JK (2003) Minireview: from anorexia to obesity – the yin and yang of body weight control. *Endocrinology* **144**, 3749–3756.
 195. Giovambattista A, Piermaria J, Suescun MO, *et al.* (2006) Direct effect of ghrelin on leptin production by cultured rat white adipocytes. *Obesity* **14**, 19–27.
 196. Barazzoni R, Zanetti M, Stebel M, *et al.* (2003) Hyperleptinemia prevents increased plasma ghrelin concentration during short-term moderate caloric restriction in rats. *Gastroenterology* **124**, 1188–1192.
 197. Shiiya T, Nakazato M, Mizuta M, *et al.* (2002) Plasma ghrelin levels in lean and obese humans and the effect of glucose on ghrelin secretion. *J Clin Endocrinol Metab* **87**, 240–244.
 198. Rigamonti AE, Pincelli AI, Corra B, *et al.* (2002) Plasma ghrelin concentrations in elderly subjects: comparison with anorexic and obese patients. *J Endocrinol* **175**, R1–R5.
 199. McCowen KC, Maykel JA, Bistrrian BR, *et al.* (2002) Circulating ghrelin concentrations are lowered by intravenous glucose or hyperinsulinemic euglycemic conditions in rodents. *J Endocrinol* **175**, R7–R11.
 200. Rosicka M, Krsek M, Jarkovska Z, *et al.* (2002) Ghrelin – a new endogenous growth hormone secretagogue. *Physiol Res* **51**, 435–441.
 201. Erdmann J, Lippl F & Schusdziarra V (2003) Differential effect of protein and fat on plasma ghrelin levels in man. *Regul Pept* **116**, 101–107.
 202. Greenman Y, Golani N, Gilad S, *et al.* (2004) Ghrelin secretion is modulated in a nutrient- and gender-specific manner. *Clin Endocrinol (Oxf)* **60**, 382–388.
 203. Figlewicz DP, Evans SB, Murphy J, *et al.* (2003) Expression of receptors for insulin and leptin in the ventral tegmental area/substantia nigra (VTA/SN) of the rat. *Brain Res* **964**, 107–115.
 204. Guan XM, Yu H, Palyha OC, *et al.* (1997) Distribution of mRNA encoding the growth hormone secretagogue receptor in brain and peripheral tissues. *Brain Res Mol Brain Res* **48**, 23–29.
 205. Wise RA (2004) Dopamine and food reward: back to the elements. *Am J Physiol Regul Integr Comp Physiol* **286**, R13.
 206. Wise RA (2004) Dopamine, learning and motivation. *Nat Rev Neurosci* **5**, 483–494.
 207. Wise RA & Rompre PP (1989) Brain dopamine and reward. *Annu Rev Psychol* **40**, 191–225.
 208. Date Y, Nakazato M, Hashiguchi S, *et al.* (2002) Ghrelin is present in pancreatic α -cells of humans and rats and stimulates insulin secretion. *Diabetes* **51**, 124–129.
 209. Kageyama H, Funahashi H, Hirayama M, *et al.* (2005) Morphological analysis of ghrelin and its receptor distribution in the rat pancreas. *Regul Pept* **126**, 67–71.
 210. Dezaki K, Sone H & Yada T (2008) Ghrelin is a physiological regulator of insulin release in pancreatic islets and glucose homeostasis. *Pharmacol Ther* **118**, 239–249.
 211. Yada T, Dezaki K, Sone H, *et al.* (2008) Ghrelin regulates insulin release and glycemia: physiological role and therapeutic potential. *Curr Diabetes Rev* **4**, 18–23.
 212. Dezaki K, Kakei M & Yada T (2007) Ghrelin uses G α 2 and activates K v channels to attenuate glucose-induced Ca $^{2+}$ signaling and insulin release in islet β -cells: novel signal transduction of ghrelin. *Diabetes* **56**, 2319–2327.
 213. Dezaki K, Hosoda H, Kakei M, *et al.* (2004) Endogenous ghrelin in pancreatic islets restricts insulin release by attenuating Ca $^{2+}$ signaling in β -cells: implication in the glycemic control in rodents. *Diabetes* **53**, 3142–3151.
 214. Rogers RC, Barnes MJ & Hermann GE (2009) Leptin ‘gates’ thermogenic action of thyrotropin-releasing hormone in the hindbrain. *Brain Res* **1295**, 135–141.
 215. Zabrocka L, Klimek J & Swierczynski J (2006) Evidence that triiodothyronine decreases rat serum leptin concentration by down-regulation of leptin gene expression in white adipose tissue. *Life Sci* **79**, 1114–1120.
 216. Medina-Gomez G, Calvo RM & Obregon MJ (2004) T $_3$ and Triac inhibit leptin secretion and expression in brown and white rat adipocytes. *Biochim Biophys Acta* **1682**, 38–47.
 217. Legradi G, Emerson CH, Ahima RS, *et al.* (1997) Leptin prevents fasting-induced suppression of prothyrotropin-releasing hormone messenger ribonucleic acid in neurons of the hypothalamic paraventricular nucleus. *Endocrinology* **138**, 2569–2576.
 218. Seoane LM, Carro E, Tovar S, *et al.* (2000) Regulation of *in vivo* TSH secretion by leptin. *Regul Pept* **92**, 25–29.
 219. Cettour-Rose P, Burger AG, Meier CA, *et al.* (2002) Central stimulatory effect of leptin on T $_3$ production is mediated by brown adipose tissue type II deiodinase. *Am J Physiol Endocrinol Metab* **283**, E980–E987.
 220. Ghamari-Langroudi M, Vella KR, Srisai D, *et al.* (2010) Regulation of thyrotropin-releasing hormone-expressing neurons in paraventricular nucleus of the hypothalamus by signals of adiposity. *Mol Endocrinol* **24**, 2366–2381.
 221. Otukonyong EE, Okutani F, Takahashi S, *et al.* (2000) Effect of food deprivation and leptin repletion on the plasma levels of estrogen (E $_2$) and NADPH-d reactivity in the ventromedial and arcuate nuclei of the hypothalamus in the female rats. *Brain Res* **887**, 70–79.
 222. Temple JL & Rissman EF (2000) Acute re-feeding reverses food restriction-induced hypothalamic–pituitary–gonadal axis deficits. *Biol Reprod* **63**, 1721–1726.
 223. Brecchia G, Bonanno A, Galeati G, *et al.* (2006) Hormonal and metabolic adaptation to fasting: effects on the hypothalamic–pituitary–ovarian axis and reproductive performance of rabbit does. *Domest Anim Endocrinol* **31**, 105–122.
 224. Barash IA, Cheung CC, Weigle DS, *et al.* (1996) Leptin is a metabolic signal to the reproductive system. *Endocrinology* **137**, 3144–3147.
 225. Bronson FH (1986) Food-restricted, prepubertal, female rats: rapid recovery of luteinizing hormone pulsing with excess food, and full recovery of pubertal development with gonadotropin-releasing hormone. *Endocrinology* **118**, 2483–2487.
 226. Chen HP, Fan J & Cui S (2006) Detection and estrogen regulation of leptin receptor expression in rat dorsal root ganglion. *Histochem Cell Biol* **126**, 363–369.
 227. Ainslie DA, Morris MJ, Wittert G, *et al.* (2001) Estrogen deficiency causes central leptin insensitivity and increased hypothalamic neuropeptide Y. *Int J Obes* **25**, 1680–1688.
 228. Chen Y & Heiman ML (2000) Chronic leptin administration promotes lipid utilization until fat mass is greatly reduced and preserves lean body mass of normal female rats. *Regul Pept* **92**, 113–119.
 229. Chakraborty S, Sachdev A, Salton SR, *et al.* (2008) Stereological analysis of estrogen receptor expression in the hypothalamic arcuate nucleus of *ob/ob* and *agouti* mice. *Brain Res* **1217**, 86–95.
 230. Diano S, Kalra SP, Sakamoto H, *et al.* (1998) Leptin receptors in estrogen receptor-containing neurons of the female rat hypothalamus. *Brain Res* **812**, 256–259.
 231. Hirose M, Minata M, Harada KH, *et al.* (2008) Ablation of estrogen receptor α (ER α) prevents upregulation of POMC by leptin and insulin. *Biochem Biophys Res Commun* **371**, 320–323.

232. Shin JH, Hur JY, Seo HS, *et al.* (2007) The ratio of estrogen receptor α to estrogen receptor β in adipose tissue is associated with leptin production and obesity. *Steroids* **72**, 592–599.
233. Konner AC, Klockener T & Bruning JC (2009) Control of energy homeostasis by insulin and leptin: targeting the arcuate nucleus and beyond. *Physiol Behav* **97**, 632–638.
234. Carvalheira JB, Siloto RM, Ignacchitti I, *et al.* (2001) Insulin modulates leptin-induced STAT3 activation in rat hypothalamus. *FEBS Lett* **500**, 119–124.
235. Carvalheira JB, Ribeiro EB, Folli F, *et al.* (2003) Interaction between leptin and insulin signaling pathways differentially affects JAK-STAT and PI 3-kinase-mediated signaling in rat liver. *Biol Chem* **384**, 151–159.
236. Carvalheira JB, Torsoni MA, Ueno M, *et al.* (2005) Cross-talk between the insulin and leptin signalling systems in rat hypothalamus. *Obes Res* **13**, 48–57.
237. Ahima RS, Saper CB, Flier JS, *et al.* (2000) Leptin regulation of neuroendocrine systems. *Front Neuroendocrinol* **21**, 263–307.
238. Finn PD, Cunningham MJ, Rickard DG, *et al.* (2001) Serotonergic neurons are targets for leptin in the monkey. *J Clin Endocrinol Metab* **86**, 422–426.
239. Orosco M, Rouch C & Gerozissis K (2000) Activation of hypothalamic insulin by serotonin is the primary event of the insulin–serotonin interaction involved in the control of feeding. *Brain Res* **872**, 64–70.
240. Buyse M, Viengchareun S, Bado A, *et al.* (2001) Insulin and glucocorticoids differentially regulate leptin transcription and secretion in brown adipocytes. *FASEB J* **15**, 1357–1366.
241. Cusin I, Sainsbury A, Doyle P, *et al.* (1995) The *ob* gene and insulin. A relationship leading to clues to the understanding of obesity. *Diabetes* **44**, 1467–1470.
242. Saladin R, De Vos P, Guerre-Millo M, *et al.* (1995) Transient increase in obese gene expression after food intake or insulin administration. *Nature* **377**, 527–529.
243. Sivitz WI, Walsh S, Morgan D, *et al.* (1998) Plasma leptin in diabetic and insulin-treated diabetic and normal rats. *Metabolism* **47**, 584–591.
244. Havrankova J, Roth J & Brownstein M (1978) Insulin receptors are widely distributed in the central nervous system of the rat. *Nature* **272**, 827–829.
245. Werther GA, Hogg A, Oldfield BJ, *et al.* (1987) Localization and characterization of insulin receptors in rat brain and pituitary gland using *in vitro* autoradiography and computerized densitometry. *Endocrinology* **121**, 1562–1570.
246. Marks JL, Porte D Jr, Stahl WL, *et al.* (1990) Localization of insulin receptor mRNA in rat brain by *in situ* hybridization. *Endocrinology* **127**, 3234–3236.
247. Shyng SL & Nichols CG (1998) Membrane phospholipid control of nucleotide sensitivity of KATP channels. *Science* **282**, 1138–1141.
248. Yang MJ, Wang F, Wang JH, *et al.* (2010) PI3K integrates the effects of insulin and leptin on large-conductance Ca^{2+} -activated K^{+} channels in neuropeptide Y neurons of the hypothalamic arcuate nucleus. *Am J Physiol Endocrinol Metab* **298**, E193–E201.
249. Heine PA, Taylor JA, Iwamoto GA, *et al.* (2000) Increased adipose tissue in male and female estrogen receptor- α knockout mice. *Proc Natl Acad Sci U S A* **97**, 12729–12734.
250. Takeda K, Toda K, Saibara T, *et al.* (2003) Progressive development of insulin resistance phenotype in male mice with complete aromatase (CYP19) deficiency. *J Endocrinol* **176**, 237–246.
251. Naaz A, Zakroczymski M, Heine P, *et al.* (2002) Effect of ovariectomy on adipose tissue of mice in the absence of estrogen receptor α (ER α): a potential role for estrogen receptor β (ER β). *Horm Metab Res* **34**, 758–763.
252. Clegg DJ, Brown LM, Woods SC, *et al.* (2006) Gonadal hormones determine sensitivity to central leptin and insulin. *Diabetes* **55**, 978–987.
253. Geer EB & Shen W (2006) Gender differences in insulin resistance, body composition and energy balance. *Genet Med* **6**, Suppl. 1, 60–75.
254. Barros RP, Machado UF, Warner M, *et al.* (2006) Muscle GLUT4 regulation by estrogen receptors ER β and ER α . *Proc Natl Acad Sci U S A* **103**, 1605–1608.
255. Brussaard HE, Gevers Leuven JA, Frolich M, *et al.* (1997) Short-term oestrogen replacement therapy improves insulin resistance, lipids and fibrinolysis in postmenopausal women with NIDDM. *Diabetologia* **40**, 843–849.
256. Stevenson JC, Crook D, Godsland IF, *et al.* (1994) Hormone replacement therapy and the cardiovascular system: nonlipid effects. *Drugs* **47**, Suppl. 2, 35–41.
257. Ding EL, Song Y, Manson JE, *et al.* (2007) Plasma sex steroid hormones and risk of developing type 2 diabetes in women: a prospective study. *Diabetologia* **50**, 2076–2084.
258. Groot LJ & Jameson JL (1995) Mechanism of thyroid hormone action. In *Endocrinology*, 3rd ed., pp. 583–601 [LJ DeGroot, editor]. Philadelphia, PA: W.B. Saunders.
259. Granner D & Pilkis S (1990) The genes of hepatic glucose metabolism. *J Biol Chem* **265**, 10173–10182.
260. Sharp GW (1979) The adenylate cyclase–cyclic AMP system in islets of Langerhans and its role in the control of insulin release. *Diabetologia* **16**, 287–296.
261. Weinstein SP, Watts J, Graves PN, *et al.* (1990) Stimulation of glucose transport by thyroid hormone in ARL 15 cells: increased abundance of glucose transporter protein and messenger ribonucleic acid. *Endocrinology* **126**, 1421–1429.
262. Meyers DS, Skwish S, Dickinson KE, *et al.* (1997) β 3-Adrenergic receptor-mediated lipolysis and oxygen consumption in brown adipocytes from cynomolgus monkeys. *J Clin Endocrinol Metab* **82**, 395–401.
263. Scheja L, Makowski L, Uysal KT, *et al.* (1999) Altered insulin secretion associated with reduced lipolytic efficiency in *ap2 -/-* mice. *Diabetes* **48**, 1987–1994.
264. Pan DA, Lillioja S, Kriketos AD, *et al.* (1997) Skeletal muscle triglyceride levels are inversely related to insulin action. *Diabetes* **46**, 983–988.
265. Pittman CS (1981) The effects of diabetes mellitus on thyroid physiology. *Thyroid Today* **4**, 1–4.
266. Ohara H, Kobayashi T, Shiraishi M, *et al.* (1974) Thyroid function of the aged as viewed from the pituitary thyroid system. *Endocrinol Jpn* **21**, 377–386.
267. Kyte J (1981) Molecular considerations relevant to the mechanism of active transport. *Nature* **292**, 201–204.
268. Lenzen S & Bailey CJ (1984) Thyroid hormones, gonadal and adrenocortical steroids and the function of the islets of Langerhans. *Endocr Rev* **5**, 411–434.
269. Lenzen S & Kloppel G (1978) Insulin secretion and the morphological and metabolic characteristics of pancreatic islets of hyperthyroid *ob/ob* mice. *Endocrinology* **103**, 1546–1555.
270. Lenzen S, Joost HG & Hasselblatt A (1976) Thyroid function and insulin secretion from the perfused pancreas in the rat. *Endocrinology* **99**, 125–129.
271. Jorns A, Tiedge M & Lenzen S (2002) Thyroxine induces pancreatic β cell apoptosis in rats. *Diabetologia* **45**, 851–855.
272. Ximenes HM, Lortz S, Jorns A, *et al.* (2007) Triiodothyronine (T_3)-mediated toxicity and induction of apoptosis in insulin-producing INS-1 cells. *Life Sci* **80**, 2045–2050.

273. Wu P (2000) Thyroid disease and diabetes. *Clin Diabetes* **18**, 38–39.
274. Franklyn JA, Sheppard MC & Ramsden DB (1983) Serum free thyroxine and free triiodothyronine concentrations in pregnancy. *BMJ* **287**, 394.
275. Franklyn JA, Wood DF, Balfour NJ, *et al.* (1987) Modulation of estrogen of thyroid effects of thyrotrophin gene expression. *J Endocrinol* **115**, 53–59.
276. Ahlquist JA, Franklyn JA, Wood DF, *et al.* (1987) Hormonal regulation of thyrotrophin synthesis and secretion. *Horm Metab Res Suppl* **17**, 86–89.
277. Fitts JM, Klein RM & Powers CA (1998) Comparison of tamoxifen effects on the actions of triiodothyronine or growth hormone in the ovariectomized-hypothyroid rat. *J Pharmacol Exp Ther* **286**, 392–402.
278. Fitts JM, Klein RM & Powers AC (2001) Estrogen and tamoxifen interplay with T₃ in male rats: pharmacologically distinct classes of estrogen responses affecting growth, bone, and lipid metabolism, and their relation to serum GH and IGF-I. *Endocrinology* **142**, 4223–4235.
279. Dahl GE, Evans NP, Moenter SM, *et al.* (1994) The thyroid gland is required for reproductive neuroendocrine responses to photoperiod in the ewe. *Endocrinology* **135**, 10–15.
280. Vasudevan N, Zhu YS, Daniel S, *et al.* (2001) Crosstalk between oestrogen receptors and thyroid hormone receptor isoforms results in differential regulation of the preproenkephalin gene. *J Neuroendocrinol* **13**, 779–790.
281. Vasudevan N, Kow LM & Pfaff DW (2001) Early membrane estrogenic effects required for full expression of slower genomic actions in a nerve cell line. *Proc Natl Acad Sci U S A* **98**, 12267–12271.
282. Vasudevan N & Pfaff DW (2007) Membrane-initiated actions of estrogens in neuroendocrinology: emerging principles. *Endocr Rev* **28**, 1–19.
283. Klieverik LP, Janssen SF, van Riel A, *et al.* (2009) Thyroid hormone modulates glucose production via a sympathetic pathway from the hypothalamic paraventricular nucleus to the liver. *Proc Natl Acad Sci U S A* **106**, 5966–5971.
284. Maratou E, Hadjidakis DJ, Kollias A, *et al.* (2009) Studies of insulin resistance in patients with clinical and subclinical hypothyroidism. *Eur J Endocrinol* **160**, 785–790.
285. Peppas M, Koliaki C, Nikolopoulos P, *et al.* (2010) Skeletal muscle insulin resistance in endocrine disease. *J Biomed Biotechnol* **2010**, 527850, (epublished 15 March 2010).
286. Alaynick WA, Way JM, Wilson SA, *et al.* (2010) ERR γ regulates cardiac, gastric, and renal potassium homeostasis. *Mol Endocrinol* **24**, 299–309.
287. Konturek SJ, Konturek JW, Pawlik T, *et al.* (2004) Brain–gut axis and its role in the control of food intake. *J Physiol Pharmacol* **55**, 137–154.
288. Stanley S, Pinto S, Segal J, *et al.* (2010) Identification of neuronal subpopulations that project from hypothalamus to both liver and adipose tissue polysynaptically. *Proc Natl Acad Sci U S A* **107**, 7024–7029.
289. Suzuki K, Simpson KE, Minnion JS, *et al.* (2010) The role of gut hormones and the hypothalamus in appetite regulation. *Endocr J* **57**, 359–372.
290. Lazar MA (1993) Thyroid hormone receptors: multiple forms, multiple possibilities. *Endocr Rev* **14**, 184–193.
291. Oppenheimer JH & Schwartz HL (1997) Molecular basis of thyroid hormone-dependent brain development. *Endocr Rev* **18**, 462–475.
292. Oppenheimer JH, Schwartz HL, Lane JT, *et al.* (1991) Functional relationship of thyroid hormone-induced lipogenesis, lipolysis, and thermogenesis in the rat. *J Clin Invest* **87**, 125–132.
293. Farsetti A, Mitsuhashi T, Desvergne B, *et al.* (1991) Molecular basis of thyroid hormone regulation of myelin basic protein gene expression in rodent brain. *J Biol Chem* **266**, 23 226–23 232.
294. Vasudevan N, Ogawa S & Pfaff D (2002) Estrogen and thyroid hormone receptor interactions: physiological flexibility by molecular specificity. *Physiol Rev* **82**, 923–944.
295. Bauer M & Whybrow PC (2001) Thyroid hormone, neural tissue and mood modulation. *World J Biol Psychiatry* **2**, 59–69.
296. Bauer M & Whybrow PC (2003) Thyroid hormone and mood modulation: new insights from functional brain imaging techniques. *Curr Psychiatry Rep* **5**, 163–165.
297. Scott RE, Wu-Peng XS, Yen PM, *et al.* (1997) Interactions of estrogen- and thyroid hormone receptors on a progesterone receptor estrogen response element (ERE) sequence: a comparison with the vitellogenin A2 consensus ERE. *Mol Endocrinol* **11**, 1581–1592.
298. Schwartz HL, Lazar MA & Oppenheimer JH (1994) Widespread distribution of immunoreactive thyroid hormone β 2 receptor (TR β 2) in the nuclei of extrapituitary rat tissues. *J Biol Chem* **269**, 24777–24782.
299. Vasudevan N, Koibuchi N, Chin WW, *et al.* (2001) Differential crosstalk between estrogen receptor ER α and ER β and the thyroid hormone receptor isoforms results in flexible regulation of the consensus ERE. *Brain Res Mol Brain Res* **95**, 9–17.
300. Vasudevan N, Davidkova G, Zhu YS, *et al.* (2001) Differential interaction of estrogen receptor and thyroid hormone receptor isoforms on the rat oxytocin receptor promoter leads to differences in transcriptional regulation. *Neuroendocrinology* **74**, 309–324.
301. Williams GR (2000) Cloning and characterization of two novel thyroid hormone receptor β isoforms. *Mol Cell Biol* **20**, 8329–8342.
302. Flamant F, Gauthier K & Samarut J (2007) Thyroid hormones signaling is getting more complex: STORMs are coming. *Mol Endocrinol* **21**, 321–333.
303. Zhu YS, Yen PM, Chin WW, *et al.* (1996) Estrogen and thyroid hormone interaction on regulation of gene expression. *Proc Natl Acad Sci U S A* **93**, 12587–12592.
304. Evans RM (1988) The steroid and hormone receptor superfamily. *Science* **240**, 889–895.
305. Mangelsdorf DJ, Thummel C, Beato M, *et al.* (1995) Overview: the nuclear receptor superfamily: the second decade. *Cell* **83**, 835–840.
306. Dellovade TL, Zhu YS, Krey L, *et al.* (1996) Thyroid hormone and estrogen interact to regulate behavior. *Proc Natl Acad Sci U S A* **93**, 12581–12586.
307. Dellovade TL, Zhu YS & Pfaff DW (1999) Thyroid hormones and estrogen affect oxytocin gene expression in hypothalamic neurons. *J Neuroendocrinol* **11**, 1–10.
308. Dellovade TL, Chan J, Vennstrom B, *et al.* (2000) The two thyroid hormone receptor genes have opposite effects on estrogen stimulated sex behaviors. *Nature Neurosci* **3**, 472–475.
309. Zhao X, Lorenc H, Stephenson H, *et al.* (2005) Thyroid hormone can increase estrogen-mediated transcription from a consensus estrogen response element in neuroblastoma cells. *Proc Natl Acad Sci U S A* **102**, 4890–4895.
310. Sonodo J, Pei L & Evans RM (2008) Nuclear receptors: decoding metabolic disease. *FEBS Lett* **582**, 2–9.
311. Jeong Y & Mangelsdorf DJ (2009) Nuclear receptor regulation of stemness and stem cell differentiation. *Exp Mol Med* **41**, 525–537.
312. Dellovade TL, Kia HK, Zhu YS, *et al.* (1999) Thyroid hormone coadministration inhibits the estrogen-stimulated

- elevation of preproenkephalin mRNA in female rat hypothalamic neurons. *Neuroendocrinology* **70**, 168–174.
313. Asarian L & Geary N (2006) Modulation of appetite by gonadal steroid hormones. *Philos Trans R Soc Lond B Biol Sci* **361**, 1251–1263.
314. Zhu YS, Dellovade T & Pfaff DW (1997) Gender-specific induction of pituitary RNA by estrogen and its modification by thyroid hormone. *J Neuroendocrinol* **9**, 395–403.
315. Morgan MA, Dellovade TL & Pfaff DW (2000) Effect of thyroid hormone administration on estrogen-induced sex behavior in female mice. *Horm Behav* **37**, 15–22.
316. Alarid ET, Preisler-Mashek MT & Solodin NM (2003) Thyroid hormone is an inhibitor of estrogen-induced degradation of estrogen receptor- α protein: estrogen-dependent proteolysis is not essential for receptor transactivation function in the pituitary. *Endocrinology* **144**, 3469–3476.
317. Adan RA, Cox JJ, Van Kats JP, *et al.* (1992) Thyroid hormone regulates the oxytocin gene. *J Biol Chem* **267**, 3771–3777.
318. la Fleur SE, Kalsbeek A, Wortel J, *et al.* (2000) Polysynaptic neural pathways between the hypothalamus, including the suprachiasmatic nucleus, and the liver. *Brain Res* **87**, 50–56.
319. Buijs RM, la Fleur SE, Wortel J, *et al.* (2003) The suprachiasmatic nucleus balances sympathetic and parasympathetic output to peripheral organs through separate preautonomic neurons. *J Comp Neurol* **464**, 36–48.
320. Bradley DJ, Young WS III & Weinberger C (1989) Differential expression of α and β thyroid hormone receptor genes in rat brain and pituitary. *Proc Natl Acad Sci U S A* **86**, 7250–7254.
321. DuSell CD, Umetani M, Shaul PW, *et al.* (2008) 27-Hydroxycholesterol is an endogenous selective estrogen receptor modulator. *Mol Endocrinol* **22**, 65–77.
322. Rangwala SM, Wang X, Calvo JA, *et al.* (2010) Estrogen-related receptor γ is a key regulator of muscle mitochondrial activity and oxidative capacity. *J Biol Chem* **285**, 22619–22629.
323. Davis PJ, Leonard JL & Davis FB (2008) Mechanisms of non-genomic actions of thyroid hormone. *Front Neuroendocrinol* **29**, 211–218.
324. Glass CK, Holloway JM, Devary OV, *et al.* (1988) The thyroid hormone receptor binds with opposite transcriptional effects to a common sequence motif in thyroid hormone and estrogen response elements. *Cell* **54**, 313–323.