

# Videodensitometric time-density curve change after alcohol septal ablation of obstructive hypertrophic cardiomyopathy

A. Nemes · A. Kalapos · V. Sasi · T. Ungi · I. Ungi ·  
T. Forster · R. Sepp

© Springer Media / Bohn Stafleu van Loghum 2012


**Keywords** Blush · Obstructive hypertrophic cardiomyopathy · Percutaneous transluminal septal myocardial ablation · Videodensitometry

A 35-year-old female patient with obstructive hypertrophic cardiomyopathy (HOCM), causing New York Heart Association Class III symptoms, underwent alcohol septal ablation [1, 2]. Before and after ablation, coronary angiograms were recorded on the left anterior descending artery, and subsequently phase-matched digital subtraction angiograms have been performed on the recordings off-line. The recently

developed computerized method for estimation of myocardial perfusion, based on the analysis of the time-density curves (TDC), was used to assess myocardial blush over a selected myocardial region of interest representing the area supplied by the ablated septal branch [3–5]. The ratio of  $G_{\max}$  (defined as maximal amplitude of the TDC) and  $T_{\max}$  (defined as the time to reach  $G_{\max}$ ) was reduced after alcohol ablation compared with pre-procedural levels indicating reduced perfusion in the ablated septal area. This new method may allow evaluation of myocardial perfusion parameters, and may assist to judge the success of alcohol ablation in hypertrophic cardiomyopathy (Fig. 1). **Q3**

A. Nemes · A. Kalapos · V. Sasi · T. Ungi · I. Ungi · T. Forster ·  
R. Sepp  
Division of Invasive Cardiology, Department of Cardiology,  
Medical Faculty, Albert Szent-Györgyi Clinical Center, University  
of Szeged, Szeged, Hungary

A. Nemes (✉)  
2nd Department of Medicine and Cardiology Center, Medical  
Faculty, Albert Szent-Györgyi Clinical Center, University of  
Szeged, Korányi fasor 6, PO Box 427, 6720, Szeged, Hungary  
e-mail: [nemes@in2nd.szote.u-szeged.hu](mailto:nemes@in2nd.szote.u-szeged.hu)



**Fig. 1** Digital subtraction angiographic and videodensitometric changes after alcohol ablation in obstructive hypertrophic cardiomyopathy. Insert 1. Coronary angiographic images of the left anterior descending coronary artery shown from the cranial-right anterior oblique orientation in an HOCM patient before (1A) and after (1B) alcohol ablation. *Arrow* indicates a sub-branch of the first septal perforator artery which was ablated during the procedure. The absence of the branch is evident on the post-procedure angiogram (1B). Insert 2. Digital subtraction angiographic images shown from the same orientation as Insert 1, before (2A) and after (2B) alcohol ablation. The region of interest (ROI), *encircled in red*, represents the area supplied by the ablated branch. Myocardial blush is represented by *greyish* opacification of the microvasculature, which is seemingly reduced after ablation (2B). Insert 3. Time-density curves based on videodensitometric evaluation of density in the region of interest from Insert 2 indicating myocardial perfusion before (3A) and after (3B) alcohol ablation. The rise and fall of density (y axis) as a function of time (x axis) is represented by a *green curve* (time-density curve, TDC). The maximal amplitude of TDC is defined as  $G_{max}$ , while the time to reach  $G_{max}$  is defined as  $T_{max}$ . Both values were automatically computed and their ratio ( $G_{max}/T_{max}$ ) was used as a parameter of myocardial perfusion changes. The ratio of  $G_{max}/T_{max}$  was reduced after alcohol ablation compared with pre-procedural levels (3.21 vs. 1.33) indicating reduced perfusion in the ablated septal area

**Acknowledgments** We gratefully acknowledge the skilled assistance of the nursing staff of Cardiac Catheterization Unit (Head: Imre Ungi, MD, PhD). Dr. Attila Nemes holds a János Bolyai Research Fellowship (Budapest, Hungary).

**References**

- Gersh BJ, Maron BJ, Bonow RO, et al. 2011 ACCF/AHA guideline for the diagnosis and treatment of hypertrophic cardiomyopathy: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol.* 2011;58:e212–60.
- Seggewiss H, Rigopoulos A, Welge D, et al. Long-term follow-up after percutaneous septal ablation in hypertrophic obstructive cardiomyopathy. *Clin Res Cardiol.* 2007;96:856–63.
- Ungi T, Ungi I, Jónás Z, et al. Myocardium selective densitometric perfusion assessment after acute myocardial infarction. *Cardiovasc Revasc Med.* 2009;10:49–54.
- Ungi T, Zimmermann Z, Balázs E, et al. Vessel masking improves densitometric myocardial perfusion assessment. *Int J Cardiovasc Imaging.* 2009;25:229–36.
- Nagy FT, Sasi V, Ungi T, et al. Correlations between myocardium selective videodensitometric perfusion parameters and corrected TIMI frame count in patients with normal epicardial coronary arteries. *Int J Cardiol.* 2012;155:498–501.