

Extended Abstract, 22<sup>nd</sup> EM Induction Workshop  
Weimar, Germany, August 24-30, 2014

---

## Theoretical electromagnetic impedance as function of ionospheric pulsation source geometry

Istvan Lempenger<sup>1</sup>, Viktor Wesztergom<sup>1</sup> and Michel Menvielle<sup>2</sup> and László Szarka<sup>1</sup> and Pál Bencze<sup>1</sup> and Attila Novak<sup>1</sup> and Judit Szendrői<sup>1</sup> and Arpad Kis<sup>1</sup> and Ahmed Lethy<sup>3</sup>

<sup>1</sup>MTA CSFK GGI, Sopron, HUNGARY, lempi@ggki.hu

<sup>2</sup>ISGI/LATMOS, Saint Maur des Fosses, FRANCE, michel.menvielle@latmos.ipsl.fr

<sup>3</sup>NRIAG, Helwan, EGYPT, ahmedellethy@gmail.com

---

### SUMMARY

The presentation introduces a theoretical study which aims to model electromagnetic impedance tensor variation due to varying ionospheric pulsation source current system geometry.

**Keywords:** source effect, pulsation current system

---

### ABSTRACT

In case of practical MT exploration the surface EM impedance is computed based on a simplification related to the nature of the source of the surface EM signals. Namely the surface fields are assumed to vary like plane waves. Well known fact that this assumption holds only in certain latitude range but lapses in equatorial and auroral latitudes. It neither stands on mid-latitudes at long periods, namely above hundreds of seconds.

In a theoretical approach, analytic computation of surface EM impedance has been performed in case of a more general source geometry. Realistic pulsation ionospheric current system model and horizontal layered media have been assumed throughout the works. The applied current system model is a temporally and spatially varying amplitude, 2D current density vector filed in the height of 120km.

The subsurface model is the simplified geological model of the site of the Széchenyi István Geophysical Observatory at Nagycenk, Hungary. The presentation provides a brief summary of the results of the analytic calculations of the source effect. The influence of the variation of the meridional and azimuthal extent, the wavelength of the source and the frequency has been estimated above a physically realistic domain of the geometrical parameter-space.

### ACKNOWLEDGEMENTS

This study was supported by the TAMOP-4.2.2.C-11/1/KONV-2012-0015 (Earth-system) project sponsored by the EU and European Social Foundation.