

Geophysical Research Abstracts
Vol. 16, EGU2014-12381, 2014
EGU General Assembly 2014
© Author(s) 2014. CC Attribution 3.0 License.



Investigating electron precipitation events using near-conjugate observations from St Johns, New Foundland, and Halley, Antarctica.

Rachael Hardman (1), Mark Clilverd (1), Craig Rodger (2), Janos Lichtenberger (3), and Donald Danskin (4)

(1) United Kingdom (rachar@bas.ac.uk), (2) Department of Physics, University of Otago, Dunedin, New Zealand, (3) Space Research Group, Department of Geophysics and Space Sciences, Eötvös University, Budapest, Hungary., (4) Geomagnetic Laboratory, Natural Resources Canada, Ottawa, Canada.

In this study we use a narrow-band radio receiver recently installed at St Johns, New Foundland, Canada, as part of the AARDDVARK network. The aerial is located close to the site of the first successful trans-Atlantic subionospheric wireless radio signal reception made by Marconi in 1901. The AARDDVARK receiver can detect near-by perturbations in subionospheric radio wave propagation conditions caused by enhanced ionization in the altitude range 50-90 km. We present the results of an investigation of electron precipitation events during 2012 and 2013 using near-conjugate observations with Halley, Antarctica. Using the signal received from the LF transmitter located in Iceland (id: NRK, 37.5kHz) we detect changes in amplitude caused by electron precipitation events that can be linked to observations made at Halley via widebeam riometer absorption, or VLF wave occurrence, or the presence of electron magnetic ion-cyclotron (EMIC) waves.