

Strong Field Molecular Ionization to Multiple Ionic States: Direct vs Indirect Pathways

Péter Sándor¹, Arthur Zhao¹, Tamás Rozgonyi², Thomas Weinacht¹

¹ Department of Physics and Astronomy, Stony Brook University, 100 Nicolls Road, Stony Brook, NY 11794-3800, USA

² Institute of Materials and Environmental Chemistry, Research Centre for Natural Sciences, Hungarian Academy of Sciences, Budapest 1025, Pusztaszeri út 59-67, Hungary

Abstract. We present velocity map imaging measurements of photoelectrons in coincidence with ions produced via strong field molecular ionization. Our measurements, in conjunction with electronic structure and Stark shift calculations, allow us to assign several features in the low energy portion of the photoelectron spectrum to different molecular electronic continua (ionic states). Furthermore, we are able to distinguish between direct and indirect ionization pathways, uncovering the role of both neutral and ionic resonances in the ionization dynamics.

1. Introduction

Strong Field Ionization plays a key role in high harmonic generation and attosecond electron dynamics (Kling & Vrakking 2008, Agostini & DiMauro 2004). It has also been used as a probe of excited state molecular dynamics (Li et al. 2010). While early works suggested that many experiments could be interpreted in terms of simply removing the most weakly bound electron (i.e. from the Highest Occupied Molecular Orbital - HOMO), recent experiments (as well as one earlier work - (Gibson et al. 1991)) have highlighted the role of ionization from multiple orbitals (Li et al. 2008, McFarland et al. 2008, Spanner et al. 2012, Kotur et al. 2012). Understanding what states of the molecular cation are populated, and how, is very important for producing and probing electron wave packets, as well as for understanding strong field ionization as a tool for molecular imaging and following neutral dynamics. Here we build upon a recent work which established direct ionization to multiple states of the molecular cation via coincidence detection of electrons and ions (Boguslavskiy et al. 2012). We make use of coincidence detection of photoelectrons and ions with velocity map imaging of the photoelectrons (Lehmann et al. 2012). Furthermore, by combining our coincidence measurements with *ab initio* electronic structure and dynamic Stark shift calculations for the molecular cations, we are able to assign peaks in the photoelectron spectrum to specific ionic states, and determine the amount of direct vs indirect ionization to each state.

2. Experiment

Our experimental apparatus consists of an amplified Ti:sapphire laser system, which produces pulses with 1 mJ pulse energy, 30 fs transform limited pulse duration (intensity FWHM), a central wavelength of 780 nm and 1 kHz repetition rate. The linearly polarized laser beam crosses the effusive molecular beam in a vacuum chamber. Here we generate charged particles, which are detected by accelerating them toward a dual stack of microchannel plates and phosphor screen with an electrostatic lens configured for velocity map imaging (VMI) of the charged particles, producing a two dimensional projection of the three dimensional charged particle velocity distribution (Eppink & Parker 1997). The hit locations on the phosphor screen are recorded and digitized by a CMOS camera. Experiments can be run in either "regular" VMI or coincidence VMI mode.

During regular velocity map imaging data acquisition, constant -950, -670 and 0 V are applied to the back, middle and front plates of the electrostatic lenses, respectively, collecting the electrons only. The number of electrons generated per laser shot is much larger than one, with the exact number depending on the experiment in question. The camera integration time is much longer than the laser repetition period (1 ms). This way, the raw images already contain $\approx 10^5$ electron hits, which results in a 2D photoelectron momentum distribution with a signal to noise ratio allowing for Abel inversion of the

image.

In coincidence velocity map imaging mode, the goal is to measure an electron and ion from the same molecule. Thus for 100% detection efficiency, one would aim to maximize the probability of ionizing one molecule per laser shot and simply discard laser shots that produced two or more ions. However, with less than 100 % detection efficiency, one has to worry about false coincidences - ions and electrons that come from separate molecules and whose partner electron or ion is not detected. Based upon a detailed analysis of the true and false coincidence rates for our detection efficiencies (see appendix I), we set the experimental parameters (laser intensity and molecular beam density) such that the number of molecules ionized per laser shot to be slightly below one - it was set to be about 0.5 for the coincidence measurements discussed in this paper. The repeller plate voltages are switched from -600 to +1000 V on the back plate, and from -420 to +700 V on the middle plate within ≈ 300 ns (10% to 90% voltage level). (The front plate stays grounded.) The switching starts immediately after an electron is detected on the MCP. After switching is done, the positively charged ion is collected. The ion collection efficiency of course strongly depends on the timing of the switching, while its detection efficiency depends on the accelerating voltage. The latter was limited by the range our high voltage switch could support (a total of 1.6 kV). The integration time of the camera is much smaller than the laser repetition period (usually 24 μ s), and synchronized with the laser pulse in such a way that once the location of the electron hit is recorded, the shutter closes. The ionic fragment, which hits the MCP microseconds later, is identified based on its time of flight. A computer algorithm calculates the coordinates of the electron hit location, and pairs it with the label of the ionic fragment. During post-processing, a 2D photoelectron momentum distribution is synthesized for each fragment from the coordinates for each electron hit. Data acquisition runs at the laser repetition rate (1 kHz); however, the rate at which coincidences occur is significantly below that (< 100 Hz). An estimate of the microchannel plate detector efficiency (as a function of particle mass and kinetic energy) for heavy (>100 a.m.u.) ions can be found in (Fraser 2002). Using equation (26) in this reference, a kinetic energy of 1 keV and an open area ratio of 55% for the MCP-s, we estimate the detection efficiencies of the cationic fragments, and use their ratio to correct the relative yields of the coincidence photoelectron spectra. For a fixed longitudinal accelerating voltage of 1 kV, the values are spread between 5 and 50 %, depending on the mass of the fragment. For electrons, using 0.6 kV for acceleration, the detection efficiency is about 50 %.

The steps for processing the data further is the same for both the regular and the coincidence VMI experiment. The 2D photoelectron momentum distributions are inverse-Abel transformed, and angularly integrated in the range of ± 18 degrees about the laser polarization axis. This value was chosen because the angular extent of the features we wished to focus on in the 2D image is within this range. Finally, the angularly integrated yield is converted to a kinetic energy distribution.

While it is difficult to directly measure the resolution of our VMI apparatus given the lack of a tunable monoenergetic source of electrons, we can estimate a lower limit

to the resolution of our VMI apparatus based upon the size of a single electron hit on the detector - the $1/e^2$ diameter of such a hit is ≈ 2 pixels. We argue that two such hits are just resolved if their center are separated by twice this width, i.e. 4 pixels. The mapping of electron velocity to camera pixel is linear (≈ 5 km/s per pixel), however, the mapping of energy is quadratic, giving nonuniform resolution across the spectrum. E.g., at 0 eV, $\Delta E \approx 1$ meV; at 1 eV, $\Delta E \approx 70$ meV, and at 2 eV, $\Delta E \approx 100$ meV.

Figures 1 and 2 show the photoelectron spectra measured in coincidence with the two most prominent ionic fragments for two different molecules from the family of halomethanes. The most prominent fragments are CH_2Br^+ in case of CH_2IBr and CH_2Cl^+ in case of CH_2BrCl . For strong field ionization of molecules with several low lying continua with comparable ionization potentials, one may expect significant structure in the low energy photoelectron spectrum. The kinetic energy of electrons ionized to the i^{th} continuum is given by:

$$K^i = nh\nu - I_p^i - U_p - E_{DSS}^i \quad (1)$$

where $h\nu$ is the energy of a single photon (typically 1.6 eV in these experiments - see discussion below), I_p^i is the ionization potential associated with the i^{th} continuum (or ionic state), U_p is the ponderomotive potential and E_{DSS}^i is the dynamic Stark-shift of the i^{th} ionic state. Based on detailed calculations/measurements of each term in this expression, we can assign each of the peaks in the coincidence spectra to a specific continuum.

3. Calculations

Excitation energies of CH_2BrI^+ at the Franck-Condon position taking into account spin-orbit coupling between electronic states are taken from (Geißler et al. 2011). The equilibrium geometry of CH_2BrCl in its ground electronic state is taken from (Rozgonyi & González 2001). The excitation energies for the five lowest excited electronic states of CH_2BrCl^+ were determined at the same level of theory as those for the other species (Geißler et al. 2011), i.e, using the multireference configuration interaction (MRCI) method implemented in the Molpro program package (Werner et al. 2012). The MRCI calculations were based on previous state-averaged complete active space self-consistent field (Roos et al. 1980) computations. The active space for both species consisted of 11 electrons on 8 active orbitals being the four non-bonding lone pairs of the halogen atoms and the C-Br and C-Cl bonding and antibonding sigma orbitals. The Douglas Kroll Hamiltonian (Douglas et al. 1974) was applied and the ANO-RCC basis sets (Widmark et al. 1990, Roos et al. 2004) were used with contractions 3s2p1d for H, 4s3p2d1f for C, 6s5p3d2f1g for Br and 5s4p2d1f for Cl atoms. The spin-orbit coupling (SOC) among the ionic states were computed using atomic mean field integrals (Heß et al. 1996). The ionization potentials for the first five ionic states for both molecules can be found in table 1. In case of both molecules, values for the ground state D_0 were taken from experimental measurements by Lago et. al. (Lago et al. 2005). While




Figure 1: Photoelectron spectra for ionization of CH_2IBr . The dashed blue line shows the spectrum measured in coincidence with CH_2Br^+ fragments, while the solid black line shows the spectrum measured in coincidence with the parent ion. Superscript on the state labels indicate the number of photons absorbed to access the specific continuum. Arrows indicate the range of peak locations allowed for the full range of ponderomotive shifts in the laser focus, whereas the shaded region highlights the expected peak locations given the arguments discussed in the text.

experimental values were not available for the excited ionic states, similar calculations for the molecule CH_2I_2 showed excellent agreement with experiment (less than 5% deviation) on the energies of the lowest four ionic states (Potts et al. 1970): the differences between calculated and measured excitation energies were less than 0.05 eV in each case.

	CH_2IBr	CH_2BrCl
D_0	9.69	10.77
D_1	10.26	11.03
D_2	10.91	11.72
D_3	11.12	11.81
D_4	13.62	14.70

Table 1: Ionization potentials in eV. Energies for D_0 are experimental values from Lago et. al., while higher lying states are calculated relative to D_0 using the MRCI method as described in the text.




Figure 2: Photoelectron spectra for ionization of CH_2BrCl . The dashed blue line shows the spectrum measured in coincidence with CH_2Cl^+ fragments, while the solid black line shows the spectrum measured in coincidence with the parent ion. Superscript on the state labels indicate the number of photons absorbed to access the specific continuum. Arrows indicate the range of peak locations allowed for the full range of ponderomotive shifts in the laser focus, whereas the shaded region highlights the expected peak locations given the arguments discussed in the text.

4. Results

The ponderomotive potential, U_p , describes the time-averaged kinetic energy of a charged particle in an electromagnetic field. It depends on the laser frequency, ω , and intensity, I , but not on any molecular parameters ($U_p = \frac{2e^2 I}{4\epsilon_0 cm_e \omega^2}$). In the short-pulse limit, the ionization potential for each electronic state is effectively increased by U_p (Freeman et al. 1987), and this has to be taken into account when making the peak assignments. We note that since not all ionization takes place at the peak intensity, there will be a distribution of ponderomotive shifts for molecules ionized at intensities around the peak intensity. However, since the ionization is a nonlinear function of intensity, most of the ionization takes place near the peak intensity (70% of the ionization yield occurs for intensities within 25% of the peak intensity), and therefore we focus on peak ponderomotive shifts in our discussion below.

We measured the peak ponderomotive shifts using the intensity dependent shifting of peaks in the photoelectron spectrum for CS_2 . We chose CS_2 because the photoelectron spectrum is much simpler than for CH_2IBr or CH_2BrCl and because we measured a linear shift in the peak locations with intensity (ponderomotive shifting), indicating that for the range of intensities used in the calibration, intermediate resonances do not play an important role in determining the peak locations. The energy difference between the




Figure 3: Photoelectron spectrum for CH_2IBr for several different laser intensities. Superscript on the state labels indicate the number of photons absorbed to access the specific continuum. Legend: laser peak intensity in TW/cm^2 (and ponderomotive potential in eV).

ground and first excited state of the cation is 2.6 eV (Potts & Fattahallah 1980), which means that for a large range of intensities it is possible to ionize only to the ground state.

Figures 1 and 2 show the photoelectron spectra for CH_2IBr and CH_2BrCl , respectively. In each figure, photoelectron spectra measured in coincidence with the parent ion (black solid line) and with the most abundant fragment (blue dashed line) can be found. Horizontal arrows indicate regions where photoelectrons coming from the lowest-lying electronic states of the ion contribute to the spectrum; the tail of each arrow indicate the appearance energy of the photoelectrons at zero field (zero ponderomotive potential), while the head of each arrow shows the same for the peak of the field (peak ponderomotive potential). The red shaded areas further highlight regions of the spectrum where the U_p is within 25% of its peak value for each state, since as noted above, most of the ionization takes place within this range of the ponderomotive shift.

5. Discussion

In assigning the peaks in the photoelectron spectrum of CH_2IBr to specific ionic states, we first note that D_4 has a significantly higher ionization potential than states D_0 to D_3 . This suggests that ionization to D_4 should be suppressed with respect to the lower states.




Figure 4: Photoelectron spectrum for CH_2BrCl for several different laser intensities. Superscript on the state labels indicate the number of photons absorbed to access the specific continuum. Legend: laser peak intensity in TW/cm^2 (and ponderomotive potential in eV).

Our measurements of the ion TOFMS are consistent with this expectation in that we measure very few fragment ions coming from ionization to D_4 , which can fragment to form I^+ and CH_2I^+ (Lago et al. 2005). Furthermore, earlier measurements performed with velocity map imaging of the ionic fragments found that CH_2Br^+ produced with kinetic energy less than 0.30 eV could be associated with dissociation on D_2 or D_3 , whereas CH_2Br^+ produced with a kinetic energy above 0.50 eV could be associated with D_4 . Measurements carried out at the same intensity as the measurements shown in this paper did not find any CH_2Br^+ with kinetic energy above 0.50 eV, consistent with the idea that there is no ionization to D_4 in the present measurements (Geißler et al. 2011). Thus, we restrict our interpretation of the spectrum to ionization to D_0 through D_3 . We also note that for both CH_2IBr and CH_2BrCl , the lowest two ionic states, D_0 and D_1 , are bound, while D_2 and D_3 are dissociative (Geißler et al. 2011, Lago et al. 2005). Finally we note that the photoelectron energy is determined at the moment of ionization, and thus transitions in the ion driven by the laser do not affect the photoelectron spectrum. Thus, we argue that peaks measured in coincidence with the parent ion must come from ionization to D_0 and D_1 . Peaks measured in coincidence with the fragment ion are more subtle, but comparison with the spectrum measured in coincidence with the parent can distinguish between different cases. Peaks in the spectrum measured in coincidence with the fragment, but which do not appear in the spectrum measured in coincidence with the parent, can be associated with direct ionization to dissociative states D_2 and D_3 ,

whereas peaks that appear in both spectra can be associated with indirect ionization to D_2 or D_3 (i.e. ionization to D_0 or D_1 followed by laser driven transitions in the ion).

Given these considerations, it is natural to assign the peaks in the CH_2BrI spectrum at 1.04 eV and ≈ 0.55 eV to D_0 and D_1 and the peak at 1.45 eV to D_2 or D_3 . We note that peak at 0.55 eV shifts with intensity (see figure 3), while the peaks at 1.04 and 1.45 eV do not. This is because the peaks at 1.04 and 1.45 eV are due to resonantly enhanced ionization (via Freeman resonances (Freeman et al. 1987, Gibson et al. 1992)), whereas the peak at 0.55 eV is not resonantly enhanced. In order to test this preliminary assignment of the peaks, we consider the energies we expect for these peaks based on the formula given above, assuming that each peak is generated near the peak intensity of the pulse and therefore experiences the peak ponderomotive shift. For this case of maximal ponderomotive shift, and considering the lowest order process that would lead to a positive photoelectron energy, we expect the peak corresponding to D_0 to be at $K = 7 \cdot 1.60 - 9.69 - 0.48 = 1.03$ eV, which compares favorably with the measured 1.04 eV. For D_2 , 7 photon ionization is not energetically allowed, and therefore we expect $K = 8 \cdot 1.60 - 10.91 - 0.48 = 1.41$ eV, which again agrees well with the experiment. For the peak assigned to D_1 , the predicted energy is $K = 7 \cdot 1.60 - 10.26 - 0.48 = 0.46$ eV. This again is in reasonable agreement with the experimental measurements, confirming our initial assignments. It is natural to look for evidence of ionization to D_3 , given the small difference in ionization potential between D_2 and D_3 . The expected location for a peak corresponding to D_3 is $K = 8 \cdot 1.60 - 11.12 - 0.48 = 1.20$ eV, at which we do see a nonzero yield, although there is not a well defined peak. Therefore it is difficult to draw a firm conclusion as to whether or not there is substantial ionization to D_3 .

Similar arguments can be made to assign the peaks in the spectrum for CH_2BrCl , taking into account that for these measurements the laser was tuned slightly to the red and thus the photon energy was 1.59 eV. Also, given the higher ionization potential of this molecule, we worked at higher intensities to get a comparable yield and thus the peak ponderomotive shift is 0.80 eV. Again, we focus our attention to the four lowest-lying states, since as in the case of CH_2IBr , D_4 is much higher in energy than the states below it. We point out that the two lowest-lying ionic states, D_0 and D_1 are not dissociative, while D_2 and D_3 are, leading mostly to the production of CH_2Cl^+ (Lago et al. 2005). Peaks corresponding to the first two of the four states mentioned are expected to be found in the spectra associated with the parent ion, as a result of direct ionization. However, we expect that some of these may also be found in the spectra of the fragment, which can be explained by post-ionization transitions of the ion from a non-dissociative to a dissociative state. Additionally, it is clear that in the spectrum of the parent, no peaks are expected to be seen corresponding to dissociative states.

In the spectrum associated with the parent ion, two narrow peaks are visible at 1.04 and 1.20 eV, which also show up in the spectrum of the fragment (see figure 2). These, based on the above arguments can be assigned to the states D_1 and D_0 , respectively. The expected kinetic energy for electrons associated with D_0 is

$K = 8 \cdot 1.59 - 10.77 - 0.80 = 1.15$ eV and for D_1 it is $K = 8 \cdot 1.59 - 11.03 - 0.80 = 0.89$ eV at the maximum ponderomotive shift. The observed appearance energies lie well within the range set between the 75% and the peak ponderomotive shift, which supports the assignment.

The spectrum associated with the fragment ion (figure 2) shows two features that are absent from the parent spectrum: a broad feature between 0.10 and 0.50 eV (centered at 0.30 eV), and a relatively narrow one centered at 0.70 eV. The peak at 0.30 eV can be associated with both D_2 and D_3 since the expected energies for the two states are close together and both fall within this broad peak. The expected energies for D_2 and D_3 are $K = 8 \cdot 1.59 - 11.72 - 0.80 = 0.20$ eV and $K = 8 \cdot 1.59 - 11.81 - 0.80 = 0.11$ eV respectively. However, the origin of the 0.70 eV peak is unclear. Looking at the intensity dependent measurements shown in figure 4, one can see that this peak and the peak at ≈ 0.30 eV in the coincidence spectrum have a common energy for a low intensity which corresponds to an intensity close to where the peak(s) appears in the spectrum. This, in conjunction with the fact that the peak is in coincidence with the fragment ion suggest that it is associated with ionization to either D_2 or D_3 . As it does not shift with intensity, it is resonantly enhanced, and the position in the spectrum corresponds to the resonant enhancement occurring at about 1/3 of the peak ponderomotive shift.

In addition to the radial distributions discussed above, the velocity map imaging measurements of the photoelectrons also provide angular distributions. While the angle-dependent yields are not the focus of the current analysis, we note that the angular distributions for the 0.30 eV and 0.70 eV peaks are the same (within the statistical uncertainty of our measurements), but different from the 1.04 eV and 1.20 eV peaks. This is consistent with the idea that the 0.30 and 0.70 eV peaks are both due to ionization to D_2/D_3 .

As noted above, resonances in both the neutral and the ion play an important role in the ionization dynamics. The intensity dependence of the photoelectron spectrum highlights the role of neutral resonances, and the comparison of photoelectron spectra in coincidence with the parent and fragment ions highlights the role of ionic resonances. In order to illustrate the role of ionic resonances and to distinguish between direct and indirect ionization to a given final ionic state, we turn to figure 5. This figure shows how the direct and the indirect ionization pathways can be associated with different peaks in the coincidence photoelectron spectra. We would like to point out that the appearance of a peak corresponding to the nondissociative state D_0 in the fragment spectrum signifies that accessing a dissociative state (most likely D_2 or D_3) took place in at least two steps (first ionization, then a transition in the ion). Hence we call this pathway 'indirect' - this is to be contrasted with the peak corresponding to D_2 , which is associated with a 'direct' multiphoton transition from S_0 (the neutral ground state) to D_2 .

As noted briefly above, the intensity dependence of the photoelectron spectra (shown in figures 3 and 4) allows us to distinguish between resonance enhanced ion-




Figure 5: Illustration of dissociative and non-dissociative pathways in the ion, and their signature in the measured photoelectron spectra associated with the parent and fragment cations of CH_2IBr . a) direct 7-photon ionization to the nondissociative continuum D_0 . b) direct 8-photon ionization to the dissociative continuum D_2 . c) 7-photon ionization to the non-dissociative continuum D_0 , followed by a post-ionization transition to a higher-lying dissociative state (D_n). Arrows point to the corresponding peaks in the spectra.

ization and non-resonant ionization. These measurements are not made in coincidence mode. The peaks which don't shift ponderomotively with intensity correspond to resonantly enhanced ionization. These are the 1.04 eV peak for D_0 and the 0.70 eV peak for D_1 in CH_2IBr ; also the 1.20 eV peak for D_0 , the 1.04 eV peak for D_1 and the 0.75 eV peak for D_2/D_3 in CH_2BrCl . However, we also point out broad features that shift to lower appearance energies as the laser peak intensity increases. One shifts from 0.50 to 0.30 eV in CH_2IBr and we assign this to a nonresonant contribution from D_1 . The other is observed in CH_2BrCl and shifts from 0.40 to 0.25 eV, and this we attribute to D_2/D_3 as was previously discussed. Based on the intensity dependent spectra, it is clear that strong field ionization in these two molecules proceeds via a mixture of resonance

enhanced and non-resonant ionization.

6. Conclusion

In conclusion, we have analyzed the velocity map imaged spectrum of photoelectrons resulting from strong field ionization in coincidence with fragment ions. Our measurements, in conjunction with electronic structure and dynamic Stark shift calculations reveal the production of multiple ionic states via strong field ionization, and show evidence for post-ionization transitions in the ion.

The authors gratefully acknowledge support from the National Science Foundation under award number 1205397 and the Hungarian National Development Agency under grant number KTIA_AIK_12-1-2012-0014.

7. Appendix I - True vs false coincidence rates

We can estimate the ratio of true and false coincidences if we assume that the number of generated electron-ion pairs (assuming one ionic fragment per electron) per laser shot exhibits a Poisson distribution with the expectation value λ and the number of occurrences k :

$$P(k, \lambda) = \frac{\lambda^k e^{-\lambda}}{k!} \quad (2)$$

If we denote electron and ion detection efficiencies by η_e and η_i , respectively, then the *true coincidence probability*, $T(\lambda)$, i.e., detecting exactly one electron with its corresponding ion, is

$$T(\lambda) = \sum_{k \geq 1} k \eta_e \eta_i (1 - \eta_e)^{k-1} (1 - \eta_i)^{k-1} \frac{\lambda^k e^{-\lambda}}{k!} \quad (3)$$

The *false coincidence probability*, $F(\lambda)$, i.e., the detected electron-ion pair does not come from the same molecule is:

$$F(\lambda) = \sum_{k \geq 2} k(k-1) \eta_e \eta_i (1 - \eta_e)^{k-1} (1 - \eta_i)^{k-1} \frac{\lambda^k e^{-\lambda}}{k!} \quad (4)$$

A desired true-to-false coincidence ratio,

$$R(\lambda) = \frac{T(\lambda)}{F(\lambda)} = \frac{1}{\lambda(1 - \eta_e)(1 - \eta_i)} \quad (5)$$

dictates the expectation value, λ , and consequently, the pulse intensity and molecular beam density that are used. In our case, we estimate $\eta_e = 0.5, \eta_i = 0.25$ and aim for $R(\lambda) \geq 5$, then we have $\lambda \approx 0.5$. In other words, by adjusting the molecular beam density and pulse intensity to obtain $\lambda \approx 0.5$, 83% of our data is true coincidence.

8. Appendix II - Dynamic Stark shift calculations

In order to estimate the dynamic Stark shift of each ionic state, E_{DSS}^i , we performed a numerical integration of the time-dependent Schrödinger equation including the molecular ground state and the five lowest ionic states of the CH_2BrCl molecule. This calculation was motivated by the fact that given the various detunings between the low lying ionic states and the laser, one cannot make the rotating wave approximation or perform adiabatic elimination. The rotating wave approximation requires the detunings to be much smaller than the laser frequency and adiabatic elimination requires that the detunings be larger compared to the Rabi frequency. Neither approximation is valid for all states, leading to rapidly oscillating terms in the Hamiltonian which one would diagonalize in order to calculate dressed state energies and Stark shifts. In order to average over these rapidly oscillating terms, as is relevant for our experimental measurements, we 'probed' the ionic state energies by looking for population transfer from the ground state as a function of UV laser frequency. The coupling between the neutral ground state and the ionic states is artificial, but allows us to probe the energy shift of the ionic states as a function of ionization pulse intensity. Only five ionic states are considered because there is a substantial energy gap to the next lowest ionic states. To model the real laser field used in the experiment, which are responsible for both the Stark shift and multiphoton ionization, we defined two fields in the calculation: a strong IR laser field inducing the Stark shift and a weak VUV field ionizing the ground state. By calculating the ionic states' population as a function of IR field strength and VUV photon energy, we can map out the Stark shift of each ionic state as a function of IR field strength.

The transition dipole moments (TDMs) between ionic states, state energies and spin-orbit couplings for the ions are based on the *ab initio* electronic structure calculations described in the calculation section, while the TDMs between the neutral and ionic states were set to 0.1 a.u. The total electronic Hamiltonian (nuclear dynamics is not considered here) consists of 3 parts, the bare Hamiltonian H_0 (not including spin-orbit coupling), spin-orbit coupling H_{SO} , and the molecule-field dipole-coupling H_{MF} :

$$H = H_0 + H_{SO} + H_{MF} \quad (6)$$

$$H_{MF} = -\vec{\mu} \cdot \vec{E} \quad (7)$$

$$(H_0 + H_{SO}) |\phi_i\rangle = \hbar\omega_i |\phi_i\rangle \quad (8)$$

The calculation is carried out in the eigenspace of $H_0 + H_{SO}$ (i.e. in the spin-orbit adiabatic basis). The wave function is written in terms of the eigenstates:

$$|\psi(t)\rangle = \tilde{a}_0(t) |\phi_0\rangle + \sum_{i \neq 0} \tilde{a}_i(t) |\phi_i\rangle \quad (9)$$




Figure 6: Calculations aimed at estimating dynamic Stark shifts for ionic states of CH_2BrCl in a strong field IR laser pulse. The graph shows population of the first five ionic states as a function of IR field strength and VUV photon energy. On the horizontal axis the frequency of the VUV radiation is plotted, while on the vertical, it is the peak intensity of the IR pulse.

Substituting (6) and (9) into the Schrödinger equation $i\hbar\frac{\partial}{\partial t}|\psi\rangle = H|\psi\rangle$ and transforming to the rotating frame $\tilde{a}_i(t) = a_i(t)e^{-i\omega_i t}$, we arrive at :

$$\dot{a}_0(t) = \frac{i}{\hbar} \sum_{i \neq 0} \mu_{0i} [\mathcal{E}_{VUV}(t)e^{i\omega_{VUV}t} + \mathcal{E}_{IR}(t)(e^{-i\omega_{IR}t} + e^{i\omega_{IR}t})] a_i(t) e^{-i\omega_{i0}t} \quad (10)$$

$$\begin{aligned} \dot{a}_{i \neq 0} = & \frac{i}{\hbar} \mu_{i0} [\mathcal{E}_{VUV}(t)e^{-i\omega_{VUV}t} + \mathcal{E}_{IR}(t)(e^{-i\omega_{IR}t} + e^{i\omega_{IR}t})] a_0(t) e^{-i\omega_{i0}t} \quad (11) \\ & + \frac{i}{\hbar} \sum_{j \neq i} \mu_{ij} \mathcal{E}_{IR}(t)(e^{-i\omega_{IR}t} + e^{i\omega_{IR}t}) a_j(t) e^{-i\omega_{ji}t} \end{aligned}$$

$$\omega_{ij} = \omega_i - \omega_j \quad i, j = 1..10$$

$$\mathcal{E}_{VUV}(t) = E_{VUV} e^{-\frac{t^2}{2\tau^2}} \quad \mathcal{E}_{IR}(t) = E_{IR} e^{-\frac{t^2}{2\tau^2}}$$

We have omitted the rapidly rotating terms, $\mathcal{E}(t)e^{\pm i(\omega_{VUV} + \omega_{i0})t}$, for $\omega_{VUV} \approx \omega_{i0}$, invoking the rotating wave approximation only for the VUV field, but not for the IR. Consequently, no simple analytical solution can be obtained. In figure 6 we show the result of a calculation for CH_2BrCl . The two-dimensional plot shows the total ionic population for a range of VUV photon energies (horizontal axis) and IR peak intensities (vertical axis). The calculation shows that population can be transferred to the lowest four ionic states when the VUV pulse is resonant with the energy difference between the neutral ground state and each of the ionic states, with energies given in table 1. As the IR field strength increases, it couples the ionic states, leading to dynamic Stark

shifts, as seen in shifts of the absorption peaks on the graph. We note that simulations were performed with the IR electric field polarized along the C-Br bond. It is along this direction that the molecule-field coupling is the largest. Nevertheless, the shifts of the absorption peaks stay below 100 meV for the intensities used in the measurements (see figure 6). Similar calculations for CH₂I Br show similar Stark shifts. Since the other terms in equation 1 are significantly larger, we argue that neglecting E_{DSS} is a good approximation.

- Agostini P & DiMauro L 2004 *Reports on Progress in Physics* **67**, 813.
- Boguslavskiy A E, Mikosch J, Gijsbertsen A, Spanner M, Patchkovskii S, Gador N, Vrakking M J J & Stolow A 2012 *Science* **335**(6074), 1336–1340.
- Douglas M & Kroll N.M 1974 *Annals of Physics* **81**, pp. 89-155
- Eppink A T J B & Parker D H 1997 *Review of Scientific Instruments* **68**(9), 3477–3484.
- Fraser G 2002 *International Journal of Mass Spectrometry* **215**(13), 13 – 30. Detectors and the Measurement of Mass Spectra.
- Freeman R R, Bucksbaum P H, Milchberg H, Darack S, Schumacher D & Geusic M E 1987 *Phys. Rev. Lett.* **59**, 1092–1095.
- Geißler D, Rozgonyi T, González-Vázquez J, González L, Marquetand P & Weinacht T C 2011 *Phys. Rev. A* **84**, 053422.
- Gibson G N, Freeman R R & McIlrath T J 1991 *Physical Review Letters* **67**(10), 1230–1233.
- Gibson G N, Freeman R R & McIlrath T J 1992 *Physical Review Letters* **69**, 1904–1907.
- Heß B.A, Marian C.M, Wahlgren U & Gropen O 1996 *Chem. Phys. Lett.* **251**, pp. 365-371
- Kling M & Vrakking M 2008 *Annu. Rev. Phys. Chem.* **59**, 463–492.
- Kotur M, Zhou C, Matsika S, Patchkovskii S, Spanner M & Weinacht T C 2012 *Phys. Rev. Lett.* **109**, 203007.
- Krausz F & Ivanov M 2009 *Rev. Mod. Phys.* **81**, 163–234.
- Lago A F, Kercher J P, Bödi A, Sztáray B, Miller B, Wurzelmann D & Baer T 2005 *The Journal of Physical Chemistry A* **109**(9), 1802–1809.
- Lehmann C S, Ram N B & Janssen M H M 2012 *Review of Scientific Instruments* **83**(9).
- Li W, Jaroń-Becker A, Hogle C, Sharma V, Zhou X, Becker A, Kapteyn H & Murnane M 2010 *Proceedings of the National Academy of Sciences* **107**(47), 20219.
- Li W, Zhou X, Lock R, Patchkovskii S, Stolow A, Kapteyn H C & Murnane M M 2008 *Science* **322**(5905), 1207–1211.
- McFarland B K, Farrell J P, Bucksbaum P H & Gühr M 2008 *Science* **322**(5905), 1232–1235.
- Potts A W & Fattahallah G H 1980 *Journal of Physics B: Atomic and Molecular Physics* **13**(13), 2545.
- Potts A W, Lempka H J, Streets D G & Price W C 1970 *Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences* **268**(1184), pp. 59–76.
- Roos B.O, Taylor P.R & Siegbahn P.E.M 1980 *Chem. Phys.* **48**, pp. 157-173
- Roos R.O, Lindh R, Malmquist P-A, Vervazov V & Widmark P-O 2004 *J. Phys. Chem. A* **108**, pp. 2851-2858
- Rozgonyi T & González L 2001 *Chem. Phys. Lett.* **350**, pp. 155-164.
- Spanner M, Patchkovskii S, Zhou C, Matsika S, Kotur M & Weinacht T C 2012 *Phys. Rev. A* **86**, 053406.
- H.-J. Werner, Knowles P.J, Knizia G., Manby F.R, Schütz M & others 2012 *MOLPRO, version 2012.1, a package of ab initio programs*
- Widmark P-O, Malmquist P-A, & Roos B.O 1990 *Theoretica Chimica Acta* **77**, pp. 291-306