

# On the Fractional Derivatives at Extreme Points

Mohammed Al-Refai  
 Department of Mathematical Sciences  
 United Arab Emirates University,  
 P.O.Box 17551, Al Ain, UAE.  
 m\_alrefai@uaeu.ac.ae

## Abstract

We correct a recent result concerning the fractional derivative at extreme points. We then establish new results for the Caputo and Riemann-Liouville fractional derivatives at extreme points.

Key words and phrases: Fractional differential equations, Caputo fractional derivative, Riemann-Liouville fractional derivative.

## 1 Introduction

In recent years several authors have discussed the existence and uniqueness results for wide classes of fractional differential equations [1, 2, 3, 4, 6, 7, 9]. The techniques implemented are mainly fixed point theorems, maximum principle and the method of lower and upper solutions. In this paper we correct a result obtained in [9] and obtain new results concerning the fractional derivatives at extreme points. These results will be of interest for many researchers, especially for those who are working in extending the method of lower and upper solutions to fractional boundary value problems [1, 7]. In the following we present some definitions and main results concerning the Caputo and Riemann-Liouville fractional derivatives.

**Definition 1.1.** Let  $f \in C[0, 1]$ ,  $\delta \geq 0$ , and  $\Gamma$  is the Euler gamma function. The left Riemann-Liouville fractional integral is defined by

$$I^\delta f(t) = \begin{cases} \frac{1}{\Gamma(\delta)} \int_0^t (t-s)^{\delta-1} f(s) ds, & \delta > 0, \\ f(t), & \delta = 0. \end{cases} \quad (1.1)$$

**Definition 1.2.** Let  $f \in C^n[0, 1]$ , the left Caputo fractional derivative is defined by

$$D_C^\delta f(t) = I^{n-\delta} \frac{d^n}{dt^n} f(t) = \begin{cases} \frac{1}{\Gamma(n-\delta)} \int_0^t (t-s)^{n-\delta-1} f^{(n)}(s) ds, & n-1 < \delta < n \in \mathbb{Z}^+, \\ f^{(n)}(t), & \delta = n \in \mathbb{Z}^+. \end{cases}$$

**Definition 1.3.** Let  $f \in C^n[0, 1]$ , the left Riemann-Liouville fractional derivative is defined by

$$D_R^\delta f(t) = \frac{d^n}{dt^n} I^{n-\delta} f(t) = \begin{cases} \frac{1}{\Gamma(n-\delta)} \frac{d^n}{dt^n} \int_0^t (t-s)^{n-\delta-1} f(s) ds, & n-1 < \delta < n \in \mathbb{Z}^+, \\ f^{(n)}(t), & \delta = n \in \mathbb{Z}^+. \end{cases}$$

It is well-known that if  $f(0) = f'(0) = \dots = f^{(n-1)}(0) = 0$ , then  $D_C^\delta f(t) = D_R^\delta f(t)$ . In general the relation between the Caputo and Riemann-Liouville fractional derivatives is given by [5, 8]

$$D_C^\delta f(t) = D_R^\delta \left( f(t) - \sum_{k=0}^{n-1} \frac{t^k}{k!} f^{(k)}(0) \right), \quad (1.2)$$

where

$$D_R^\delta t^k = \frac{\Gamma(k+1)}{\Gamma(k-\delta+1)} t^{k-\delta}. \quad (1.3)$$

## 2 Main Results

We first show that the following result claimed in [9] is not correct. The following is claimed as Theorem 2.2 of [9].

• *Let a function  $f \in C^2(0,1) \cap C[0,1]$ , attain its minimum over the interval  $[0,1]$  at the point  $t_0 \in (0,1]$ . Then  $D_C^\delta f(t_0) \geq 0$ , for all  $1 < \delta \leq 2$ .*

As a counter example we consider  $f(t) = t(t - \frac{1}{2})(t - 1)$ ,  $0 \leq t \leq 1$ . Direct calculations imply that  $f(t)$  has absolute minimum value at  $t_0 = \frac{3+\sqrt{3}}{6} < 1$ . For  $1 < \delta < 2$ , we have

$$D_C^\delta t^3 = \frac{\Gamma(4)}{\Gamma(4-\delta)} t^{3-\delta}, \quad D_C^\delta t^2 = \frac{\Gamma(3)}{\Gamma(3-\delta)} t^{2-\delta} \quad \text{and} \quad D_C^\delta t = 0.$$

Thus,

$$D_C^{1.1} f(t_0) = \frac{(3 + \sqrt{3})^{1.9}}{6^{0.9} \Gamma(2.9)} - \frac{3^{0.1} (3 + \sqrt{3})^{0.9}}{2^{0.9} \Gamma(1.9)} = -0.4277 \dots < 0,$$

which contradicts the result in Theorem 2.2 of [9]. We correct the above result by imposing more conditions on  $f$ . We have

**Theorem 2.1.** *Let  $f \in C^2[0,1]$  attain its minimum at  $t_0 \in (0,1)$ , then*

$$D_C^\delta f(t_0) \geq \frac{t_0^{-\delta}}{\Gamma(2-\delta)} \left[ (\delta-1)(f(0) - f(t_0)) - t_0 f'(0) \right], \quad \text{for all } 1 < \delta < 2. \quad (2.1)$$

*Proof.* We define the auxiliary function  $h(t) = f(t) - f(t_0)$ ,  $t \in [0,1]$ . Then  $h(t)$  satisfies the following in  $[0,1]$

$$h(t) \geq 0, \quad h(t_0) = h'(t_0) = 0, \quad h''(t_0) \geq 0 \quad \text{and} \quad D_C^\delta h(t) = D_C^\delta f(t).$$

Integration by parts of

$$D_C^\delta h(t_0) = \frac{1}{\Gamma(2-\delta)} \int_0^{t_0} (t_0 - s)^{1-\delta} h''(s) ds,$$

yields

$$\Gamma(2-\delta) D_C^\delta h(t_0) = (t_0 - s)^{1-\delta} h'(s) \Big|_0^{t_0} - (\delta-1) \int_0^{t_0} (t_0 - s)^{-\delta} h'(s) ds. \quad (2.2)$$

Since  $h'(t_0) = 0$  and  $h''(t_0)$  is bounded, there exists  $\mu_1(t) \in C[0,1]$  such that  $h'(t) = (t_0 - t)\mu_1(t)$ . We have for  $1 < \delta < 2$

$$\lim_{t \rightarrow t_0} \frac{h'(t)}{(t_0 - t)^{\delta-1}} = \lim_{t \rightarrow t_0} \frac{(t_0 - t)\mu_1(t)}{(t_0 - t)^{\delta-1}} = \lim_{t \rightarrow t_0} (t_0 - t)^{2-\delta} \mu_1(t) = 0.$$

Hence

$$\Gamma(2 - \delta) D_C^\delta h(t_0) = -t_0^{1-\delta} h'(0) - (\delta - 1) \int_0^{t_0} (t_0 - s)^{-\delta} h'(s) ds. \quad (2.3)$$

Since  $h(t_0) = h'(t_0) = 0$  and  $h''(t_0)$  is bounded, there exists  $\mu_2(t) \in C[0, 1]$  such that  $h(t) = (t_0 - t)^2 \mu_2(t)$ . Thus

$$\int_0^{t_0} (t_0 - s)^{-\delta-1} h(s) ds = \int_0^{t_0} (t_0 - s)^{-\delta+1} \mu_2(s) ds,$$

is bounded and

$$\lim_{t \rightarrow t_0} \frac{h(t)}{(t_0 - t)^\delta} = \lim_{t \rightarrow t_0} \frac{(t_0 - t)^2 \mu_2(t)}{(t_0 - t)^\delta} = \lim_{t \rightarrow t_0} (t_0 - t)^{2-\delta} \mu_2(t) = 0.$$

Integrating Eq. (2.3) by parts and using the above result together with  $h(t) \geq 0$  on  $[0, 1]$  yields

$$\begin{aligned} \Gamma(2 - \delta) D_C^\delta h(t_0) &= -t_0^{1-\delta} h'(0) - (\delta - 1) \left[ (t_0 - s)^{-\delta} h(s) \Big|_0^{t_0} - \delta \int_0^{t_0} (t_0 - s)^{-\delta-1} h(s) ds \right], \\ &= -t_0^{1-\delta} h'(0) - (\delta - 1) \left[ -t_0^{-\delta} h(0) - \delta \int_0^{t_0} (t_0 - s)^{-\delta-1} h(s) ds \right] \\ &= -t_0^{1-\delta} h'(0) + (\delta - 1) t_0^{-\delta} h(0) + \delta(\delta - 1) \int_0^{t_0} (t_0 - s)^{-\delta-1} h(s) ds \\ &\geq -t_0^{1-\delta} h'(0) + (\delta - 1) t_0^{-\delta} h(0) = -t_0^{1-\delta} f'(0) + (\delta - 1) t_0^{-\delta} (f(0) - f(t_0)) \end{aligned}$$

and the result is obtained.  $\square$

**Corollary 2.1.** *Let  $f \in C^2[0, 1]$  attain its minimum at  $t_0 \in (0, 1)$ , and  $f'(0) \leq 0$ . Then  $D_C^\delta f(t_0) \geq 0$ , for all  $1 < \delta < 2$ .*

*Proof.* By Theorem 2.1 there holds  $D_C^\delta f(t_0) \geq \frac{1}{\Gamma(2-\delta)} \left[ (\delta - 1) t_0^{-\delta} (f(0) - f(t_0)) - t_0^{1-\delta} f'(0) \right]$ . Since  $f(t_0) \leq f(0)$ ,  $t_0 > 0$  and  $f'(0) \leq 0$ , we obtain  $D_C^\delta f(t_0) \geq 0$ .  $\square$

The following result is obtained as Theorem 1 of [7].

• *Let a function  $f \in W_t^1((0, T)) \cap C([0, T])$  attain its maximum over the interval  $[0, T]$  at the point  $\tau = t_0, t_0 \in (0, T)$ . Then*

$$D_C^\delta f(t_0) \geq 0, \quad 0 < \delta < 1,$$

where  $W_t^1((0, T))$  denotes the space of functions  $f \in C^1((0, T])$  such that  $f' \in L((0, T))$  and  $L((0, T))$  being the set of functions Lebesgue integrable on  $(0, T)$ .

By substituting  $g = -f$ , we have the following result.

• *Let a function  $g \in W_t^1((0, T)) \cap C([0, T])$  attain its minimum over the interval  $[0, T]$  at the point  $\tau = t_0, t_0 \in (0, T)$ . Then  $D_C^\delta g(t_0) \leq 0$ ,  $0 < \delta < 1$ .*

The following result is a simple generalization to the above one for  $t \in (0, 1)$ .

**Theorem 2.2.** *Let  $f \in C^1[0, 1]$  attain its minimum at  $t_0 \in (0, 1)$ , then*

$$D_C^\delta f(t_0) \leq \frac{t_0^{-\delta}}{\Gamma(1 - \delta)} [f(t_0) - f(0)] \leq 0, \quad \text{for all } 0 < \delta < 1. \quad (2.4)$$

*Proof.* We define the auxiliary function  $h(t) = f(t) - f(t_0), t \in [0, 1]$ . Then  $h(t) \geq 0$ , on  $[0, 1]$ ,  $h(t_0) = h'(t_0) = 0$  and  $h(t) = (t_0 - t)\mu_3(t)$  for some  $\mu_3(t) \in C[0, 1]$ . Integration by parts of

$$D_C^\delta h(t_0) = \frac{1}{\Gamma(1 - \delta)} \int_0^{t_0} (t_0 - s)^{-\delta} h'(s) ds,$$

yields

$$\Gamma(1 - \delta) D_C^\delta h(t_0) = (t_0 - s)^{-\delta} h(s) \Big|_0^{t_0} - \delta \int_0^{t_0} (t_0 - s)^{-\delta-1} h(s) ds. \quad (2.5)$$

For  $0 < \delta < 1$ , we have  $\int_0^{t_0} (t_0 - s)^{-\delta-1} h(s) ds = \int_0^{t_0} (t_0 - s)^{-\delta} \mu_3(s) ds$  is bounded and

$$\lim_{t \rightarrow t_0} \frac{h(t)}{(t_0 - t)^\delta} = \lim_{t \rightarrow t_0} (t_0 - t)^{1-\delta} \mu_3(t) = 0.$$

Thus

$$\Gamma(1 - \delta) D_C^\delta h(t_0) = -t_0^{-\delta} h(0) - \delta \int_0^{t_0} (t_0 - s)^{-\delta-1} h(s) ds \leq -t_0^{-\delta} h(0) = -t_0^{-\delta} (f(0) - f(t_0)),$$

and the result is obtained.

In the following we present analogous results concerning the Riemann-Liouville fractional derivative.

**Theorem 2.3.** *Let  $f \in C^2[0, 1]$  attain its minimum at  $t_0 \in (0, 1)$ , then*

$$D_R^\delta f(t_0) \geq \frac{t_0^{-\delta}}{\Gamma(2 - \delta)} (\delta - 1) f(t_0) \text{ for all } 1 < \delta < 2. \quad (2.6)$$

Moreover, if  $f(t) \geq 0$  in  $[0, 1]$ , then  $D_R^\delta f(t_0) \geq 0$ .

*Proof.* From Eq.'s (1.2)-(1.3) we have for  $1 < \delta < 2$

$$D_R^\delta f(t) = \frac{t^{-\delta}}{\Gamma(2 - \delta)} \left[ (1 - \delta) f(0) + t f'(0) \right] + D_C^\delta f(t).$$

Applying the result in Eq. (2.1) yields

$$\begin{aligned} D_R^\delta f(t_0) &\geq \frac{t_0^{-\delta}}{\Gamma(2 - \delta)} \left[ (1 - \delta) f(0) + t f'(0) \right] + \frac{t_0^{-\delta}}{\Gamma(2 - \delta)} \left[ (\delta - 1) (f(0) - f(t_0)) - t_0 f'(0) \right] \\ &= \frac{t_0^{-\delta}}{\Gamma(2 - \delta)} [(\delta - 1) f(t_0)]. \end{aligned}$$

If  $f(t) \geq 0$  then  $f(t_0) \geq 0$  and finally  $D_R^\delta f(t_0) \geq 0$ . □

□

**Theorem 2.4.** *Let  $f \in C^1[0, 1]$  attain its minimum at  $t_0 \in (0, 1)$ , then*

$$D_R^\delta f(t_0) \leq \frac{t_0^{-\delta}}{\Gamma(1 - \delta)} f(t_0), \text{ for all } 0 < \delta < 1. \quad (2.7)$$

Moreover, if  $f(t_0) \leq 0$ , then  $D_R^\delta f(t_0) \leq 0$ .

*Proof.* From Eq.'s (1.2)-(1.3) we have for  $0 < \delta < 1$

$$D_R^\delta f(t) = \frac{t^{-\delta}}{\Gamma(1-\delta)}f(0) + D_C^\delta f(t).$$

Using the result in Eq. (2.4) we obtain

$$D_R^\delta f(t_0) \leq \frac{t_0^{-\delta}}{\Gamma(1-\delta)}f(0) - \frac{t_0^{-\delta}}{\Gamma(1-\delta)}(f(0) - f(t_0)) = \frac{t_0^{-\delta}}{\Gamma(1-\delta)}f(t_0),$$

and  $D_R^\delta f(t_0) \leq 0$  provided  $f(t_0) \leq 0$ . □

**Remark 2.1.** *Analogous results for the fractional derivatives at absolute maximum points are obtained by applying the above results on  $-f(t)$ .*

## References

- [1] M. Al-Refai, M. Hajji, Monotone iterative sequences for nonlinear boundary value problems of fractional order, *Nonlinear Analysis Series A: Theory, Methods and Applications*, 74(2011), 3531-3539.
- [2] S. Abbas and M. Benchohra, Upper and lower solutions method for impulsive partial hyperbolic differential equations with fractional order, *Nonlinear Analysis: Hybrid Systems*, 4(2010), 406-413.
- [3] XiWang Dong, JinRong Wang, Yong Zhou, On nonlocal problems for fractional differential equations in Banach spaces, *Opuscula Mathematica*, **3**, 31(2011).
- [4] Rahmat Khan, Existence and approximation of solutions to three-point boundary value problems for fractional differential equations, *Electronic Journal of Qualitative Theory of Differential Equations*, 58(2011), 1-8.
- [5] Changpin Li, Weihua Deng, Remarks on fractional derivatives, *Applied Mathematics and Computation*, 187(2007), 777-784.
- [6] Sihua Liang, Jihui Zhang, Positive solutions for boundary value problems of nonlinear fractional differential equations, *Nonlinear Analysis*, 71(2009), 5545-5550.
- [7] Yury Luchko, Maximum principle for the generalized time-fractional diffusion equation, *Journal of Mathematical Analysis and Applications*, 351(2009), 218-223.
- [8] I. Podlubny, *Fractional Differential Equations*, Academic Press, New york, (1999).
- [9] A. Shi and S. Zhang, Upper and lower solutions method and a fractional differential equation boundary value problem, *Electronic Journal of Qualitative Theory of Differential Equations*, 30(2009), 1-13.

(Received April 11, 2012)