

Isolation and selection of *Bacillus* strains with high potential probiotic that used in catfish farming (*Pangasianodon hypophthalmus*)

Nguyen Van Minh^{1*}, Nguyen Hoang Tuan Duy¹, Do Phuong Quynh¹, Tran Thi A Ni², Tran Kien Duc³, Duong Nhat Linh¹

¹Faculty of Biotechnology, Ho Chi Minh City Open University, Vietnam

²MIDOLI Co., Ltd, Vietnam

³Ho Chi Minh City University of Science, Vietnam

*Corresponding author: minh.nv@ou.edu.vn

ARTICLE INFO

ABSTRACT

DOI:10.46223/HCMCOUJS.tech.en.10.1.355.2020

Received: August 19th, 2019

Revised: December 27th, 2019

Accepted: December 31st, 2019

Keywords:

Bacillus subtilis, probiotic, *Edwardsiella ictaluri*, *Perionyx excavatus*, *Pangasianodon hypophthalmus*

In this study, we isolated 28 strains of *Bacillus* spp. from water samples, catfish pond mud samples and earthworm manure (*Perionyx excavatus*). By the cross-streak agar methods, 22 *Bacillus* strains showed the inhibition ability to *Edwardsiella ictaluri*, which caused Bacillary Necrosis *Pangasius* (BNP) in catfish (*Pangasianodon hypophthalmus*). Both *Bacillus* sp. Q₁₆ and Q₁₁₁ strains showed the highest inhibition to *E. ictaluri* by the double-layer agar methods. Finally, two *Bacillus* strains (Q₁₆, Q₁₁₁) were selected as a source of potential probiotic because of the ability of extracellular enzyme secretion (protease, amylase, cellulase) strong growth at 0,1-1% salt concentrations, survival within the pH range 6-8, resistance to low pH and low bile salts, inability to produce haemolysin enzyme, sensitivity to eight antibiotics in the three impacting groups (inhibition of wall synthesis, inhibition mechanism of protein synthesis, inhibition of nucleic acid synthesis). Two *Bacillus* strains (Q₁₆, Q₁₁₁) were identified that they belong to *Bacillus subtilis* by biochemical method and 16S rRNA gene sequencing method. This study indicated that two *Bacillus* strains (Q₁₆, Q₁₁₁) isolated from catfish pond can be applied as high potential probiotics that used to farm catfish.

1. Introduction

In recent years, the Catfish job (*Pangasianodon hypophthalmus*) at Mekong Delta is more developed and yield is being enhanced. However, the farmers carry out intensive farming constantly to increase yield and profit, therefore, the problem of catfish disease occurs more often and more damages. One of the common diseases in catfish, bacterial disease is the most effective in catfish farming, especially purulent kidney liver disease caused by *Edwardsiella*

ictaluri (Dang & Nguyen, 2012). Although the chemicals and antibiotics cure and protect catfish, the disadvantages of them increase antibiotic-resistance bacteria and decrease the effects of antibiotics on humans and animals (Moriarty, 1997). One of the solutions to this problem is using biological products in biological control disease (Sudha, Chauhan, Dixit, Babu, & Jamil, 2010). Many pieces of research showed *Bacillus* have probiotic activity and biological control (Chao et al., 2012; Moriarty, 2006) in aquaculture. Gram-positive bacteria are used worldwide as probiotics. The wide applications belong to endospore-forming members of *Bacillus* genera (Huynh, Le, & Cutting, 2005), in which *Bacillus subtilis* is commonly used in aquaculture. Probiotics have been shown resistance to diseases, and they are excellent preventive tools against pathogens. Probiotics play an important role in creating resistance to infectious diseases and in producing antibacterial materials that prevent pathogenic bacteria from getting into organisms. Some products demonstrated the ability of probiotics in the protection for aquatic animals against pathogenic infection such as *Bacillus* spp. against *Streptococcus iniae* (Cha, Rahimnejad, Yang, Kim, & Lee, 2013). Probiotics have proven their effectiveness in improving water quality. They also enhanced the decomposition of organic matter, reduced nitrogen and phosphorus concentrations, and controlled ammonia, nitrite, and hydrogen sulfide (Cha et al., 2013; Ma, Cho, & Oh, 2009). Nguyen et al. (2013) reported that isolated bacteria from earthworms have the potential to make probiotic products and biological control some bacteria's diseases on the aquatic animals. In this study, some *Bacillus* spp. strains were isolated and selected from catfish gut, water, muddy water pond samples and from earthworm manure (*P. excavates*), they are potential to make the probiotic products that could be applied to catfish farming.

2. Materials and methods

2.1. Materials

Bacillus spp. strains (Labeled as Q) were isolated from water, catfish muddy water pond and gut samples in Thoai Son district, An Giang province and Lap Vo district, Dong Thap province. *Bacillus* spp. strains (Labeled as F) were isolated from earthworm manure (*P. excavates*), which were provided by Microbiology Lab at Ho Chi Minh City Open University. *E. ictaluri* was provided by Research Institute for Aquaculture No.2. Catfish were provided by hatchery Binh Thanh 1 of An Giang Fishers Association center.

2.2. Methods

2.2.1. Isolation *Bacillus* spp.

Bacillus spp. were isolated and purified on Nutrient Agar media (NA). Each strain was presumptive identification based on the Bergey method. The criteria of *Bacillus* spp. identification were positive bacteria, spore, positive Oxidase and positive catalase based on Bergey (1994, as cited in Holt, Krieg, Sneath, Staley, & Williams, 1994).

2.2.2. Antagonistic test to *E. ictaluri*

Each *Bacillus* spp. strain was tested the ability to resist to *E. ictaluri* by Cross-Streak method. (Lemos, Toranzo, & Barja, 1985). Double-layer agar method base on (Jock, Völksch, Mansvelt, & Geider, 2002). The result was recorded in 24, 48 and 72h and repeated 3 times.

2.2.3. Production extracellular enzyme test

This experiment was tested on some extracellular enzymes such as protease (caseinase, gelatinase), amylase, lipase, cellulase. Qualitative protease enzyme based on Montville (1983) (Montville, 1983), amylase and lipase enzymes based on Hankin and Anagnostakis (1975) (Hankin & Anagnostakis, 1975), cellulase enzyme based on Samanta, Pal, and Sem (1989) (Samanta et al., 1989).

2.2.4. Resistance saline and pH test

Bacteria trial was cultured on TSB media (Tryptic Soy Broth) in 30°C/24h, dilution with NaCl 0,85% at 10⁸CFU/mL concentration based on McFarland 0.5. In resistance saline test, 1% bacteria broth was cultured on TSB media with different NaCl concentrations: 0,1; 0,2; 0,4; 0,6; 0,8 and 1% (w/w), in 30°C/24h. In the resistance pH test, the range of pH was examined from 4 to 10. Bacteria could tolerance saline and pH when could growth in media.

2.2.5. Resistance to acid gastric test

Bacteria trial were cultured on DSM media (Difco Sporulation Medium) 37°C/48h. The bacteria broth at 10⁸CFU/mL concentration, 0,1mL bacteria broth was transferred into 10mL TSB media (Tryptic Soy Broth) at pH = 2, pH = 3 in shake cultured in 30°C, 200rpm/min. At 0h, 1h, 2h, 1mL broth was certificated cold 6000rpm/5 mins, kept and washed sediment by NaCl 0,85% and dilution to appropriate concentration in NaCl 0,85%. The sample was cultured on TSA media (Tryptic Soy Agar) in 3 different consecutive concentrations, in 37°C/24-48h. Count and analyze the living bacteria trial follow the formula: % living bacteria: $N_i/N_x \times 100$, $N_i = \log \text{CFU/mL}$ after culture time, $N_x = \log \text{CFU/mL}$ at 0h (Cukrowska et al., 2009).

2.2.6. Resistance to bile salt test

Bacteria broth was prepared at 10⁸CFU/mL concentration. 0,1mL bacteria broth was transferred into 10mL TSB media with bile salt in 0,5, 1 and 2% concentration, shake culture in 37°C, 200rpm/min. At 0h, 1h, 2h, 1mL broth sample was removed and analyze living bacteria.

2.2.7. Production hemolysin enzyme test

Isolated bacteria were cultured on BA (Blood Agar), this media was added 5% sheep blood, in 30°C/24. This experiment was conducted with control (non-hemolysin) (Van, 2006).

2.2.8. Minimum Inhibitory Concentration (MIC)

The antibiotic was dilution in MHB (Mueller Hinton Broth). Each bacterium was isolated & cultured on MHB in 37°C/24h and diluted in NaCl 0,85% at the final concentration is 10⁶ CFU/mL. The antibiotic was diluted in 2 consecutive concentration in MHB media with line concentration: 128; 64; 32; 16; 8; 4; 2; 1; 0,5; 0,25; 0,125; 0,0625, transferred into each tube 0,5mL antibiotic and 0,5mL bacteria brotha at 10⁶CFU/mL, in 37°C/16-18h. The control sample was conducted without antibiotics. The result was based on MIC Breakpoint for Bacillus genus, according to CLSI (Schwalbe, Steele, & Goodwin, 2007).

2.2.9. Identification

Selection bacteria were identified by the biochemical method Bergey (as cited in Holt, Krieg, Sneath, Staley, & Williams, 1994) and the molecular method based on 16S rRNA was conducted by Macrogen Company, Korea.

3. Results

3.1. Isolation of *Bacillus* spp.

We isolated 6 *Bacillus* spp. from earthworm manure (*P. excavates*) (Labeled as F) and 22 *Bacillus* spp. from water, catfish pond mud and gut (Labeled as Q). Its characteristics are Gram-positive, catalase-positive, rod-shaped, intracellular spore.




Figure 1. (A) Gram-stained image of *Bacillus* Q32. (B) *Bacillus* Q32 colony characteristics on NA after 24 hours of culture

3.2. The result of antagonistic

There are 22 *Bacillus* spp. Strain (F₁₁, F₂, F₂₇, F₃₃, Q₁₁, Q₁₁₁, Q₁₂, Q₁₃, Q₁₆, Q₁₆₂, Q₂₁, Q₂₃, Q₂₅, Q₂₄₀, Q₂₇₀, Q₀, Q₂₉, Q₃, Q₃₀, Q₃₂, Q₂, Q₆) that could resist to *E. ictaluri* at 24h, 48h, and 72h from 28 strain were tested by the cross-streak method and the double-layer agar method. The results were shown in Figure 2 and Figure 1. 2 strains *Bacillus* sp. Q₁₆ and Q₁₁₁ showed the highest resistance to *E. ictaluri* with the inhibition zone over 30mm. According to the research of Ho et al. (2017) showed the *B. pumilus* (47B), *B. amyloliquefaciens* (48C, 51G, 39B) and *B. megaterium* (4A, 62D) showed high inhibition activity against *E. ictaluri*, with inhibition zone over 10mm. Therefore, *Bacillus* sp. Q₁₆ and Q₁₁₁ strains were selected for the next experiment.




Figure 2. *Bacillus* strains resistance *E. ictaluri* at 24h, 48h, and 72h




Figure 3. Bacillus strains resistance *E.ictaluri*

3.3. Production extracellular enzyme test

The result showed *Bacillus* sp. Q₁₆ and Q₁₁₁ could produce extracellular enzymes such as amylase enzyme, gelatinase, caseinase, cellulase and couldn't produce lipase enzyme (Figure 4). In addition, enzyme production activity of *Bacillus* was also Joo, Hur, Han, and Kim (2007), showed that strains of *Bacillus* spp. can produce many extracellular enzymes such as amylase, cellulase, lipase.




Figure 4. Extracellular enzyme production test

(A: Gelatinase, B: Cellulase, C: Caseinase, D: Amylase)

3.4. Resistance to saline and pH test

Bacillus sp. Q₁₁₁ and Q₁₆ could resistance and strong growth in concentration saline to 1%. In the pH test, 2 strains could strong growth in pH from 6 to 8. Ho et al. (2017) showed Bacillus could growth in pH from 5 to 10.

3.5. Resistance to acid gastric test

The result of Bacillus sp. Q₁₆ and Q₁₁₁ was shown in Figure 5, Figure 6 and Figure 7.

In pH = 3 and 2, Bacillus sp. Q₁₁₁ và Q₁₆ could strong growth in 2h. Therefore, 2 strains could resistance to acid gastric. This result was similar to the research of Sudha et al. (2010) and Hyronimus, Marrec, Sassi, and Deschamps (2000) showed that Bacillus spp. could live in pH = 2, 3.




Figure 5. Living bacterial of strains test with pH = 2




Figure 6. Living bacterial of strains test with pH = 3



Figure 7. Acid gastric tolerance of *Bacillus* sp. Q16

3.6. Resistance to bile salt test

The result was shown in Figure 8, 9, 10.

The result showed *Bacillus* sp. Q₁₆ could growth in bile salt 0,5% and the living rate was decreased extremely at 1% and 2%. But *Bacillus* sp. Q₁₁₁ could growth after 2h at 0,5% and 1%. In 2% concentration bile salt after 2h culture, living rate of *Bacillus* sp. Q₁₁₁ was more than *Bacillus* sp. Q₁₆. In research of Sudha et al. (2010) showed *Bacillus coagulans* could live in media with 2% concentration bile salt.




Figure 8. Living bacterial of strains test in bile salt 0,5%




Figure 9. Living bacterial of strains test in bile salt 1%




Figure 10. Living bacterial of strains test in bile salt 2%




Figure 11. Bile salt tolerance of *Bacillus* sp. Q₁₁₁ at 2%

3.7. Hemolysin test

Bacillus sp. Q₁₆ và Q₁₁₁ didn't have hemolysin (γ) (Figure 12). This was the first step to recognize the characteristic disease, to apply safely to the human when it infected through food (Shafiqur, Shakila, Niamul, & Manjurul, 2009).




Figure 12. Hemolysin test of strains *Bacillus* sp. Q₁₆ và Q₁₁₁

3.8. Sensitive antibiotic test

Bacillus sp. Q₁₁₁ and Q₁₆ were sensitive to 8 antibiotics (penicillin, cefotaxime, vancomycin, chloramphenicol, erythromycin, gentamycin, tetracycline, ciprofloxacin) including in 3 impacting groups.

3.9. Identification

Based on Bergey (1994 as cited in Holt, Krieg, Sneath, Staley, & Williams, 1994), *Bacillus* sp. Q₁₆ and Q₁₁₁ were similar to *Bacillus subtilis* is 100% (20/20 test) by the biochemical method.

By molecular method based on sequence 16S rRNA, the sequence of 2 strains was compared to Genbank NCBI BLAST showed *Bacillus* sp. Q₁₆ and Q₁₁₁ were similar 100% with *Bacillus subtilis* and *Bacillus amyloliquefaciens*. According to Kwon et al. (2009), sequence 16S rRNA of *B. subtilis* and *B. amyloliquefaciens* couldn't distinguish. Therefore, we conducted 3 test biochemical α -D-glucosidase production, CMC hydrolysis, L-tryptophan-aminopeptidase production to distinguish *B. subtilis* and *B. amyloliquefaciens* (Cowan & Steel, 1993). The test biochemical result of *Bacillus* sp. Q₁₆ and Q₁₁₁ showed that 2 strains similar to *Bacillus subtilis* (positive α -D-glucosidase, positive CMC hydrolysis, negative L-tryptophan-aminopeptidase). Based on the biochemical method and molecular method, *Bacillus* sp. Q₁₆ and Q₁₁₁ were identified as *Bacillus subtilis*.

We have to study some important experiments such as testing of evaluation about *Bacillus* safety, the ability of *Bacillus* strains to protect catfish against *Edwardsiella ictaluri*, optimization of the media of *Bacillus* strains to applicate for manufacturing the probiotics.

4. Discussion

Many studies show that *Bacillus* was used as a biological control in aquaculture. *Bacillus* strains isolated from soil or channel catfish intestine were screened for their antagonism against *Edwardsiella ictaluri* and *Aeromonas hydrophila*, the causative agents of enteric septicemia of catfish (ESC) and Motile *Aeromonas* Septicaemia (MAS). Many studies proved that the *Bacillus* spp. can secrete antibacterial compounds belonging to the peptide, lipopeptide, and bacteriocin (Abriouel, Franz, Omar, & Gálvez, 2011). Aly, Ahmed, Ghareeb, and Mohamed (2008) showed that *B. subtilis* secretes antagonism to *A. hydrophila* and *Pseudomonas fluorescens*. Vo, Van, and Nguyen (2013) researched the probiotic products including a mixture of strains of *B. circulans* B3, *B. subtilis* N26.3, *P. acidilactici* LA61 used at a concentration of 1×10^7 CFU/g and catfish were fed in 4 weeks that could enhance the resistance and resist to purulent liver disease caused by *E. ictaluri* in catfish. According to Nguyen et al. (2013) selected two strains of *Bacillus* spp. (Q₁₆ and Q₁₁₁) were the most resistant to *E. ictaluri*, and through the safety assessment and the protection ability test for pangasius under the infective condition with *E. ictaluri* showed that both strains *Bacillus* spp. (Q₁₆ and Q₁₁₁) were safe and able to protect the host.

5. Conclusion

In this study, we selected 2 *Bacillus* spp. (Q₁₆ and Q₁₁₁) that have potential probiotic, resistance to *E. ictaluri*. Additionally, there could produce 3 extracellular enzymes (amylase, protease, cellulase) which resist to saline, pH, acid gastric, bile salt, that couldn't produce

hemolysin enzyme resistance to the antibiotic. Therefore, this study results had the potential to make/produce probiotics, applications in aquaculture.

ACKNOWLEDGMENTS

This work was funded by Ho Chi Minh Open University.

References

- Abriouel, H., Franz, C. M. A. P., Omar, N. B., & Gálvez, A. (2011). Diversity and applications of *Bacillus bacteriocins*. *FEMS Microbiology Reviews*, 35(1), 201-232.
- Aly, S. M., Ahmed, Y. A. G., Ghareeb, A. A. A., & Mohamed, M. F. (2008). Studies on *Bacillus subtilis* and *Lactobacillus acidophilus*, as potential probiotics, on the immune response and resistance of *Tilapia nilotica* (*Oreochromis niloticus*) to challenge infections. *Fish & Shellfish Immunology*, 25(1/2), 128-36. doi:10.1016/j.fsi.2008.03.013
- Barrow, G., & Feltham, R. (1993). *Cowan and Steel's manual for the identification of medical bacteria*. Cambridge, UK: Cambridge University Press.
- Cha, J. H., Rahimnejad, S., Yang, S. Y., Kim, K. W., & Lee, K. J. (2013). Evaluations of *Bacillus* spp. as dietary additives on growth performance, innate immunity and disease resistance of olive flounder (*Paralichthys olivaceus*) against *Streptococcus iniae* and as water additives. *Aquaculture*, 402-403, 50-57. doi:10.1016/j.aquaculture.2013.03.030
- Chao, R., Abel, C., Malachi, A. W., Nancy, C., Bui, C. T. D., Joseph, C. N., ... Mark, R. L. (2012). Identification of *Bacillus* strains for biological control of catfish pathogens. *PLOS One*, 7(9), 1-9.
- Cowan, S. T., & Steel, K. J. (1993). *Cowan and Steel's manual for identification of medical bacteria* (3rd ed.) (pp. 199-241). Cambridge, UK: Cambridge University Press.
- Cukrowska, B., Motyl, I., Kozáková, H., Schwarzer, M., Górecki, R. K., Klewicka, E., ... Libudzisz, Z. (2009). Probiotic *Lactobacillus* Strains: In vitro and in vivo Studies, *Folia Microbiologica*, 54(6), 533-537.
- Dang, O. T. H., & Nguyen, P. T. (2012). Trial of *Edwardsiella ictaluri* on *Pangasius hypophthalmus* by Erythromycin Thiocynate antibiotic. *Can Tho University Journal of Science*, 2012(22c), 146-154.
- Hankin, L., & Anagnostakis, S. L. (1975). The use of solid media for detection of enzyme production by fungi. *Cryptogamie, Mycologie*, 67(3), 597- 607.
- Ho, T. T. T., Nguyen, H. T., Nguyen, N. T., Ong, M. Q., Korntip, K., Sasimamas, U., & Nontawith, A. (2017). Identification and characterization of potential probiotic *Bacillus* spp. for application in striped catfish (*Pangasianodon hypophthalmus* [Sauvage, 1878]). *Journal of Fisheries and Environment*, 41(2), 20-32.
- Holt, J. G., Krieg, N. R., Sneath, P. H. A., Staley, J. T., & Williams S. T. (1994). *Bergey's manual of determinative bacteriology* (9th ed.). Philadelphia, PA: Lippincott Williams & Wilkins.
- Huynh, A. H., Le, H. D., & Cutting, S. M. (2005). The use of bacterial spore formers as probiotics. *FEMS Microbiol Reviews*, 29(4), 813-835. doi:10.1016/j.femsre.2004.12.001

- Hyronimus, B., Marrec, C. L., Sassi, A. H., & Deschamps, A. (2000). Acid and bile tolerance of spore-forming lactic acid bacteria. *International Journal of Food Microbiology*, 61(2-3), 193-197. doi:10.1016/s0168-1605(00)00366-4
- Jock, S., Völksch, B., Mansvelt, L., & Geider, K., (2002). Characterization of Bacillus strains from apple and pear trees in South Africa antagonistic to Erwinia amylovora. *FEMS Microbiology Letters*, 211(2), 247-252. doi:10.1111/j.1574-6968.2002.tb11232.x
- Joo, M., Hur, S., Han, Y., & Kim, J. (2007). Isolation, identification and characterization of Bacillus strains from the traditional Korean soybean-fermented food, Chungkookjang. *Journal of Applied Biological Chemistry*, 50(4), 202-210.
- Kwon, G. H., Lee, H. A., Park, J. Y., Kim, J. S., Park, C. S., Kwon, D. Y., ... Kim, J. H. (2009). Development of a RAPD-PCR method for identification of Bacillus species isolated from Cheonggukjang. *International Journal of Food Microbiology*, 129(3), 282-287. doi:10.1016/j.ijfoodmicro.2008.12.013
- Lemos, M. L., Toranzo, A. E., & Barja, J. L. (1985). Antibiotic activity of epiphytic bacteria isolated from intertidal seaweeds. *Microbial Ecology*, 11(2), 149-163. doi:10.1007/BF02010487
- Ma, C. W., Cho, Y. S., & Oh, K. H. (2009). Removal of pathogenic bacteria and nitrogens by Lactobacillus spp. JK-8 and JK-11. *Aquaculture*, 287(3/4), 266-270. doi:10.1016/j.aquaculture.2008.10.061
- Montville, T. J. (1983). Dual-substrate plate diffusion assay for proteases. *Applied and Environmental Microbiology*, 45(1), 200-204.
- Moriarty, D. J. W. (1997). The role of microorganisms in aquaculture ponds. *Aquaculture*, 151(1-4), 333-349.
- Moriarty, D. J. W. (2006). Safety of aquaculture probiotics. *Global Aquaculture Advocate*, 86-87.
- Nguyen, M. V., Nguyen, D. H. T., Do, Q. P., Vo, N. N. Y., Duong, L. N., Nguyen, T. T. N., & Le, P. H. (2013). Biocontrol of pathogenic Edwardsiella ictaluri by Bacillus spp. isolated from catfish pond. *Journal of Science and Technology*, 51(5C), 508-512.
- Samanta, R., Pal, D., & Sem, S. P. (1989). Production of hydrolases by N₂ fixing microorganisms. *Biochemie and Physiologie der pflanzen*, 185(1/2), 75-81.
- Sánchez-Ortiz, A. C., Luna-González, A., Campa-Córdova, A. I., Escamilla-Montes, R., Flores-Miranda, M. C., & Mazon-Suastegui, J. M. (2015). Isolation and characterization of potential probiotic bacteria from pustulosebark (Anadara tuberculosa) suitable for shrimp farming. *Latin American Journal of Aquatic Research*, 43(1), 123-136.
- Schwalbe, R., Steele, M. L., & Goodwin, A. C. (2007). *Antimicrobial susceptibility testing protocols*. Boca Raton, FL: CRC Press.
- Shafiqur, R., Shakila, N. K., Niamul, N. M., & Manjurul, K. M. (2009). Application of probiotic bacteria: A novel approach towards ensuring food safety in shrimp aquaculture. *Journal of Bangladesh Academy of Sciences*, 33(1), 139-144.

- Sudha, M. R., Chauhan, P., Dixit, K., Babu, S., & Jamil, K. (2010). Molecular typing and probiotic attributes of a new strain of *Bacillus coagulans* - Unique IS-2: A potential biotherapeutic agent. *Genetic Engineering and Biotechnology Journal GEBJ*, 7, 1-20.
- Vo, M. S., Van, T. T., & Nguyen, T. N. T. (2013). Dietary administration of the probiotic enhanced innate immune responses and disease resistance of the striped catfish against *Edwardsiella ictaluri*. *Journal of Agriculture and Rural Development*, 14, 81-89.