

ON MULTIPLE BORSUK NUMBERS IN NORMED SPACES

ZSOLT LÁNGI AND MÁRTON NASZÓDI

ABSTRACT. Hujter and Lángi defined the k -fold Borsuk number of a set S in Euclidean n -space of diameter $d > 0$ as the smallest cardinality of a family \mathcal{F} of subsets of S , of diameters strictly less than d , such that every point of S belongs to at least k members of \mathcal{F} .

We investigate whether a k -fold Borsuk covering of a set S in a finite dimensional real normed space can be extended to a completion of S . Furthermore, we determine the k -fold Borsuk number of sets in not angled normed planes, and give a partial characterization for sets in angled planes.

1. INTRODUCTION

In 1933, Borsuk [5] posed the problem whether any set S of diameter $d > 0$ in Euclidean n -space \mathbb{R}^n is the union of $n + 1$ sets of diameters less than d . A proof of the affirmative answer for $n = 2$ appeared in [5], and for $n = 3$ in [6] (for finite S , see [9], [11]). Sixty years after the problem appeared, Kahn and Kalai [14] proved that for large values of n the answer is negative. For surveys on Borsuk's problem, see [3, 18].

Boltyanski [1] gave a characterization of bounded sets according to their *Borsuk number* (that is, the least number of smaller diameter pieces that they can be partitioned into) in the Euclidean plane: *Let $\emptyset \neq S \subset \mathbb{R}^2$ be a bounded set that is not a singleton. Then the Borsuk number of S is 3 if S has a unique completion (see Definition 2.1) and 2 otherwise.*

Grünbaum [8] was the first to consider the Borsuk numbers of sets with respect to a metric distinct from the Euclidean, and determined the Borsuk numbers of sets in the plane equipped with the ℓ_∞ norm. The problem was solved for arbitrary normed planes in [4]:

Theorem 1.1 (Boltyanski-Soltan). *Let S be a compact set in the normed plane with unit ball \mathbf{B} . Then the Borsuk number of S is*

- $a(S) = 4$ if, and only if, \mathbf{B} and S are homothetic parallelograms;

1991 *Mathematics Subject Classification.* 52C17, 05C15, 52A21.

Key words and phrases. Borsuk's problem, diameter, unique completion, covering, bodies of constant width, complete, multiple chromatic number.

The authors gratefully acknowledge the support of the János Bolyai Research Scholarship of the Hungarian Academy of Sciences, and the Hung. Nat. Sci. Found. (OTKA) grants: K72537 and PD104744.

- $a(S) = 3$ if, and only if, $a(S) \neq 4$, there is a unique completion C of S with respect to \mathbf{B} , and S satisfies the supporting line property: for any pair of parallel supporting lines of C , S has a point on at least one;
- $a(S) = 2$ otherwise.

As a generalization of Borsuk's problem, Hujter and Lángi [12] defined the k -fold Borsuk number, $a_k(S)$, of a set S of diameter $d > 0$ as the smallest cardinality of a family \mathcal{F} of subsets of S , of diameter strictly less than d , such that every point of S belongs to at least k members of \mathcal{F} . Among other results, they determined the k -fold Borsuk numbers of any set in the Euclidean plane.

Motivated by Boltyanski's result, we investigate whether a (k -fold) Borsuk covering of a set S can be extended to a completion of S . Theorem 1 states that such an extension is possible in certain *Minkowski spaces* (ie. finite dimensional real normed spaces) provided that S has a unique completion. The class of these Minkowski spaces include Euclidean n -space for all n . This result has been known in the Euclidean plane [1] but is new in higher dimensional Euclidean spaces. In Theorem 2, we extend this result to not angled Minkowski planes (see Definition 3.2).

In Theorems 3, 4 and 5, we find the k -fold Borsuk numbers of sets in not angled normed planes, and of sets that cannot be completed uniquely to a Reuleaux polygon in angled planes.

2. DEFINITIONS AND NOTATIONS

We denote the closed unit ball centered at a point $x \in \mathbb{R}^n$ of a Minkowski space by $\mathbf{B}(x)$, and its boundary, the unit sphere by $\mathbf{S}(x)$. For a set A , the intersection of unit balls centered at the points of A is denoted as

$$\mathbf{B}A = \bigcap_{x \in A} \mathbf{B}(x).$$

Definition 2.1. A bounded set C in an n -dimensional Minkowski space is *complete*, if no set of the same diameter properly contains C . (Note that a complete set is clearly compact and convex.) A set S is a *set of unique completion* if there is a unique complete set C containing S of the same diameter as S .

Proposition 2.2. *Let S be a set of unit diameter in an n -dimensional Minkowski space. Then*

- S is complete if, and only if $S = \mathbf{B}S$,
- S is a set of unique completion if, and only if, $\mathbf{B}S = \mathbf{B}^2S$ ie. $\mathbf{B}S$ is complete.

The first statement is due to Eggleston [7], where it is called the *spherical intersection property*, the second is due to Moreno (Corollary 3 in [16]). Note that in the second case the completion of S is $\mathbf{B}S$.

We define the *distance of a set A* of a Minkowski space and a point x as $d_{\mathbf{B}}(x, A) = \inf\{d_{\mathbf{B}}(x, a) : a \in A\}$, where $d_{\mathbf{B}}(x, a)$ is the distance of the points a and x in the normed space with unit ball \mathbf{B} .

3. EXTENDING A BORSUK COVERING IN CERTAIN MINKOWSKI SPACES

Our goal is to extend a Borsuk covering of a closed set S of unique completion in a Minkowski space to its unique completion \mathbf{BS} . In general, a Borsuk covering of a compact set may not extend to any of its completions: consider a pair of points which in Euclidean space have many completions, all of whose Borsuk number is above two.

3.1. Extension of a Borsuk covering in certain Minkowski spaces. We define the following ‘‘Lens Cutting Condition’’ which holds in certain Minkowski spaces:

(LCC) *For any two distinct points u and v in \mathbb{R}^n with $d_{\mathbf{B}}(u, v) \leq 1$ and $x \in \mathbb{S}(u) \cap \mathbb{S}(v)$ and $\varepsilon > 0$, there is a $w \in \mathbb{R}^n$ such that $x \notin \mathbf{B}(w)$ but $\mathbf{B}(w) \supset \mathbf{B}(u) \cap \mathbf{B}(v) \setminus \varepsilon \mathbf{B}(x)$.*

Remark 3.1. It is not hard to see that (LCC) holds in all Euclidean spaces.

Theorem 1. *If (LCC) holds in a Minkowski space then any k -fold Borsuk covering of a closed set of unique completion extends to a k -fold Borsuk covering of its completion.*

Proof. We prove the Theorem for $k = 1$, the general case follows from the same argument. Let $S = Q_1 \cup \dots \cup Q_k$ be a Borsuk covering of a closed set S of unique completion by closed sets of diameter at most $r < 1$. Note that a Borsuk covering of the boundary of a set may be extended to the set in a straightforward way (cf. also Remark 4.3). Thus, we will define sets $Q'_1 \cup \dots \cup Q'_m = \text{bd } \mathbf{BS}$ that form a Borsuk covering of the boundary of the completion \mathbf{BS} of S .

For all i , Q'_i will contain $Q_i \cap \text{bd } \mathbf{BS}$ and some more points of $\text{bd } \mathbf{BS}$. For an $x \in (\text{bd } \mathbf{BS})$ we take the index i such that $d(x, Q_i)$ is minimal (if it is not unique, we take all such i), and include x into Q'_i . Clearly, Q'_i is closed.

Note that for any $x \in \mathbf{BS} \setminus S$ we have that

(*) *there are no two distinct points $u, v \in \mathbf{BS}$ with $d_{\mathbf{B}}(x, u) = d_{\mathbf{B}}(x, v) = 1$.*

Suppose the contrary. Then $S \subseteq \mathbf{B}^2 S \subseteq \mathbf{B}(u) \cap \mathbf{B}(v)$. On the other hand, $\mathbf{B}^2 S$ is the intersection of all unit balls that contain S , and hence by (LCC), $\mathbf{B}^2 S \subseteq (\mathbf{B}(u) \cap \mathbf{B}(v)) \setminus \{x\}$, contradicting $x \in \mathbf{BS} = \mathbf{B}^2 S$.

The family of the sets q'_i is a Borsuk partition of $\text{bd } \mathbf{BS}$. Indeed, let $x, y \in Q'_i$. If x or y is in S then clearly, $d(x, y) < 1$. If both are in $Q'_i \setminus S$ then, by (*), $d(x, y) < 1$. \square

3.2. Extension of a Borsuk covering in certain Minkowski planes. It is not difficult to see that a strictly convex normed plane (that is, when the unit disk \mathbf{B} is strictly convex) satisfies (LCC), and thus has the extension property of Theorem 1. Next, we consider a class of Minkowski planes that is wider than the class of strictly convex planes, and where (LCC) does not hold, but the extension property still does. The following definition is from [3] (cf. also [4]).

Definition 3.2. A normed plane with unit ball \mathbf{B} is *angled*, if for some non-collinear points a, b, c , we have $[a, b] \cup [b, c] \subset \mathbb{S}$.

Theorem 2. *Let S be a set of unique completion in a not angled normed plane, and let C be the completion of S . Then any k -fold Borsuk covering \mathcal{F} of S can be extended to a k -fold Borsuk covering of C .*

From this point on throughout this section, we assume that the Minkowski plane we work with is not angled.

The following monotonicity lemma appeared in [15].

Lemma 3.3 (Lassak). *Let $t \mapsto p(t)$ (with $t \in [0, 1]$) be a simple, closed, continuous curve, defining the boundary of a complete body of diameter one in a Minkowski plane. Let $p = p(0)$, and let t_1 and t_2 be the smallest and the largest values of t such that $\text{dist}_{\mathbf{B}}(p, p(t)) = 2$. Then the function $t \mapsto \text{dist}_{\mathbf{B}}(p, p(t))$ is*

- *strictly increasing on the interval $[0, t_1]$,*
- *equal to one on $[t_1, t_2]$, and*
- *strictly decreasing on $[t_2, 1]$.*

Corollary 3.4. *Let C be a complete body of diameter one in a Minkowski plane. Then, for any $p \in \text{bd } C$ we have the following.*

- *The set of points of C at unit distance from p is a connected arc of $\mathbb{S}(p) \cap \text{bd } C$.*
- *If $\|q - p\|_{\mathbf{B}} = \|r - p\|_{\mathbf{B}}$ for some $q, r \in \text{bd } C$, then the arc of $\text{bd } C$, connecting q and r and not containing p , belongs to the circle $\mathbb{S}(p)$.*

Lemma 3.5. *If C is a complete body in a Minkowski plane, and $[a, b]$, $[c, d]$ are two disjoint diameters of C such that a, b, c, d are in counterclockwise order in $\text{bd } C$, then $[a, d]$, $[b, c] \subset \text{bd } C$ and they are parallel.*

Proof. Consider the quadrangle $Q = \text{conv}\{a, b, c, d\}$. Observe that as $[a, b]$ and $[c, d]$ are diameters of C , neither C nor $Q \subseteq C$ contains a translate of neither $[a, b]$ nor $[c, d]$ in its interior. Thus, $[a, c]$ and $[b, d]$ are parallel, and they belong to $\text{bd } C$. \square

Lemma 3.6 is a straightforward consequence of Theorems 33.7 and 33.9 of [3].

Lemma 3.6. *Let S be a compact set of unique completion in a not angled normed plane, and let C be its completion. Then, for any parallel supporting lines L and L' of C , L or L' contains a point of S . In other words, S satisfies the supporting line property (see page 2).*

Lemma 3.7. *Let S be a compact set of diameter one and of unique completion, C . Then, for any point $x \in (\text{bd } C) \setminus S$, there is an open circle arc of radius one, containing p and being contained in $\text{bd } C$, such that its endpoints and its center belong to S .*

Proof. Let $x \in (\text{bd } C) \setminus S$. Then, since $\mathbf{B}S = C$, there is a point $p \in S$ such that $x \in \mathbb{S}(p)$. Clearly, $[p, x]$ is a diameter of C , and thus, $p \in \text{bd } C$. Let L and L' be a pair of parallel supporting lines of C such that $x \in L$ and $p \in L'$. For simplicity, we imagine these lines as vertical such that L is to the left of L' . Let $[a, b] = C \cap L$ and $[c, d] = C \cap L'$, and note that these segments might be degenerate. Without loss of generality, we assume that a, b, c and d are in this counterclockwise order in $\text{bd } C$.

First, we show that at least one of a and c belongs to S . Indeed, consider a sequence of supporting lines L_m of C , with positive slopes, such that the limit of $L_m \cap C$ is $\{a\}$. For any m , let L'_m be the supporting line of C , parallel to and different from L_m . Clearly, the limit of $L'_m \cap C$ is $\{c\}$. Now, by Lemma 3.6, we have that for any m , L_m or L'_m contains a point of S . Thus, the observation follows from the compactness of S . We may show similarly that at least one of b and d belongs to S .

Now we prove the assertion. If both $[a, x]$ and $[x, b]$ contain a point of S , then we may observe that $[a, b] \subset \mathbb{S}(p)$, and thus, our lemma follows. Assume that exactly one of these segments, say $[x, b]$, contains a point of S . Then $a \notin S$, and thus, $c \in S$. Let G be the arc of $(\text{bd } C) \cap \mathbb{S}(c)$, starting at x and above the line connecting x and c . If G does not contain a point of S , then for some point $c' \in L' \setminus [c, d]$, we have $S \subset \mathbf{B}(c')$; or in other words, $c' \in \mathbf{B}S = C$; a contradiction. Thus, G contains a point of S , which yields the assertion.

We are left with the case that $[a, b] \cap S = \emptyset$, which, in particular, implies that $c, d \in S$. Note that if $c \neq d$, then, by Lemma 3.5, for any $y \in \text{relint}[c, d]$, we have $C \cap \mathbb{S}(y) = [a, b]$. Thus, moving y slightly to the right, we can find a point y' such that $S \subset \mathbf{B}(y')$, but $[a, b] \cap \mathbf{B}(y') = \emptyset$. This yields that $C \not\subset \mathbf{B}(y')$, or in other words that $y' \notin \mathbf{B}C = C$, contradicting $y' \in \mathbf{B}S = C$. Thus, we obtain that $c = d$. In this case, similarly like in the previous paragraph, one can show that both arcs of $(\text{bd } C) \cap \mathbf{B}(c)$, starting at x , contain a point of S , and the assertion readily follows. \square

Proof of Theorem 2. Note that it suffices to extend \mathcal{F} to a k -fold Borsuk covering of $\text{bd } C$.

Let $\mathcal{F} = \{Q_1, Q_2, \dots, Q_m\}$ be a k -fold Borsuk covering of S . Without loss of generality, we may assume that S is compact. Let ε be chosen in such a way that the diameter of every member of \mathcal{F} is at most $1 - 3\varepsilon$. Now, for every i , we set $Q_i^* = Q_i + \varepsilon B$, and observe that $\mathcal{F}^* = \{Q_1^*, Q_2^*, \dots, Q_m^*\}$ is still a k -fold Borsuk covering of S .

Consider the connected components of $\text{bd } C \setminus S$. By Lemma 3.7, they are open circle arcs of unit radius, with their centers contained in S . Note that \mathcal{F}^* is a k -fold covering of any such arc not longer than 2ε . Since $\text{bd } C$ has a bounded length, there are only finitely many arcs that are not covered k -fold by \mathcal{F}^* . Thus, by induction, it suffices to prove that \mathcal{F}^* can be extended to cover k -fold at least one such arc.

Consider an arc G that is not covered by \mathcal{F}^* k -fold. Let $p \in S$ denote the center, and $q, r \in S$ denote the endpoints of G . If, for every $x \in G$, p is the only point of C at unit distance from x , then we can apply the argument in the proof of Theorem 1. Thus, assume that for some $x \in G$ and $p' \in C$ with $p \neq p'$, we have $x \in \mathbb{S}(p')$, where without loss of generality, we may assume that, say, $[p, r]$ and $[p', x]$ are disjoint. Note that since $[p, r]$ and $[p', x]$ are diameters of C , we have that $[p, p']$ and $[x, r]$ are parallel, and are contained in $\text{bd } C$.

Let L and L' be the line containing $[r, x]$ and $[p, p']$, respectively. Observe that the points diametrically opposite to any point in the relative interior of $L \cap C$ are the points of $L' \cap C$. Let y be the endpoint of $L \cap \text{bd } C$ closer to x than to r . If $q \in L$, then

we may add the segment $[q, x]$ to any Q_i^* containing q , and $[x, r]$ to any Q_i^* containing r . Thus, we may assume that y is a point of G .

Consider the case that the points diametrically opposite to y are only the points of $L' \cap C$. Then we may add the segment $[y, r]$ to any Q_i^* containing r . On the other hand, note that if some $u \in \text{bd } C$ is diametrically opposite to any point of the arc between y and q , then it is diametrically opposite to q as well. Thus, we may add this arc to any Q_i^* containing q .




FIGURE 1. An illustration for the proof of Theorem 2

Finally, assume that there is some point $p'' \notin L' \cap C$ that is diametrically opposite to y (cf. Figure 1). Then, clearly, the points p', p, p'', y are in this cyclic order in $\text{bd } C$, and $[y, p'']$ and $[q, p]$ are disjoint diameters of C , which yields, by Lemma 3.5, that $[p, p'']$ and $[q, y]$ are parallel, and both are contained in $\text{bd } C$. Thus, $\text{bd } C$, and also $\mathbb{S}(p)$, contains an angle, which contradicts the conditions of the theorem. \square

Corollary 3.8. *Let S be a set of unique completion in a not angled normed plane, and let C be the completion of S . Then for any value of k , $a_k(S) = a_k(C)$.*

4. THE MULTIPLE BORSUK NUMBERS OF SETS IN A NOT ANGLED NORMED PLANE

We start with three observations, which, for sets in a Euclidean space, appeared as Remarks 1–3 in [12]. Their proofs are straightforward modifications of those in [12], and hence we omit them.

Remark 4.1. The sequence $a_k(S)$ is sub-additive for every set S in any normed (or metric) space. More precisely, for any positive integers k, l , we have $a_{k+l}(S) \leq a_k(S) + a_l(S)$.

Remark 4.2. Let S be a set of diameter $d > 0$ in a normed (or metric) space. Then for every set S of diameter $d > 0$ and every $k \geq 1$, we have $a_k(S) \geq 2k$. Furthermore, for every value of k , if $a(S) = 2$, then $a_k(S) = 2k$, and if $a(S) > 2$, then $a_k(S) > 2k$.

Remark 4.3. Let $S \subset \mathbb{R}^n$ be a set of positive diameter in a normed space. Then for every value of k , $a_k(S) = a_k(\text{bd } S)$.

Let S be a bounded set in a normed plane. By Theorem 1.1, if S is not a set of unique completion then $a(S) = 2$, which yields that for any k , $a_k(S) = 2k$. Combined with Corollary 3.8, it yields that it suffices to characterize the k -fold Borsuk numbers of complete sets. To do this, we need a generalization of the notion of Reuleaux polygons for normed planes (cf. also [21], [19] and [10]).

Definition 4.4. Let C be a complete set in a normed plane. If C is the intersection of finitely many translates of \mathbf{B} , we say that C is a *Reuleaux polygon*. If m is the smallest number such that C is the intersection of m translates of \mathbf{B} , then we say that C has m sides.

Theorem 3. Let C be a complete set of diameter one in a normed plane, which is not a Reuleaux polygon. Then for every k , $a_k(C) = 2k + 1$.

Proof. Clearly, by Remark 4.2, for every k , we have $a_k(C) \geq 2k + 1$. Thus, we need to show that if C is not a Reuleaux polygon, then C , or equivalently, $\text{bd } C$, can be covered k -fold by $2k + 1$ subsets of smaller diameters.

To do this, we prove the existence of $2k + 1$ diameters $[p_i, p_{2k+1+i}]$, where $i = 1, 2, \dots, 4k + 2$ of C , such that for any $j \neq 2k + 1 + i$, $[p_i, p_j]$ is not a diameter of C . Observe that from this, the assertion follows. Indeed, by Lemma 3.3, we have that any two of these diameters intersect. Thus, we may label their endpoints in such a way that $p_1, p_2, \dots, p_{4k+2}$ are in counterclockwise order in $\text{bd } C$. Let A_i be the arc of $\text{bd } C$, connecting p_i and p_{2k+i} and not containing p_{2k+1+i} . Then A_i is of diameter less than one, and the arcs A_{1+ks} , where $s = 1, 2, \dots, 2k + 1$, form a k -fold Borsuk covering of $\text{bd } C$.

For simplicity, for any point $x \in \text{bd } C$, we set $G(x) = C \cap \mathbb{S}(x) \subset \text{bd } C$. We choose the required diameters as follows. Let $[p_1, p_{2k+2}]$ be an arbitrary diameter of C . Let q_1, r_1 and q_{2k+2}, r_{2k+2} be the endpoints of the arcs $G(p_1)$ and $G(p_{2k+2})$, respectively. It follows from Lemma 3.3 that $G(p_{2k+2}) \subseteq G(q_1) \cup G(r_1)$ and $G(p_1) \subseteq G(q_{2k+2}) \cup G(r_{2k+2})$. Then, as no finitely many unit circle arcs cover $\text{bd } C$, $X_2 = \text{bd } C \setminus (G(q_1) \cup G(r_1) \cup G(q_{2k+2}) \cup G(r_{2k+2})) \neq \emptyset$.

Observe that for any $x \in X_2$, $\|x - p_1\|_{\mathbf{B}}$ and $\|x - p_{2k+2}\|_{\mathbf{B}}$ are strictly less than one, and any point diametrically opposite to x is also contained in X_2 . Let $p_2 \in X_2$ arbitrary. Since C is complete, there is some $p_{2k+3} \in \text{bd } C$ such that $[p_2, p_{2k+3}]$ is a diameter of C . Then $p_{2k+3} \in X_2$; that is, $\|p_{2k+3} - p_1\|_{\mathbf{B}}$ and $\|p_{2k+3} - p_{2k+2}\|_{\mathbf{B}}$ are strictly less than one. Let us define $q_2, r_2, q_{2k+3}, r_{2k+3}$ similarly as for p_1 and p_{2k+2} . Now, set $X_3 = X_2 \setminus (G(q_2) \cup G(r_2) \cup G(q_{2k+3}) \cup G(r_{2k+3}))$. Since $\text{bd } C$ is not covered by finitely many unit circle arcs, we have $X_3 \neq \emptyset$. Thus, using the argument as for

$[p_2, p_{2k+3}]$, we can find a diameter $[p_3, p_{2k+4}]$ with $p_3, p_{2k+4} \in X_3$, satisfying the required conditions. Since C is not a Reuleaux polygon, repeating this procedure we may choose the required $2k + 1$ diameters for any value of k . \square

Theorem 4. *If C is an m -sided Reuleaux polygon of diameter one in the not angled norm with unit disk \mathbf{B} , then*

- (1) m is an odd integer,
- (2) if $m = 2s + 1$, then the k -fold Borsuk number of C is $a_k(C) = 2k + \lceil \frac{k}{s} \rceil$.

Proof. Let G_i , where $i = 1, 2, \dots, m$, be unit circle arcs that cover $\text{bd } C$, and let p_i, q_i and r_i be the center and the two endpoints of G_i , respectively. Clearly, we may assume that no G_i is a proper subset of any unit circle arc in $\text{bd } C$.

We label the points in such a way that in counterclockwise order, q_i is the starting and r_i is the endpoint of G_i , and the points q_1, q_2, \dots, q_m are in this counterclockwise order in $\text{bd } C$. For simplicity, we call the G_i s the *sides*, and their endpoints the *vertices* of C . Note that r_1, r_2, \dots, r_m are in this counterclockwise order as well, as otherwise $G_i \subset G_j$ for some $i \neq j$, which contradicts the assumption that C is m -sided. By Lemma 3.3, we have that p_1, p_2, \dots, p_m are also in this counterclockwise order.

Since C is complete, $p_i \in \text{bd } C$ for every value of i . Furthermore, since m is the minimal number of unit circle arcs that cover $\text{bd } C$, there is no point that belongs to more than two arcs. We observe also that if p_i is in the relative interior of a segment $[x, y] \subset \text{bd } C$, then, by Lemma 3.5, $G_i = [q_i, r_i]$ is a segment. Thus, replacing G_i by, say $\mathbb{S}(x) \cap C$, we still have a family of m unit circle arcs that cover $\text{bd } C$. This implies that, without loss of generality, we may assume that no p_i is in the relative interior of a segment on $\text{bd } C$.

Consider, first, the case that two consecutive sides, say G_i and G_{i+1} overlap. Then q_i, q_{i+1}, r_i and r_{i+1} are in this counterclockwise order in $\text{bd } C$. Thus, $[p_i, r_i]$ and $[p_{i+1}, q_{i+1}]$ are disjoint diameters, which yields, by Lemma 3.5, that $[p_i, p_{i+1}], [q_{i+1}, r_i] \subset \text{bd } C$, and that they are parallel. Hence, for any two overlapping sides of C , the common part is a straight line segment.

Now we show that the intersection of any two consecutive sides of C contains the center of exactly one side. Consider the sides G_i and G_{i+1} .

Case 1, G_i and G_{i+1} do not overlap. Then $r_i = q_{i+1}$. Observe that $p_i, p_{i+1} \in \mathbb{S}(r_i) \cap C$. Let G be the arc of $\text{bd } C$ connecting p_i and p_{i+1} and not containing r_i . We show that there is a point in the relative interior of G which is diametrically opposite only to r_i . Note that since C is a Reuleaux polygon, it yields that in this case $C \cap \mathbb{S}(r_i)$ must be a side of C .

Let p be an arbitrary relative interior point of G , and assume that $C \cap \mathbb{S}(p)$ contains not only r_i , but some other point x as well. Without loss of generality, we may assume that $x \in G_i$, which yields that $[p_i, r_i]$ and $[p, x]$ are disjoint diameters of C . Thus, by Lemma 3.5, $[p, p_i], [r_i, x] \subset \text{bd } C$, and they are parallel. Since here p is an arbitrary relative interior point of G , we have that either $G = [p_i, p_{i+1}]$ or there is some relative interior point z of G such that $G = [p_i, z] \cup [z, p_{i+1}]$. Observe that $G \subset C \cap \mathbb{S}(r_i)$, and

hence, as \mathbf{B} is not angled, it follows that $G = [p_i, p_{i+1}]$. Furthermore, for some point $x \in C$, we have that $[r_i, x] \subset \text{bd } C$, and that $[r_i, x]$ and $[p_i, p_{i+1}]$ are parallel. This means that $[r_i, x]$ belongs to both $\mathbb{S}(p_i)$ and $\mathbb{S}(p_{i+1})$, which contradicts our assumption that G_i and G_{i+1} do not overlap.

Case 2, G_i and G_{i+1} overlap; or in other words, $r_i \neq q_{i+1}$. Then, similarly like in Case 1, we have that $[r_i, q_{i+1}], [p_i, p_{i+1}] \subset \text{bd } C$, and they are parallel. Let L and L' denote the line containing $[p_i, p_{i+1}]$ and $[r_i, q_{i+1}]$, respectively. Observe that $\mathbb{S}(p_i)$ and $\mathbb{S}(p_{i+1})$ both contain $C \cap L'$, and thus, we have $C \cap L' = [r_i, q_{i+1}]$. Furthermore, note that, for any point p in the relative interior of $[p_i, p_{i+1}]$, the points of C diametrically opposite to p are exactly the points of $[r_i, q_{i+1}]$. Thus, the center of any side of C containing p is a point of $[q_{i+1}, r_i]$. Since we chose the sides of C in such a way that no center is contained in a straight line segment in $\text{bd } C$, we have that only q_{i+1} or r_i can be the center of a side, and also that $L \cap C = [p_i, p_{i+1}]$.

Suppose, for contradiction, that both q_{i+1} and r_i are centers, and let these sides be G_j and G_{j+1} . Then, we have $[p_i, p_{i+1}] \subseteq G_j \cap G_{j+1}$, and, similarly like in the previous paragraph, we may obtain that $[p_i, p_{i+1}] = G_j \cap G_{j+1}$. Thus, $q_{j+1} = p_i$ and $r_j = p_{i+1}$. Since $q_i \neq q_{i+1} = p_j$ and $q_j \neq q_{j+1} = p_i$, it follows that $[q_{i+1}, q_j]$ and $[p_i, q_i]$ are disjoint diameters of C . Hence, by Lemma 3.5, we have that $[q_i, q_{i+1}]$ and $[q_j, p_i]$ are parallel and contained in $\text{bd } C$. Thus, $[q_i, q_{i+1}]$ and $[q_{i+1}, r_i]$ are both contained in $\mathbb{S}(p_i)$, which contradicts our assumption that the normed plane is not angled.

We have shown that the intersection of any two consecutive sides contains the center of exactly one side. Since any point of $\text{bd } C$ belongs to at most two sides of C , these intersections are pairwise disjoint. As the number of centers is equal to the number of intersections, it follows that the center of every side of C is contained in one of these intersections. In fact, we showed a bit more: every center is the vertex of some other side.

For every value of i , consider a point z_i that belongs to G_i but no other side of C . Note that since no point of $\text{bd } C$ belongs to more than two sides of C , this is possible, and also that, by Lemma 3.5, the segments $[p_i, z_i]$, where $i = 1, 2, \dots, m$, are pairwise intersecting diameters of C . Clearly, the $2m$ points p_i and z_j form an alternating sequence S in $\text{bd } C$, and each of the two open arcs of $\text{bd } C$, starting at, say, p_1 and ending at z_1 , contains exactly $m - 1$ points. Since the subsequence of S in any of the above two arcs, starts with some z_i and ends with some p_j , we have that $m - 1$ is an even number, and thus, m is odd.

Now we prove the second part. Let $m = 2s + 1$. According to the previous paragraph, we have that for every i , $p_i \in G_{i+s} \cap G_{i+s+1}$. First, we show that the points z_i can be chosen in such a way that the set $Z = \{z_i : i = 1, 2, \dots, m\}$ contains no diametrically opposite pair.

Assume that for every i , z_i belongs to only G_i , but Z contains a diametrically opposite pair, say z_i and z_j . Then $j = i - s$ or $j = i + s$. Without loss of generality, we may assume that z_i and z_{i+s} are diametrically opposite. From this, by Lemma 3.5, we obtain that $[z_i, p_{i+s}]$ and $[z_{i+s}, p_i]$ are parallel and are contained in $\text{bd } C$. Let L be the

line containing $[p_{i+s}, z_i]$. Note that p_{i+s} is an endpoint of $L \cap C$, and let x be the other endpoint. Observe that $p_{i-s} \notin L$, as otherwise $z_{i+s} \in G_{i-s}$, which is a contradiction. In addition, x is not diametrically opposite to z_{i-s} . Indeed, if $[x, z_{i-s}]$ is a diameter, then $[x, z_{i-s}]$ and $[p_{i-s}, p_i]$ are disjoint diameters, and thus Lemma 3.5 yields that $[x, p_{i-s}], [z_{i-s}, p_i] \subset \text{bd } C$, which contradicts our assumption that the normed plane is not angled (cf. Figure 2). Now we choose any point $y \in (\text{bd } C) \setminus L$ sufficiently close to x , and replace z_i by y . Then, clearly, y is diametrically opposite to neither z_{i-s} nor z_{i+s} . Thus, to choose a subset Z that does not contain diametrically opposite points, we start with any set and then, applying the argument of this paragraph, we may replace the points one by one to reduce the number of diametrically opposite points.




FIGURE 2. An illustration for the proof of Theorem 4

We constructed a subset $Z = \{z_i : i = 1, 2, m\}$ such that for every i , z_i belongs only to G_i , and Z contains no diametrically opposite pair. Let A_i denote the closed arc of $\text{bd } C$, which, in counterclockwise order, starts at z_i and ends at z_{i+s} . Observe that by Lemma 3.3 and the choice of Z , no such arc contains a diametrically opposite pair. On the other hand, the sets A_{j_s} , where $j = 0, 1, \dots, 2k + \lceil \frac{k}{s} \rceil - 1$ and the indices are taken mod m , covers $\text{bd } C$ k -fold, and thus, they are a k -fold Borsuk covering of $\text{bd } C$. This proves that $a_k(C) \leq 2k + \lceil \frac{k}{s} \rceil$.

To prove the other direction, we note that the k -fold Borsuk coverings of the set $\{p_i : i = 1, 2, \dots, m\}$ can be identified with the k -fold vertex-colorings of a $(2s + 1)$ -cycle. Since it is known (cf. [20]) that the k -fold chromatic number of such a cycle is $2k + \lceil \frac{k}{s} \rceil$, the assertion follows. \square

From Theorems 1.1, 3 and Remark 4.2, we immediately obtain the following.

Theorem 5. *Let S be a set of positive diameter in a normed plane \mathbf{B} .*

- If S is not a set of unique completion, or S does not satisfy the supporting line property, then for every value of k , $a_k(S) = 2k$.
- If S is a set of unique completion that satisfies the supporting line property (see page 2) and the completion of S is not a Reuleaux polygon, then for every k , $a_k(S) = 2k + 1$.

5. REMARKS AND QUESTIONS

Remark 5.1. We note that our results cannot be extended to angled planes. For example, Theorem 2 fails if the unit disk \mathbf{B} is a parallelogram. Besides, any centrally symmetric polygon with $4m$ sides is a Reuleaux polygon with $2m$ sides in its norm (and thus, it has even sides according to our definition).

Remark 5.2. The k -fold Borsuk number of an o -symmetric polygon P with $2m$ sides in its own norm is $a_k(P) = 2k + \lceil \frac{2k}{m-1} \rceil$.

Proof. Let the vertices of the polygon be p_1, p_2, \dots, p_{2m} in counterclockwise order. Then p_i is diametrically opposite to p_{i+k-1}, p_{i+k} and p_{i+k+1} . Thus, the inequality $a_k(P) \geq 2k + \lceil \frac{2k}{m-1} \rceil$ follows from the Pigeonhole Principle. On the other hand, if G_i denotes the shorter arc in $\text{bd } P$, connecting the midpoints of $[p_i, p_{i+1}]$ and $[p_{i+k-1}, p_{i+k}]$, then, clearly, G_i contains the vertices of no diameter of P . Thus, the arcs $G_{i+t(k-1)}$, where $t = 1, 2, \dots, 2k + \lceil \frac{2k}{m-1} \rceil$, form a k -fold Borsuk-covering of $\text{bd } P$. \square

Remark 5.3. It is proven in [4] that in any angled normed plane there is a complete set of Borsuk number two. In other words, for a normed plane, the result in [1] about the Borsuk numbers of sets in the Euclidean plane holds in the same form if, and only if the plane is not angled. According to our results, the same can be observed about the multiple Borsuk numbers of sets.

Remark 5.4. In any angled normed plane, there is a Borsuk covering of a set of unique completion, satisfying the supporting line property (see page 2), that cannot be completed to a Borsuk covering of its completion.

Proof. If the norm is a parallelogram norm, the remark trivially follows. Hence, we may assume that the unit disk \mathbf{B} is not a parallelogram, and that its boundary contains $[x, y] \cup [y, z]$ and $[-x, -y] \cup [-y, -z]$. Without loss of generality, we may assume that the lines, containing $[x, y]$ and $[y, z]$, intersect \mathbf{B} in $[x, y]$ and $[y, z]$, respectively.

Let C be the truncation of \mathbf{B} with a line connecting the relative interior points w_1 and w_2 of $[x, y]$ and $[y, z]$, respectively. Clearly, the unique completion of C is \mathbf{B} , and C satisfies the supporting line property. Let w be the midpoint of $[w_1, w_2]$. Let u_1 and u_2 be relative interior points of $[-x, -y]$ and $[-y, -z]$, respectively (cf. Figure 3). Then the shorter arcs of $\text{bd } C$ connecting w to u_1 , u_1 to u_2 , and u_2 to w , is a Borsuk covering of $\text{bd } C$. On the other hand, y cannot be added to any of these arcs, which yields that this covering cannot be extended to any Borsuk covering of \mathbf{B} . \square




FIGURE 3. A Borsuk covering may not be extended in an angled plane

Note that if \mathbf{B} is a parallelogram, then the only complete sets of unit diameter are the translates of \mathbf{B} (cf. [22], [17] or [13]).

Remark 5.5. Let S be a compact set with $a(S) = 3$ in the normed plane where \mathbf{B} is a parallelogram. Then $a_k(S) = 3k$ for every k .

Proof. Without loss of generality, let \mathbf{B} be the unique completion of S . By the supporting line property, S contains at least two consecutive vertices of \mathbf{B} . Furthermore, since \mathbf{B} is the unique completion, S contains a point of the opposite side of \mathbf{B} . Thus, S contains three points at pairwise normed distances equal to $\text{diam } S$, which yields $a_k(S) \geq 3k$. By sub-additivity, we have $a_k(S) \leq 3k$, and the assertion readily follows. \square

REFERENCES

- [1] V.G. Boltyanski, *On the partition of plane figures into pieces of smaller diameters (in Russian)*, Colloq. Math. **21** (1970), 253-263.
- [2] V. G. Boltyanskii and I. C. Gohberg, *The Decomposition of Figures into Smaller Parts*, translated from Russian, The University of Chicago Press, Chicago, 1980.
- [3] V. Boltyanski, H. Martini and P.S. Soltan, *Excursions into Combinatorial Geometry*, Springer-Verlag, Berlin Heidelberg, 1997.
- [4] V.G. Boltyanski and V. Soltan, *Borsuk's problem (in Russian)*, Mat. Zametki **22** (1977), 621-631.
- [5] K. Borsuk, *Drei Sätze über die n -dimensionale euklidische Sphäre*, Fundamenta Math. **20** (1933), 177-190.
- [6] H.G. Eggleston, *Covering a three-dimensional set with sets of smaller diameter*, J. London Math. Soc. **30** (1955), 11-24.
- [7] H.G. Eggleston, *Sets of constant width in finite dimensional Banach spaces*, Israel J. Math. **3** (1965), 163-172.
- [8] B. Grünbaum, *Borsuk's partition conjecture in Minkowski planes*, Bull. Res. Council Israel, F1, N1 (1957), 25-30.
- [9] B. Grünbaum, *A simple proof of Borsuk's conjecture in three dimensions*, Proc. Cambridge Philos. Soc. **53** (1957), 776-778.
- [10] P.C. Hammer, *Convex curves of constant Minkowski breadth*, Proc. Symp. Pure Math. **7**, Amer. Math. Soc., Providence RI (1963), 291-304.

- [11] A. Heppes, *On the partitioning of a three-dimensional point set into sets of smaller diameter (Hungarian)*, Magyar Tud. Akad. Mat. Fiz. Oszt. Közl. **7** (1957), 413-416.
- [12] M. Hujter and Z. Lángi, *On the multiple Borsuk numbers of sets*, Israel J. Math., DOI:10.1007/s11856-013-0048-1, arXiv:1206.0892
- [13] A. Joós and Z. Lángi, *On the relative distances of seven points in a plane convex body*, J. Geom. **87** (2007), 83-95.
- [14] J. Kahn, and G. Kalai, *A counterexample to Borsuk's conjecture*, Bull. Amer. Math. Soc. **29** (1993), 60–62.
- [15] M. Lassak, *On relatively equilateral polygons inscribed in a convex body*, Publ. Math. Debrecen **65** (2004), 133-148.
- [16] J.P. Moreno, *Porosity and unique completion in strictly convex spaces*, Math. Z. **267**(1-2) (2011), 173-184.
- [17] M. Naszódi and B. Visy, *Sets with a unique extension to a set of constant width*, Discrete geometry, Monogr. Textbooks Pure Appl. Math., vol. 253, Dekker, New York, 2003, pp. 373–380. MR 2034729 (2004k:52005)
- [18] A.M. Raigorodskii, *Around Borsuks hypothesis*, Journal of Mathematical Sciences **154** (2008), no. 4, 604–623 (English).
- [19] G.T. Sallee, *The maximal set of constant width in a lattice*, Pacific J. Math. **28** (1969), 669-674.
- [20] S. Stahl, *n-tuple colorings and associated graphs*, J. Combin. Theory Ser. B **20** (1976), 185–203.
- [21] A.C. Thompson, *Minkowski Geometry*, Encyclopedia of Mathematics and its Applications **63**, Cambridge University Press, Cambridge, 1996.
- [22] D. Yost, *Irreducible convex sets*, Mathematika **38** (1991), no. 1, 134–155. MR 1116691 (92h:52006)

ZSOLT LÁNGI, DEPT. OF GEOMETRY, BUDAPEST UNIVERSITY OF TECHNOLOGY AND ECONOMICS, BUDAPEST, EGRY JÓZSEF U. 1., HUNGARY, 1111

E-mail address: zlangi@math.bme.hu

MÁRTON NASZÓDI, DEPT. OF GEOMETRY, LORAND EÖTVÖS UNIVERSITY, PÁZMÁNY PÉTER SÉTÁNY 1/C BUDAPEST, HUNGARY 1117

E-mail address: marton.naszodi@math.elte.hu