

A new approach to constant term identities and Selberg-type integrals

Gyula Károlyi*

School of Mathematics and Physics, The University of Queensland, Brisbane, QLD 4072, Australia

Zoltán Lóránt Nagy

Alfréd Rényi Institute of Mathematics, Reáltanoda utca 13–15, Budapest, 1053 Hungary

Fedor V. Petrov^{1,2}

Steklov Institute of Mathematics, Fontanka 27, 191023 St. Petersburg, Russia

Vladislav Volkov²

Saint-Petersburg State University, Universitetsky prospekt 28, 198504 St. Petersburg, Russia

Abstract

Selberg-type integrals that can be turned into constant term identities for Laurent polynomials arise naturally in conjunction with random matrix models in statistical mechanics. Built on a recent idea of Karasev and Petrov we develop a general interpolation based method that is powerful enough to establish many such identities in a simple manner. The main consequence is the proof of a conjecture of Forrester related to the Calogero–Sutherland model. In fact we prove a more general theorem, which includes Aomoto’s constant term identity at the same time. We also demonstrate the relevance of the method in additive combinatorics.

Keywords: Aomoto’s constant term identity, Calogero–Sutherland model, Combinatorial Nullstellensatz, Erdős–Heilbronn conjecture, Forrester’s conjecture, Hermite interpolation, Selberg integral

1. Introduction

Perhaps the most famous constant term identity is the one associated with the name of Freeman Dyson. In his seminal paper [13] dated back to 1962, Dyson proposed to replace Wigner’s classical Gaussian-based random matrix models by what now is known as the circular ensembles. The study of their joint eigenvalue probability density functions led Dyson to the following conjecture. Consider the family of Laurent polynomials

$$\mathcal{D}(\mathbf{x}; \mathbf{a}) := \prod_{1 \leq i \neq j \leq n} \left(1 - \frac{x_i}{x_j}\right)^{a_i}$$

parametrized by a sequence $\mathbf{a} = (a_1, \dots, a_n)$ of nonnegative integers, where $\mathbf{x} = (x_1, \dots, x_n)$ is a sequence of indeterminates. Denoting by $\text{CT}[\mathcal{L}(\mathbf{x})]$ the constant term of the Laurent polynomial $\mathcal{L} = \mathcal{L}(\mathbf{x})$, Dyson’s hypothesis can be formulated as the identity

$$\text{CT}[\mathcal{D}(\mathbf{x}; \mathbf{a})] = \frac{(a_1 + a_2 + \dots + a_n)!}{a_1! a_2! \dots a_n!} =: \binom{|\mathbf{a}|}{\mathbf{a}},$$

*Corresponding author. On leave from Eötvös University, Budapest.

Email addresses: karolyi@cs.elte.hu (Gyula Károlyi), nagyzoztanlorant@gmail.com (Zoltán Lóránt Nagy), fedyapetrov@gmail.com (Fedor V. Petrov), vladvolkov239@gmail.com (Vladislav Volkov)

¹Also at St. Petersburg State University.

²Also at B.N. Delone International Laboratory “Discrete and Computational Geometry”.

where $|\mathbf{a}| = a_1 + a_2 + \dots + a_n$. Using the shorthand notation $\mathcal{D}(\mathbf{x}; k)$ for the equal parameter case $\mathbf{a} = (k, \dots, k)$, the constant term of $\mathcal{D}(\mathbf{x}; k)$ for $k = 1, 2, 4$ corresponds to the normalization factor of the partition function for the circular orthogonal, unitary and symplectic ensemble, respectively.

Dyson's conjecture was confirmed by Gunson [unpublished]³ and Wilson [49] in the same year. The most elegant proof, based on Lagrange interpolation, is due to Good [22].

Let q denote yet another independent variable. In 1975 Andrews [5] suggested the following q -analogue of Dyson's conjecture: The constant term of the Laurent polynomial

$$\mathcal{D}_q(\mathbf{x}; \mathbf{a}) := \prod_{1 \leq i < j \leq n} \left(\frac{x_i}{x_j} \right)_{a_i} \left(\frac{qx_j}{x_i} \right)_{a_j}$$

must be the q -multinomial coefficient

$$\begin{bmatrix} |\mathbf{a}| \\ \mathbf{a} \end{bmatrix} := \frac{(q)_{|\mathbf{a}|}}{(q)_{a_1} (q)_{a_2} \dots (q)_{a_n}},$$

where $(t)_k = (1-t)(1-tq)\dots(1-tq^{k-1})$. Note that the slight asymmetry of the function \mathcal{D}_q disappears when one considers $\mathcal{D} = \mathcal{D}_1$; specializing at $q = 1$, Andrews' conjecture gives back that of Dyson.

Despite several attempts [27, 46, 47] the problem remained unsolved until 1985, when Zeilberger and Bressoud [53] found a tour de force combinatorial proof; see also [9]. Shorter proofs are due to Gessel and Xin [21] and Cai [10]. Recently an idea of Karasev and Petrov [35] led to a very short proof by Károlyi and Nagy [37], which we consider as a precursor to the present paper.

Constant term identities like these and their generalizations are intimately related to Selberg's integral formula [44]. Colloquially referred to as the Selberg integral, it asserts

$$\begin{aligned} S_n(\alpha, \beta, \gamma) &:= \int_0^1 \dots \int_0^1 \prod_{i=1}^n t_i^{\alpha-1} (1-t_i)^{\beta-1} \prod_{1 \leq i < j \leq n} |t_i - t_j|^{2\gamma} dt_1 \dots dt_n \\ &= \prod_{j=0}^{n-1} \frac{\Gamma(\alpha + j\gamma)\Gamma(\beta + j\gamma)\Gamma(1 + (j+1)\gamma)}{\Gamma(\alpha + \beta + (n+j-1)\gamma)\Gamma(1 + \gamma)}, \end{aligned}$$

where the complex parameters α, β, γ satisfy

$$\Re(\alpha) > 0, \quad \Re(\beta) > 0, \quad \Re(\gamma) > -\min\{1/n, \Re(\alpha)/(n-1), \Re(\beta)/(n-1)\}.$$

The continued interest in the Selberg integral, demonstrated for example by the most recent article [43], is due to its role in random matrix theory, statistical mechanics, special function theory among other fields; see the comprehensive exposition [19].

The Selberg integral is well-known to be equivalent to Morris's constant term identity [42]

$$\text{CT} \left[\prod_{j=1}^n (1-x_j)^a (1-1/x_j)^b \mathcal{D}(\mathbf{x}; k) \right] = \prod_{j=0}^{n-1} \frac{(a+b+kj)!(kj+k)!}{(a+kj)!(b+kj)!k!}, \quad (1.1)$$

or in a more compact form,

$$\text{CT}[\mathcal{M}(\mathbf{x}; a, b, k)] = M(n; a, b, k),$$

where the parameters a, b, k are nonnegative integers. The equivalence is established, via a suitable change of variables, by an application of a theorem of Carlson [11] and the residue theorem. This method can be employed to reduce Selberg-type integrals to constant term identities.

³Gunson's proof is similar to that of Wilson, cf. [13]. A related conjecture of Dyson is proved by Gunson in [23].

Introducing an extra $t_1 \cdots t_m$ factor into the integrand, Aomoto [4] in 1987 proved an extension of the Selberg integral. Based on the fundamental theorem of calculus, it yields besides Anderson's [3] one of the simplest known proofs of the Selberg integral itself. Turned into a constant term identity, Aomoto's integral reads as

$$\text{CT} \left[\prod_{j=1}^n (1-x_j)^{a+\chi(j \leq m)} (1-1/x_j)^b \mathcal{D}(\mathbf{x}; k) \right] = \prod_{j=0}^{n-1} \frac{(a+b+kj+\chi(j \geq n-m))!(kj+k)!}{(a+kj+\chi(j \geq n-m))!(b+kj)!k!}, \quad (1.2)$$

where $\chi(S)$ is equal to 1 if the statement S is true and 0 otherwise.

Intimately related to the theory of random matrices, in particular the Dyson Brownian motion model [14], is the Calogero-Sutherland quantum many body system for spinless quantum particles on the unit circle interacting via the $1/r^2$ two-body potential, see [18, Chapter 11]. Generalizations to include internal degrees of freedom of the particles were formulated in the early 1990's. In his 1995 paper [17] Forrester initiated the study of the analogue of the Selberg integral for the corresponding exact multicomponent ground-state wavefunction. Presented in the form of the constant term for the Laurent polynomial

$$\mathcal{F}(\mathbf{x}; n_0; a, b, k) = M(\mathbf{x}; a, b, k) \prod_{n_0 < i \neq j \leq n} \left(1 - \frac{x_i}{x_j} \right),$$

the normalization factor for the most interesting two-component case can be determined by the conjectured identity

$$\text{CT} [\mathcal{F}(\mathbf{x}; n_0; a, b, k)] = M(n_0; a, b, k) \times \prod_{j=0}^{n-n_0-1} \frac{(j+1)(a+b+kn_0+(k+1)j)!(kn_0+(k+1)j+k)!}{(a+kn_0+(k+1)j)!(b+kn_0+(k+1)j)!k!}.$$

A q -analogue of this hypothesis which extends the q -Morris ex-conjecture [42] was formulated and studied in [7]. Despite several further attempts [8, 20, 25, 31, 32, 33, 34], these conjectures have been resolved only in some particular cases. The main achievement in the present paper is the proof of these identities, and in a form that also includes Aomoto's formula (1.2); see Theorem 6.2 for the precise formulation. Along the way we develop a method with a wide range of possible applications, some of which are given as instructive examples.

A new proof of the Dyson conjecture given in [35] and the subsequent proof of the Zeilberger–Bressoud identity presented in [37] are based on a quick application of the following explicit version of the Combinatorial Nullstellensatz [1] found independently by Lasoń [40] and by Karasev and Petrov [35].

Lemma 1.1. *Let \mathbb{F} be an arbitrary field and $F \in \mathbb{F}[x_1, x_2, \dots, x_n]$ a polynomial of degree $\deg(F) \leq d_1 + d_2 + \dots + d_n$. For arbitrary subsets C_1, C_2, \dots, C_n of \mathbb{F} with $|C_i| = d_i + 1$, the coefficient of $\prod x_i^{d_i}$ in F is*

$$\sum_{c_1 \in C_1} \sum_{c_2 \in C_2} \cdots \sum_{c_n \in C_n} \frac{F(c_1, c_2, \dots, c_n)}{\phi'_1(c_1)\phi'_2(c_2)\cdots\phi'_n(c_n)},$$

where $\phi_i(z) = \prod_{c \in C_i} (z - c)$.

One principal aim of the present paper is to turn this idea into a method, which has the power to reduce seemingly difficult evaluations to simple combinatorial problems. To this end, in the next section we present a somewhat abstract framework, which allows us to extend the previous lemma to multisets via Hermite interpolation. In Section 3 we demonstrate the strength of the method in additive combinatorics by providing a new proof of an extension of the Erdős–Heilbronn conjecture, which is devoid of the heavy technical details that were needed previously. This is followed in Section 4 by an application to a problem of Kadell [29] in algebraic combinatorics, where the amount of reduction of former complexities is even more voluminous. In Section 5, which can be viewed as a prelude to the main result, we reestablish (1.1) using our method, thereby giving a short proof of the Selberg integral itself. Besides formulating our main result, in Section 6 we point out how a slight modification yields, modulo some routine computation, a one-page derivation of the q -Morris identity. The same idea with more delicate combinatorics leads to the solution of the problem of Forrester in the concluding section. Finally we mention that the method developed here can be successfully applied to prove Kadell's orthogonality conjectures [30], see [36].

2. On the Combinatorial Nullstellensatz

Alon's Nullstellensatz [1] describes effectively the structure of polynomials which vanish on a finite Cartesian product over an arbitrary field. It implies the following non-vanishing criterion. Let F be a polynomial as in Lemma 1.1. If the coefficient of $\prod x_i^{d_i}$ in F is non-zero, then F cannot vanish on a set $C_1 \times C_2 \times \dots \times C_n$, where $|C_i| > d_i$ for every i . Note that this is also an immediate consequence of Lemma 1.1. A standard application of the polynomial method to prove a combinatorial hypothesis works as follows. Assuming the falsity of the hypothesis, build a polynomial whose values are all zero over a large Cartesian product, then compute the coefficient of the appropriate leading term. If that coefficient is not zero, the criterion leads to the desired contradiction. The difficulty often lies in the computation of that coefficient. This is where the power of Lemma 1.1 comes into the picture, which is clearly demonstrated in the next section. An extension of the non-vanishing criterion for the case when C_i are multisets, along with some applications, was obtained recently by Kós and Rónyai [39]; see also [38]. Here we generalize Lemma 1.1 in a similar spirit.

Let $\mathfrak{V}_1, \dots, \mathfrak{V}_n$ be vector spaces over the same field \mathbb{F} . For each i , fix a basis \mathfrak{B}_i in \mathfrak{V}_i and fix the corresponding basis $\otimes \mathfrak{B}_i$ in the tensor product space $\otimes \mathfrak{V}_i$. Consider arbitrary non-empty subsets $A_i \subseteq \mathfrak{B}_i$, labelled vectors $a_i \in A_i$, and linear functionals $\eta_i \in \text{Hom}(\mathfrak{V}_i, \mathbb{F})$ that satisfy the conditions $\eta_i(a_i) = 1$ and $\eta_i(b) = 0$ for every $b \in A_i \setminus \{a_i\}$. Our tool will be the following straightforward observation.

Lemma 2.1. *Assume that the tensor $F \in \otimes \mathfrak{V}_i$ satisfies the following condition: if $b_i \in \mathfrak{B}_i$ and the coordinate of F at $\otimes b_i$ does not vanish, then either $b_i = a_i$ for every i or $b_i \in A_i \setminus \{a_i\}$ for at least one index i . Then the coordinate of F at $\otimes a_i$ equals $(\otimes \eta_i)(F)$. \square*

We will apply this lemma in the following situation:

$$\mathfrak{V}_i = \mathbb{F}[x_i], \quad \mathfrak{B}_i = \{1, x_i, x_i^2, \dots\}, \quad A_i = \{1, x_i, \dots, x_i^{d_i}\}, \quad a_i = x_i^{d_i}.$$

Moreover we will assume that the value of $\eta_i \in \text{Hom}(\mathfrak{V}_i, \mathbb{F})$ at $f \in \mathbb{F}[x_i]$ is the same as the coefficient of $x_i^{d_i}$ in f if $\deg(f) \leq d_i$. Now $\mathbb{F}[x_1, \dots, x_n]$, as a vector space over \mathbb{F} , can be identified with $\otimes \mathfrak{V}_i$ via the unique isomorphism, which extends the correspondence

$$x_1^{k_1} \dots x_n^{k_n} \longleftrightarrow x_1^{k_1} \otimes \dots \otimes x_n^{k_n}, \quad k_i \in \{0, 1, 2, \dots\}.$$

An important feature of this identification is the following.

Lemma 2.2. *Assume that linear functionals $\vartheta_i \in \text{Hom}(\mathfrak{V}_i, \mathbb{F})$ are given in the form $\vartheta_i(f) = f^{(m_i)}(c_i)$ for some elements $c_i \in \mathbb{F}$ and nonnegative integers m_i . Then for any polynomial $G \in \mathbb{F}[x_1, \dots, x_n]$,*

$$(\otimes \vartheta_i)(G) = \frac{\partial^{m_1 + \dots + m_n} G}{\partial x_1^{m_1} \dots \partial x_n^{m_n}}(c_1, \dots, c_n).$$

Proof. Indeed, identifying $\mathbb{F} \otimes \dots \otimes \mathbb{F}$ with \mathbb{F} via the correspondence $\alpha_1 \otimes \dots \otimes \alpha_n \longleftrightarrow \alpha_1 \dots \alpha_n$, for any monomial $G = x_1^{k_1} \dots x_n^{k_n} \in \otimes \mathfrak{B}_i$ we obtain

$$(\otimes \vartheta_i)(G) = \otimes (\vartheta_i(x_i^{k_i})) = \prod_{i=1}^n k_i(k_i - 1) \dots (k_i - m_i + 1) \cdot (\otimes c_i^{k_i - m_i}) = \frac{\partial^{m_1 + \dots + m_n} G}{\partial x_1^{m_1} \dots \partial x_n^{m_n}}(c_1, \dots, c_n).$$

The general statement follows by linearity. \square

Let $F \in \mathbb{F}[x_1, \dots, x_n]$. We say that no monomial majorizes $\prod x_i^{d_i}$ in F if every monomial $\prod x_i^{k_i}$ with a non-zero coefficient in F satisfies either $k_i = d_i$ for every i or $k_i < d_i$ for some i . This is certainly the case if $\deg(F) \leq d_1 + \dots + d_n$. Such a polynomial F obviously satisfies the condition in Lemma 2.1. As a warm-up exercise we reestablish Lemma 1.1 in a slightly stronger form using this language.

Theorem 2.3. Let $F \in \mathbb{F}[x_1, \dots, x_n]$ be a polynomial such that no monomial majorizes $M = \prod x_i^{d_i}$ in F . Let C_1, \dots, C_n be arbitrary subsets of \mathbb{F} such that $|C_i| = d_i + 1$ for every i . Then the coefficient of M in F can be evaluated as

$$\sum_{c_1 \in C_1} \sum_{c_2 \in C_2} \cdots \sum_{c_n \in C_n} \prod_{i=1}^n \kappa(C_i, c_i) F(c_1, c_2, \dots, c_n),$$

where $\kappa(C_i, c_i) = \left(\prod_{c \in C_i \setminus \{c_i\}} (c_i - c) \right)^{-1}$. Consequently, if the above coefficient is not zero, then there exists a system of representatives $c_i \in C_i$ such that $F(c_1, c_2, \dots, c_n) \neq 0$.

Proof. Define the linear functionals $\eta_i \in \text{Hom}(\mathfrak{A}_i, \mathbb{F})$ by $\eta_i(f) = \sum_{c_i \in C_i} \kappa(C_i, c_i) f(c_i)$. According to the Lagrange interpolation formula, $\eta_i(f)$ is equal to the coefficient of $x_i^{d_i}$ in f for any $f \in \mathbb{F}[x_i]$ with $\deg(f) \leq d_i$. Since each η_i is a linear combination of linear functionals of the form $\vartheta_i(f) = f^{(0)}(c_i)$, the claim follows easily from Lemmas 2.1 and 2.2. \square

Extending the notion of the 0/1-valued characteristic function of a set, a finite multiset C in \mathbb{F} can be represented by a multiplicity function $\omega : \mathbb{F} \rightarrow \{0, 1, 2, \dots\}$ with finite sum $|C| := \sum_{x \in \mathbb{F}} \omega(x)$. We denote by $\text{supp}(C) := \{c \in \mathbb{F} \mid \omega(c) \neq 0\}$ the supporting set of C and, with a slight abuse of notation, write $c \in C$ if $c \in \text{supp}(C)$. A finite union of multisets is understood as the sum of the corresponding multiplicity functions. An appropriate generalization of Theorem 2.3 for multisets can be formulated as follows.

Theorem 2.4. Let $F \in \mathbb{F}[x_1, \dots, x_n]$ be a polynomial such that no monomial majorizes $M = \prod x_i^{d_i}$ in F . Let C_1, \dots, C_n be arbitrary multisets in \mathbb{F} with corresponding multiplicity functions $\omega_1, \dots, \omega_n$ such that $|C_i| = d_i + 1$ for every i . Assume that either $\text{char}(\mathbb{F}) = 0$ or $\text{char}(\mathbb{F}) \geq \omega_i(c)$ for every index i and $c \in \mathbb{F}$. Then the coefficient of M in F can be evaluated as

$$[M]F = \sum_{c_1 \in C_1} \sum_{m_1 < \omega_1(c_1)} \cdots \sum_{c_n \in C_n} \sum_{m_n < \omega_n(c_n)} \prod_{i=1}^n \kappa(C_i, c_i, m_i) \frac{\partial^{m_1 + \dots + m_n} F}{\partial x_1^{m_1} \dots \partial x_n^{m_n}}(c_1, \dots, c_n),$$

where

$$\kappa(C_i, c_i, m_i) = \frac{1}{m_i! \cdot (\omega_i(c_i) - 1 - m_i)!} \cdot \left(\frac{1}{\prod_{c \in C_i \setminus \{c_i\}} (x - c)^{\omega_i(c)}} \right) \Bigg|_{x=c_i}.$$

Consequently, if $[M]F \neq 0$, then there exists a system of representatives $c_i \in C_i$ and multiplicities $m_i < \omega_i(c_i)$ such that

$$\frac{\partial^{m_1 + \dots + m_n} F}{\partial x_1^{m_1} \dots \partial x_n^{m_n}}(c_1, \dots, c_n) \neq 0.$$

Remarks. 1. We tacitly assume that the m_i 's are nonnegative integers. 2. When each ω_i is a 0/1-valued function, the statement reduces to Theorem 2.3. 3. It is possible to derive this result, in a slightly weaker form, from the earlier works of Kós et al. [38, 39]. We preferred this more direct approach.

Proof. To construct the linear functionals η_i we replace Lagrange interpolation by Hermite interpolation. For $c_i \in C_i$, $0 \leq m_i < \omega_i(c_i)$, let $g(C_i, c_i, m_i)$ denote the unique polynomial of degree less than $|C_i|$, provided by the Chinese Remainder Theorem, to the system of simultaneous congruences

$$\begin{aligned} g(C_i, c_i, m_i)(x_i) &\equiv (x_i - c_i)^{m_i} / m_i! \pmod{(x_i - c_i)^{\omega_i(c_i)}}, \\ g(C_i, c_i, m_i)(x_i) &\equiv 0 \pmod{(x_i - c)^{\omega_i(c)}} \quad (c \in C_i \setminus \{c_i\}) \end{aligned}$$

in \mathfrak{A}_i . That is, $g(C_i, c_i, m_i)$ is the unique polynomial $g \in \mathbb{F}[x_i]$ of degree less than or equal to d_i , which satisfies $g^{(m_i)}(c_i) = 1$ and $g^{(m')}(u) = 0$ otherwise if $m' < \omega_i(u)$, $u \in \mathbb{F}$. Denote by $\kappa(C_i, c_i, m_i)$ the coefficient of $x_i^{d_i}$ in $g(C_i, c_i, m_i)$. Then Lemmas 2.1 and 2.2 can be applied as before for the linear functionals $\eta_i \in \text{Hom}(\mathfrak{A}_i, \mathbb{F})$ given by

$$\eta_i(f) = \sum_{c_i \in C_i} \sum_{m_i < \omega_i(c_i)} \kappa(C_i, c_i, m_i) f^{(m_i)}(c_i).$$

To compute the coefficients $\kappa(C_i, c_i, m_i)$, write $p_i(x_i) = \prod_{c \in C_i \setminus \{c_i\}} (x_i - c)^{\omega_i(c)}$. That is, there exist polynomials $h_i, r_i \in \mathbb{F}[x_i]$ with $\deg(h_i) < \omega_i(c_i)$ and $\deg(r_i) \leq d_i - \omega_i(c_i)$ such that

$$h_i(x_i) = \frac{g(C_i, c_i, m_i)(x_i)}{p_i(x_i)} = \frac{(x_i - c_i)^{m_i}}{m_i! p_i(x_i)} + (x_i - c_i)^{\omega_i(c_i)} \frac{r_i(x_i)}{p_i(x_i)}$$

with $\kappa(C_i, c_i, m_i)$ being the coefficient of $x_i^{\omega_i(c_i)-1}$ in $h_i(x_i)$. Expanding both the left- and the right-hand side as a formal power series in the variable $x_i - c_i$ one finds that $\kappa(C_i, c_i, m_i)$ is the coefficient of $(x_i - c_i)^{\omega_i(c_i)-m_i-1}$ in $1/(m_i! p_i(x_i))$. The result follows by an application of Taylor's formula. \square

3. An application to additive theory

Let $\mathcal{S} = \{S_{ij} \mid 1 \leq i < j \leq n\}$ be a family of subsets of the cyclic group $\mathbb{Z}_p := \mathbb{Z}/p\mathbb{Z}$ of prime order p . For a collection of sets $A_1, \dots, A_n \subseteq \mathbb{Z}_p$, consider the following restricted sumset:

$$\bigwedge_{\mathcal{S}} A_i = \{a_1 + \dots + a_n \mid a_i \in A_i, a_j - a_i \notin S_{ij} \text{ for } i < j\}.$$

For the special case when $A_i \equiv A$ and $S_{ij} \equiv \{0\}$, Dias da Silva and Hamidoune [12] proved

$$\left| \bigwedge_{\mathcal{S}} A_i \right| \geq \min \{p, n|A| - n^2 + 1\},$$

thus establishing a long-standing conjecture of Erdős and Heilbronn [16]. Their proof exploited the properties of cyclic spaces of derivations on exterior product spaces and the representation theory of symmetric groups; see [2] for another proof based on the polynomial method. A far reaching generalization was obtained by Hou and Sun [26]. Here we use Lemma 1.1 to reestablish their result in a short and elegant manner, thereby also providing a simplified proof to the Dias da Silva–Hamidoune theorem. Note that although our formulation below is slightly different, it is still equivalent to [26, Theorem 1.1].

Theorem 3.1. *Let A_1, \dots, A_n be subsets of a field \mathbb{F} such that $|A_i| = k$ for $1 \leq i \leq n$ and assume that $S_{ij} \subseteq \mathbb{F}$ satisfy $|S_{ij}| \leq s$ for $1 \leq i < j \leq n$. If either $\text{char}(\mathbb{F}) = 0$ or*

$$\text{char}(\mathbb{F}) > \max \{n\lceil s/2 \rceil, n(k-1) - n(n-1)\lceil s/2 \rceil\},$$

then

$$\left| \bigwedge_{\mathcal{S}} A_i \right| \geq n(k-1) - n(n-1)\lceil s/2 \rceil + 1.$$

Proof. Since posing extra restrictions cannot increase the size of the sumset, we will assume that s is even and $|S_{ij}| = s = 2t$ holds for every pair $i < j$. We may also assume that $k-1 \geq (n-1)t$. We proceed by way of contradiction. Suppose that $\bigwedge_{\mathcal{S}} A_i$ is contained in a set C of size $n(k-1) - n(n-1)t$, and consider the polynomial

$$\prod_{e \in C} (x_1 + \dots + x_n - e) \times \prod_{i < j} \left(\prod_{e \in S_{ij}} (x_j - x_i - e) \right).$$

This polynomial of degree $n(k-1)$ vanishes on the Cartesian product $A_1 \times \dots \times A_n$. According to Lemma 1.1, the coefficient of the monomial $\prod x_i^{k-1}$ must be zero. This coefficient remains the same if we slightly modify the polynomial and consider

$$F(\mathbf{x}) = \prod_{e=\binom{n}{2}t+1}^{n(k-1)-\binom{n}{2}t} (x_1 + \dots + x_n - e) \times \prod_{i < j} \left(\prod_{e=-t}^{t-1} (x_j - x_i - e) \right)$$

instead, keeping all leading terms intact. This coefficient is easy to compute when one applies Lemma 1.1 with $C_i \equiv \{0, 1, \dots, k-1\}$. Indeed, if $F(\mathbf{c}) \neq 0$ for some $\mathbf{c} \in C_1 \times \dots \times C_n$, then $|c_j - c_i| \geq t$ for every pair $i < j$. Accordingly,

$$\binom{n}{2}t \leq c_1 + \dots + c_n \leq n(k-1) - \binom{n}{2}t,$$

thus it must be $c_1 + \dots + c_n = \binom{n}{2}t$ and the numbers c_1, \dots, c_n , in some order, must coincide with the numbers $0, t, 2t, \dots, (n-1)t$. Moreover, it must be the natural order, for if $c_i > c_j$ for some $i < j$, then $c_i - c_j \geq t + 1$. Thus the computation of the coefficient reduces to the evaluation of

$$\frac{F(c_1, c_2, \dots, c_n)}{\phi'_1(c_1)\phi'_2(c_2)\dots\phi'_n(c_n)}$$

at the point $\mathbf{c} = (0, t, 2t, \dots, (n-1)t)$. After some cancellations this leads to the value

$$(-1)^{\binom{n}{2}t} \times \frac{(n(k-1) - n(n-1)t)!}{(t!)^n} \times \prod_{i=1}^n \frac{(it)!}{(k-1 - (i-1)t)!}$$

which is not zero in view of the assumption on the characteristic of the field. This contradiction completes the proof. \square

The tightness of the bound is demonstrated by the choice

$$A_i \equiv \{0, 1, \dots, k-1\}, S_{ij} \equiv \{-t+1, -t+2, \dots, t-1\}.$$

3.1. Further examples

The alert reader must have already extracted from the above argument the following general statement about restricted sumsets, which is rather folklore, cf. [2, Theorem 2.1].

Theorem 3.2. *Let d_i, s_{ij} denote non-negative integers, and let A_1, \dots, A_n and S_{ij} ($1 \leq i < j \leq n$) be subsets of a field \mathbb{F} with $|A_i| = d_i + 1$, $|S_{ij}| = s_{ij}$. Assume that $N = \sum d_i - \sum s_{ij} \geq 0$. If the coefficient of the monomial $\prod x_i^{d_i}$ in the polynomial*

$$F_0(\mathbf{x}) = (x_1 + \dots + x_n)^N \prod_{i < j} (x_j - x_i)^{s_{ij}} \in \mathbb{F}[x_1, \dots, x_n]$$

is non-zero, then $|\bigwedge_{i,j} A_i| > N$.

In the proof of Theorem 3.1 we applied Lemma 1.1 in the case $d_i \equiv k-1$, $s_{ij} \equiv 2t$ to obtain the coefficient $[x_1^{d_1} \dots x_n^{d_n}]F_0$ in a simple product form. Similar arguments work in the following cases. The first example concerns a related result of Sun and Yeh, cf. [48, Theorem 1.1], which involves only a minor modification.

Example 3.3. *Let $d_i = k-i$, $s_{ij} \equiv 2t-1$. Then $N = n(k-1) - n(n-1)t$ and*

$$[x_1^{d_1} \dots x_n^{d_n}]F_0 = (-1)^{\binom{n}{2}t} \times \frac{N!}{(t!)^n n!} \times \prod_{i=1}^n \frac{(it)!}{(k-1 - (i-1)t)!}.$$

Proof. Apply Lemma 1.1 to the modified polynomial

$$F(\mathbf{x}) = \prod_{e=\binom{n}{2}t+1}^{n(k-1)-\binom{n}{2}t} (x_1 + \dots + x_n - e) \times \prod_{i < j} \left(\prod_{e=1-t}^{t-1} (x_j - x_i - e) \right)$$

with the choice $C_i = \{0, 1, \dots, k-1\} \setminus \{jm \mid 0 \leq j < i-1\}$. Once again $F(\mathbf{c}) \neq 0$ for $\mathbf{c} \in C_1 \times \dots \times C_n$ if and only if $c_i = (i-1)t$ for every $1 \leq i \leq n$, and the slight changes in the computation are easy to detect. \square

The next example considers the Alon–Nathanson–Ruzsa theorem [2]. Although our approach is not significantly different from the original proof, we include it for it represents an atypical application of Lemma 1.1, when more than one $\mathbf{c} \in C_1 \times \cdots \times C_n$ contributes to a non-zero summand.

Example 3.4. Let the d_i be arbitrary, $s_{ij} \equiv 1$. Then $N = d_1 + \cdots + d_n - \binom{n}{2}$ and

$$[x_1^{d_1} \cdots x_n^{d_n}]F_0 = \frac{N!}{d_1! \cdots d_n!} \prod_{i < j} (d_j - d_i).$$

Proof. Replace the polynomial F_0 by

$$F(\mathbf{x}) = \prod_{e=\binom{n}{2}+1}^{d_1+\cdots+d_n} (x_1 + \cdots + x_n - e) \times \prod_{i < j} (x_j - x_i)$$

and apply Lemma 1.1 with $C_i = \{0, 1, \dots, d_i\}$. Consider an element $\mathbf{c} \in C_1 \times \cdots \times C_n$ whose coordinates c_i are mutually different. Then $F(\mathbf{c}) \neq 0$ only if $\{c_1, \dots, c_n\} = \{0, \dots, n-1\}$. That is, there is a permutation $\pi = \pi_c \in \mathfrak{S}(n)$ such that $c_i = \pi_c(i) - 1$. For such a \mathbf{c} ,

$$F(\mathbf{c}) = (-1)^N N! \times \text{sign}(\pi_c) \prod_{i < j} (j - i), \quad \phi'(c_i) = (-1)^{d_i - c_i} c_i! (d_i - c_i)!.$$

Since $\binom{d_i}{c_i} = 0$ for $d_i < c_i$, it is enough to prove that

$$\sum_{\pi \in \mathfrak{S}(n)} \text{sign}(\pi) \prod_{i=1}^n \binom{d_i}{\pi(i) - 1} = \prod_{i < j} \frac{d_j - d_i}{j - i}.$$

To establish this identity, notice that both sides are completely antisymmetric polynomials of minimum possible degree $n(n-1)/2$ in the variables d_i , which attain the same value at $(d_1, \dots, d_n) = (0, \dots, n-1)$. \square

Remark. A more direct proof goes as follows. Write $x^{[k]} = x(x-1)\cdots(x-k+1)$ and consider the polynomials

$$F(\mathbf{x}) = \left(\sum_{i=1}^n x_i - \binom{n}{2} \right)^{[N]} \prod_{i < j} (x_j - x_i), \quad F^*(\mathbf{x}) = \sum_{k_1+\cdots+k_n=d_1+\cdots+d_n} \frac{N!}{\prod k_i!} \prod_{i < j} (k_j - k_i) \prod_{i=1}^n x_i^{[k_i]}.$$

It is enough to prove that $F - F^*$ vanishes on the Cartesian product of the sets $C_i = \{0, 1, \dots, d_i\}$, for then $[x_1^{d_1} \cdots x_n^{d_n}](F - F^*) = 0$ by Lemma 1.1 and therefore

$$[x_1^{d_1} \cdots x_n^{d_n}]F_0 = [x_1^{d_1} \cdots x_n^{d_n}]F = [x_1^{d_1} \cdots x_n^{d_n}]F^* = \frac{N!}{d_1! \cdots d_n!} \prod_{i < j} (d_j - d_i)$$

as claimed. For the proof, notice that $\mathbf{c} \in C_1 \times \cdots \times C_n$ implies $F(\mathbf{c}) = F^*(\mathbf{c}) = 0$ unless $c_i = d_i$ for every i , in which case $F(\mathbf{c}) = F^*(\mathbf{c})$ follows from the very choice of the coefficients in F^* . This argument can be extended to show that in fact $F = F^*$.

Our final example originates in Xin [50], where it appears in the form of the constant term identity

$$\text{CT} \left[x_1^{-a_1} \cdots x_n^{-a_n} (x_1 + \cdots + x_n)^{a_1 + \cdots + a_n} \prod_{i \neq j} (1 - x_j/x_i)^{a_i} \right] = \binom{|\mathbf{a}|}{\mathbf{a}}, \quad (3.1)$$

see also [20]. Here the full capacity of Theorem 2.4 can be exploited with a minimum amount of computation.

Example 3.5. Let $d_i = na_i$, $s_{ij} = a_i + a_j$. Then $N = a_1 + \cdots + a_n$ and

$$[x_1^{d_1} \cdots x_n^{d_n}]F_0 = (-1)^{\sum_{i < j} a_i} \binom{|\mathbf{a}|}{\mathbf{a}}.$$

Proof. For the proof we may assume that $\text{char}(\mathbb{F}) = 0$. Choose an arbitrary set $B = \{b_1, \dots, b_n\} \subset \mathbb{F}$ so that $b_1 + \dots + b_n = 0$, and consider the multisets C_1, \dots, C_n with $\text{supp}(C_i) = B$ and multiplicity functions given by $\omega_i(b_j) = a_i + \chi(j = i)$; then $|C_i| = d_i + 1$. We apply Theorem 2.4 to the polynomial

$$F(\mathbf{x}) = (-1)^{\sum_{i < j} a_i} F_0(\mathbf{x}) = (x_1 + \dots + x_n)^{a_1 + \dots + a_n} \prod_{i \neq j} (x_i - x_j)^{a_i}.$$

There is only one non-zero summand in the summation formula for $[x_1^{d_1} \dots x_n^{d_n}]F$. Indeed, suppose that

$$\frac{\partial^{m_1 + \dots + m_n} F}{\partial x_1^{m_1} \dots \partial x_n^{m_n}}(c_1, \dots, c_n) \neq 0$$

for some $\mathbf{c} \in C_1 \times \dots \times C_n$ with $0 \leq m_i < \omega_i(c_i)$. First we show that the coordinates c_i are mutually different. Assume that on the contrary, $c_i = c_j$ for some $i \neq j$. Then $m_i + m_j \leq \omega_i(c_i) + \omega_j(c_j) \leq a_i + a_j - 1$. This implies that the polynomial $H := \prod (\partial/\partial x_i)^{m_i} F$ is divisible by $x_i - x_j$, a contradiction.

Thus, $\{c_1, \dots, c_n\} = \{b_1, \dots, b_n\}$. Note that $m_i \leq a_i$. If $\sum m_i < \sum a_i$, then H is divisible by $\sum x_i$, a contradiction. Accordingly, $m_i = a_i$, $c_i = b_i$ for every i . Moreover, all the $a_1 + \dots + a_n$ partial derivatives must be applied to the term $(x_1 + \dots + x_n)^{a_1 + \dots + a_n}$ in F . After all, we get

$$[x_1^{d_1} \dots x_n^{d_n}]F = \prod_{i=1}^n \kappa(C_i, b_i, a_i) \frac{\partial^{a_1 + \dots + a_n} F}{\partial x_1^{a_1} \dots \partial x_n^{a_n}}(b_1, \dots, b_n) = \binom{|\mathbf{a}|}{\mathbf{a}},$$

for $\prod_{i \neq j} (b_i - b_j)^{a_i} = \prod_i \prod_{c \in B \setminus \{b_i\}} (b_i - c)^{a_i}$. □

Remarks. 1. A connection between restricted sumsets and Morris's constant term identity was made recently by Zhou [55]. 2. Let $h_r(\mathbf{x}) = \sum_{1 \leq j_1 < \dots < j_r \leq n} x_{j_1} \dots x_{j_r}$, denote the complete symmetric function of degree r . Following Good's method [22] one gets the following generalization of (3.1), also implicit in [50]:

$$\text{CT} \left[x_1^{-ra_1} \dots x_n^{-ra_n} h_r(x_1, \dots, x_n)^{a_1 + \dots + a_n} \prod_{i \neq j} (1 - x_j/x_i)^{a_i} \right] = \binom{|\mathbf{a}|}{\mathbf{a}}.$$

It would be interesting to obtain a proof of this identity based on the Combinatorial Nullstellensatz.

4. On a problem of Kadell

The aforementioned idea of Aomoto led Kadell [29] to discover and prove the following Dyson-type identity. Fix $m < n$. For $1 \leq r \leq n$ and an m -element subset M of $\{1, 2, \dots, n\}$, consider the Laurent polynomial

$$\mathcal{K}_{r,M}(\mathbf{x}; \mathbf{a}) = \left(1 + \sum_{v \notin M} a_v\right) \prod_{s \in M} \left(1 - \frac{x_r}{x_s}\right) \mathcal{D}(\mathbf{x}; \mathbf{a}).$$

Note that $\mathcal{K}_{r,M}(\mathbf{x}; \mathbf{a}) = 0$ if $r \in M$. Then, according to [29, Theorem 1],

$$\text{CT} \left[\sum_{r=1}^n \sum_{|M|=m} \mathcal{K}_{r,M}(\mathbf{x}; \mathbf{a}) \right] = n \binom{n-1}{m} (1 + |\mathbf{a}|) \binom{|\mathbf{a}|}{\mathbf{a}}. \quad (4.1)$$

Kadell suggested that each non-zero function $\mathcal{K}_{r,M}(\mathbf{x}; \mathbf{a})$ must have the same contribution to the constant term. He formulated an even more general hypothesis (see [29, Conjecture 2]), which was established recently by Zhou [54] based on the first layer formulas for Dyson-coefficients [41, Theorem 1.7]. Kadell also suggested the following q -analogue of his hypothesis.

Conjecture 4.1 ([29, Conjecture 3]). *Let $M \subset \{1, 2, \dots, n\}$ and $\{r_s \mid s \in M\} \cap M = \emptyset$. Then*

$$\text{CT} \left[\prod_{1 \leq i < j \leq n} \left(\frac{x_i}{x_j} \right)_{a_i^*} \left(\frac{qx_j}{x_i} \right)_{a_j^*} \right] = \frac{1 - q^{1+|\mathbf{a}|}}{1 - q^{1 + \sum_{v \notin M} a_v}} \binom{|\mathbf{a}|}{\mathbf{a}},$$

where, with a slight abuse of notation, $a_i^* = a_i^*(j) = a_i + \chi(j \in M, i = r_j)$ and $a_j^* = a_j^*(i) = a_j + \chi(i \in M, j = r_i)$.

Zhou [54] pointed out that this conjecture already fails for $n = 3$, $|M| = m = 1$, and proved a meaningful q -analogue of [29, Conjecture 2], which is too technical to be recalled here in detail. Our main result in this section is the proof of the following special case of Conjecture 4.1 corresponding to $M = \{1, \dots, m\}$ and $r_s \equiv n$ that, unexpectedly, does not seem to be implied by Zhou's result.

Theorem 4.2. *Let $m < n$. Then*

$$\text{CT} \left[\prod_{1 \leq i < j \leq n} \binom{x_i}{x_j}_{a_i} \binom{qx_j}{x_i}_{a_j^*} \right] = \frac{1 - q^{1+|a|}}{1 - q^{1 + \sum_{v=m+1}^n a_v}} \begin{bmatrix} |a| \\ \mathbf{a} \end{bmatrix},$$

where $a_n^* = a_n + \chi(i \leq m)$ and $a_j^* = a_j$ otherwise.

Specializing at $q = 1$ and taking into account the symmetry of the Dyson product we obtain the following special case of [29, Conjecture 2], which already implies (4.1).

Corollary 4.3. *Let $m < n$, $M \subset \{1, \dots, n\}$ with $|M| = m$ and $r \in \{1, \dots, n\} \setminus M$. Then*

$$\text{CT} \left[\prod_{s \in M} \left(1 - \frac{x_r}{x_s} \right) \mathcal{D}(\mathbf{x}; \mathbf{a}) \right] = \frac{1 + |a|}{1 + \sum_{v \notin M} a_v} \begin{bmatrix} |a| \\ \mathbf{a} \end{bmatrix}.$$

As a final remark we mention that the $m = 1$ case of this corollary in conjunction with the Zeilberger–Bressoud theorem immediately implies Sills' [45, Theorem 1.1]: For $1 \leq r \neq s \leq n$,

$$\text{CT} [(x_r/x_s) \mathcal{D}(\mathbf{x}; \mathbf{a})] = \frac{-a_s}{1 + |a| - a_s} \begin{bmatrix} |a| \\ \mathbf{a} \end{bmatrix}.$$

In general, one may use the inclusion-exclusion principle to obtain a formula for the constant term of

$$\left(x_r^m / \prod_{s \in M} x_s \right) \mathcal{D}(\mathbf{x}; \mathbf{a}),$$

in agreement with [41, Theorem 1.7].

Proof of Theorem 4.2. Note that if $a_i = 0$ for some $i < n$, then we may omit all factors that include the variable x_i without affecting the constant term. Accordingly, we may assume that each a_i , with the possible exception of a_n , is a positive integer. Clearly the constant term equals the coefficient of the monomial

$$\prod_{i=1}^n x_i^{|a| - a_i + \chi(i \leq m)}$$

in the homogeneous polynomial

$$F(\mathbf{x}) = \prod_{1 \leq i < j \leq n} \left(\prod_{k=0}^{a_i-1} (x_j - x_i q^k) \times \prod_{k=1}^{a_j^*} (x_i - x_j q^k) \right),$$

where

$$a_j^* = \begin{cases} a_j + 1 & \text{if } j = n \text{ and } i \leq m, \\ a_j & \text{otherwise.} \end{cases}$$

To express this coefficient we apply Lemma 1.1 with $\mathbb{F} = \mathbb{Q}(q)$. Once again, the aim is to choose the sets C_i so that $F(\mathbf{c}) = 0$ for all but one element $\mathbf{c} \in C_1 \times \dots \times C_n$. This can be easily achieved as follows. Let $C_i = \{q^{a_i} \mid a_i \in B_i\}$, where

$$B_i = \{0, 1, \dots, |a| - a_i + \chi(i \leq m)\}.$$

The sets C_i clearly have the right cardinalities. Now assume that $c_i = q^{\alpha_i} \in C_i$ and $F(\mathbf{c}) \neq 0$. Then all the α_i are distinct. Moreover,

$$\alpha_j \geq \alpha_i + a_i + \chi(j < i) + \chi(i = n, j \leq m)$$

holds for $\alpha_j > \alpha_i$. Next consider the unique permutation $\pi \in \mathfrak{S}_n$ for which

$$0 \leq \alpha_{\pi(1)} < \alpha_{\pi(2)} < \cdots < \alpha_{\pi(n)} \leq |\mathbf{a}| - a_{\pi(n)} + \chi(\pi(n) \leq m).$$

We obtain the chain of inequalities

$$|\mathbf{a}| - a_{\pi(n)} = \sum_{i=1}^{n-1} a_{\pi(i)} \leq \sum_{i=1}^{n-1} (\alpha_{\pi(i+1)} - \alpha_{\pi(i)}) = \alpha_{\pi(n)} - \alpha_{\pi(1)} \leq |\mathbf{a}| - a_{\pi(n)} + 1.$$

Notice that the first inequality is strict if π is not the identity permutation, while the second inequality is strict if $\pi(n) > m$. Suppose that $\pi(n) \neq n$. Since $\pi \neq \text{id}$, it must be $\pi(n) \leq m$. Consider the index i with $\pi(i) = n$. Then $\alpha_{\pi(i+1)} - \alpha_{\pi(i)} = a_{\pi(i)} + 1$, which implies $\pi(i+1) > m$. Therefore there must be an index $i+1 \leq j < n$ such that $\pi(j) > m$ and $\pi(j+1) \leq m$. For such a j we have $\alpha_{\pi(j+1)} - \alpha_{\pi(j)} \geq a_{\pi(j)} + 1$, resulting in

$$|\mathbf{a}| - a_{\pi(n)} = \sum_{i=1}^{n-1} a_{\pi(i)} \leq \sum_{i=1}^{n-1} (\alpha_{\pi(i+1)} - \alpha_{\pi(i)}) - 2 \leq |\mathbf{a}| - a_{\pi(n)} - 1,$$

a contradiction. Thus we can conclude that $\pi(n) = n$, implying $\pi = \text{id}$ and $\alpha_i = a_1 + \cdots + a_{i-1}$ for every i .

It only remains to substitute these values into

$$\frac{F(c_1, c_2, \dots, c_n)}{\phi'_1(c_1)\phi'_2(c_2)\cdots\phi'_n(c_n)},$$

which is quite a routine calculation. Therefore we only recall that substituting the same values in the same formula working with

$$F(\mathbf{x}) = \prod_{1 \leq i < j \leq n} \left(\prod_{k=0}^{a_i-1} (x_j - x_i q^k) \times \prod_{k=1}^{a_j} (x_i - x_j q^k) \right)$$

and $B_i = \{0, 1, \dots, |\mathbf{a}| - a_i\}$ yields the q -Dyson constant term $\text{CT}[\mathcal{D}_q(\mathbf{x}; \mathbf{a})]$, see [37]. The changes are easily detected, and noting $\alpha_i + a_i = \alpha_{i+1}$, $\alpha_n + a_n = |\mathbf{a}|$ we find that the constant term in question is indeed

$$\prod_{i=1}^m \frac{q^{\alpha_i} - q^{\alpha_n + a_n + 1}}{q^{\alpha_i} - q^{|\mathbf{a}| - a_i + 1}} \left[\begin{matrix} |\mathbf{a}| \\ \mathbf{a} \end{matrix} \right] = \frac{1 - q^{1+|\mathbf{a}| - \alpha_1}}{1 - q^{1+|\mathbf{a}| - \alpha_{m+1}}} \left[\begin{matrix} |\mathbf{a}| \\ \mathbf{a} \end{matrix} \right],$$

as claimed. □

5. A new proof of the Selberg integral

Due to its equivalence to the Selberg integral, it will be enough to establish Morris's constant term identity (1.1). Making the Laurent polynomial homogeneous by the introduction of a new variable does not affect the constant term. Thus, we are to determine the constant term of the Laurent polynomial

$$\mathcal{M}(x_0, \mathbf{x}; a, b, k) := \prod_{j=1}^n \left(1 - \frac{x_j}{x_0}\right)^a \left(1 - \frac{x_0}{x_j}\right)^b \prod_{1 \leq i \neq j \leq n} \left(1 - \frac{x_i}{x_j}\right)^k,$$

which is the same as the coefficient of $x_0^{na} \prod_{i=1}^n x_i^{(n-1)k+b}$ in the homogeneous polynomial

$$\prod_{j=1}^n (x_0 - x_j)^a (x_j - x_0)^b \times \prod_{1 \leq i \neq j \leq n} (x_j - x_i)^k.$$

As in Section 3, we modify this polynomial without affecting this leading coefficient and consider

$$F(x_0, \mathbf{x}) = \prod_{j=1}^n \prod_{e=-a}^{b-1} (x_j - x_0 - e) \times \prod_{1 \leq i < j \leq n} \prod_{e=-k}^{k-1} (x_j - x_i - e). \quad (5.1)$$

To apply Lemma 2.4 efficiently, we choose sets $C_i = \{0, 1, \dots, (n-1)k + b\}$ for $1 \leq i \leq n$ and multiset

$$C_0 = \{0\} \cup \bigcup_{\ell=0}^{n-1} \{k\ell + 1, k\ell + 2, \dots, k\ell + a\}.$$

They have the right cardinality, the latter one being an ordinary set if $k \geq a$. Consider $c_i \in C_i$ and $m_i < \omega_i(c_i)$. Note that $m_i = 0$ for $1 \leq i \leq n$. We proceed to prove that

$$\frac{\partial^{m_0 + \dots + m_n} F}{\partial x_0^{m_0} \dots \partial x_n^{m_n}}(c_0, \dots, c_n) = \frac{\partial^{m_0} F}{\partial x_0^{m_0}}(c_0, \dots, c_n) = 0$$

for all but one such selection.

Lemma 5.1. *If $c_0 \neq 0$, then*

$$\frac{\partial^{m_0} F}{\partial x_0^{m_0}}(c_0, \dots, c_n) = 0. \quad (5.2)$$

Proof. Write $S_\ell = \{k\ell + 1, k\ell + 2, \dots, k\ell + a\}$. Since $m_0 < \omega_0(c_0)$, there is an index $0 \leq u \leq n - \omega_0(c_0)$ such that $c_0 \in S_u \cap S_{u+1} \cap \dots \cap S_{u+m_0}$. That is,

$$(u + m_0)k + 1 \leq c_0 \leq uk + a.$$

Accordingly, if c_j lies in the interval $[uk, (u + m_0)k + b]$ for some $1 \leq i \leq n$, then $c_0 - a \leq c_j \leq c_0 + b - 1$ and there is a term of the form $x_j - x_0 - e$ in F which attains 0 when evaluated at the point (c_0, \mathbf{c}) . It follows, that (5.2) holds if more than m_0 of such c_i lie in the interval $[uk, (u + m_0)k + b]$.

Otherwise either at least $u + 1$ of c_1, \dots, c_n lie in the interval $[0, ku - 1]$, or at least $n - m_0 - u$ of them lie in the interval $[(u + m_0)k + b + 1, (n - 1)k + b]$. In either case there is a pair of indices $1 \leq i < j \leq n$ such that $|c_j - c_i| < k$, meaning that there is a term of the form $x_j - x_i - e$ in F which attains 0 when evaluated at the point (c_0, \mathbf{c}) , and once again we arrive at (5.2). \square

Thus we only have to consider the case when $c_0 = 0$; then $\omega_0(c) = 1$ and $m_0 = 0$. If

$$\frac{\partial^{m_0} F}{\partial x_0^{m_0}}(c_0, \dots, c_n) = F(c_0, \mathbf{c}) \neq 0,$$

then $c_1, \dots, c_n \in [b, (n-1)k + b]$ and $|c_j - c_i| \geq k$ for each pair $1 \leq i < j \leq n$. Therefore the numbers c_1, \dots, c_n , in some order, must coincide with the numbers $b, k + b, \dots, (n-1)k + b$. Moreover it must be the natural order, for if $c_i > c_j$ for some $i < j$, then $c_i - c_j \geq k + 1$. It only remains to evaluate

$$\prod_{i=0}^n \kappa(C_i, c_i, 0) F(c_0, \mathbf{c})$$

at the point $(c_0, \mathbf{c}) = (0, b, k + b, \dots, (n-1)k + b)$. Since $\omega_i(c_i) = 0$ for each i , we simply have

$$\kappa(C_i, c_i, 0) = \frac{1}{\prod_{c \in C_i \setminus \{c_i\}} (c_i - c)^{\omega_i(c)}}$$

and one easily recovers (1.1).

Remark. For the sake of simplicity, we tacitly assumed that the parameters a, b, k are positive integers. It is not difficult to modify the above proof to suit the remaining cases and we leave it to the reader. Alternatively, one can easily reduce the $k = 0$ case to the Chu-Vandermonde identity, whereas the $\min\{a, b\} = 0$ case is just the equal parameter case of Dyson's identity.

6. Interlude

Replace the polynomial in (5.1) by

$$\prod_{j=1}^n \prod_{e=1}^a (x_0 - q^e x_j) \prod_{e=0}^{b-1} (x_j - q^e x_0) \times \prod_{1 \leq i < j \leq n} \prod_{e=0}^{k-1} (x_j - q^e x_i) \prod_{e=1}^k (x_i - q^e x_j).$$

Also replace the multisets C_i by multisets which consist of powers of q whose exponents belong to C_i , and with the same multiplicities. Repeating the proof given in the previous section almost verbatim one obtains without any difficulty the following version of the q -Morris constant term identity:

$$\text{CT} \left[\prod_{j=1}^n (qx_j)_a (1/x_j)_b \mathcal{D}_q(\mathbf{x}; k) \right] = \prod_{j=0}^{n-1} \frac{(q)_{a+b+kj} (q)_{kj+k}}{(q)_{a+kj} (q)_{b+kj} (q)_k}.$$

Although the identity conjectured in Morris's thesis [42] reads slightly differently as

$$\text{CT} \left[\prod_{j=1}^n (x_j)_a (q/x_j)_b \mathcal{D}_q(\mathbf{x}; k) \right] = \prod_{j=0}^{n-1} \frac{(q)_{a+b+kj} (q)_{kj+k}}{(q)_{a+kj} (q)_{b+kj} (q)_k}, \quad (6.1)$$

the two are easily seen to be equivalent, for each monomial of degree zero has the same coefficient in the Laurent polynomials $\prod_{j=1}^n (qx_j)_a (1/x_j)_b$ and $\prod_{j=1}^n (x_j)_a (q/x_j)_b$. Morris's conjecture was established independently in [24] and [28] via the proof of a q -Selberg integral proposed by Askey [6], followed by a more elementary proof in [52].

The above argument relates to the one given in the previous section in a similar way as the derivation of the q -analogue of Dyson's conjecture in [37] relates to the original version of Karasev and Petrov's proof [35] for the Dyson product. One may say that applications of Lemma 1.1 (or its generalization Theorem 2.4) allows one to prove an appropriate q -analogue practically along the same lines as the original identity, even without the need to modify the corresponding polynomial. This works also the other way around: the way (6.1) is formulated gives a hint of an alternative proof of (1.1) which involves a slightly different modification along with a slightly different choice of the multisets C_i . Our preference was given to the modification, which allowed a more simple choice for the C_i as well as to keep the natural order of the variables x_0, x_1, \dots, x_n for the q -analogue in the following sense.

All the constant term identities and their q -analogues studied in this paper can be formulated in the following context. Let $\mathbf{B} = (\beta_{ij})$ denote an $(n+1) \times (n+1)$ matrix with rows and columns numbered from 0 to n , corresponding to the natural order of the variables. It is assumed that the entries are non-negative integers and all the diagonal entries are zero. Associated to such a matrix is the Laurent polynomial

$$\mathcal{L}(x_0, \mathbf{x}; \mathbf{B}) = \prod_{0 \leq i \neq j \leq n} \left(1 - \frac{x_i}{x_j} \right)^{\beta_{ij}}$$

and its q -analogue

$$\mathcal{L}_q(x_0, \mathbf{x}; \mathbf{B}) = \prod_{0 \leq i < j \leq n} \left(\frac{x_i}{x_j} \right)_{\beta_{ij}} \left(\frac{qx_j}{x_i} \right)_{\beta_{ji}}.$$

Thus, one can write $\mathcal{D}(\mathbf{x}; \mathbf{a}) = \mathcal{L}(x_0, \mathbf{x}; \mathbf{B}_{\mathcal{D}})$ and $\mathcal{M}(x_0, \mathbf{x}; a, b, k) = \mathcal{L}(x_0, \mathbf{x}; \mathbf{B}_{\mathcal{M}})$ with the matrices

$$\mathbf{B}_{\mathcal{D}} = \left(\begin{array}{c|cccc} 0 & 0 & 0 & 0 & \dots & 0 \\ \hline 0 & 0 & a_1 & a_1 & \dots & a_1 \\ 0 & a_2 & 0 & a_2 & \dots & a_2 \\ 0 & a_3 & a_3 & 0 & \dots & a_3 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & a_n & a_n & a_n & \dots & 0 \end{array} \right) \quad \text{and} \quad \mathbf{B}_{\mathcal{M}} = \left(\begin{array}{c|cccc} 0 & b & b & b & \dots & b \\ \hline a & 0 & k & k & \dots & k \\ a & k & 0 & k & \dots & k \\ a & k & k & 0 & \dots & k \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ a & k & k & k & \dots & 0 \end{array} \right)$$

corresponding to the Dyson resp. Morris constant term identities, whereas $\mathcal{D}_q(\mathbf{x}; \mathbf{a}) = \mathcal{L}_q(x_0, \mathbf{x}; \mathbf{B}_{\mathcal{D}})$. Note that simultaneous permutation of the rows and columns of \mathbf{B} according to the same element of \mathfrak{S}_{n+1} has no effect on $\text{CT}[\mathcal{L}(x_0, \mathbf{x}; \mathbf{B})]$. Generally it is not the case for $\text{CT}[\mathcal{L}_q(x_0, \mathbf{x}; \mathbf{B})]$, but as we explained in relation to the q -Morris identity, one may always apply the cyclic permutation

$$n \rightarrow n-1 \rightarrow \dots \rightarrow 1 \rightarrow 0 \rightarrow n$$

or any of its powers without affecting the constant term.

Theorem 4.2 concerns $\text{CT}[\mathcal{L}_q(x_0, \mathbf{x}; \mathbf{B}_{\mathcal{K}})]$ for the matrix

$$\mathbf{B}_{\mathcal{K}} = \left(\begin{array}{c|cccc|cccc} 0 & 0 & 0 & \dots & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & a_1 & \dots & a_1 & a_1 & \dots & a_1 & a_1 \\ 0 & a_2 & 0 & \dots & a_2 & a_2 & \dots & a_2 & a_2 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & a_m & a_m & \dots & 0 & a_m & \dots & a_m & a_m \\ \hline 0 & a_{m+1} & a_{m+1} & \dots & a_{m+1} & 0 & \dots & a_{m+1} & a_{m+1} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & a_{n-1} & a_{n-1} & \dots & a_{n-1} & a_{n-1} & \dots & 0 & a_{n-1} \\ 0 & a_n + 1 & a_n + 1 & \dots & a_n + 1 & a_n & \dots & a_n & 0 \end{array} \right).$$

Applying the above mentioned cyclic permutations to $\mathbf{B}_{\mathcal{K}}$, after rearranging indices we obtain the following more general form of Theorem 4.2.

Theorem 6.1. Fix an arbitrary integer $r \in \{1, 2, \dots, n\}$. Then Conjecture 4.1 is valid with the choice of $M = \{r + 1, \dots, r + m\}$ and $r_s \equiv r$, where indices are understood modulo n .

Aomoto's identity (1.2) and Forrester's conjecture are related to the matrices

$$\mathbf{B}_{\mathcal{A}} = \left(\begin{array}{c|ccc|ccc} 0 & b & \dots & b & b & \dots & b \\ a & 0 & \dots & k & k & \dots & k \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ a & k & \dots & 0 & k & \dots & k \\ \hline a+1 & k & \dots & k & 0 & \dots & k \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ a+1 & k & \dots & k & k & \dots & 0 \end{array} \right) \quad \text{and} \quad \mathbf{B}_{\mathcal{F}} = \left(\begin{array}{c|ccc|ccc} 0 & b & \dots & b & b & \dots & b \\ a & 0 & \dots & k & k & \dots & k \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ a & k & \dots & 0 & k & \dots & k \\ \hline a & k & \dots & k & 0 & \dots & k+1 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ a & k & \dots & k & k+1 & \dots & 0 \end{array} \right),$$

where the last m resp. $n - n_0$ rows/columns are separated. In the first case we rearranged the matrix so that a q -analogue can be formulated within our framework. Our main result concerns the overlay of these matrices when $m \geq n - n_0$, that is, the matrix

$$\mathbf{B}_{\mathcal{AF}} = \left(\begin{array}{c|ccc|ccc|ccc} 0 & b & \dots & b & b & \dots & b & b & \dots & b \\ a & 0 & \dots & k & k & \dots & k & k & \dots & k \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ a & k & \dots & 0 & k & \dots & k & k & \dots & k \\ \hline a+1 & k & \dots & k & 0 & \dots & k & k & \dots & k \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ a+1 & k & \dots & k & k & \dots & 0 & k & \dots & k \\ \hline a+1 & k & \dots & k & k & \dots & k & 0 & \dots & k+1 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ a+1 & k & \dots & k & k & \dots & k & k+1 & \dots & 0 \end{array} \right) \begin{array}{c} 0 \\ \hline 1 \\ \vdots \\ \hline n-m \\ \hline n-m+1 \\ \vdots \\ \hline n_0 \\ \hline n_0+1 \\ \vdots \\ \hline n \end{array}.$$

Theorem 6.2. *Let n be a positive integer. For arbitrary nonnegative integers a, b, k and $m, n_0 \leq n \leq m + n_0$,*

$$\text{CT}[\mathcal{L}_q(x_0, \mathbf{x}; \mathbf{B}_{\mathcal{AF}})] = \prod_{j=0}^{n-1} \frac{(q)_{a+b+kj+\chi(j>n_0)(j-n_0)+\chi(j \geq n-m)}(q)_{kj+\chi(j>n_0)(j-n_0)+k}}{(q)_{a+kj+\chi(j>n_0)(j-n_0)+\chi(j \geq n-m)}(q)_{b+kj+\chi(j>n_0)(j-n_0)}(q)_k} \times \prod_{j=1}^{n-n_0} \frac{1-q^{(k+1)j}}{1-q^{k+1}}.$$

When $m = 0$, this proves Baker and Forrester's [7, Conjecture 2.1], and further specializing at $q = 1$, Forrester's original conjecture as well. The $n_0 = n$ case gives the following q -analogue of Aomoto's identity.

Corollary 6.3. *Let n be a positive integer. For arbitrary nonnegative integers a, b, k and $m \leq n$,*

$$\text{CT}[\mathcal{L}_q(x_0, \mathbf{x}; \mathbf{B}_{\mathcal{A}})] = \prod_{j=0}^{n-1} \frac{(q)_{a+b+kj+\chi(j \geq n-m)}(q)_{kj+k}}{(q)_{a+kj+\chi(j \geq n-m)}(q)_{b+kj}(q)_k}.$$

A more general version of this identity, which involves an additional parameter attached to b , was established in Kadell's paper [28]. An elementary proof was claimed recently by Xin and Zhou [51]. Replacing k by $k + 1$ we obtain that Theorem 6.2 is valid for arbitrary $m \leq n$ when $n_0 = 0$. Although the condition $n \leq m + n_0$ is crucial to our proof given in the next section, it does not seem to be necessary.

Conjecture 6.4. *Theorem 6.2 remains valid without the restriction $n \leq m + n_0$.*

7. Proof of the conjecture of Forrester

Clearly $\text{CT}[\mathcal{L}_q(x_0, \mathbf{x}; \mathbf{B})]$ equals the coefficient of $\prod_j x_j^{B_j}$, where $B_j = \sum_i \beta_{ij}$, in the polynomial

$$F_q(x_0, \mathbf{x}; \mathbf{B}) := \prod_{0 \leq i < j \leq n} \left(\prod_{t=0}^{\beta_{ij}-1} (x_j - q^t x_i) \times \prod_{t=1}^{\beta_{ji}} (x_i - q^t x_j) \right).$$

Claim 7.1. *Suppose that $c_i = q^{\alpha_i}$ for some integers α_i such that $F_q(c_0, \mathbf{c}; \mathbf{B}) \neq 0$. Let $j > i$. Then $\alpha_j \geq \alpha_i$ implies $\alpha_j \geq \alpha_i + \beta_{ij}$, and $\alpha_i > \alpha_j$ implies $\alpha_i \geq \alpha_j + \beta_{ji} + 1$. Both statements are valid even if the corresponding entry in \mathbf{B} is zero. The same is true with F_q replaced by any of its partial derivatives in which $m_i = m_j = 0$. \square*

We are to apply Theorem 2.4 with the polynomial $F = F_q(\cdot; \mathbf{B}_{\mathcal{AF}})$. As in Section 5, we will assume that the parameters a, b, k are positive integers and leave the rest to the reader.

7.1. The choice for the multisets C_i

Write $\gamma_i = \beta_{in}$ for $0 \leq i < n$ and let $\Delta_t = \sum_{i=0}^t \gamma_i$. Thus,

$$\gamma_0 = b, \gamma_1 = \dots = \gamma_{n_0} = k, \gamma_{n_0+1} = \dots = \gamma_{n-1} = k + 1$$

and $\beta_{ij} = \gamma_{\min\{i,j\}}$ for $1 \leq i \neq j \leq n$. Consider the intervals $I_t = [\Delta_t - \gamma_t + 1, \Delta_t] = [\Delta_{t-1} + 1, \Delta_t]$, where here and thereafter $[u, v]$ stands for the set of integers ℓ satisfying $u \leq \ell \leq v$. The intervals $I_0 := [0, b], I_1, \dots, I_{n-1}$ are mutually disjoint. The multisets C_i are defined in the form $C_i = \{q^\alpha \mid \alpha \in A_i\}$, where for $1 \leq j \leq n$

$$A_j = \{0\} \cup \bigcup_{t=0}^{n-1} [\Delta_t - \gamma_{\min\{t,j\}} + 1, \Delta_t] \subseteq \bigcup_{t=0}^{n-1} I_t = [0, \Delta_{n-1}]$$

is an ordinary set and

$$A_0 = \{0\} \cup \bigcup_{t=0}^{n-1} [\Delta_t - b + 1, \Delta_t - b + \beta_{t+1,0}]$$

is a multiset. Then $|C_i| = |A_i| = B_i + 1$ holds for every $0 \leq i \leq n$. We are to show that

$$\frac{\partial^{m_0+\dots+m_n} F}{\partial x_0^{m_0} \dots \partial x_n^{m_n}}(c_0, \dots, c_n) = \frac{\partial^{m_0} F}{\partial x_0^{m_0}}(c_0, \dots, c_n) = 0$$

for all but one selection of elements $c_i \in C_i$ and multiplicities $m_i < \omega_i(c_i)$, namely when $c_0 = 1$, $c_i = q^{\Delta_{i-1}}$ for $1 \leq i \leq n$, and all the multiplicities are zero.

7.2. The combinatorics

Consider such a selection and write $c_i = q^{\alpha_i}$. Note that $\omega_1(c_1) = \cdots = \omega_n(c_n) = \omega_0(q^0) = 1$. The above statement is verified by the juxtaposition of the following two lemmas.

Lemma 7.2. *Let $\alpha_0 = 0$. If $F(c_0, c_1, \dots, c_n) \neq 0$, then $\alpha_i = \Delta_{i-1}$ for every $1 \leq i \leq n$.*

Lemma 7.3. *If $\alpha_0 \neq 0$, then $(\partial^{m_0} F / \partial x_0^{m_0})(c_0, \dots, c_n) = 0$.*

One key to each is the following consequence of Claim 7.1.

Lemma 7.4. *Suppose that $(\partial^{m_0} F / \partial x_0^{m_0})(c_0, \dots, c_n) \neq 0$. Then for every $1 \leq t \leq n-1$ there is at most one index $1 \leq i \leq n$ such that $\alpha_i \in I_t$.*

Proof. Assume that, on the contrary, there is a pair $1 \leq i \neq j \leq n$ such that $\alpha_i, \alpha_j \in I_t$. Let $\alpha_j \geq \alpha_i$, then it follows from Claim 7.1 that $\alpha_j - \alpha_i \geq k$. The length of I_t is $\gamma_t \in \{k, k+1\}$. Thus, it must be $\gamma_t = k+1$, $\alpha_i = \Delta_t - k$ and $\alpha_j = \Delta_t$. Consequently, $t > n_0$, $i < j$ and $i \leq n_0$. Therefore $\Delta_t - \gamma_{\min\{t, i\}} + 1 = \Delta_t - k + 1$ and $\alpha_i \notin A_i$, a contradiction. \square

Proof of Lemma 7.2. For every $1 \leq i \leq n$ we have $\alpha_i \geq \alpha_0$, therefore $\alpha_i \geq \beta_{0i} = b$ by Claim 7.1. Moreover, $k > 0$ implies that $\alpha_1, \dots, \alpha_n$ are all distinct, thus it follows from Lemma 7.4 that each of the intervals I_0, I_1, \dots, I_{n-1} contains precisely one of them. Let $\pi \in \mathfrak{S}_n$ denote the unique permutation for which $\alpha_{\pi(1)} < \cdots < \alpha_{\pi(n)}$, then $\alpha_{\pi(i)} \in I_{i-1}$. By Claim 7.1 we have

$$\alpha_{\pi(i+1)} \geq \alpha_{\pi(i)} + \beta_{\pi(i), \pi(i+1)} + \chi(\pi(i) > \pi(i+1)).$$

Consequently,

$$\alpha_{\pi(n_0+1)} \geq b + kn_0 + \sum_{i=1}^{n_0} \chi(\pi(i) > \pi(i+1)) \geq \Delta_{n_0} + \sum_{i=1}^{n_0} \chi(\pi(i) > \pi(i+1)).$$

Since $\alpha_{\pi(n_0+1)} \leq \Delta_{n_0}$, it follows that $\alpha_{\pi(1)} = b$, $\pi(1) < \cdots < \pi(n_0+1)$, and $\beta_{\pi(i), \pi(i+1)} = k$ for $1 \leq i \leq n_0$. This in turn implies that $\pi(n_0) \leq n_0$, thus $\pi(i) = i$ and $\alpha_i = \Delta_{i-1}$ for $1 \leq i \leq n_0$.

Now for $n_0 < i < n$ we have $\pi(i), \pi(i+1) > n_0$ and thus $\beta_{\pi(i), \pi(i+1)} = k+1$. Restricting π to the set $[n_0+1, n]$ and starting with $\alpha_{\pi(n_0+1)} = \Delta_{n_0}$, a similar argument completes the proof. \square

Proof of Lemma 7.3. Assume that, contrary to the statement, $(\partial^{m_0} F / \partial x_0^{m_0})(c_0, \dots, c_n) \neq 0$. Write $S_t = [\Delta_t - b + 1, \Delta_t - b + \beta_{t+1,0}]$. Since $\alpha_0 \neq 0$ and $m_0 < \omega_0(c_0)$, there is an index $0 \leq u \leq n - \omega_0(c_0)$ such that $\alpha_0 \in S_u \cap S_{u+1} \cap \cdots \cap S_{u+m_0}$. That is,

$$\Delta_{u+m_0} - b + 1 \leq \alpha_0 \leq \Delta_u - b + \beta_{u+1,0}.$$

Accordingly, if α_j lies in the interval

$$T_{uj} = [\Delta_u - b + \beta_{u+1,0} - \beta_{j0}, \Delta_{u+m_0}]$$

for some $1 \leq j \leq n$, then $\alpha_0 - \beta_{j0} \leq \alpha_j \leq \alpha_0 + \beta_{0j} - 1$ and there is a term of the form $x_j - q^t x_0$ or $x_0 - q^t x_j$ in F which attains 0 when evaluated at the point (c_0, \mathbf{c}) . There cannot be more than m_0 such terms. It is implied by Lemma 7.4 that at most $n-1-u-m_0$ of the distinct numbers $\alpha_1, \dots, \alpha_n$ can lie in the interval $[\Delta_{u+m_0} + 1, \Delta_{n-1}]$.

It follows that at least $u+1$ of the numbers α_j satisfy $\alpha_j \leq \Delta_u - b + \beta_{u+1,0} - \beta_{j0} - 1$. This is clearly impossible if $u+1 \leq n-m$, for then $\Delta_u - b + \beta_{u+1,0} - \beta_{j0} - 1 \leq uk - 1$ in view of $n-m \leq n_0$, and on the other hand the difference between any two such α_j is at least k in view of Claim 7.1. Thus, $u \geq n-m$ and $\beta_{u+1,0} = a+1$. Consider

$$\alpha_{\nu(1)} < \cdots < \alpha_{\nu(u+1)} \leq \Delta_u - b + \beta_{u+1,0} - \beta_{\nu(u+1),0} - 1 \leq \Delta_u - b.$$

If $u \leq n_0$, then it must be $\alpha_{\nu(i)} = (i-1)k$ and $\nu(1) < \cdots < \nu(u+1)$, but then $\nu(u+1) \geq u+1 > n-m$, $\beta_{\nu(u+1),0} = a+1$, implying $\alpha_{\nu(u+1)} \in T_{u, \nu(u+1)}$, which is absurd. This means that $u \geq n_0+1$. It is easy to see that $\alpha_{\nu(i+1)} - \alpha_{\nu(i)} \geq \gamma_{\nu(i)}$ for $i \leq u$, thus $\alpha_{\nu(u+1)} \geq \sum_{i=1}^u \gamma_{\nu(i)} \geq \Delta_u - b$. Therefore $\sum_{i=1}^u \gamma_{\nu(i)} = \Delta_u - b$, which implies that $\{\nu(1), \dots, \nu(u)\} \supseteq \{1, \dots, n_0\}$. Consequently, $\nu(u+1) \geq n_0+1 > n-m$, which leads to a contradiction as before. \square

7.3. The computation

It only remains to evaluate

$$\frac{F_q(q^0, q^{\Delta_0}, \dots, q^{\Delta_{n-1}}; \mathbf{B})}{\psi_0 \psi_1 \dots \psi_n}, \quad (7.1)$$

where

$$\psi_j = \prod_{\alpha \in A_j \setminus \{\Delta_{j-1}\}} (q^{\Delta_{j-1}} - q^\alpha)$$

for $j = 1, \dots, n$, and with the shorthand notation $\Delta_u^v = \gamma_u + \dots + \gamma_v = \Delta_v - \Delta_{u-1}$,

$$\psi_0 = \prod_{t=0}^{n-1} \prod_{\alpha=\Delta_t'+1}^{\Delta_t'+\beta_{t+1,0}} (1 - q^\alpha) = \prod_{j=1}^n [\Delta_1^{j-1} + 1, \Delta_1^{j-1} + \beta_{j0}]_q. \quad (7.2)$$

From now on, $[u, v]_q := (1 - q^u) \dots (1 - q^v) = (q)_v / (q)_{u-1}$, with $[u, u]_q$ abbreviated as $[u]_q$. Both the numerator and the denominator in (7.1) is the product of factors in the form $\pm q^u (1 - q^v)$ with some non-negative integers u, v . More precisely, collecting factors of a similar nature together we find that the numerator is the product of the factors

$$(-1)^{\gamma_0} \times q^{1+\dots+(\gamma_0-1)} \times [\Delta_1^{j-1} + 1, \Delta_0^{j-1} + \beta_{j0}]_q \quad \text{for } 1 \leq j \leq n, \quad (7.3)$$

$$(-1)^{\gamma_i} \times q^{\Delta_{i-1}+\dots+(\Delta_{i-1}+\gamma_i-1)} \times [\Delta_i^{j-1} - \gamma_i + 1, \Delta_i^{j-1}]_q \quad \text{for } 1 \leq i < j \leq n, \quad (7.4)$$

and

$$q^{\gamma_i \Delta_{i-1}} \times [\Delta_i^{j-1} + 1, \Delta_i^{j-1} + \gamma_i]_q \quad \text{for } 1 \leq i < j \leq n. \quad (7.5)$$

In the denominator, besides (7.2) we have the factors

$$(-1) \times [\Delta_{j-1}]_q \times \psi_{j<} \times \psi_{j=} \times \psi_{j>} \quad \text{for } 1 \leq j \leq n, \quad (7.6)$$

where

$$\psi_{j<} = \prod_{t=0}^{j-2} (-1)^{\gamma_t} \times q^{(\Delta_t - \gamma_t + 1) + \dots + \Delta_t} \times [\Delta_{t+1}^{j-1}, \Delta_{t+1}^{j-1} + \gamma_t - 1]_q, \quad (7.7)$$

$$\psi_{j=} = (-1)^{\gamma_{j-1}-1} \times q^{(\Delta_{j-1} - \gamma_{j-1} + 1) + \dots + (\Delta_{j-1} - 1)} \times [1, \gamma_{j-1} - 1]_q, \quad (7.8)$$

and

$$\psi_{j>} = \prod_{t=j}^{n-1} q^{\gamma_t \Delta_{j-1}} \times [\Delta_t^j - \gamma_j + 1, \Delta_t^j]_q. \quad (7.9)$$

Now the powers of -1 and q cancel out due to the simple identity

$$n\gamma_0 + \sum_{1 \leq i < j \leq n} \gamma_i = n + \sum_{0 \leq t < j-1 \leq n-1} \gamma_t + \sum_{1 \leq j \leq n} (\gamma_{j-1} - 1)$$

and the somewhat more subtle

$$\begin{aligned} n \binom{\gamma_0}{2} + \sum_{1 \leq i < j \leq n} \left(2\gamma_i \Delta_{i-1} + \binom{\gamma_i}{2} \right) \\ = \sum_{0 \leq t < j-1 \leq n-1} \left(\gamma_t \Delta_t - \binom{\gamma_t}{2} \right) + \sum_{j=1}^n \left((\gamma_{j-1} - 1) \Delta_{j-1} - \binom{\gamma_{j-1} - 1}{2} \right) + \sum_{0 \leq j-1 < t \leq n-1} \gamma_j \Delta_{j-1}. \end{aligned}$$

It remains to deal with the factors of the form $[u, v]_q$. Those from (7.4) and (7.9) cancel out. Those from (7.3) and (7.2) yield

$$\prod_{j=1}^n \frac{(q)_{\Delta_0^{j-1} + \beta_{j0}}}{(q)_{\Delta_1^{j-1} + \beta_{j0}}} = \prod_{j=0}^{n-1} \frac{(q)_{a+b+kj+\chi(j>n_0)(j-n_0)+\chi(j \geq n-m)}}{(q)_{a+kj+\chi(j>n_0)(j-n_0)+\chi(j \geq n-m)}}. \quad (7.10)$$

As for the rest, the contribution from (7.5) and (7.7) with the substitution $t + 1 = i$ gives

$$\begin{aligned}
\prod_{1 \leq i < j \leq n} \frac{[\Delta_i^{j-1} + 1, \Delta_i^{j-1} + \gamma_i]_q}{[\Delta_i^{j-1}, \Delta_i^{j-1} + \gamma_{i-1} - 1]_q} &= \prod_{1 \leq i < j \leq n} \frac{[\Delta_i^{j-1}, \Delta_i^{j-1} + \gamma_i]_q \cdot [\Delta_i^{j-1} + \gamma_{i-1}]_q}{[\Delta_i^{j-1}, \Delta_i^{j-1} + \gamma_{i-1}]_q \cdot [\Delta_i^{j-1}]_q} \\
&= \prod_{j=2}^n \frac{[\Delta_1^{j-1}, \Delta_1^{j-1} + \gamma_1]_q}{[\Delta_1^{j-1}, \Delta_1^{j-1} + \gamma_0]_q} \times \Psi \times \prod_{1 \leq i < j \leq n} \frac{[\Delta_{i-1}^{j-1}]_q}{[\Delta_i^{j-1}]_q} \\
&= \prod_{j=2}^n \frac{(q)_{\Delta_1^{j-1} + \gamma_1}}{(q)_{\Delta_1^{j-1} + \gamma_0}} \times \Psi \times \prod_{j=2}^n \frac{[\Delta_{j-1}]_q}{1 - q^{\gamma_{j-1}}}
\end{aligned} \tag{7.11}$$

in the first place, where the factor

$$\Psi = \prod_{j=n_0+2}^n [\Delta_{n_0+1}^{j-1} + \gamma_{n_0+1}]_q = \prod_{j=2}^{n-n_0} (1 - q^{(k+1)j})$$

only occurs when $n_0 > 0$. Combining (7.11) with the contribution of the factors $[\Delta_{j-1}]_q = 1 - q^{\Delta_{j-1}}$ from (7.6) and the factors $[1, \gamma_{j-1} - 1]_q = (q)_{\gamma_{j-1}-1}$ from (7.8), shifting indices we obtain

$$\prod_{j=1}^{n-1} \frac{(q)_{\Delta_1^j + \gamma_1}}{(q)_{\Delta_1^j + \gamma_0}} \times \prod_{j=0}^{n-1} \frac{1}{(q)_{\gamma_j}} \times \left(\prod_{j=2}^{n-n_0} (1 - q^{(k+1)j}) \right)^{\chi(n_0 > 0)},$$

in agreement with

$$\frac{(q)_{k_j + \chi(j > n_0)(j - n_0) + k}}{(q)_{b + k_j + \chi(j > n_0)(j - n_0)} (q)_k} \times \prod_{j=1}^{n-n_0} \frac{1 - q^{(k+1)j}}{1 - q^{k+1}}. \tag{7.12}$$

Putting together (7.10) and (7.12) completes the proof of Theorem 6.2.

Remark. For all the identities considered in this paper, the formulas exhibit, apart from some minor deviations, quite a similar pattern, and it is more or less clear from the above argument, why it is so. We do not elaborate on this here, but the motivated reader may come up with other families of matrices \mathbf{B} for which a similar proof strategy might work. We believe that the details given above can be useful in such a quest.

7.4. A rationality result

It is possible to prove Theorem 6.2 based solely on Lemma 1.1; in fact this is how our result was originally obtained. It involves the same combinatorics applied when $k \geq a + 1$, in which case A_0 is an ordinary set. The extension of the result that includes all non-negative integers k depends on the following rationality lemma, inspired by [20, Proposition 2.4].

Lemma 7.5. *Fix nonnegative integers r_i, s_i for $1 \leq i \leq n$, satisfying $\sum r_i = \sum s_i$. There is a rational function $Q = Q(z) \in \mathbb{Q}(q)(z)$ that depends only on n and the numbers r_i, s_i , such that*

$$\text{CT} \left[\frac{x_1^{r_1} \cdots x_n^{r_n}}{x_1^{s_1} \cdots x_n^{s_n}} \mathcal{D}_q(\mathbf{x}; k) \right] = Q(q^k) \frac{(q)_{nk}}{(q)_k^n}.$$

Expanding the degree zero part of

$$\prod_{j=1}^n (qx_j)_{a + \chi(j \leq m)} (1/x_j)_b \prod_{n_0 < i < j \leq n} (1 - q^k x_i/x_j)(1 - q^{k+1} x_j/x_i)$$

into a sum of monomial terms and applying the above lemma to each such term individually, we find that there is a rational function $R \in \mathbb{Q}(q)(z)$ depending only on the parameters n, m, n_0, a, b such that

$$\text{CT}[\mathcal{L}_q(x_0, \mathbf{x}; \mathbf{B}_{\mathcal{AF}})] = R(q^k) \frac{(q)_{nk}}{(q)_k^n}.$$

Reorganizing the formula in Theorem 6.2 in the form

$$P(q^k) \frac{(q)_{nk}}{(q)_k^n}$$

with a function $P \in \mathbb{Q}(q)(z)$ which also depends only on n, m, n_0, a, b , the theorem established for $k \geq a + 1$ yields $P = R$, which in turn implies the full content of the result.

It only remains to prove Lemma 7.5, and this is executed with yet another application of Lemma 1.1. Since a similar — in fact more general — result was found recently by Doron Zeilberger and his able computer [15], we only give a brief account. As the $k = 0$ case is trivial, we will assume $k > 0$.

Proof of Lemma 7.5. The constant term of $\mathcal{D}_q(\mathbf{x}; k)$ equals the coefficient of $\prod x_i^{(n-1)k}$ in the polynomial

$$F(\mathbf{x}) = \prod_{1 \leq i < j \leq n} \left(\prod_{t=0}^{k-1} (x_j - q^t x_i) \times \prod_{t=1}^k (x_i - q^t x_j) \right).$$

Set $C_i = \{q^{\alpha_i} \mid \alpha_i \in [0, (n-1)k]\}$. Then $F(\mathbf{c}) = 0$ for every $\mathbf{c} \in C_1 \times \cdots \times C_n$ except when $c_i = q^{(i-1)k}$ for every i . According to Lemma 1.1,

$$\text{CT}[\mathcal{D}_q(\mathbf{x}; k)] = \frac{F(q^0, q^k, \dots, q^{(n-1)k})}{\psi_1 \psi_2 \dots \psi_n} = \frac{(q)_{nk}}{(q)_k^n} \quad \text{where} \quad \psi_i = \prod_{0 \leq j \leq (n-1)k, j \neq (i-1)k} (q^{(i-1)k} - q^j).$$

We compare this product to the constant term in the lemma, which equals the coefficient of $\prod x_i^{(n-1)k + s_i}$ in the polynomial $F^*(\mathbf{x}) = x_1^{r_1} \dots x_n^{r_n} F(\mathbf{x})$. Accordingly we set $C_i^* = \{q^{\alpha_i} \mid \alpha_i \in [0, (n-1)k + s_i]\}$ and note that for $\mathbf{c} \in C_1^* \times \cdots \times C_n^*$ we have $F^*(\mathbf{c}) \neq 0$ if and only if the exponents α_i are all distinct and

$$\alpha_{\pi(i+1)} \geq \alpha_{\pi(i)} + k + \chi(\pi(i) > \pi(i+1))$$

holds for $1 \leq i \leq n-1$ with the unique permutation $\pi = \pi_{\mathbf{c}} \in \mathfrak{S}_n$ satisfying $\alpha_{\pi(1)} < \cdots < \alpha_{\pi(n)}$. Consequently, $\alpha_i = (\pi^{-1}(i) - 1)k + \epsilon_i$ for some $\epsilon_i = \epsilon_i(\mathbf{c}) \in [0, s_{\pi(i)}]$.

Set $C = \{\mathbf{c} \in C_1^* \times \cdots \times C_n^* \mid F^*(\mathbf{c}) \neq 0\}$, and write $s = \max s_i$. It follows that

$$|C| \leq n! \binom{s+n}{n}.$$

Moreover, the set $\mathcal{S} = \{(\pi_{\mathbf{c}}, \epsilon_1(\mathbf{c}), \dots, \epsilon_n(\mathbf{c})) \mid \mathbf{c} \in C\}$ is independent of k ; it depends only on n and the numbers s_i . It follows from Lemma 1.1 that, using the notation $\tau = \pi^{-1}$,

$$\text{CT} \left[\frac{x_1^{r_1} \dots x_n^{r_n}}{x_1^{s_1} \dots x_n^{s_n}} \mathcal{D}_q(\mathbf{x}; k) \right] = \sum_{\mathbf{c} \in C} \prod_{i=1}^n q^{((\tau(i)-1)k + \epsilon_i)r_i} \frac{F(\dots, q^{(\tau(i)-1)k + \epsilon_i}, \dots)}{\psi_1^* \psi_2^* \dots \psi_n^*}$$

where

$$\psi_{\pi(i)}^* = \prod_{0 \leq j \leq (n-1)k + s_{\pi(i)}, j \neq (i-1)k + \epsilon_{\pi(i)}} (q^{(i-1)k + \epsilon_{\pi(i)}} - q^j).$$

One readily checks that for each $\Sigma = (\pi, \epsilon_1, \dots, \epsilon_n) \in \mathcal{S}$ there exist rational functions $Q_i \in \mathbb{Q}(q)(z)$ that depend only on n , the numbers r_j, s_j and the sequence Σ , such that

$$\prod_{i=1}^n q^{((\tau(i)-1)k + \epsilon_i)r_i} = Q_0(q^k), \quad \frac{\psi_i}{\psi_{\pi(i)}^*} = Q_i(q^k) \quad \text{and} \quad \frac{F(\dots, q^{(\tau(i)-1)k + \epsilon_i}, \dots)}{F(q^0, q^k, \dots, q^{(n-1)k})} = Q_{n+1}(q^k).$$

The result follows. □

Acknowledgments

We gratefully acknowledge the support of the Australian Research Council, ERC Advanced Research Grant No. 267165, OTKA Grants 81310 and 100291, RFBR Grants 11-01-00677-a, 13-01-00935-a and 13-01-12422-ofi-m, President of Russia Grant MK-6133.2013.1, and Russian Government Mega-Grant Project 11.G34.31.0053.

References

- [1] N. Alon, Combinatorial Nullstellensatz, *Combin. Probab. Comput.* 8 (1999) 7–29.
- [2] N. Alon, M.B. Nathanson, I.Z. Ruzsa, The polynomial method and restricted sums of congruence classes, *J. Number Theory* 56 (1996) 404–417.
- [3] G.W. Anderson, A short proof of Selberg’s generalized beta formula, *Forum Math.* 3 (1991) 415–417.
- [4] K. Aomoto, Jacobi polynomials associated with Selberg integrals, *SIAM J. Math. Anal.* 18 (1987) 545–549.
- [5] G.E. Andrews, Problems and prospects for basic hypergeometric functions, in: R.A. Askey (Ed.), *Theory and Application of Special Functions*, Academic Press, New York, 1975, pp. 191–224.
- [6] R. Askey, Some basic hypergeometric extensions of integrals of Selberg and Andrews, *SIAM J. Math. Anal.* 11 (1980) 938–951.
- [7] T.H. Baker, P.J. Forrester, Generalizations of the q -Morris constant term identity, *J. Combin. Theory Ser. A* 81 (1998) 69–87.
- [8] W. Baratta, Some properties of Macdonald polynomials with prescribed symmetry, *Kyushu J. Math.* 64 (2010) 323–343.
- [9] D.M. Bressoud, I.P. Goulden, Constant term identities extending the q -Dyson theorem, *Trans. Amer. Math. Soc.* 291 (1985) 203–228.
- [10] T.W. Cai, Macdonald symmetric functions of rectangular shapes, arXiv:1308.3821.
- [11] F. Carlson, *Sur une Classe de Séries de Taylor*, Ph.D. Thesis, Uppsala Univ., 1914.
- [12] J.A. Dias da Silva, Y.O. Hamidoune, Cyclic spaces for Grassmann derivatives and additive theory, *Bull. London Math. Soc.* 26 (1994) 140–146.
- [13] F.J. Dyson, Statistical theory of energy levels of complex systems. I, *J. Math. Phys.* 3 (1962) 140–156.
- [14] F.J. Dyson, A Brownian-motion model for the eigenvalues of a random matrix, *J. Math. Phys.* 3 (1962) 1191–1198.
- [15] S.B. Ekhad, D. Zeilberger, How to extend Károlyi and Nagy’s BRILLIANT proof of the Zeilberger–Bressoud q -Dyson theorem in order to evaluate ANY coefficient of the q -Dyson product, *Personal Journal of Shalosh B. Ekhad and Doron Zeilberger*. See also arXiv:1308.2983.
- [16] P. Erdős, R.L. Graham, *Old and New Problems and Results in Combinatorial Number Theory*, L’Enseignement Mathématique, Geneva, 1980.
- [17] P.J. Forrester, Normalization of the wavefunction for the Calogero–Sutherland model with internal degrees of freedom, *Int. J. Mod. Phys. B* 9 (1995) 1243–1261.
- [18] P.J. Forrester, *Log-Gases and Random Matrices*, London Math. Soc. Monographs, vol. 34, Princeton University Press, 2010.
- [19] P.J. Forrester, S.O. Warnaar, The importance of the Selberg integral, *Bull. Amer. Math. Soc. (N.S.)* 45 (2008) 489–534.
- [20] I.M. Gessel, L. Lv, G. Xin, Y. Zhou, A unified elementary approach to the Dyson, Morris, Aomoto, and Forrester constant term identities, *J. Combin. Theory Ser. A* 115 (2008) 1417–1435.
- [21] I.M. Gessel, G. Xin, A short proof of the Zeilberger–Bressoud q -Dyson theorem, *Proc. Amer. Math. Soc.* 134 (2006) 2179–2187.
- [22] I.J. Good, Short proof of a conjecture by Dyson, *J. Math. Phys.* 11 (1970) 1884.
- [23] J. Gunson, Proof of a conjecture by Dyson in the statistical theory of energy levels, *J. Math. Phys.* 3 (1962) 752–753.
- [24] L. Habsieger, Une q -intégrale de Selberg et Askey, *SIAM J. Math. Anal.* 19 (1988) 1475–1489.
- [25] S. Hamada, Proof of Baker–Forrester’s constant term conjecture for the cases $N_1 = 2, 3$, *Kyushu J. Math.* 56 (2002) 243–266.
- [26] Q.-H. Hou, Z.-W. Sun, Restricted sums in a field, *Acta Arith.* 102 (2002) 239–249.
- [27] K.W.J. Kadell, A proof of Andrews’s q -Dyson conjecture for $n = 4$, *Trans. Amer. Math. Soc.* 290 (1985) 127–144.
- [28] K.W.J. Kadell, A proof of Askey’s conjectured q -analogue of Selberg’s integral and a conjecture of Morris, *SIAM J. Math. Anal.* 19 (1988) 969–986.
- [29] K.W.J. Kadell, Aomoto’s machine and the Dyson constant term identity, *Methods Appl. Anal.* 5 (1998) 335–350.
- [30] K.W.J. Kadell, A Dyson constant term orthogonality relation, *J. Combin. Theory Ser. A* 89 (2000) 291–297.
- [31] J. Kaneko, On Forrester’s generalization of Morris constant term identity, in: *q -series From a Contemporary Perspective* (South Hadley, MA, 1998), Contemporary Mathematics, vol. 254, Amer. Math. Soc., Providence, 2000, pp. 271–282.
- [32] J. Kaneko, Forrester’s constant term conjecture and its q -analogue, in: *Physics and Combinatorics* (Nagoya, 2000), World Sci. Publ., River Edge, NJ, 2001, pp. 49–62.
- [33] J. Kaneko, Forrester’s conjectured constant term identity. II, *Ann. Combin.* 6 (2002) 383–397.
- [34] J. Kaneko, On Baker–Forrester’s constant term conjecture, *J. Ramanujan Math. Soc.* 18 (2003) 349–367.
- [35] R.N. Karasev, F.V. Petrov, Partitions of nonzero elements of a finite field into pairs, *Israel J. Math.* 192 (2012) 143–156.
- [36] Gy. Károlyi, A. Lascoux, S.O. Warnaar, Constant term identities and Poincaré polynomials, *Trans. Amer. Math. Soc.*, to appear.
- [37] Gy. Károlyi, Z.L. Nagy, A simple proof of the Zeilberger–Bressoud q -Dyson theorem, *Proc. Amer. Math. Soc.*, to appear.
- [38] G. Kós, T. Mészáros, L. Rónyai, Some extensions of Alon’s Nullstellensatz, *Publ. Math. Debrecen* 79 (2011) 507–519.
- [39] G. Kós, L. Rónyai, Alon’s Nullstellensatz for multisets, *Combinatorica* 32 (2012) 589–605.
- [40] M. Lasoń, A generalization of Combinatorial Nullstellensatz, *Electron. J. Combin.* 17 (2010) #N32, 6 pages.
- [41] L. Lv, G. Xin, Y. Zhou, A family of q -Dyson style constant term identities, *J. Combin. Theory Ser. A* 116 (2009) 12–29.
- [42] W.G. Morris, *Constant Term Identities for Finite and Affine Root Systems: Conjectures and Theorems*, Ph.D. Thesis, Univ. Wisconsin–Madison, 1982.

- [43] D. Ostrovsky, Selberg integral as a meromorphic function, *Int. Math. Res. Not. IMRN* (2013) 3988–4028.
- [44] A. Selberg, Bemerkninger om et multipelt integral, *Norsk Mat. Tidsskr.* 26 (1944) 71–78.
- [45] A.V. Sills, Disturbing the Dyson conjecture, in a *generally* GOOD way, *J. Combin. Theory Ser. A* 113 (2006) 1368–1380.
- [46] R.P. Stanley, The q -Dyson conjecture, generalized exponents, and the internal product of Schur functions, in: *Combinatorics and Algebra* (Boulder, 1983), *Contemporary Mathematics*, vol. 34, Amer. Math. Soc., Providence, 1984, pp. 81–94.
- [47] R.P. Stanley, The stable behavior of some characters of $SL(n, \mathbb{C})$, *Lin. Multilin. Alg.* 16 (1984) 3–27.
- [48] Z.-W. Sun, Y.-N. Yeh, On various restricted sumsets, *J. Number Theory* 114 (2005) 209–220.
- [49] K.G. Wilson, Proof of a conjecture by Dyson, *J. Math. Phys.* 3 (1962) 1040–1043.
- [50] G. Xin, A residue theorem for Malcev–Neumann series, *Adv. Appl. Math.* 35 (2005) 271–293.
- [51] G. Xin, Y. Zhou, A Laurent series proof of the Habsieger–Kadell q -Morris identity, arXiv:1302.6642.
- [52] D. Zeilberger, A Stembridge–Stanton style elementary proof of the Habsieger–Kadell q -Morris identity, *Discrete Math.* 79 (1990) 313–322.
- [53] D. Zeilberger, D.M. Bressoud, A proof of Andrews’ q -Dyson conjecture, *Discrete Math.* 54 (1985) 201–224.
- [54] Y. Zhou, On Kadell’s two conjectures for the q -Dyson product, *Electron. J. Combin.* 18(2) (2011) #P2, 15 pages.
- [55] Y. Zhou, New extensions to the sumsets with polynomial restrictions, arXiv:1202.3190.