

1 **Estrogen receptor alpha and beta differentially mediate C5aR agonist evoked**  
2 **Ca<sup>2+</sup>-influx in neurons through L-type voltage-gated Ca<sup>2+</sup> channels**

3 Imre Farkas<sup>a</sup>, Miklós Sárvári<sup>a</sup>, Máté Aller<sup>b</sup>, Noriko Okada<sup>c</sup>, Hidechika Okada<sup>d</sup>, István  
4 Likó<sup>e</sup>, Zsolt Liposits<sup>a,f</sup>

5 <sup>a</sup>Laboratory of Endocrine Neurobiology and <sup>b</sup>Laboratory of Cellular Pharmacology,  
6 Institute of Experimental Medicine, Hungarian Academy of Sciences, Budapest,  
7 Hungary, <sup>c</sup>Department of Immunology, Nagoya City University, Nagoya, Japan,  
8 <sup>d</sup>Choju Medical Institute, Toyohashi, Japan, <sup>e</sup>Gedeon Richter Plc., Budapest,  
9 Hungary, <sup>f</sup>Department of Neuroscience, Faculty of Information Technology, Pázmány  
10 Péter Catholic University, Budapest, Hungary

11

12 **Disclosure summary:** The authors have nothing to disclose

13

14 **Corresponding author and to whom reprint requests should be addressed:**

15 Imre Farkas, PhD

16 Laboratory of Endocrine Neurobiology, Institute of Experimental Medicine,  
17 Hungarian Academy of Sciences, Szigony. u. 43., H-1083 Budapest, Hungary

18 Tel.: 36-1-210-9400; Fax: 36-1-210-9944

19 Email: [farkas@koki.hu](mailto:farkas@koki.hu)

20

21 **Abstract**

22 Complement C5a is associated primarily with inflammation. The widespread  
23 expression of its receptors, C5aR and C5L2 in neuronal cells, however, suggests  
24 additional regulatory roles for C5a in the CNS. C5aR agonist (PL37-MAP) evokes  
25  $\text{Ca}^{2+}$ -influx in GT1-7 neuronal cell line and the  $\text{Ca}^{2+}$ -influx is regulated by estradiol.  
26 In the present study, we examined further the mechanism of  $\text{Ca}^{2+}$ -influx and the  
27 contribution of the two estrogen receptor (ER) isotypes, ER $\alpha$  and ER $\beta$ , to estrogenic  
28 modulation of intracellular  $\text{Ca}^{2+}$ -content. GT1-7 neurons were treated with isotype  
29 selective ER agonists for 24h then C5aR agonist evoked  $\text{Ca}^{2+}$ -responses were  
30 measured by  $\text{Ca}^{2+}$ -imaging. Transcriptional changes were followed by real-time PCR.  
31 We found that not only estradiol (100pM), but the ER $\alpha$  selective agonist PPT  
32 (100pM) enhanced the PL37-MAP-evoked  $\text{Ca}^{2+}$ -influx (E2: 215%, PPT: 175%,  
33 compared to the PL37-MAP-evoked  $\text{Ca}^{2+}$ -influx). In contrast, the ER $\beta$  selective  
34 agonist DPN (100pM) significantly reduced the  $\text{Ca}^{2+}$ -influx (32%). Attenuated  $\text{Ca}^{2+}$ -  
35 response (25%) was observed in Ca-free environment and depletion of the  $\text{Ca}^{2+}$ -pool  
36 by CPA eliminated the remaining elevation in the  $\text{Ca}^{2+}$ -content, demonstrating that  
37 the majority of  $\text{Ca}^{2+}$  originated from the extracellular compartment. L-type voltage-  
38 gated  $\text{Ca}^{2+}$ -channel (L-VGCC) blocker nifedipine abolished the  $\text{Ca}^{2+}$ -influx, while R-  
39 type  $\text{Ca}^{2+}$ -channel blocker SNX-482 had no effect, exemplifying the predominant role  
40 of L-VGCC in this process. Acute pre-treatments (8min) with ER agonists did not  
41 affect the evoked  $\text{Ca}^{2+}$ -influx, revealing that the observed effects of estrogens were  
42 genomic. Therefore, we checked estrogenic regulation of C5a receptors and L-VGCC  
43 subunits. ER agonists increased C5aR mRNA expression, whereas they differentially  
44 regulated C5L2. Estradiol decreased transcription of  $\text{Ca}_v1.3$  L-VGCC subunit. Based  
45 on these results we propose that estradiol may differentially modulate C5a-induced

46 Ca<sup>2+</sup>-influx via L-VGCCs in neurons depending on the expression of the two ER  
47 isotypes.

48

49 **Keywords:** GT1-7 neuron, complement C5a receptor, estrogen receptor alpha,  
50 estrogen receptor beta; voltage-gated calcium channel

51

52 **Abbreviations:**

53 C – complement system

54 C5aR – “classical” complement 5a receptor (G-protein coupled)

55 C5L2 – “second” complement 5a receptor (non-G-protein coupled)

56 CPA – cyclopiazonic acid

57 DPN – Diarylpropionitrile

58 E2 – 17β-estradiol

59 ERα – estrogen receptor alpha

60 ERβ – estrogen receptor beta

61 GnRH – gonadotropin-releasing hormone

62 HBSS – Hanks’ Balanced Salt Solution

63 L-VGCC – L-type voltage-gated calcium channel

64 PPT – 4,4',4''-(4-Propyl-[1H]-pyrazole-1,3,5-triyl)trisphenol

65 PVN – paraventricular nucleus

66 SON – supraoptic nucleus

67

68

69        1. **Introduction**

70            The complement (C) system is an ancient immune pathway comprised of  
71 numerous elements activated in a cascade leading to the elimination of pathogens  
72 (Speth et al., 2008). During C activation, the C5a anaphylatoxin, a 74 amino-acid long  
73 fragment of the fifth component of C is released when C5 is cleaved by the C5  
74 convertase. C5a binds to two receptors, the “classical” G-protein coupled C5aR  
75 (CD88) and the non-G-protein coupled receptor C5L2 (GPR77). Both receptors are  
76 expressed on immune and non-immune cell types. C5a binding to C5aR leads to  
77 various events such as increased intracellular calcium level and activation of  
78 intracellular signaling cascades resulting in functional responses e.g. recruiting and  
79 activation of inflammatory cells, degranulation, delayed or enhanced apoptosis,  
80 phagocytosis, histamine release, and chemotaxis (Fujita et al., 2004; Guo and Ward,  
81 2005). C5L2 may function as a decoy receptor regulating the inflammatory response  
82 resulted from the C5a/C5aR binding (Bamberg et al., 2010; Woodruff et al., 2011).

83            Expression of C5aR has been demonstrated in astrocytes, microglia and  
84 neurons of the central nervous system (CNS) (Woodruff et al., 2010). The cellular  
85 expression pattern of C5L2 is similar to that seen for C5aR. Pyramidal neurons in the  
86 hippocampus and the cortex, Purkinje cells of the cerebellum and neuroblastoma cells  
87 express C5aR (Farkas et al., 1999; Farkas et al., 1998a; Farkas et al., 1998b). Function  
88 of the C5aR in neurons remains elusive. A C5aR-related apoptotic pathway and the  
89 role of this receptor in neurodegenerative diseases such as Alzheimer’s disease have  
90 been suggested (Farkas et al., 2003; Fonseca et al., 2009; Fonseca et al., 2011). In  
91 contrast, the neuroprotective role of C5a has also been demonstrated (Woodruff et al.,  
92 2010).

93 C5aR has recently been identified in hypothalamic neurons, including  
94 gonadotropin-releasing hormone (GnRH)-producing cells, immortalised GnRH-  
95 producing GT1-7 neurons and neurons of the paraventricular (PVN) and supraoptic  
96 (SON) nuclei (Farkas et al., 2008). GT1-7 neurons establish neuronal network with  
97 co-ordinated activity and produce GnRH in a pulsatile fashion (Liposits et al., 1991;  
98 Wetsel et al., 1992). Pulsatility and volume of the secretion is in strong correlation  
99 with synchronised firing (Moenter et al., 2003; Thiery and Pelletier, 1981; Wilson et  
100 al., 1984). Various factors released during inflammation can play role in the function  
101 of these cells (Karsch et al., 2002). Cannabinoids also affect GnRH neurons by  
102 utilising the retrograde endocannabinoid signaling mechanism (Farkas et al., 2010).

103 The estrogen receptor alpha and beta (ER $\alpha$  and ER $\beta$ ) are expressed in  
104 numerous hypothalamic neurons, such as GnRH cells and the neurons of PVN and  
105 SON (Hrabovszky et al., 2004; Shughrue et al., 1997; Shughrue and Merchenthaler,  
106 2001), GT1-7 cells express both ER subtypes (Roy et al., 1999). 17 $\beta$ -estradiol (E2)  
107 can modulate the electric function of GT1-7 cells and exert both negative and positive  
108 feedback on the firing (Christian et al., 2005; Farkas et al., 2007).

109 Our previous experiments have shown that administration of a C5aR agonist  
110 results in robust calcium (Ca<sup>2+</sup>) influx in GnRH neurons. In addition, E2 pre-treatment  
111 elevates this Ca<sup>2+</sup>-response suggesting that the signal transduction pathways related to  
112 the C5aRs and the ERs, respectively, can modulate each other (Farkas et al., 2008).  
113 Change in the intracellular Ca<sup>2+</sup>-milieu can heavily affect firing properties of the  
114 neurons. Firing can be fine-tuned for example by the opening and closing of the L-  
115 type voltage-gated Ca<sup>2+</sup>-channels (L-VGCCs). Ca<sub>v</sub>1.2 and Ca<sub>v</sub>1.3 subunits of the L-  
116 VGCCs are strongly involved in spontaneous firing and pacemaking (Zuccotti et al.,  
117 2011). In the present study, therefore, we investigated further, how C5aR and ER

118 subtypes interact using the immortalised GnRH-producing GT1-7 neurons as a  
119 neuronal model with Ca<sup>2+</sup>-imaging and quantitative real-time PCR methods. Genomic  
120 and non-genomic actions of ER subtypes on the C5aR-mediated Ca<sup>2+</sup>-influx were  
121 examined. Potential sources of the increased Ca<sup>2+</sup>-content were also studied, including  
122 the role of various VGCCs and the intracellular Ca<sup>2+</sup>-pool.  
123

124 **2. Materials and methods**

125 *2.1 Cell culture*

126 GnRH-producing immortalised GT1-7 neurons were cultured in Dulbecco  
127 Modified Eagle Medium (DMEM) containing high-glucose and supplemented with  
128 10% fetal calf serum (FCS) and 5% horse serum (HS). Prior to ER agonist treatment  
129 the culturing medium was replaced with a steroid/thyroid- and phenol red-free one  
130 and cells were cultured in this medium for 24hrs. Subsequently, the cells were treated  
131 with 17 $\beta$ -estradiol (E2, SIGMA), the highly potent ER $\beta$  receptor agonist DPN  
132 (Diarylpropionitrile, Tocris) and the ER $\alpha$  receptor agonist PPT (4,4',4''-(4-Propyl-  
133 [1H]-pyrazole-1,3,5-triyl)trisphenol, Tocris) at various concentrations (100pM -  
134 20nM) for 24h and then used for calcium imaging and RT-PCR experiments. In the  
135 experiments examining effect of acute treatment, the E2, DPN, and PPT were added  
136 to the cells 8min before starting the calcium imaging recording.

137

138 *2.2 Calcium imaging*

139 Cultured GT1-7 cells were loaded with the calcium-sensitive fluorescent dye  
140 Fura-2 AM (1 $\mu$ M; Molecular Probes, Eugene, Oregon) in loading buffer Hanks'  
141 Balanced Salt Solution (HBSS) containing 0.1% DMSO and Pluronic-F127 (1 $\mu$ M,  
142 Molecular Probes) for 1.5h at room temperature (RT). After washing with HBSS, the  
143 experiments were carried out at RT. The antisense homology box peptide fragment of  
144 the C5a (RAARISLGPRCIKAFTE) was synthesised in multiple antigenic peptide  
145 form (termed PL37-MAP, 2.5 $\mu$ M). Sequence of the PL37 is a fragment sequence of  
146 the C5a, representing a "strong" antisense homology box region in the C5a (Baranyi  
147 et al., 1995). Our previous works applying both C5a and PL37, respectively,  
148 demonstrated that PL37 is a potent agonist of the C5aR preserving the biological

149 activity of the C5a and triggering responses similar to that of the C5a (Baranyi et al.,  
150 1996; Fujita et al., 2004). The peptide was pipetted directly onto the cells in HBSS  
151 after a 1min baseline recording and then the diluted peptide remained in the HBSS  
152 during recording. In the case of E2, DPN, PPT pre-treatment, the cells were pre-  
153 treated with them as described in the “Cell culture” section and all of the rinsing and  
154 extracellular solutions contained the same concentration of E2, DPN and PPT. After  
155 the 1min baseline recording the PL37-MAP peptide was introduced into the bath fluid  
156 containing E2, DPN, and PPT and then the diluted peptide remained in the HBSS-E2  
157 or HBSS-DPN or HBSS-PPT mixture during recording.

158         When PL37-MAP was applied in  $\text{Ca}^{2+}$ -free extracellular solution (phosphate  
159 buffered salt solution=PBS, pH 7.4), the HBSS was changed to PBS just before  
160 starting the recording, except when the cells were treated with cyclopiazonic acid  
161 (CPA, 10 $\mu\text{M}$ , Tocris). CPA is a specific blocker of the  $\text{Ca}^{2+}$ -ATP-ase of the  
162 intracellular  $\text{Ca}^{2+}$ -store endoplasmic reticulum and depletes these  $\text{Ca}^{2+}$ -stores. CPA  
163 was applied to the GT1-7 neurons in PBS 30min before starting the measurements.

164         The VGCC blockers nifedipine (10 $\mu\text{M}$ , SIGMA) and SNX-482 (100nM) were  
165 added to the HBSS just before starting the  $\text{Ca}^{2+}$ -imaging measurement and remained  
166 in the HBSS during recording.

167         The experiments were carried out with an ARGUS HiSCA  $\text{Ca}^{2+}$ -imaging  
168 system (Hamamatsu Photonics, Hamamatsu, Japan) or with an Olympus BX50WI  
169 microscope equipped with a Polychrome II monochromator (TILL Photonics), a  
170 cooled CCD camera (Photometrics Quantix, Tucson, AZ, USA), and controlled by the  
171 Axon Imaging Workbench 6.0 software (Axon Instruments, Union City, CA, USA).  
172 The ratio of the fluorescent signals obtained at excitation wavelengths of 340 and 380  
173 nm was used to determine changes in the intracellular  $\text{Ca}^{2+}$ -concentration.



174 The surface density of the cultured cells was 500,000-750,000 cells/cm<sup>2</sup>, the  
175 magnification of the objective lens used was 40x, and the area of the calcium imaging  
176 acquisition was 0.038 mm<sup>2</sup>. Before starting the calcium imaging recordings, an area  
177 was chosen where at least 7 individual cells without overlaps could be clearly  
178 observed and measured (7-15 neurons depending on the surface density of the cells in  
179 the area measured).

180

### 181 *2.3 Total RNA isolation from GT1-7 cells*

182 Total RNA was isolated from GT1-7 cells using the RNeasy Mini Kit  
183 (QIAGEN, Hilden, Germany). RNA analytics included capillary electrophoresis using  
184 Agilent 2100 Bioanalyzer (Santa Clara, CA). All RNA samples displayed RNA  
185 integrity numbers (RIN) above 8.5.

186

### 187 *2.4 Quantitative real-time PCR*

188 Inventoried TaqMan assays were selected to study in depth the regulation of  
189 genes of our interest by quantitative real-time PCR. Each assay consisted of a FAM  
190 dye-labeled TaqMan MGB probe and two PCR primers. Every assay was optimised  
191 by the manufacturer to run under universal thermal cycling conditions with a final  
192 reaction concentration of 250nM for the probe and 900nM for each primer. Reverse  
193 transcription and real-time PCR were run as described earlier (Sarvari et al., 2010).  
194 RealTime StatMiner (Integromics, Granada, Spain) software and relative  
195 quantification against calibrator samples ( $\Delta\Delta C_t$ ) were used for analysis of Applied  
196 Biosystems TaqMan gene expression assays. Two house-keeping genes (Gapdh, Hprt)  
197 were applied as internal controls. The geometric mean of Ct values of Gapdh and  
198 Hprt1 was used for subsequent  $\Delta C_t$  calculation (Vandesompele et al., 2002). Relative

199 quantity (RQ) represents the change in the expression of a given gene in response to a  
200 treatment compared to basal (control) expression of the given gene. We considered  
201 changes with  $RQ > 1.5$  as up-regulation or  $RQ < 0.67$  as down-regulation (21).

202

### 203 *2.5 Statistical analysis*

204  $Ca^{2+}$ -imaging recordings using the fluorescence ratio obtained at 340 and 380  
205 nm wavelengths were baseline corrected, then the area-under-curve data of the  
206 records representing the net  $Ca^{2+}$ -influx were analyzed. Group data of the cells ( $n \geq 7$ )  
207 were expressed as mean  $\pm$  standard error (SEM). Statistical significance was analyzed  
208 using ANOVA followed by Newman-Keuls (NK) test (GraphPad Software Inc.,  
209 USA), and considered at  $p < 0.05$ .

210

211 **3. Results**

212 *3.1 Estrogens differentially modulate the Ca<sup>2+</sup>-influx evoked by the C5aR agonist*

213 The C5aR agonist peptide PL37-MAP (2.5µM) triggered robust Ca<sup>2+</sup>-influx in  
214 GT1-7 neurons (Fig. 1a). Onset of the elevation of intracellular Ca<sup>2+</sup>-concentration  
215 was within 1-1.5min after application of the peptide. When the neurons were pre-  
216 treated with E2 (100pM, 24h), the evoked Ca<sup>2+</sup>-influx started earlier (in less than 30  
217 seconds) and the amplitude of the records was higher than that of the recordings  
218 evoked by the PL37-MAP alone (Fig. 1b). Pre-treatment with the selective ERα  
219 agonist PPT (100pM, 24h) elevated the PL37-MAP-triggered Ca<sup>2+</sup>-influx. The  
220 elevation was similar to the increased Ca<sup>2+</sup>-influx measured with E2 (Fig. 1c). In  
221 contrast, when the cells were pre-treated with the selective ERβ agonist DPN (100pM,  
222 24h), the elevation of the Ca<sup>2+</sup>-content was much lower and started in 1.5-2min after  
223 introducing the PL37-MAP peptide into the bath fluid (Fig. 1d).

224 Histogram of the normalised area-under-curve data representing the net  
225 changes in the intracellular free Ca<sup>2+</sup> concentration showed significant increase when  
226 PL37-MAP was applied to GT1-7 neurons in the presence of either E2 (100pM, 24h)  
227 or PPT (100pM, 24h), revealing facilitation of the PL37-MAP-evoked Ca<sup>2+</sup>-influx by  
228 both E2 and PPT. On the other hand, DPN (100pM, 24h) significantly decreased (Fig.  
229 1e) the Ca<sup>2+</sup>-influx evoked by PL37-MAP (E2: 226.6 ± 27.52%; PPT: 159.1 ±  
230 16.74%; DPN 18.6 ± 10.47% of the Ca<sup>2+</sup>-influx evoked by PL37-MAP alone;  
231 p<0.001). Application of higher concentrations of ER agonists resulted in similar  
232 effect except in the case of PPT (Fig. 1e). E2 (20nM, 24h) increased the PL37-MAP-  
233 evoked Ca<sup>2+</sup>-influx whereas DPN (20nM, 24h) attenuated it significantly (E2: 169.7 ±  
234 16.67%; DPN 10.9 ± 3.65% of the Ca<sup>2+</sup>-influx evoked by PL37-MAP alone;  
235 p<0.001).

236

237 *3.2 The effects of ER agonists on the Ca<sup>2+</sup>-influx are genomic*

238 In order to determine whether the observed effect of ER agonists on the PL37-  
239 MAP-evoked Ca<sup>2+</sup>-influx was genomic, PL37-MAP was applied after an acute (8min)  
240 application of E2, PPT, and DPN. Ca<sup>2+</sup>-imaging experiments demonstrated that short  
241 administration of ER agonists failed to influence the change in the intracellular free  
242 Ca<sup>2+</sup>-concentration triggered by PL37-MAP (Fig. 1f). Examination of the area-under-  
243 curve data showed no significant differences (E2: 115.1 ± 13.78%; PPT: 132.2 ±  
244 13.97%; DPN 146.9 ± 45.22% of the Ca<sup>2+</sup>-influx evoked by PL37-MAP alone). The  
245 results revealed that genomic effects of the E2, DPN, and PPT were necessary to  
246 modulate the Ca<sup>2+</sup>-response.

247 *3.3 The major Ca<sup>2+</sup>-source is extracellular*

248 Potential sources of the increase in the intracellular Ca<sup>2+</sup>-content were also  
249 investigated with Ca<sup>2+</sup>-imaging on the GT1-7 neurons (Fig. 2a-g). Application of  
250 PL37-MAP in Ca<sup>2+</sup>-free extracellular fluid (PBS) resulted in a significantly lower  
251 Ca<sup>2+</sup>-response than in HBSS, which was independent from the ER agonist pre-  
252 treatment (p<0.001). Administration of PL37-MAP resulted in a significantly  
253 attenuated elevation in Ca<sup>2+</sup>-concentration (25.3 ± 4.91% of the Ca<sup>2+</sup>-increase evoked  
254 by PL37-MAP in Ca<sup>2+</sup>-containing extracellular fluid). Nevertheless, the effect of the  
255 PL37-MAP was not entirely eliminated in PBS, suggesting, that intracellular Ca<sup>2+</sup>-  
256 sources were also activated during the process. Therefore, the effect of PL37-MAP  
257 was also examined when CPA (10µM, 30min) was present in the PBS. Depletion of  
258 the Ca<sup>2+</sup>-store in the endoplasmic reticulum by CPA resulted in ablation of the  
259 increase in the intracellular free Ca<sup>2+</sup>-content (3.2 ± 0.66% of the Ca<sup>2+</sup>-increase  
260 evoked by PL37-MAP in Ca<sup>2+</sup>-containing extracellular fluid). Application of PL37-

261 MAP in PBS in the presence of E2, DPN, or PPT (20nM, 24h) demonstrated  
262 significant decrease of the Ca<sup>2+</sup>-response (E2: 16.9 ± 1.63%; PPT: 22.9 ± 5.82%;  
263 DPN: 27.5 ± 5.45% of the Ca<sup>2+</sup>-increase evoked by PL37-MAP in Ca<sup>2+</sup>-containing  
264 extracellular fluid), however, these data did not differ from the Ca<sup>2+</sup>-increase evoked  
265 by PL37-MAP in Ca<sup>2+</sup>-free solution.

### 266 *3.4 L-type but not the R-type voltage-gated Ca<sup>2+</sup>-channels are involved in the PL37-* 267 *MAP-evoked Ca<sup>2+</sup>-influx*

268 In order to investigate which Ca<sup>2+</sup>-channel was involved in the Ca<sup>2+</sup>-influx  
269 triggered by C5aR activation, PL37-MAP was applied to GT1-7 neurons in the  
270 presence of blockers of various VGCCs. The L- and the R-type Ca<sup>2+</sup>-channels have  
271 been reported as the most abundant ones in the GT1-7 cells (Watanabe et al., 2004),  
272 therefore these two channels were examined. Application of nifedipine (10 µM), the  
273 inhibitor of the L-VGCC, resulted in a significantly reduced Ca<sup>2+</sup>-influx evoked by  
274 PL37-MAP (Fig. 3a-f). Pre-treatment the cells with E2, PPT or DPN (20nM, 24h) did  
275 not modify effect of nifedipine (nifedipine alone: 39.7 ± 3.29%; E2: 35.8 ± 6.94%;  
276 PPT: 32.4 ± 4.66%; DPN: 34.2 ± 6.48% of the data measured with PL37-MAP alone;  
277 p<0.001).

278 In contrast to nifedipine, SNX-482 (100nM), the inhibitor of the R-type Ca<sup>2+</sup>-  
279 channels, had no effect on the changes in the intracellular Ca<sup>2+</sup>-concentration evoked  
280 by PL37-MAP (Fig. 4a-b, f). In the presence of SNX-482, the triggered Ca<sup>2+</sup>-influx  
281 did not differ from the one measured with PL37-MAP alone (SNX: 81.6 ± 13.95 %).  
282 In addition, block of the R-type Ca<sup>2+</sup>-channels did not influence effect of the pre-  
283 treatment with E2, PPT or DPN (20 nM, 24 h) (Figs. 4c-f). The Ca<sup>2+</sup>-influx increased  
284 upon E2 whereas decreased upon DPN pre-treatment significantly (E2: 139.5 ±  
285 14.63%, PPT: 86.6 ± 5.33%, DPN: 43.3 ± 6.15%, p<0.001).

286

287 *3.5 Estrogenic modulation of the transcription of the C5a receptors and the Ca<sub>v</sub>1.3*  
288 *subunit of the L-type Ca<sup>2+</sup>-channel in GT1-7 cells*

289 We examined the effects of E2 and isotype selective ER agonists on the  
290 transcription of genes encoding C5a receptors C5aR and C5L2, and L-VGCC subunits  
291 Ca<sub>v</sub>1.2, and Ca<sub>v</sub>1.3, by real-time PCR. We demonstrated estrogenic regulation of the  
292 C5a receptor genes (Table 1). C5ar1 (the classical C5aR) was up-regulated by the  
293 three ER agonists. C5L2 was regulated differentially, E2 increased while PPT  
294 decreased its transcription. Cacna1d (gene for Ca<sub>v</sub>1.3) was regulated only by E2  
295 whereas Cacna1c (Ca<sub>v</sub>1.2) showed no estrogenic regulation.

296

297 **4. Discussion**

298 In the present study, we examined further estrogenic modulation of the C5aR  
299 agonist-evoked  $Ca^{2+}$ -response using the GnRH-producing GT1-7 cell line as a  
300 neuronal model and applying isotype selective ER agonists. We demonstrated that i)  
301  $ER\alpha$  and  $ER\beta$  agonists differentially modulated the C5aR agonist-evoked  $Ca^{2+}$ -influx,  
302 ii) estrogenic modulation was dependent on genomic effects, iii)  $Ca^{2+}$ -influx was  
303 mediated primarily through L-VGCC, iv) estrogens up-regulated C5aR mRNA  
304 expression while differentially regulated C5L2.

305 *4.1 Estrogens differentially modulate the PL37-MAP-evoked  $Ca^{2+}$ -influx*

306 Our present results showed that the C5aR agonist-evoked  $Ca^{2+}$ -influx was  
307 differentially mediated by various ER agonists in GT1-7 neurons. Expression of  
308 C5aR, and  $ER\alpha$  and  $ER\beta$  has long been reported in various types of neurons (Farkas  
309 et al., 2003; Farkas et al., 2008; Hrabovszky et al., 2004; Hrabovszky et al., 2000;  
310 Hrabovszky et al., 2001; Shughrue et al., 1997; Shughrue and Merchenthaler, 2001;  
311 Stahel et al., 1997a; Stahel et al., 1997b; Wilson et al., 2002; Woodruff et al., 2010).  
312 Differential modulation by  $ER\alpha$  and  $ER\beta$  could be important, because  $Ca^{2+}$ -influx  
313 evoked by the activation of C5aR can differentially affect functions of a neuronal cell,  
314 such as firing pattern, shape of after-hyperpolarisation and depolarising after-  
315 potentials, neurotransmitter release, plasticity, gene transcription, and vulnerability  
316 (Berridge, 1998; Zuccotti et al., 2011).

317 In the present experiments DPN decreased the  $Ca^{2+}$ -influx evoked by PL37-  
318 MAP. Nevertheless, the amplitude of the inward ion current in the  $ER\beta$ -expressing  
319 GnRH neurons from slices obtained from E2 substituted mice was higher than those  
320 from ovariectomized mice (Farkas et al., 2008). The reasons of the discrepancy may  
321 lie in the differences between the two models.

322 4.2 *Estrogenic modulation of the evoked Ca<sup>2+</sup>-influx is dependent on genomic effects*

323 Estrogenic modulation of the Ca<sup>2+</sup>-signal evoked by the activated C5aR was  
324 genomic rather than rapid in our experiments. Numerous estradiol-regulated genes  
325 have already been identified in GT1-7 neurons by expression profiling (Varju et al.,  
326 2009). Majority of the responding genes were up-regulated in these cells, including  
327 potassium channel subunits and transporters, transcription factors, molecules related  
328 to cell death, immune response, neurotransmitter, hormone and neuropeptide  
329 receptors, regulators of G-protein signaling. Those results support our present data  
330 showing up-regulation of C5aR.

331 In our present experiments, we found no acute effect of E2 on the Ca<sup>2+</sup>-influx  
332 evoked in the GT1-7 cells. In another model, published from another laboratory (Sun  
333 et al., 2010), both genomic and rapid changes resulted from the E2 administration  
334 were reported by potentiating the Ca<sup>2+</sup>-current in GnRH neurons in the acute brain  
335 slice in 5min. This report described, however, that percentage of the responding cells  
336 depended upon the concentration of E2 and only doses of E2 much higher than used  
337 in our experiments could evoke response with high rate of success. In, addition, the  
338 GnRH neurons presented an “all or none” ability to respond to acutely administered  
339 E2 (Sun et al., 2010). The observed discrepancies may reflect differences in the basic  
340 physiology and regulation of GnRH neurons integrated within the preoptic brain slice  
341 preparation versus the immortalised GT1-7 neurons cultured *in vitro*.

342 4.3 *L-type but not R-type Ca<sup>2+</sup>-channels are involved in the PL37-MAP-evoked Ca<sup>2+</sup>-*  
343 *influx*

344 In GT1-7 cells, the two major VGCCs are the L- and R-type channels  
345 (Watanabe et al., 2004). Both of them play a critical role in the regulation of Ca<sup>2+</sup>-  
346 dependent GnRH-release. In addition, R-type Ca<sup>2+</sup>-channels have been reported to be



347 involved in the release of neurotransmitters in calyx-type synapses of the medial  
348 nucleus of the trapezoid body, oxytocin neurons, and adrenal chromaffin cells  
349 (Albillos et al., 2000; Wang et al., 1999; Wu et al., 1998; Wu et al., 1999). R-type  
350 channels are responsible for the dendritic  $\text{Ca}^{2+}$ -influx induced by action potentials in  
351 CA1 pyramidal neurons of the hippocampus (Magee and Johnston, 1995; Sabatini and  
352 Svoboda, 2000). L-VGCCs contribute to  $\text{Ca}^{2+}$ -dependent gene transcription and can  
353 modulate firing properties of neurons (Gomez-Ospina et al., 2006; Zuccotti et al.,  
354 2011). Our results have now revealed that the  $\text{Ca}^{2+}$ -ion current resulted from the  
355 activation of the C5aR in GT1-7 neurons passed through the L-VGCCs but not via R-  
356 type channels. In addition, differential modulation of the  $\text{Ca}^{2+}$ -influx by ER agonists  
357 affected the function of the L-type channel. These data suggest that physiological  
358 functions of the L-VGCC such as regulation of the GnRH release, parameters of the  
359 firing, and various gene transcriptional events are affected by C5aR activation.  
360 Modulation of the L-VGCC by E2 and DPN has recently been reported in GnRH  
361 neurons of the acute brain slice of the mice demonstrating changes in the ion current  
362 via these channels under physiological conditions (Sun et al., 2010). Similarly,  
363 modulation of the T-type voltage-gated  $\text{Ca}^{2+}$ -channels by estradiol was also  
364 demonstrated (Bosch et al., 2009; Qiu et al., 2006; Zhang et al., 2009).

365         It is an intriguing question, how the activation of C5aR can regulate opening  
366 of a VGCC. One possibility is a change in the threshold level by phosphorylation of  
367 the L-VGCC, occurring as a downstream event of the C5aR activation. G-protein-  
368 coupled receptor activation can result in activation of diverse pathways involving  
369 enzymes such as protein kinase A or protein kinase C yielding phosphorylation of the  
370 L-VGCC, in particular its  $\text{Ca}_v1.2$  or  $\text{Ca}_v1.3$  subunits (Dai et al., 2009; Dolphin, 2009).  
371 This phosphorylation could eventually modify various electric parameters of the

372 neurons such as open probability of the channel or the threshold level. Nevertheless,  
373 the existence of a C5aR-related phosphorylation of L-VGCC requires further  
374 examination. Other G-protein mediated mechanisms can also be involved such as  
375 direct G-protein related modulation of the L-VGCCs (Currie, 2010; Tedford and  
376 Zamponi, 2006). In this paradigm, the L-VGCC molecule possesses residues  
377 interacting directly with the  $\beta\gamma$  subunits of the G-protein. The interaction is  
378 membrane-delimited, i.e. involves a second messenger molecule that remains  
379 associated with the plasma membrane, rather than diffusing to the channel via a  
380 cytoplasmic pathway (Hille, 1994).

381 Our  $\text{Ca}^{2+}$ -imaging measurements showed that in addition to the extracellular  
382 sources, intracellular  $\text{Ca}^{2+}$ -stores were also involved in the elevation of the  $\text{Ca}^{2+}$ -  
383 content evoked by the activation of the C5aR. The intracellular  $\text{Ca}^{2+}$ -stores could be  
384 triggered to release  $\text{Ca}^{2+}$  directly via the C5aR-related signal transduction pathway  
385 (Nishiura et al., 2010). Another possible pathway for this action is that C5aR-  
386 activation opens the L-VGCCs first and then the L-VGCCs activate the intracellular  
387  $\text{Ca}^{2+}$ -stores via a putative coupling (Kim et al., 2007; Kolarow et al., 2007), however,  
388 these opportunities require further elaboration.

389

#### 390 *4.4 C5aR agonist and estrogens modulate function of L-VGCC crucial in firing*

391 The characteristic firing pattern is a crucial feature of the hormone secreting  
392 neurons. Pulsatile release of the GnRH, for example, is indispensable for the proper  
393 function of the reproductive system (Moenter et al., 2003). In addition, the  
394 synchronous firing of the GnRH-producing neurons correlates with this pulsatility  
395 (Moenter et al., 2003). The  $\text{Ca}^{2+}$ -channels mediates how the neurons fire, therefore,  
396 effects disturbing the intracellular  $\text{Ca}^{2+}$ -milieu could have an effect on the firing

397 properties of the GnRH cells and consequently, the pulsatile secretion of GnRH. The  
398 L-VGCC is considered as one of the key mediators of the firing pattern (Zuccotti et  
399 al., 2011). Its  $Ca_v1.2$  and  $Ca_v1.3$  subunits expressed in neurons and endocrine cells,  
400 such as pancreatic beta, adrenal chromaffin cells (Catterall et al., 2005) and contribute  
401 to the spontaneous firing and pacemaking of the neurons (Zuccotti et al., 2011).  
402 However, literature data show activity-dependent differences between them.  $Ca_v1.3$  is  
403 more effective at low levels of activity such as during interburst intervals, and  $Ca_v1.2$   
404 is more efficient at high levels of activity such as during interspike intervals in the  
405 bursts (Zhang et al., 2006). It has recently been reported that E2 decreases mRNA  
406 expression of the  $Ca_v1.2$ , but has no effect on the  $Ca_v1.3$  subunit in the hippocampus  
407 of aged female rats (Brewer et al., 2009). Our real-time PCR measurements revealed  
408 that the  $Ca_v1.3$  subunit was down-regulated by E2 in GT1-7 cells. This discrepancy  
409 might originate from the cell-type differences in the two experimental models.

410         The point of convergence of the signals coming from C5aR and ERs, was not  
411 the expression of the  $Ca^{2+}$ -channel, because the  $Ca_v1.2$  and  $Ca_v1.3$  subunits were not  
412 regulated by DPN and PPT. Therefore, our present data suggest that any differential  
413 transcriptional regulation of elements of C5a/C5aR signaling which might be involved  
414 in the explored differential effects of the ER agonists on the evoked  $Ca^{2+}$ -influx  
415 should be upstream of L-VGCC. Possible candidates could be the regulator molecules  
416 of G-protein signaling (RGS2, RGS9 and RGS 10) which were earlier shown to be  
417 regulated by E2 in the GT1-7 cells (Varju et al., 2009).

#### 418 *4.5 C5aR is up-regulated by ER agonists*

419         In our studies, the expression of classical C5aR was up-regulated by E2 and  
420 the used isotype selective ER agonists, suggesting that neurons could respond more  
421 effectively to the inflammatory mediator C5a in the presence of estrogens. In contrast,

422 the expression of C5L2 was differentially regulated by ER agonists, displaying the  
423 up-regulation of this receptor by E2 and the attenuation of its expression by PPT. As  
424 the decoy receptor modulates the performance of the classical C5aR, the elucidation  
425 of this inverse regulatory trend warrants further investigation.

426         The results of the present *in vitro* study raise the questions of how activation  
427 of C5aR could occur by its ligand, the C5a in hormone secreting neurons *in vivo*. The  
428 hypophysiotrophic axonal projections of numerous hormone secreting neurons  
429 terminate outside the blood brain barrier, suggesting that these neurons are capable of  
430 monitoring C5a released either in the hypothalamus or the blood. Since several rodent  
431 hypothalamic neurons have been shown to express functional C5a receptors (Farkas et  
432 al., 2008), it is reasonable to assume that the inflammatory mediator C5a can alter the  
433 physiological properties and cellular functions of these neurons. In these mechanisms,  
434 the Ca<sup>2+</sup>-influx occurring via L-VGCCs might have a pivotal role.

435

436

437 **5. Conclusions**

438 Summing up, this study provided evidence that C5aR-mediated  $\text{Ca}^{2+}$ -signaling  
439 can be differentially modulated via  $\text{ER}\alpha$  and  $\text{ER}\beta$ . In addition, estrogens potentiate the  
440 sensitivity of GT1-7 neurons for C5a by up-regulation of C5aR through  $\text{ER}\alpha$  and  
441  $\text{ER}\beta$ . C5aR activation leads to  $\text{Ca}^{2+}$ -influx through L-VGCCs. Although the  
442 transcription of the  $\text{Ca}_v1.3$  L-VGCC subunit is regulated by E2, the isotype specific  
443 ligands had no effect, indicating that these subunits are not the primary targets of  $\text{ER}\alpha$   
444 and  $\text{ER}\beta$  agonist actions upon C5a/C5aR signaling. The significance of the present  
445 findings relates to the better understanding of the differential impact of estrogens on  
446 the C5a-evoked response of neurons which express  $\text{ER}\alpha$  and/or  $\text{ER}\beta$ .

447

448

449 **Acknowledgement**

450 We thank Drs. Tibor Zelles and Lajos Baranyi for their invaluable advices. We are  
451 grateful to Dr. Pamela Mellon for providing us the GT1-7 neuronal cell line.

452 This research was supported by grants from the Hungarian Scientific Research Found  
453 (OTKA T73002). The research leading to these results has received funding from the  
454 European Community's Seventh Framework Programme (FP7/2007-2013) under  
455 grant agreement No. 245009.

456

457

458 **References**

- 459 Albillos, A., Neher, E., Moser, T., 2000. R-Type Ca<sup>2+</sup> channels are coupled to the  
460 rapid component of secretion in mouse adrenal slice chromaffin cells. *J Neurosci* 20,  
461 8323-8330.
- 462 Bamberg, C.E., Mackay, C.R., Lee, H., Zahra, D., Jackson, J., Lim, Y.S., Whitfeld,  
463 P.L., Craig, S., Corsini, E., Lu, B., Gerard, C., Gerard, N.P., 2010. The C5a receptor  
464 (C5aR) C5L2 is a modulator of C5aR-mediated signal transduction. *J Biol Chem* 285,  
465 7633-7644.
- 466 Baranyi, L., Campbell, W., Ohshima, K., Fujimoto, S., Boros, M., Okada, H., 1995.  
467 The antisense homology box: a new motif within proteins that encodes biologically  
468 active peptides. *Nat Med* 1, 894-901.
- 469 Baranyi, L., Campbell, W., Okada, H., 1996. Antisense homology boxes in C5a  
470 receptor and C5a anaphylatoxin: a new method for identification of potentially active  
471 peptides. *J Immunol* 157, 4591-4601.
- 472 Berridge, M.J., 1998. Neuronal calcium signaling. *Neuron* 21, 13-26.
- 473 Bosch, M.A., Hou, J., Fang, Y., Kelly, M.J., Ronnekleiv, O.K., 2009. 17Beta-estradiol  
474 regulation of the mRNA expression of T-type calcium channel subunits: role of  
475 estrogen receptor alpha and estrogen receptor beta. *J Comp Neurol* 512, 347-358.
- 476 Brewer, L.D., Dowling, A.L., Curran-Rauhut, M.A., Landfield, P.W., Porter, N.M.,  
477 Blalock, E.M., 2009. Estradiol reverses a calcium-related biomarker of brain aging in  
478 female rats. *J Neurosci* 29, 6058-6067.
- 479 Catterall, W.A., Perez-Reyes, E., Snutch, T.P., Striessnig, J., 2005. International  
480 Union of Pharmacology. XLVIII. Nomenclature and structure-function relationships  
481 of voltage-gated calcium channels. *Pharmacol Rev* 57, 411-425.

482 Christian, C.A., Mobley, J.L., Moenter, S.M., 2005. Diurnal and estradiol-dependent  
483 changes in gonadotropin-releasing hormone neuron firing activity. *Proc Natl Acad Sci*  
484 *U S A* 102, 15682-15687.

485 Currie, K.P., 2010. G protein modulation of CaV2 voltage-gated calcium channels.  
486 *Channels (Austin)* 4, 497-509.

487 Dai, S., Hall, D.D., Hell, J.W., 2009. Supramolecular assemblies and localized  
488 regulation of voltage-gated ion channels. *Physiol Rev* 89, 411-452.

489 Dolphin, A.C., 2009. Calcium channel diversity: multiple roles of calcium channel  
490 subunits. *Curr Opin Neurobiol* 19, 237-244.

491 Farkas, I., Baranyi, L., Kaneko, Y., Liposits, Z., Yamamoto, T., Okada, H., 1999. C5a  
492 receptor expression by TGW neuroblastoma cells. *Neuroreport* 10, 3021-3025.

493 Farkas, I., Baranyi, L., Liposits, Z.S., Yamamoto, T., Okada, H., 1998a. Complement  
494 C5a anaphylatoxin fragment causes apoptosis in TGW neuroblastoma cells.  
495 *Neuroscience* 86, 903-911.

496 Farkas, I., Baranyi, L., Takahashi, M., Fukuda, A., Liposits, Z., Yamamoto, T.,  
497 Okada, H., 1998b. A neuronal C5a receptor and an associated apoptotic signal  
498 transduction pathway. *J Physiol* 507 ( Pt 3), 679-687.

499 Farkas, I., Kallo, I., Deli, L., Vida, B., Hrabovszky, E., Fekete, C., Moenter, S.M.,  
500 Watanabe, M., Liposits, Z., 2010. Retrograde endocannabinoid signaling reduces  
501 GABAergic synaptic transmission to gonadotropin-releasing hormone neurons.  
502 *Endocrinology* 151, 5818-5829.

503 Farkas, I., Takahashi, M., Fukuda, A., Yamamoto, N., Akatsu, H., Baranyi, L.,  
504 Tateyama, H., Yamamoto, T., Okada, N., Okada, H., 2003. Complement C5a  
505 receptor-mediated signaling may be involved in neurodegeneration in Alzheimer's  
506 disease. *J Immunol* 170, 5764-5771.



507 Farkas, I., Varju, P., Liposits, Z., 2007. Estrogen modulates potassium currents and  
508 expression of the Kv4.2 subunit in GT1-7 cells. *Neurochem Int* 50, 619-627.

509 Farkas, I., Varju, P., Szabo, E., Hrabovszky, E., Okada, N., Okada, H., Liposits, Z.,  
510 2008. Estrogen enhances expression of the complement C5a receptor and the C5a-  
511 agonist evoked calcium influx in hormone secreting neurons of the hypothalamus.  
512 *Neurochem Int* 52, 846-856.

513 Fonseca, M.I., Ager, R.R., Chu, S.H., Yazan, O., Sanderson, S.D., LaFerla, F.M.,  
514 Taylor, S.M., Woodruff, T.M., Tenner, A.J., 2009. Treatment with a C5aR antagonist  
515 decreases pathology and enhances behavioral performance in murine models of  
516 Alzheimer's disease. *J Immunol* 183, 1375-1383.

517 Fonseca, M.I., Chu, S.H., Berci, A.M., Benoit, M.E., Peters, D.G., Kimura, Y.,  
518 Tenner, A.J., 2011. Contribution of complement activation pathways to  
519 neuropathology differs among mouse models of Alzheimer's disease. *J*  
520 *Neuroinflammation* 8, 4.

521 Fujita, E., Farkas, I., Campbell, W., Baranyi, L., Okada, H., Okada, N., 2004.  
522 Inactivation of C5a anaphylatoxin by a peptide that is complementary to a region of  
523 C5a. *J Immunol* 172, 6382-6387.

524 Gomez-Ospina, N., Tsuruta, F., Barreto-Chang, O., Hu, L., Dolmetsch, R., 2006. The  
525 C terminus of the L-type voltage-gated calcium channel Ca(V)1.2 encodes a  
526 transcription factor. *Cell* 127, 591-606.

527 Guo, R.F., Ward, P.A., 2005. Role of C5a in inflammatory responses. *Annu Rev*  
528 *Immunol* 23, 821-852.

529 Hille, B., 1994. Modulation of ion-channel function by G-protein-coupled receptors.  
530 *Trends Neurosci* 17, 531-536.

531 Hrabovszky, E., Kallo, I., Steinhauser, A., Merchenthaler, I., Coen, C.W., Petersen,  
532 S.L., Liposits, Z., 2004. Estrogen receptor-beta in oxytocin and vasopressin neurons  
533 of the rat and human hypothalamus: Immunocytochemical and in situ hybridization  
534 studies. *J Comp Neurol* 473, 315-333.

535 Hrabovszky, E., Shughrue, P.J., Merchenthaler, I., Hajszan, T., Carpenter, C.D.,  
536 Liposits, Z., Petersen, S.L., 2000. Detection of estrogen receptor-beta messenger  
537 ribonucleic acid and 125I-estrogen binding sites in luteinizing hormone-releasing  
538 hormone neurons of the rat brain. *Endocrinology* 141, 3506-3509.

539 Hrabovszky, E., Steinhauser, A., Barabas, K., Shughrue, P.J., Petersen, S.L.,  
540 Merchenthaler, I., Liposits, Z., 2001. Estrogen receptor-beta immunoreactivity in  
541 luteinizing hormone-releasing hormone neurons of the rat brain. *Endocrinology* 142,  
542 3261-3264.

543 Karsch, F.J., Battaglia, D.F., Breen, K.M., Debus, N., Harris, T.G., 2002. Mechanisms  
544 for ovarian cycle disruption by immune/inflammatory stress. *Stress* 5, 101-112.

545 Kim, S., Yun, H.M., Baik, J.H., Chung, K.C., Nah, S.Y., Rhim, H., 2007. Functional  
546 interaction of neuronal Cav1.3 L-type calcium channel with ryanodine receptor type 2  
547 in the rat hippocampus. *J Biol Chem* 282, 32877-32889.

548 Kolarow, R., Brigadski, T., Lessmann, V., 2007. Postsynaptic secretion of BDNF and  
549 NT-3 from hippocampal neurons depends on calcium calmodulin kinase II signaling  
550 and proceeds via delayed fusion pore opening. *J Neurosci* 27, 10350-10364.

551 Liposits, Z., Merchenthaler, I., Wetsel, W.C., Reid, J.J., Mellon, P.L., Weiner, R.I.,  
552 Negro-Vilar, A., 1991. Morphological characterization of immortalized hypothalamic  
553 neurons synthesizing luteinizing hormone-releasing hormone. *Endocrinology* 129,  
554 1575-1583.

555 Magee, J.C., Johnston, D., 1995. Characterization of single voltage-gated Na<sup>+</sup> and  
556 Ca<sup>2+</sup> channels in apical dendrites of rat CA1 pyramidal neurons. *J Physiol* 487 ( Pt  
557 1), 67-90.

558 Moenter, S.M., DeFazio, A.R., Pitts, G.R., Nunemaker, C.S., 2003. Mechanisms  
559 underlying episodic gonadotropin-releasing hormone secretion. *Front*  
560 *Neuroendocrinol* 24, 79-93.

561 Nishiura, H., Tokita, K., Li, Y., Harada, K., Woodruff, T.M., Taylor, S.M., Nsiama,  
562 T.K., Nishino, N., Yamamoto, T., 2010. The role of the ribosomal protein S19 C-  
563 terminus in Gi protein-dependent alternative activation of p38 MAP kinase via the  
564 C5a receptor in HMC-1 cells. *Apoptosis* 15, 966-981.

565 Qiu, J., Bosch, M.A., Jamali, K., Xue, C., Kelly, M.J., Ronnekleiv, O.K., 2006.  
566 Estrogen upregulates T-type calcium channels in the hypothalamus and pituitary. *J*  
567 *Neurosci* 26, 11072-11082.

568 Roy, D., Angelini, N.L., Belsham, D.D., 1999. Estrogen directly respresses  
569 gonadotropin-releasing hormone (GnRH) gene expression in estrogen receptor-alpha  
570 (ERalpha)- and ERbeta-expressing GT1-7 GnRH neurons. *Endocrinology* 140, 5045-  
571 5053.

572 Sabatini, B.L., Svoboda, K., 2000. Analysis of calcium channels in single spines using  
573 optical fluctuation analysis. *Nature* 408, 589-593.

574 Sarvari, M., Kallo, I., Hrabovszky, E., Solymosi, N., Toth, K., Liko, I., Molnar, B.,  
575 Tihanyi, K., Liposits, Z., 2010. Estradiol replacement alters expression of genes  
576 related to neurotransmission and immune surveillance in the frontal cortex of middle-  
577 aged, ovariectomized rats. *Endocrinology* 151, 3847-3862.

578 Shughrue, P.J., Lane, M.V., Merchenthaler, I., 1997. Comparative distribution of  
579 estrogen receptor-alpha and -beta mRNA in the rat central nervous system. *J Comp*  
580 *Neurol* 388, 507-525.

581 Shughrue, P.J., Merchenthaler, I., 2001. Distribution of estrogen receptor beta  
582 immunoreactivity in the rat central nervous system. *J Comp Neurol* 436, 64-81.

583 Speth, C., Prodinger, W.M., Wurzner, R., Stoiber, H., Dierich, M.P., 2008.  
584 Complement, In: Paul, W.E. (Ed.), *Fundamental immunology*, 6th ed, Lippincott  
585 Williams and Wilkins, Philadelphia, pp. 1047-1078.

586 Stahel, P.F., Frei, K., Eugster, H.P., Fontana, A., Hummel, K.M., Wetsel, R.A., Ames,  
587 R.S., Barnum, S.R., 1997a. TNF-alpha-mediated expression of the receptor for  
588 anaphylatoxin C5a on neurons in experimental *Listeria meningoenzephalitis*. *J*  
589 *Immunol* 159, 861-869.

590 Stahel, P.F., Kossmann, T., Morganti-Kossmann, M.C., Hans, V.H., Barnum, S.R.,  
591 1997b. Experimental diffuse axonal injury induces enhanced neuronal C5a receptor  
592 mRNA expression in rats. *Brain Res Mol Brain Res* 50, 205-212.

593 Sun, J., Chu, Z., Moenter, S.M., 2010. Diurnal in vivo and rapid in vitro effects of  
594 estradiol on voltage-gated calcium channels in gonadotropin-releasing hormone  
595 neurons. *J Neurosci* 30, 3912-3923.

596 Tedford, H.W., Zamponi, G.W., 2006. Direct G protein modulation of Cav2 calcium  
597 channels. *Pharmacol Rev* 58, 837-862.

598 Thiery, J.C., Pelletier, J., 1981. Multiunit activity in the anterior median eminence and  
599 adjacent areas of the hypothalamus of the ewe in relation to LH secretion.  
600 *Neuroendocrinology* 32, 217-224.

601 Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A.,  
602 Speleman, F., 2002. Accurate normalization of real-time quantitative RT-PCR data by

603 geometric averaging of multiple internal control genes. *Genome Biol* 3,  
604 RESEARCH0034.

605 Varju, P., Chang, K.C., Hrabovszky, E., Merchenthaler, I., Liposits, Z., 2009.  
606 Temporal profile of estrogen-dependent gene expression in LHRH-producing GT1-7  
607 cells. *Neurochem Int* 54, 119-134.

608 Wang, G., Dayanithi, G., Newcomb, R., Lemos, J.R., 1999. An R-type Ca(2+) current  
609 in neurohypophysial terminals preferentially regulates oxytocin secretion. *J Neurosci*  
610 19, 9235-9241.

611 Watanabe, M., Sakuma, Y., Kato, M., 2004. High expression of the R-type voltage-  
612 gated Ca<sup>2+</sup> channel and its involvement in Ca<sup>2+</sup>-dependent gonadotropin-releasing  
613 hormone release in GT1-7 cells. *Endocrinology* 145, 2375-2383.

614 Wetsel, W.C., Valenca, M.M., Merchenthaler, I., Liposits, Z., Lopez, F.J., Weiner,  
615 R.I., Mellon, P.L., Negro-Vilar, A., 1992. Intrinsic pulsatile secretory activity of  
616 immortalized luteinizing hormone-releasing hormone-secreting neurons. *Proc Natl*  
617 *Acad Sci U S A* 89, 4149-4153.

618 Wilson, M.E., Rosewell, K.L., Kashon, M.L., Shughrue, P.J., Merchenthaler, I., Wise,  
619 P.M., 2002. Age differentially influences estrogen receptor-alpha (ERalpha) and  
620 estrogen receptor-beta (ERbeta) gene expression in specific regions of the rat brain.  
621 *Mech Ageing Dev* 123, 593-601.

622 Wilson, R.C., Kesner, J.S., Kaufman, J.M., Uemura, T., Akema, T., Knobil, E., 1984.  
623 Central electrophysiologic correlates of pulsatile luteinizing hormone secretion in the  
624 rhesus monkey. *Neuroendocrinology* 39, 256-260.

625 Woodruff, T.M., Ager, R.R., Tenner, A.J., Noakes, P.G., Taylor, S.M., 2010. The role  
626 of the complement system and the activation fragment C5a in the central nervous  
627 system. *Neuromolecular Med* 12, 179-192.

628 Woodruff, T.M., Nandakumar, K.S., Tedesco, F., 2011. Inhibiting the C5-C5a  
629 receptor axis. *Mol Immunol*.

630 Wu, L.G., Borst, J.G., Sakmann, B., 1998. R-type Ca<sup>2+</sup> currents evoke transmitter  
631 release at a rat central synapse. *Proc Natl Acad Sci U S A* 95, 4720-4725.

632 Wu, L.G., Westenbroek, R.E., Borst, J.G., Catterall, W.A., Sakmann, B., 1999.  
633 Calcium channel types with distinct presynaptic localization couple differentially to  
634 transmitter release in single calyx-type synapses. *J Neurosci* 19, 726-736.

635 Zhang, C., Bosch, M.A., Rick, E.A., Kelly, M.J., Ronnekleiv, O.K., 2009. 17Beta-  
636 estradiol regulation of T-type calcium channels in gonadotropin-releasing hormone  
637 neurons. *J Neurosci* 29, 10552-10562.

638 Zhang, H., Fu, Y., Altier, C., Platzer, J., Surmeier, D.J., Bezprozvanny, I., 2006.  
639 Ca<sub>v</sub>1.2 and Ca<sub>v</sub>1.3 neuronal L-type calcium channels: differential targeting and  
640 signaling to pCREB. *Eur J Neurosci* 23, 2297-2310.

641 Zuccotti, A., Clementi, S., Reinbothe, T., Torrente, A., Vandael, D.H., Pirone, A.,  
642 2011. Structural and functional differences between L-type calcium channels: crucial  
643 issues for future selective targeting. *Trends Pharmacol Sci* 32, 366-375.

644

645

646 **Figure legends**

647 **Figure 1. Ca<sup>2+</sup>-influx evoked by the C5aR agonist PL37-MAP (PL37) in the**  
648 **presence of E2, DPN, or PPT in the GT1-7 neurons.** a) The PL37-MAP induced  
649 robust Ca<sup>2+</sup>-influx in the cells. b) Pre-treatment of the GT1-7 cells with E2 (100pM,  
650 24h) potentiated the Ca<sup>2+</sup>-influx significantly. Onset of the response started earlier and  
651 the amplitude of it was higher than without pre-treatment. c) Pre-treatment with the  
652 PPT (100pM, 24h) resulted in an elevated Ca<sup>2+</sup>-influx. d) DPN pre-treatment (100pM,  
653 24h), however, attenuated the Ca<sup>2+</sup>-influx evoked by the PL37-MAP, significantly. e)  
654 The histogram shows the area-under-curve values, representing the net Ca<sup>2+</sup>-influx in  
655 the neurons, in the percentage of the Ca<sup>2+</sup>-influx measured with the PL37-MAP alone  
656 (PL37+100 pM E2: 226,6±27,52%; PL37+100 pM PPT: 159,1±16,74%; PL37+100  
657 pM DPN: 18,6±10,47%; PL37+20 nM E2: 169,7±16,67%; PL37+20 nM PPT:  
658 80,2±9,73%; PL37+20 nM DPN: 10,9±3,65%) f) Acute (8min) pre-treatments with  
659 the E2, DPN, or PPT showed no significant effect on the PL37-MAP-evoked Ca<sup>2+</sup>-  
660 influx, demonstrating that the observed effect of E2, DPN, or PPT was genomic  
661 (PL37+E2: 115,1±13,78%; PL37+PPT: 132,1±13,97%; PL37+DPN: 146,9±45,22%).  
662 \*=p<0.05, \*\*=p<0.01. Arrow shows the onset of the administration of the PL37-  
663 MAP.

664

665 **Figure 2. Ca<sup>2+</sup>-influx evoked by the PL37-MAP (PL37) in the presence of CPA,**  
666 **E2, DPN, or PPT in the GT1-7 neurons in Ca<sup>2+</sup>-free extracellular solution.** a-f)  
667 Calcium imaging recordings show that extracellular calcium sources play important  
668 role in the Ca<sup>2+</sup>-influx. g) The histogram reveals that the Ca<sup>2+</sup>-free extracellular  
669 environment decreased the Ca<sup>2+</sup>-response significantly, demonstrating that majority of  
670 the elevation in the Ca<sup>2+</sup>-concentration was from the extracellular source. This change

671 was independent from the pre-treatment with E2, DPN or PPT. CPA, a depletor of the  
672  $\text{Ca}^{2+}$ -pool in the endoplasmic reticulum, however, eliminated the remaining  $\text{Ca}^{2+}$ -  
673 response in the neurons, showing that the endoplasmic reticulum was the intracellular  
674 source of the remaining  $\text{Ca}^{2+}$ -response (PL37 in  $\text{Ca}^{2+}$ -free:  $25.3 \pm 4.91\%$ ; PL37+CPA  
675 in  $\text{Ca}^{2+}$ -free:  $3.2 \pm 0.66\%$ ; PL37+E2 in  $\text{Ca}^{2+}$ -free:  $16.9 \pm 1.63\%$ ; PL37+PPT in  $\text{Ca}^{2+}$ -  
676 free:  $22.9 \pm 5.82\%$ ; PL37+DPN in  $\text{Ca}^{2+}$ -free:  $27.5 \pm 5.45\%$ ).  $*=p<0.05$ ,  $**=p<0.01$ .  
677 Arrow shows the onset of the administration of the PL37-MAP.

678

679 **Figure 3.  $\text{Ca}^{2+}$ -influx evoked by the PL37-MAP in the presence of nifedipine**  
680 **(blocker of L-type  $\text{Ca}^{2+}$ -channels), E2, DPN, or PPT in the GT1-7 neurons. a-b)**  
681 Calcium imaging recordings show that comparing to the control, nifedipine (Nif)  
682 eliminated the  $\text{Ca}^{2+}$ -influx significantly. c-e) This decrease was independent from the  
683 presence of the E2, DPN, or PPT, demonstrating role of the L-type  $\text{Ca}^{2+}$ -channels. f)  
684 The histogram of the area-under-curve data shows the values expressed in the  
685 percentage of the  $\text{Ca}^{2+}$ -influx measured with the PL37-MAP alone (PL37+Nif:  
686  $39.7 \pm 3.29\%$ ; PL37+E2+Nif:  $35.8 \pm 6.94\%$ ; PL37+PPT+Nif:  $32.4 \pm 4.66\%$ ;  
687 PL37+DPN+Nif:  $32.4 \pm 4.66\%$ ).  $*=p<0.05$ . Arrow shows the onset of the  
688 administration of the PL37-MAP.

689

690 **Figure 4.  $\text{Ca}^{2+}$ -influx evoked by the PL37-MAP in the presence of SNX-482**  
691 **(blocker of R-type  $\text{Ca}^{2+}$ -channels), E2, DPN, or PPT in the GT1-7 neurons. a-e)**  
692 The ratiometric graphs revealed that SNX-482 (SNX) showed no effect on the  $\text{Ca}^{2+}$ -  
693 influx evoked by the PL37-MAP, and it exerted no influence on the effect of the ER  
694 agonists, demonstrating that the R-type  $\text{Ca}^{2+}$ -channel plays no role in these processes.  
695 f) The histogram of the area-under-curve data shows the values expressed in the




696 percentage of the Ca<sup>2+</sup>-influx measured with the PL37-MAP alone (PL37+SNX:  
697 81.6±13.95%; PL37+E2+SNX: 139.5±14.63%; PL37+PPT+SNX: 86.6±5.33%;  
698 PL37+DPN+SNX: 43.3±6.15%). \*= $p < 0.05$ . Arrow shows the onset of the  
699 administration of the PL37-MAP.  
700

701 **Table legend**

702 **Table 1. Transcriptional modulation of L-type Ca<sup>2+</sup>-channel subunits and C5a**  
703 **receptors.** Transcription of L-type Ca<sup>2+</sup>-channel Ca<sub>v</sub>1.2 (Cacna1c) and Ca<sub>v</sub>1.3  
704 (Cacna1d) subunits, and C5a receptors CD88 (C5ar1) and C5L2 (Gpr77) was  
705 followed by real-time PCR. Table shows the arithmetic mean and standard deviation  
706 of relative quantities (RQ) from two independent experiments. Arrows show direction  
707 of regulation of the respective transcript where change is significant. The lack of  
708 evidence for regulation is denoted with “-“.

709

710



711

Fig. 1.

712





Fig. 2.

713

714





Fig. 3.

715

716



717

718

Fig. 4.

719  
720

<b>Gene symbol and physiologic name</b>	<b>E2</b>		<b>PPT</b>		<b>DPN</b>	
Cacna1c (L-type Ca <sup>2+</sup> -channel Ca <sub>v</sub> 1.2 subunit)	0.867±0.164	-	0.905±0.073	-	0.748±0.052	-
Cacna1d (L-type Ca <sup>2+</sup> -channel Ca <sub>v</sub> 1.3 subunit)	0.413±0.326	↓	0.960±0.023	-	0.795±0.117	-
C5ar1 (C5aR)	2.476±0.715	↑	2.039±0.241	↑	1.938±0.743	↑
Gpr77 (C5L2)	1.634±0.336	↑	0.562±0.101	↓	0.955±0.577	-

721  
722  
723  
724  
725

Table 1.